

SIGN 158

British guideline on the management of asthma

A national clinical guideline

First published 2003

Revised edition published July 2019



British
Thoracic
Society

NHS
SCOTLAND

Key to evidence statements and recommendations

Levels of evidence

- 1⁺⁺ | High-quality meta-analyses, systematic reviews of RCTs, or RCTs with a very low risk of bias
- 1⁺ | Well-conducted meta-analyses, systematic reviews, or RCTs with a low risk of bias
- 1⁻ | Meta-analyses, systematic reviews, or RCTs with a high risk of bias
- 2⁺⁺ | High-quality systematic reviews of case-control or cohort studies
High-quality case-control or cohort studies with a very low risk of confounding or bias and a high probability that the relationship is causal
- 2⁺ | Well-conducted case-control or cohort studies with a low risk of confounding or bias and a moderate probability that the relationship is causal
- 2⁻ | Case-control or cohort studies with a high risk of confounding or bias and a significant risk that the relationship is not causal
- 3 | Non-analytic studies, eg case reports, case series
- 4 | Expert opinion

Grades of recommendation

Note: The grade of recommendation relates to the strength of the supporting evidence on which the evidence is based. It does not reflect the clinical importance of the recommendation.

- A** | At least one meta-analysis, systematic review, or RCT rated as 1⁺⁺, and directly applicable to the target population; *or*
A body of evidence consisting principally of studies rated as 1⁺, directly applicable to the target population, and demonstrating overall consistency of results
- B** | A body of evidence including studies rated as 2⁺⁺, directly applicable to the target population, and demonstrating overall consistency of results; *or*
Extrapolated evidence from studies rated as 1⁺⁺ or 1⁺
- C** | A body of evidence including studies rated as 2⁺, directly applicable to the target population and demonstrating overall consistency of results; *or*
Extrapolated evidence from studies rated as 2⁺⁺
- D** | Evidence level 3 or 4; *or*
Extrapolated evidence from studies rated as 2⁺

Good-practice points

- ✓ | Recommended best practice based on the clinical experience of the guideline development group.



NHS Evidence has accredited the process used by Scottish Intercollegiate Guidelines Network to produce clinical guidelines. Accreditation is applicable to guidance produced using the processes described in SIGN 50: a guideline developer's handbook, 2011 edition (www.sign.ac.uk/sign-50).

More information on accreditation can be viewed at www.evidence.nhs.uk

Healthcare Improvement Scotland (HIS) is committed to equality and diversity and assesses all its publications for likely impact on the six equality groups defined by age, disability, gender, race, religion/belief and sexual orientation.

SIGN guidelines are produced using a standard methodology that has been **equality impact assessed** to ensure that these equality aims are addressed in every guideline. This methodology is set out in the current version of SIGN 50, our guideline manual, which can be found at www.sign.ac.uk/sign-50.html. The EQIA assessment of the manual can be seen at www.sign.ac.uk/assets/sign50eqia.pdf. The full report in paper form and/or alternative format is available on request from the Healthcare Improvement Scotland Equality and Diversity Officer.

Every care is taken to ensure that this publication is correct in every detail at the time of publication. However, in the event of errors or omissions corrections will be published in the web version of this document, which is the definitive version at all times. This version can be found on our web site www.sign.ac.uk

Scottish Intercollegiate Guidelines Network
British Thoracic Society

British guideline on the management of asthma

A national clinical guideline

First published 2003

Revised edition published July 2019

978-1-909103-70-2

First published 2003

Revised edition published July 2019

Scottish Intercollegiate Guidelines Network

Gyle Square, 1 South Gyle Crescent
Edinburgh EH12 9EB

www.sign.ac.uk

British Thoracic Society

17 Doughty Street
London WC1N 2PL

www.brit-thoracic.org.uk

SIGN and BTS consent to the photocopying of this guideline for the purpose of implementation in the NHS in England, Wales, Northern Ireland and Scotland.

Contents

1	Introduction	1
1.1	The need for a guideline	1
1.2	Remit of the guideline	1
1.3	Statement of intent	4
2	Key recommendations	6
2.1	Diagnosis	6
2.2	Monitoring	7
2.3	Supported self management	7
2.4	Non-pharmacological management	7
2.5	Pharmacological management	7
2.6	Inhaler devices	8
2.7	Acute asthma	8
2.8	Difficult asthma	9
2.9	Asthma in pregnancy	9
2.10	Occupational asthma	9
3	Diagnosis	10
3.1	Definition and overarching principles	10
3.2	Predictive value of individual symptoms, signs and diagnostic tests	11
3.3	Practical approach to diagnosis	20
3.4	Organisation of diagnostic services	28
3.5	Wheezing in preschool children and the future risk of developing persistent asthma	29
4	Monitoring asthma	30
4.1	Targeting care	31
4.2	Monitoring current asthma symptom control	31
4.3	Predicting future risk of asthma attacks	32
4.4	Physiological measures	36
4.5	Other approaches	37
5	Supported self management	38
5.1	Effectiveness of supported self management	38
5.2	Components of a self-management programme	38
5.3	Self management in specific patient groups	42
5.4	Adherence and concordance	45
5.5	Implementation in practice	48
6	Non-pharmacological management	50
6.1	Primary prevention	50
6.2	Secondary prevention	55

7	Pharmacological management	62
7.1	Intermittent reliever therapy	63
7.2	Regular preventer therapy	63
7.3	Initial add-on therapy	70
7.4	Additional controller therapies	72
7.5	Specialist therapies	73
7.6	Deceasing treatment	82
7.7	Specific management issues	82
8	Inhaler devices	85
8.1	Technique and training	85
8.2	β_2 agonist delivery	85
8.3	Inhaled corticosteroids for stable asthma	86
8.4	Prescribing devices	86
8.5	Use and care of spacers	87
8.6	Environmental impact of metered-dose inhalers	87
9	Management of acute asthma	89
9.1	Lessons from asthma deaths and near-fatal asthma	89
9.2	Acute asthma in adults	91
9.3	Treatment of acute asthma in adults	94
9.4	Further investigation and monitoring	100
9.5	Asthma management protocols and proformas	100
9.6	Hospital discharge and follow up	101
9.7	Acute asthma in children	102
9.8	Initial treatment of acute asthma in children	104
9.9	Second-line treatment of acute asthma in children	109
10	Difficult asthma	113
10.1	Defining and assessing difficult asthma	113
10.2	Factors contributing to difficult asthma	113
11	Asthma in adolescents	117
11.1	Definitions	117
11.2	Prevalence of asthma in adolescents	117
11.3	Diagnosis and assessment	117
11.4	Risk factors	118
11.5	Comorbidities and modifiable behaviours	119
11.6	Asthma attacks and the risk of hospital admission	120
11.7	Long-term outlook and entry into the workplace	120
11.8	Non-pharmacological management	120
11.9	Pharmacological management	121
11.10	Inhaler devices	121
11.11	Organisation and delivery of care	122
11.12	Patient education and self management	123

12	Asthma in pregnancy	126
12.1	Natural history and management of stable asthma	126
12.2	Management of acute asthma in pregnancy	127
12.3	Drug therapy in pregnancy	128
12.4	Management during labour	131
12.5	Drug therapy for breastfeeding mothers	132
13	Occupational asthma	133
13.1	Incidence	133
13.2	At-risk populations	133
13.3	Diagnosis	133
13.4	Management of occupational asthma	136
14	Organisation and delivery of care	138
14.1	Care pathways	138
14.2	Educating clinicians	138
14.3	Asthma clinics	139
14.4	Telehealthcare	140
14.5	School-based interventions	143
14.6	Ethnicity/culture-based interventions	143
14.7	Lay-led interventions	144
14.8	Pharmacist-led interventions	144
15	Provision of information	145
15.1	Checklist of information for patients and carers	145
15.2	Publications from SIGN	147
15.3	Sources of further information	148
16	The evidence base	150
16.1	Systematic literature review	150
16.2	Recommendations for research	150
17	Development of the guideline	152
17.1	Introduction	152
17.2	Guideline development group	152
17.3	Acknowledgements	154
17.4	Consultation and peer review	154
Abbreviations		156
Annexes		158
References		172

1 Introduction

1.1 The need for a guideline

Asthma is a common condition which produces a significant workload for general practice, hospital outpatient clinics and inpatient services. It is clear that much of this morbidity relates to poor management, particularly around the use of preventative medicine.

1.1.1 Background

In 1999 the British Thoracic Society (BTS) and the Scottish Intercollegiate Guidelines Network (SIGN) agreed to jointly produce a comprehensive new asthma guideline, both having previously published guidance on asthma. The original BTS guideline dated back to 1990 and the SIGN guidelines to 1996. Both organisations recognised the need to develop the new guideline using evidence-based methodology explicitly. The joint process was further strengthened by collaboration with Asthma UK, the Royal College of Physicians of London, the Royal College of Paediatrics and Child Health, the General Practice Airways Group (now Primary Care Respiratory Society UK), and the British Association of Accident and Emergency Medicine (now the College of Emergency Medicine). The outcome of these efforts was the British Guideline on the Management of Asthma, developed using SIGN methodology¹ and published in 2003.²

1.1.2 Updating the evidence

Between 2004 and 2012 sections within the guideline were updated annually. Subsequently, updating moved to a biennial basis, beginning with the 2014 update. This edition of the guideline was issued in 2019. All updates were published on the BTS and SIGN websites. A list of the key questions addressed in this update is given in Annex 1. Any updates to the guideline in the period between scheduled updates will be noted on the SIGN and BTS websites.

A summary of the search histories for each section is given in Annex 2. It is hoped that this asthma guideline continues to serve as a basis for high quality management of both acute and chronic asthma and a stimulus for research into areas of management for which there is little evidence (*see section 16.2*).

1.2 Remit of the guideline

1.2.1 Overall objectives

This guideline provides recommendations based on current evidence for best practice in the management of asthma. It makes recommendations on management of adults, including pregnant women, and adolescents and children with asthma. In sections 7 and 8 on pharmacological management and inhaler devices, respectively, and in section 4.3 on predicting future risk of asthma attacks, each recommendation has been graded and the supporting evidence assessed for adults and adolescents over 12 years old, children 5–12 years, and children under 5 years. Further information on managing asthma in adolescents (10–19 years of age as defined by the World Health Organization)³ is given in section 11.

The guideline considers diagnosis of asthma and management in all patients with a diagnosis of asthma, although there is less evidence available for people at either age extreme. The guideline does not cover patients whose primary diagnosis is not asthma, for example those with chronic obstructive pulmonary disease (COPD) or cystic fibrosis, but patients with these conditions can also have asthma. Under these circumstances many of the principles set out in this guideline will apply to the management of their asthma symptoms.

1.2.2 Target users of the guideline

This guideline will be of particular interest to healthcare professionals involved in the care of people with asthma including general practitioners, consultants and specialists in respiratory medicine, nurses, pharmacists and other allied health professionals with an interest in respiratory care. The guideline will also be of interest to people with asthma, their parents and carers; those who interact with people with asthma outside of the NHS, such as teachers; voluntary organisations with an interest in asthma; and those planning the delivery of services in the NHS in England, Wales, Northern Ireland and Scotland.

1.2.3 Summary of updates to the guideline, by section

Guideline section		Year of update
2	Key recommendations	2014, 2016, 2019
3	Diagnosis	2008, 2011, 2016
4	Monitoring asthma	2008, 2011, 2019
5	Supported self management	2004, 2008, 2014, 2016, 2019
6	Non-pharmacological management	2008, 2014, 2016, 2019
7	Pharmacological management	2004, 2005, 2006, 2008, 2009, 2011, 2014, 2016, 2019
8	Inhaler devices	2005, 2014, 2019
9	Management of acute asthma	2004, 2009, 2014, 2016, 2019
10	Difficult asthma	2008, 2014, 2016
11	Asthma in adolescents	2011
12	Asthma in pregnancy	2005, 2008, 2009, 2014
13	Occupational asthma	2005, 2008, 2014, 2016
14	Organisation and delivery of care	2008, 2014, 2016

1.2.4 Summary of updates to the 2019 edition of the guideline, by section

The table below lists all the sections and subsections of the guideline that were updated in 2019. This update includes a complete revision of the section on monitoring, and updates to sections including supported self management, non-pharmacological management of asthma, pharmacological management of asthma, inhaler devices and management of acute asthma.

Guideline section		Year of update
2	Key recommendations	Updated: 2.2 Monitoring Minor update: 2.1 Diagnosis, 2.4 Non-pharmacological management, 2.10 Occupational asthma
3	Diagnosis	Minor update: 3.3.3 Low probability of asthma based on initial structured clinical assessment, 3.4 Organisation of diagnostic services
4	Monitoring asthma	New: Table 7, 4.1 Targeting care, 4.2 Monitoring current asthma symptom control, 4.3 Predicting future risk of asthma attacks, 4.3.1 Adults, 4.3.2 School-aged children, 4.3.3 Preschool children, 4.3.4 People with severe asthma, 4.4 Physiological measures, 4.4.1 Spirometry and peak expiratory flow, 4.4.2 Fractional exhaled nitric oxide, 4.4.3 Eosinophils, 4.5 Other approaches
5	Supported self management	New: 5.2.3 Increasing inhaled corticosteroids to abort an asthma attack Updated: 5.4.3 Interventions to improve medication adherence Minor update: 5.2.2 Personalised asthma action plans, Table 11
6	Non-pharmacological management	New: 6.2.8 Vitamin D Updated: 6.2.1 House dust mite avoidance, 6.2.2 Other allergens, 6.2.3 Smoking, 6.2.4 Air pollution, 6.2.13 Air ionisers, 6.2.14 Breathing exercises
7	Pharmacological management	Updated: 7.1.1 Frequency of dosing of inhaled SABA, Table 12 Categorisation of inhaled corticosteroids by dose – adults, Table 13 Categorisation of inhaled corticosteroids by dose – children, 7.3.5 Single combination inhaler for maintenance and reliever therapy, 7.4 Additional controller therapies, 7.5 Specialist therapies; 7.5.2 Other approaches, 7.5.3 Continuous or frequent use of oral steroids, 7.5.4 Monoclonal antibody, 7.5.6 Immunotherapy for asthma, 7.5.7 Bronchial thermoplasty, Figure 2 Summary of management in adults, Figure 3 Summary of management in children, Minor updates: 7.1 Intermittent reliever therapy, 7.5.5 Other agents, 7.7.2 Exercise-induced asthma Deleted: Patients on oral steroids not previously tried on inhaled therapy
8	Inhaler devices	New: 8.6 Environmental impact of metered-dose inhalers
9	Management of acute asthma	Updated: 9.3.5 Magnesium sulphate, 9.3.12 Critical care settings (adults), 9.9.5 Critical care settings (children), 9.9.6 Non-invasive ventilation Minor update: 9.3.2 β_2 agonist bronchodilators, 9.9.3 Intravenous magnesium sulphate
13	Occupational asthma	Minor update: 13.3 Diagnosis
15	Provision of information	New: 15.1 Checklist of information for patients and carers, 15.2 Publications from SIGN
16	The evidence base	Updated: 16.2 Recommendations for research
	Annexes	New: Annex 1 Key questions addressed in this update Updated: Annexes 3 – 9 Management of acute asthma

1.3 Statement of intent

This guideline is not intended to be construed or to serve as a standard of care. Standards of care are determined on the basis of all clinical data available for an individual case and are subject to change as scientific knowledge and technology advance and patterns of care evolve. Adherence to guideline recommendations will not ensure a successful outcome in every case, nor should they be construed as including all proper methods of care or excluding other acceptable methods of care aimed at the same results. The ultimate judgement must be made by the appropriate healthcare professional(s) responsible for clinical decisions regarding a particular clinical procedure or treatment plan. This judgement should only be arrived at through a process of shared decision making with the patient, covering the diagnostic and treatment choices available. It is advised, however, that significant departures from the national guideline or any local guidelines derived from it should be fully documented in the patient's medical records at the time the relevant decision is taken.

1.3.1 Influence of financial and other interests

It has been recognised that financial interests in, or close working relationships with, pharmaceutical companies may have an influence on the interpretation of evidence from clinical studies.

It is not possible to completely eliminate any possible bias from this source, nor even to quantify the degree of bias with any certainty. SIGN requires that all those involved in the work of guideline development should declare all financial interests, whether direct or indirect, annually for as long as they are actively working with the organisation. By being explicit about the influences to which contributors are subjected, SIGN acknowledges the risk of bias and makes it possible for guideline users or reviewers to assess for themselves how likely it is that the conclusions and guideline recommendations are based on a biased interpretation of the evidence.

Signed copies are retained by the SIGN Executive and a register of interests is available in the supporting material section for this guideline at www.sign.ac.uk

1.3.2 Patient version

Patient versions of this guideline are available from the SIGN website, www.sign.ac.uk

1.3.3 Prescribing of licensed medicines outwith their marketing authorisation

Recommendations within this guideline are based on the best clinical evidence. Some recommendations may be for medicines prescribed outwith the marketing authorisation (MA) also known as product licence. This is known as 'off label' use.

Medicines may be prescribed 'off label' in the following circumstances:

- for an indication not specified within the marketing authorisation
- for administration via a different route
- for administration of a different dose
- for a different patient population.

An unlicensed medicine is a medicine which does not have MA for medicinal use in humans.

Generally 'off label' prescribing of medicines becomes necessary if the clinical need cannot be met by licensed medicines within the marketing authorisation. Such use should be supported by appropriate evidence and experience.⁴

"Prescribing medicines outside the conditions of their marketing authorisation alters (and probably increases) the prescribers' professional responsibility and potential liability."⁴

The General Medical Council (GMC) recommends that when prescribing a medicine 'off label', doctors should:

- be satisfied that there is no suitably licensed medicine that will meet the patient's need
- be satisfied that there is sufficient evidence or experience of using the medicine to show its safety and efficacy
- take responsibility for prescribing the medicine and for overseeing the patient's care, including monitoring the effects of the medicine and any follow-up treatment, or ensure that arrangements are made for another suitable doctor to do so
- make a clear, accurate and legible record of all medicines prescribed and, when not following common practice, the reasons for prescribing an unlicensed medicine.

Non-medical prescribers should ensure that they are familiar with the legislative framework and their own professional prescribing standards.

Prior to any prescribing, the licensing status of a medication should be checked in the summary of product characteristics (www.medicines.org.uk). The prescriber must be competent, operate within the professional code of ethics of their statutory bodies and the prescribing practices of their employers.⁵

1.3.4 Additional advice on the use of new and existing medicines and treatments

The National Institute for Health and Care Excellence (NICE) develops technology appraisals that make recommendations on the use of new and existing medicines and treatments within the NHS in England and Wales.

The Scottish Medicines Consortium (SMC) provides advice to NHS boards and their Area Drug and Therapeutics Committees about the status of all newly-licensed medicines, all new formulations of existing medicines and new indications for established products, within NHSScotland.

Until 1 October 2017, Healthcare Improvement Scotland reviewed Multiple Technology Appraisals produced by NICE and provided advice about their applicability in NHSScotland. If Healthcare Improvement Scotland has advised that Multiple Technology Appraisal guidance was applicable in Scotland, NHSScotland should take account of this and ensure that recommended medicines and treatment are made available to meet clinical need where appropriate.

NICE Multiple Technology Appraisals deemed valid for NHSScotland supersede extant SMC advice as they are generally underpinned by a larger and more recent evidence base.

Practitioners should be aware of this additional advice on medicines and treatments recommended in this guideline and that recommendations made by these organisations and restrictions on their use may differ between England and Wales and Scotland.

2 Key recommendations

The following recommendations were highlighted by the guideline development group as the key clinical recommendations that should be prioritised for implementation. The grade of recommendation relates to the strength of the supporting evidence on which the recommendation is based. It does not reflect the clinical importance of the recommendation.

2.1 Diagnosis

C Compare the results of diagnostic tests undertaken whilst a patient is asymptomatic with those undertaken when a patient is symptomatic to detect variation over time.

D Carry out quality-assured spirometry using the lower limit of normal to demonstrate airway obstruction, provide a baseline for assessing response to initiation of treatment and exclude alternative diagnoses.

- Obstructive spirometry with positive bronchodilator reversibility increases the probability of asthma.
- Normal spirometry in an asymptomatic patient does not rule out the diagnosis of asthma.

D Undertake a structured clinical assessment to assess the initial probability of asthma. This should be based on:

- a history of recurrent episodes (attacks) of symptoms, ideally corroborated by variable peak flow when symptomatic and asymptomatic
- symptoms of wheeze, cough, breathlessness and chest tightness that vary over time
- recorded observation of wheeze heard by a healthcare professional
- personal/family history of other atopic conditions (in particular, atopic eczema/dermatitis, allergic rhinitis)
- no symptoms/signs to suggest alternative diagnoses.

✓ In patients with a high probability of asthma:

- record the patient as likely to have asthma and commence a carefully monitored initiation of treatment (typically six weeks of inhaled corticosteroids)
- assess the patient's status with a validated symptom questionnaire, ideally corroborated by lung function tests (FEV₁ at clinic visits or by domiciliary serial peak flows to capture times with/without symptoms)
- with a good symptomatic and objective response to treatment, confirm the diagnosis of asthma and record the basis on which the diagnosis was made
- if the response is poor or equivocal, check inhaler technique and adherence, arrange further tests and consider alternative diagnoses.

2.2 Monitoring

>12 yrs 5-12 yrs <5 yrs

D	B	D	Assess risk of future asthma attacks at every asthma review by asking about history of previous attacks, objectively assessing current asthma control, and reviewing reliever use.
---	---	---	--

2.3 Supported self management

A	All people with asthma (and/or their parents or carers) should be offered self-management education, which should include a written personalised asthma action plan and be supported by regular professional review.
A	Prior to discharge, inpatients should receive written personalised asthma action plans, given by healthcare professionals with expertise in providing asthma education.
D	Adherence to long-term asthma treatment should be routinely and regularly addressed by all healthcare professionals within the context of a comprehensive programme of accessible proactive asthma care.

2.4 Non-pharmacological management

B	People with asthma and parents of children with asthma should be advised about the dangers of smoking and second-hand tobacco smoke exposure, and be offered appropriate support to stop smoking.
B	Weight loss interventions (including dietary and exercise-based programmes) should be considered for overweight and obese adults and children with asthma to improve asthma control.
A	Breathing exercise programmes (including face-to-face physiotherapist-taught methods and audiovisual programmes) can be offered to adults with asthma as an adjuvant to pharmacological treatment to improve quality of life and reduce symptoms.

2.5 Pharmacological management

✓	Before initiating a new drug therapy practitioners should check adherence with existing therapies, check inhaler technique, and eliminate trigger factors.		
A	A	A	Inhaled corticosteroids are the recommended preventer drug for adults and children for achieving overall treatment goals.
A			The first choice as add-on therapy to inhaled corticosteroids in adults is an inhaled long-acting β_2 agonist, which should be considered before increasing the dose of inhaled corticosteroids.
D	D		<p>If asthma control remains suboptimal after the addition of an inhaled long-acting β_2 agonist then:</p> <ul style="list-style-type: none"> increase the dose of inhaled corticosteroids from low dose to medium dose in adults or from very low dose to low dose in children (5–12 years), if not already on these doses. consider adding a leukotriene receptor antagonist.

2.6 Inhaler devices

- | | | | |
|----------|--|---|---|
| B | ✓ | ✓ | Prescribe inhalers only after patients have received training in the use of the device and have demonstrated satisfactory technique. |
| ✓ | Generic prescribing of inhalers should be avoided as this might lead to people with asthma being given an unfamiliar inhaler device which they are not able to use properly. | | |
| ✓ | In young children, a pMDI and spacer is the preferred method of delivery of β_2 agonists and inhaled corticosteroids. A face mask is required until the child can breathe reproducibly using the spacer mouthpiece. Where this is ineffective a nebuliser may be required. | | |

2.7 Acute asthma

2.7.1 Adults

- | | |
|----------|--|
| D | Refer to hospital any patients with features of acute severe or life-threatening asthma. |
| C | Give controlled supplementary oxygen to all hypoxaemic patients with acute severe asthma titrated to maintain an SpO₂ level of 94–98%. Do not delay oxygen administration in the absence of pulse oximetry but commence monitoring of SpO₂ as soon as it becomes available. |
| A | Use high-dose inhaled β_2 agonists as first-line agents in patients with acute asthma and administer as early as possible. Reserve intravenous β_2 agonists for those patients in whom inhaled therapy cannot be used reliably. |
| A | Give steroids in adequate doses to all patients with an acute asthma attack. |

2.7.2 Children

- | | |
|----------|---|
| ✓ | Children with life-threatening asthma or SpO ₂ <94% should receive high-flow oxygen via a tight-fitting face mask or nasal cannula at sufficient flow rates to achieve normal saturations of 94–98%. |
| A | Inhaled β_2 agonists are the first-line treatment for acute asthma in children. |
| A | Give oral steroids early in the treatment of acute asthma attacks in children. |

2.7.3 All patients

- | | |
|---|--|
| ✓ | It is essential that the patient's primary care practice is informed within 24 hours of discharge from the emergency department or hospital following an asthma attack. Ideally this communication should be directly with a named individual responsible for asthma care within the practice. |
|---|--|

2.8 Difficult asthma

- D** Patients with difficult asthma should be systematically evaluated, including:
- confirmation of the diagnosis of asthma, and
 - identification of the mechanism of persisting symptoms and assessment of adherence to therapy.

2.9 Asthma in pregnancy

B Women should be advised of the importance of maintaining good control of their asthma during pregnancy to avoid problems for both mother and baby.

C Counsel women with asthma regarding the importance and safety of continuing their asthma medications during pregnancy to ensure good asthma control.

2.10 Occupational asthma

B In patients with adult onset, or reappearance of childhood asthma, healthcare professionals should consider that there may be an occupational cause.

- ✓ Adults with suspected asthma or unexplained airflow obstruction should be asked:
- Are you the same, better, or worse on days away from work?
 - Are you the same, better, or worse on holiday?
- Those with positive answers should be investigated for occupational asthma.

3 Diagnosis

The diagnosis of asthma is a clinical one. The absence of consistent gold-standard diagnostic criteria means that it is not possible to make unequivocal evidence-based recommendations on how to make a diagnosis of asthma.

Section 3.1 defines asthma and highlights overarching principles, section 3.2 describes the diagnostic accuracy of individual symptoms, signs and diagnostic tests, and section 3.3 describes a pragmatic approach to establishing a diagnosis of asthma based on current evidence and the collective experience of the guideline development group.

3.1 Definition and overarching principles

3.1.1 Definition

Central to all definitions is the presence of symptoms (more than one of wheeze, breathlessness, chest tightness, cough) and of variable airflow obstruction. More recent descriptions of asthma, in both children and adults, have included airway hyper-responsiveness and airway inflammation as components of the disease reflecting a developing understanding of the diverse subtypes (phenotypes and endotypes) of asthma and their underpinning mechanisms.⁶

3.1.2 Tests influence the probability of asthma but do not prove a diagnosis

There is no single diagnostic test for asthma. Building on the definitions in section 3.1.1, diagnosis is based on clinical assessment (*see section 3.3*) supported by objective tests that seek to demonstrate variable airflow obstruction or the presence of airway inflammation (*see section 3.2*). Both clinical assessment of symptoms and signs and objective tests have significant false positive and false negative rates (*see Table 1*).

Objective tests influence the probability of a diagnosis of asthma, but the magnitude of that influence depends on the probability prior to testing as well as the predictive value of the test. Therefore, in a patient with a very high probability of asthma prior to testing, the results of a diagnostic test with a substantial false negative rate will have minimal influence. In contrast, in a patient with an intermediate or low probability of asthma, a positive diagnostic test may significantly shift the probability towards an asthma diagnosis (*see section 3.3*).

3.1.3 Asthma status and the outcome of diagnostic tests for asthma vary over time

Diagnostic tests are typically performed at a single point in time whereas asthma status varies over time. Patients on primary care asthma registers who have not received prescriptions for a year are considered to be 'inactive',⁷ and there is evidence that some patients shift from 'inactive' to 'active' status (and vice versa) over time.⁸⁻¹⁰

Objective tests performed when patients are asymptomatic or during an 'inactive' period may result in false negatives. For example, in primary care patients with intermittent asthma symptoms, spirometry confirmed obstruction in 16–39% of patients,¹¹⁻¹³ and bronchodilator reversibility was demonstrated in only 15–17% of patients.^{12, 14, 15} In contrast, in a population admitted to hospital with a physician-diagnosed asthma attack, 83% had obstructive lung function.¹⁶ In a prospective

2+
3

longitudinal study in primary care, fractional exhaled nitric oxide (FeNO) was only positive in 40% of people with diagnosed asthma at 12 months, and one in five were falsely negative.⁹

2+
3

Time may, however, be used to advantage if objective signs and tests when a patient is symptomatic are compared with measurements when they are asymptomatic. In the event of diagnostic uncertainty it may be helpful to repeat investigations.

C Compare the results of diagnostic tests undertaken whilst a patient is asymptomatic with those undertaken when a patient is symptomatic to detect variation over time.

3.2 Predictive value of individual symptoms, signs and diagnostic tests

The individual symptoms and signs and the diagnostic tests and thresholds typically used in clinical practice and their performance in diagnostic studies are shown in Table 1.

These data, however, need to be interpreted with caution. The performance of the diagnostic tests, as assessed by reported sensitivities/specificities and positive and negative predictive values (PPV/NPV), vary widely. This reflects methodological considerations such as the use of different reference (gold) standards and variation in defined thresholds for tests, as well as the diverse clinical contexts for these studies (*see Table 1*). The majority of studies assessing diagnostic test accuracy recruited patients from secondary care clinics; the predictive value of tests in people presenting to primary care with undifferentiated respiratory symptoms is less well reported.

The predictive value of objective tests is often poor and reinforces the need for test results to be used in conjunction with a structured clinical assessment to assess the probability of asthma in an individual presenting with respiratory symptoms suggestive of asthma (*see section 3.3.1*).

3.2.1 Symptoms and signs

The predictive value of isolated symptoms or signs is poor (*see Table 1*). In adults, isolated symptoms of cough, wheeze and shortness of breath are neither sensitive nor specific for asthma.¹⁷ Almost all children with asthma have intermittent cough, wheeze and/or exercise induced symptoms, but only about a quarter of children with these symptoms have asthma.¹⁸⁻²⁰ Enquiring about the episodic nature of symptoms (for example acute attacks) as opposed to current symptoms may improve the predictive value.^{18, 19, 21, 22}

2++
2+

Wheezing is one of a number of respiratory noises that occur in children. Parents often use the term wheezing as a non-specific label to describe any abnormal respiratory noise. It is important to distinguish wheezing – a continuous, high-pitched musical sound coming from the chest – from other respiratory noises, such as stridor or rattly breathing.²³

Wheeze heard by a healthcare professional on auscultation is an important sign that increases the probability of asthma.²²

2++

Combinations of symptoms and signs are clinically more helpful than isolated symptoms, especially in children. For example, two thirds of children with a cluster of cough, wheeze, chest tightness, dyspnoea, and exercise symptoms have asthma. Asthma is very unlikely if a child does not have at least some of these symptoms and signs.¹⁸⁻²⁰

2+

3.2.2 Spirometry and bronchodilator reversibility

Spirometry is the investigation of choice for identification of airflow obstruction and is widely available, including in primary care, although training is required to obtain reliable recordings and to interpret the results, particularly in children. The probability of asthma, differential diagnosis (*see Tables 4 and 5*) and approach to investigation is different in patients with and without airflow obstruction at the time baseline spirometry is undertaken.

Confirmation of an asthma diagnosis hinges on demonstration of airflow variability over short periods of time. A normal spirogram obtained when the patient is asymptomatic does not, therefore, exclude the diagnosis of asthma.¹¹⁻¹³ Alternative reasons for obstructive spirometry, for example chronic obstructive pulmonary disease (COPD) in adults, must also be considered. In a population of adults presenting to primary care with new respiratory symptoms, only a third of those with obstructive spirometry had asthma and almost two thirds had COPD. Only a quarter of those subsequently thought to have asthma had obstructive spirometry at the time of assessment.²⁴

2++
2+
3

Measuring lung function in children under five years of age is difficult and requires techniques which are not widely available outwith specialist centres. For developmentally mature children over five years of age conventional lung function testing is possible in most settings with an operator trained and experienced in undertaking paediatric spirometry. As in adults, normal results on testing, especially if performed when the child is asymptomatic, do not exclude a diagnosis of asthma.²⁵ Asthma severity classified by symptoms and use of medicines correlates poorly with single measurements of forced expiratory volume in one second (FEV₁) and other spirometric indices: FEV₁ is often normal in children with persistent asthma, and abnormal results may be seen in children with other respiratory diseases.^{25, 26}

2+

In children, the relationship between asthma symptoms and lung function tests, including bronchodilator reversibility, is complex. Measures of gas trapping (residual volume and the ratio of residual volume to total lung capacity) may be superior to measurement of expiratory flow at detecting airways obstruction especially in asymptomatic children.^{25, 27}



Operators should be trained to undertake quality-assured spirometry and be experienced in providing tests in the relevant age groups.

The FEV₁/forced vital capacity (FVC) ratio changes with age. In young children it can be as high as 90% so use of the commonly used fixed ratio of 70% will substantially underestimate airflow limitation. Conversely, in adults over 40 years, levels below 70% may be normal and use of a 70% threshold will overestimate obstruction. Accordingly, use of lower limits of normal is now recommended and is becoming easily available through software built into spirometers.^{6, 28-30} Detailed data about normal values for different age groups is available from the report of the European Respiratory Society Global Lung function Initiative.³¹

2+

From a practical perspective, the spirometers widely used in clinical practice provide the lower and upper limits of the normal range of spirometry parameters (although they usually use the fixed ratio to generate the automated interpretation reports).

In adults with obstructive spirometry, an improvement in FEV₁ of 12% or more in response to either β_2 agonists or corticosteroid treatment trials, together with an increase in volume of 200 ml or more, is regarded as a positive test,³² although some people with COPD can have significant reversibility.³³ An improvement of greater than 400 ml in FEV₁ strongly suggests underlying asthma. In children, an improvement in FEV₁ of 12% or more is regarded as a positive test.³²

D Carry out quality-assured spirometry using the lower limit of normal to demonstrate airway obstruction, provide a baseline for assessing response to initiation of treatment and exclude alternative diagnoses.

- **Obstructive spirometry with positive bronchodilator reversibility increases the probability of asthma.**
- **Normal spirometry in an asymptomatic patient does not rule out the diagnosis of asthma.**

3.2.3 Tests of variability in lung function

Peak expiratory flow monitoring

Peak expiratory flow (PEF) should be recorded as the best of three forced expiratory blows from total lung capacity with a maximum pause of two seconds before blowing. The patient can be standing or sitting. Further blows should be done if the largest two PEF are not within 40 L/min.³⁴

Peak expiratory flow is best used to provide an estimate of variability of airflow from multiple measurements made over at least two weeks. Increased variability may be evident from twice-daily readings. More frequent readings will result in a better estimate³⁵ but the improved precision is likely to be achieved at the expense of reduced patient compliance.³⁶ Use of electronic meters and diaries with time and date stamps can overcome problems of compliance and accuracy when recording peak flows in paper diaries.³⁷

Peak expiratory flow variability is usually calculated as the difference between the highest and lowest PEF expressed as a percentage of the average PEF,³⁸⁻⁴⁰ although one study showed that three or more days a week with significant variability was more sensitive and specific than calculating mean differences.⁴¹

The upper limit of the normal range for variability is around 20% using four or more PEF readings per day^{39, 40, 42} but may be lower using twice-daily readings.⁴³ Studies have shown sensitivities of between 3% and 46% for identifying physician-diagnosed asthma.^{17, 38, 44} One limitation of these epidemiological studies is that it is not always clear whether the participants were symptomatic at the time of the monitoring. Peak expiratory flow charting when asthma is 'inactive' is unlikely to confirm variability; one study showed that significant PEF variability was associated with respiratory symptoms in the previous week.³⁸

Peak expiratory flow records from frequent readings taken at work and away from work are useful when considering a diagnosis of occupational asthma (see section 13.3.1). A computer-generated analysis of occupational records which provides an index of the work effect is available.⁴⁵

In children, serial measures of peak-flow variability and FEV₁ show poor concordance with disease activity and do not reliably rule the diagnosis of asthma in or out.²⁶

Direct challenge tests

The most widely-used method of measuring airway responsiveness relies on measuring response in terms of change in FEV₁ a set time after inhalation of increasing concentrations of histamine or methacholine. The agent can be delivered by breath-activated dosimeter, via a nebuliser using tidal breathing, or via a hand-held atomiser.⁴⁶ The response is usually quantified as the provocative concentration (PC₂₀) or dose (PD₂₀) of bronchoconstrictor required to cause a 20% fall in FEV₁ calculated by linear interpolation of the log concentration or dose response curve. A PC₂₀ of 8 mg/ml or less is regarded as positive.^{38, 47, 48}

Two thirds, or more, of adults with a positive methacholine challenge have asthma and the false negative rate is less than 10%.¹⁷ Tests of airway responsiveness are of little value in patients with established airflow obstruction as the specificity is low.^{49, 50}

Methacholine challenge tests in schoolchildren only marginally increase the diagnostic accuracy after the symptom history is taken into account.⁵¹ However, in a child, a negative methacholine test, which has a high negative predictive value, makes a diagnosis of asthma improbable.²⁹

3

Indirect challenge tests

Other potentially helpful tests of variability in lung function include indirect challenges such as exercise and inhaled mannitol.⁵² A positive response to these indirect stimuli, such as a fall in FEV₁ of greater than 15%, is a specific marker of asthma but the tests are less sensitive than challenges using methacholine and histamine, particularly in patients tested while on treatment.^{52, 53}

In children, a positive exercise challenge test (as opposed to a history of exercise-induced symptoms) is highly predictive of asthma with a false positive rate of less than 10%.¹⁷ A negative response to an exercise challenge test is helpful in excluding asthma in children with exercise-related breathlessness.⁵⁴

3

C Referral for challenge tests should be considered in adults with no evidence of airflow obstruction on initial assessment in whom other objective tests are inconclusive but asthma remains a possibility.

- ✓ A peak flow recorded when symptomatic (eg during the assessment of an asthma attack) may be compared with a peak flow when asymptomatic (eg after recovery from an asthma attack) in order to confirm variability.
- ✓ In adults, serial peak-flow records may demonstrate variability in symptomatic patients, but should be interpreted with caution and with regard to the clinical context. There is no evidence to support the routine use of peak-flow monitoring in the diagnosis of asthma in children.
- ✓ Serial peak flows (at least four readings a day) are the initial investigation of choice in suspected occupational asthma.

3.2.4 Tests to detect eosinophilic airway inflammation or atopy

Fractional exhaled nitric oxide (FeNO)

A positive FeNO test suggests eosinophilic inflammation and provides supportive, but not conclusive, evidence for an asthma diagnosis. There is overlap between the levels seen in normal non-asthmatic populations and in people with atopic asthma.²⁸ There are some important confounders.

4

FeNO levels are:⁵⁵⁻⁵⁷

- increased in patients with allergic rhinitis exposed to allergen, even without any respiratory symptoms
- increased by rhinovirus infection in healthy individuals, but this effect is inconsistent in people with asthma
- increased in men; tall people; and by consumption of dietary nitrates
- lower in children
- reduced in cigarette smokers
- reduced by inhaled or oral steroids.

2+

4

In steroid-naïve adults, a FeNO level of 40 parts per billion (ppb) or more is regarded as positive; in schoolchildren a FeNO level of 35 ppb or more is regarded as a positive test.³²

In eight studies in adults recruited from secondary care with symptoms suggestive of asthma, sensitivities for FeNO ranged from 43–88% and specificities from 60–92%. The PPV and NPV ranged from 54–95% and 65–93%, respectively (see Table 1).¹⁷ On this basis, approximately one in five people with a positive FeNO test will not have asthma (false positives), and conversely one in five people with a negative FeNO test will have asthma (false negatives). There are no data from primary care populations.

2++

It is feasible to measure FeNO in children from the age of three to four years.⁵⁸ In children, FeNO is closely linked with atopic status, age and height.^{59, 60}

D Use measurement of FeNO (if available) to find evidence of eosinophilic inflammation. A positive test increases the probability of asthma but a negative test does not exclude asthma.

Tests of atopic status

Positive skin-prick tests,⁶¹ blood eosinophilia $\geq 4\%$,⁶² or a raised allergen-specific immunoglobulin E (IgE) to a range of common aeroallergens^{63, 64} increase the probability of asthma in schoolchildren and adults.^{32, 61} The PPV for individual tests are, however, poor (see Table 1). Non-atopic wheezing is as frequent as atopic wheezing in school-aged children.⁶⁵

2++

4

D Use a previous record of skin-prick tests, blood eosinophilia of 4% or more, or a raised allergen-specific IgE to corroborate a history of atopic status, but do not offer these tests routinely as a diagnostic test for asthma.

Sputum eosinophils

Eosinophilic airway inflammation in adults can be assessed non-invasively using the induced sputum differential eosinophil count.^{55, 66} Sputum induction is feasible in school-aged children but is technically demanding and time consuming and remains a research tool.^{67, 68} Experience with induced sputum is limited to a few centres and more research needs to be done before any recommendations can be made on its use as a diagnostic test in clinical practice.

Table 1: Summary of individual diagnostic tests

Strategy	Description*	Parameter*	Range of predictive values* (Note that a single value indicates data from a single study)				Comments**
			Sens ⁱ	Spec ⁱⁱ	PPV ⁱⁱⁱ	NPV ^{iv}	
Clinical assessment							
Symptoms and signs	The commonest symptoms assessed were cough and wheeze and, in adults, shortness of breath.	Cough in adults	16–66%	26–64%	8–44%	18–92%	As isolated symptoms cough, wheeze and shortness of breath are neither sensitive, nor specific for asthma. Most children with asthma have intermittent cough, wheeze and exercise-induced symptoms, but only about a quarter of children with these symptoms have asthma. Note that the single study in preschool children compared current symptoms with a diagnosis of asthma two years later.
		Wheeze in adults	9–76%	34–87%	10–81%	28–94%	
		Dyspnoea in adults	11–73%	38–71%	41–59%	26–70%	
		Cough in schoolchildren ²⁰	63%	75%	14%	97%	
		Wheeze in children ²⁰	59%	93%	34%	97%	
		Cough in preschool children	88%	7%	76%	15%	
Symptom variability		Wheeze in preschool children	54%	57%	80%	27%	Asking about episodic symptoms improves the positive predictive values in children compared with current symptoms.
		Shortness of breath in preschool children	76%	52%	84%	40%	
		Episodic symptoms in adults	9–40%	36–91%	14–86%	18–93%	
		Diurnal symptoms in adults	30–56%	36–83%	48–76%	18–67%	
		Symptoms after exercise in adults	5–40%	32–93%	5–81%	58–84%	
Combinations of symptoms (typically cough, wheeze, chest tightness, dyspnoea, exercise symptoms)		Episodic symptoms in children ^{18, 19}	36–93%	35–93%	40–94%	62–90%	Combinations of symptoms are clinically more helpful than isolated symptoms, especially in children. For example, two thirds of children with a cluster of cough, wheeze, chest tightness, dyspnoea and exercise symptoms have asthma. Asthma is unlikely if a child does not have at least some of these symptoms.
		Symptoms after exercise in children ^{18, 19}	82–94%	59–73%	54–86%	79–91%	
		Nocturnal symptoms in children ^{18, 19}	57–84%	58–78%	64–85%	57–82%	
		Symptom scores in adults	60%	66%	44–94%	66–97%	
		Symptom scores in children ^{18–20}	45–83%	85–97%	80%	51%	
		Symptoms of cough and wheeze in preschool children	49%	59%			

Table 1: Summary of individual diagnostic tests (continued)

Strategy	Description*	Parameter*	Range of predictive values* (Note that a single value indicates data from a single study)				Comments**
			Sens ⁱ	Spec ⁱⁱ	PPV ⁱⁱⁱ	NPV ^{iv}	
Clinical assessment (continued)							
History of atopy	Personal/family history of atopic/allergic diseases	Personal history of atopy in adults	54-55%	68-74%	46-76%	45-79%	Past history (personal or family) of atopic disease has poor sensitivity and specificity for asthma.
		Personal history of rhinitis/eczema in preschool children	47-62%	20-75%	72-86%	14-30%	
		Family history of atopy in adults	26-60%	56-83%	44-74%	38-70%	
		Family history of atopy in children	43-44%	57-70%	51-77%	24-62%	
Strategies for demonstrating airway obstruction							
Spirometry	Regard a FEV ₁ /FVC ratio of less than 70% as a positive test for obstructive airway disease.	Obstructive spirometry in adults Obstructive spirometry in children (5-18 yrs)	23-47% 52%	31-100% 73%	45-100% 75%	18-73% 49%	In the four larger studies (adults and children), the NPV was between 18% and 54% which means that more than half of patients being investigated who have normal spirometry will have asthma (ie false negatives).
Strategies for demonstrating variability in airway obstruction							
Bronchodilator reversibility	In adults, regard an improvement in FEV ₁ of ≥12% and ≥200 ml as a positive test. In children regard an improvement in FEV ₁ of ≥12% as a positive test.	Bronchodilator reversibility in adults Bronchodilator reversibility in schoolchildren (using a threshold of 9% change in FEV ₁) ⁶⁹	17-69% 50%	55-81% 86%	53-82%	22-68%	In these secondary care populations, about one in three people with a positive reversibility test will not have asthma (the cohorts all included people with COPD); and at least one in three people with a negative bronchodilator reversibility test will have asthma.

Table 1: Summary of individual diagnostic tests (continued)

Strategy	Description*	Parameter*	Range of predictive values* (Note that a single value indicates data from a single study)				Comments**
			Sens ⁱ	Spec ⁱⁱ	PPV ⁱⁱⁱ	NPV ^{iv}	
Strategies for demonstrating variability in airway obstruction (continued)							
Challenge tests	Regard a PC ₂₀ value of 8 mg/ml or less as a positive test.	Methacholine challenge in adults. Methacholine challenge in children ^{29, 51, 70}	51–100% 47–86%	39–100% 36–97%	60–100% 20%	46–100% 94%	Challenge tests are a good indicator for those with a definitive diagnosis of asthma already (based upon clinical judgment, signs and symptoms and response to antiasthma therapy)
	Fall in FEV ₁ ≥15% at cumulative dose of ≤635 mg is positive	Mannitol in adults Mannitol in children	56% 63%	75% 81%	80%	49%	These data are from a single study in adults and children with symptoms of asthma on questionnaire.
	Exercise challenge (range of thresholds used: 8-20% fall in FEV ₁)	Exercise challenge in adults Exercise challenge in children	26–80% 69–72%	100% 69–72%	100% 90–99%	0% 5–73%	The studies in adults had very small sample sizes. The larger study in children had a false positive rate of 1% (PPV 99%).
Peak-flow charting	Monitor peak flows for 2–4 weeks, calculate mean variability. Regard ≥20% variability as a positive test.	PEF charting in adults in a population study - using mean variability of >20% - using mean variability of >15% - using diurnal variation >15% on >3 days/week PEF charting in children - using variation >12.3% (95 th centile)	46% 3–5% 20%	80% 98–99% 97%	97% 60–67% 82%	10% 60% 64%	It is not clear whether the patients in these studies were symptomatic at the time of the charting, and results may not reflect clinical use in symptomatic populations. One study concluded that the number of days with diurnal variation was more accurate than calculating the mean variation.
Strategies for detecting eosinophilic inflammation or atopy							
FeNO	Adults: Regard a FeNO level of 40 ppb or more as a positive test Children 5–16yrs: regard a FeNO level of 35 ppb or more as a positive test	FeNO in adults FeNO in schoolchildren	43–88% 57%	60–92% 87%	54–95% 90%	65–93% 49%	These studies are all in secondary care populations. Approximately one in five adults with a positive FeNO test will not have asthma (ie false positives) and one in five adults with a negative FeNO test will have asthma (ie false negatives).

Table 1: Summary of individual diagnostic tests (continued)

Strategy	Description*	Parameter*	Range of predictive values* (Note that a single value indicates data from a single study)				Comments**
			Sens ⁱ	Spec ⁱⁱ	PPV ⁱⁱⁱ	NPV ^{iv}	
Strategies for detecting eosinophilic inflammation or atopy (continued)							
Blood eosinophils	Suggested thresholds for blood eosinophils: Adults >4.15% Children ≥4% ⁶²	Blood eosinophils in adults Blood eosinophils in children	15–36% 55–62%	39–100% 67–84%	39–100% 56–69%	27–65% 73%	Elevated blood eosinophil level is poorly predictive. The threshold varies in these studies from 4.0–6.3%.
IgE		Any allergen-specific IgE >0.35 kU/L in adults Total IgE in adults >100 kU/L	54–93% 57%	67–73% 78%	5–14% 5%	95–99% 99%	A normal IgE substantially reduces the probability of asthma in adults with a false negative rate of less than 1 in 10, although a positive result is poorly predictive.
Skin-prick testing		Any positive test (wheal ≥3 mm) in adults Any positive test (wheal ≥3 mm) in children	61–62% 44–79%	63–69% 56–92%	14–81% 65–92%	39–96% 36–79%	

Notes:

* Data derived from NICE evidence tables unless otherwise specified.¹⁷ Only studies reporting sensitivity, specificity, PPV and NPV are included here.

** Comments have been added by the guideline development group as an aid to interpretation of the data presented.

i Sensitivity (Sens) is the probability of a test being positive when asthma is present.

ii Specificity (Spec) is the probability of a test being negative when asthma is absent.

iii Positive predictive value (PPV) is the proportion of patients with a positive test who actually have asthma (100 minus the PPV is the proportion of patients with a false positive test).

iv Negative predictive value (NPV) is the proportion of patients with a negative test who do not have asthma (100 minus the NPV is the proportion of patients with asthma but in whom test was negative).

Reference tests

In most of the studies, the reference test was spirometry plus either bronchodilator reversibility or a challenge test, although some studies also included a 'typical history of attacks' or diurnal variation, or used physician diagnosis. Studies evaluating methacholine challenge tests used physician diagnosis or bronchodilator reversibility and/or diurnal peak-flow variability. In children, the reference tests used were physician-diagnosed asthma plus spirometry, or documented history of wheeze on at least two occasions, and variability in FEV₁ over time or on exercise testing.

3.3 Practical approach to diagnosis

The diagnosis of asthma in children and adults is based on the recognition of a characteristic pattern of respiratory symptoms, signs and test results (*see Table 2*) and the absence of any alternative explanation for these.

At present, there is no definitive evidence to inform the most appropriate choice of algorithm for making a diagnosis of asthma in clinical settings. There are pragmatic observational studies which can inform the clinical process of making a diagnosis,^{9, 12, 13, 15, 71} or which compare outcomes of diagnostic tests in different settings,^{16, 21, 72} and some potentially useful algorithms,^{11, 20-22, 72} or symptom questionnaires in children have been derived.^{18, 19} This section and the associated diagnostic algorithm (*see Figure 1*), therefore, represent consensus opinion, building on the overarching principles defined in section 3.1. It is further informed by the evidence available from these pragmatic studies combined with data from the diagnostic studies described in section 3.2. There is an urgent need for diagnostic accuracy studies and implementation research to confirm, prospectively, the diagnostic accuracy of retrospectively-derived algorithms and to define the optimal approach to making a diagnosis in different clinical practice settings.

All studies evaluating diagnostic approaches have used a clinical assessment, sometimes using diagnostic,⁷² or standard morbidity questions,^{15, 71} as the basis for the diagnostic process.^{11-15, 22, 71, 72} A number of studies have highlighted the diagnostic significance of episodic symptoms and confirmed wheezing as important predictors of asthma.^{13, 16, 18-20, 22} Studies also illustrate the importance of observing events over time and documenting the basis on which a diagnosis is made.^{9, 72}

In adults, absence of smoking and young age of onset are typically included in algorithms designed to distinguish asthma from COPD.^{12, 14, 15, 71, 72}

3.3.1 Initial structured clinical assessment

The predictive value of individual symptoms or signs is poor (*see Table 1*), and a structured clinical assessment including all information available from the history, examination and historical records should be undertaken. The clinical features that influence the probability that episodic respiratory symptoms are due to asthma are summarised in Table 2.

Alternative explanations for the symptoms or signs and/or the possibility of comorbid conditions such as COPD in adults with a smoking history, obesity, and dysfunctional breathing, which can produce features that mimic asthma, must be considered (*see Tables 4 and 5*). For working adults with airflow obstruction, occupational asthma should be considered and suitable screening questions asked (*see section 13.3*).

Table 2: Factors to consider in an initial structured clinical assessment

<p>Episodic symptoms (see sections 3.2.1 and 3.2.2)^{13, 18, 19, 21, 22 62, 73, 74}</p> <p>More than one of the symptoms of wheeze, breathlessness, chest tightness and cough occurring in episodes with periods of no (or minimal) symptoms between episodes. Note that this excludes cough as an isolated symptom in children.⁷⁵ For example:</p> <ul style="list-style-type: none"> • a documented history of acute attacks of wheeze, triggered by viral infection or allergen exposure with symptomatic and objective improvement with time and/or treatment • recurrent intermittent episodes of symptoms triggered by allergen exposure as well as viral infections and exacerbated by exercise and cold air, and emotion or laughter in children • in adults, symptoms triggered by taking non-steroidal anti-inflammatory medication or beta blockers. <p>An historical record of significantly lower FEV₁ or PEF during symptomatic episodes compared with asymptomatic periods provides objective confirmation of the obstructive nature of the episodic symptoms.</p>
<p>Wheeze confirmed by a healthcare professional on auscultation (see section 3.2.1)²²</p> <p>It is important to distinguish wheezing from other respiratory noises, such as stridor or rattly breathing.</p> <p>Repeatedly normal examination of chest when symptomatic reduces the probability of asthma.</p>
<p>Evidence of diurnal variability^{18, 19, 22, 32, 73}</p> <p>Symptoms which are worse at night or in the early morning.</p>
<p>Atopic history (see section 3.2.4)^{17, 22, 62, 74, 76, 77}</p> <p>Personal history of an atopic disorder (ie eczema or allergic rhinitis) or a family history of asthma and/or atopic disorders, potentially corroborated by a previous record of raised allergen-specific IgE levels, positive skin-prick tests to aeroallergens or blood eosinophilia.</p>
<p>Absence of symptoms, signs or clinical history to suggest alternative diagnoses (including but not limited to COPD, dysfunctional breathing, obesity) (see section 3.3.3)</p>

D Undertake a structured clinical assessment to assess the initial probability of asthma. This should be based on:

- a history of recurrent episodes (attacks) of symptoms, ideally corroborated by variable peak flows when symptomatic and asymptomatic
- symptoms of wheeze, cough, breathlessness and chest tightness that vary over time
- recorded observation of wheeze heard by a healthcare professional
- personal/family history of other atopic conditions (in particular, atopic eczema/dermatitis, allergic rhinitis)
- no symptoms/signs to suggest alternative diagnoses.

3.3.2 High probability of asthma based on initial structured clinical assessment

Adults and children with a typical clinical assessment including recurrent episodes of symptoms (attacks), wheeze heard by a healthcare professional, historical record of variable airflow obstruction and a positive history of atopy (*see Table 2*) and without any features to suggest an alternative diagnosis (*see Tables 4 and 5*) have a high probability of asthma.²² If there is doubt, the diagnosis should be considered as being of intermediate probability and further investigations will be needed (*see section 3.3.4*). 2+

Obstructive spirometry and a positive bronchodilator test provide objective evidence of variable airflow obstruction,¹⁷ and further increase the probability of asthma.^{20, 22, 24} However, as spirometry has a false negative rate of at least 50%,¹⁷ normal spirometry does not rule out asthma.²⁴ If the patient is symptomatic peak-flow charting, if performed correctly, may provide objective evidence of variability. 2++
2+

- ✓ In patients with a high probability of asthma:
- record the patient as likely to have asthma and commence a carefully monitored initiation of treatment (typically six weeks of inhaled corticosteroids) (*see Table 3*)
 - assess the patient's status with a validated symptom questionnaire, ideally corroborated by lung function tests (FEV₁ at clinic visits or by domiciliary serial peak flows to capture times with/without symptoms)
 - with a good symptomatic and objective response to treatment, confirm the diagnosis of asthma and record the basis on which the diagnosis was made
 - if the response is poor or equivocal, check inhaler technique and adherence, arrange further tests and consider alternative diagnoses.

3.3.3 Low probability of asthma based on initial structured clinical assessment

Adults and children who do not have any of the typical features on initial structured clinical assessment (*see Table 2*) or who have symptoms suggestive of an alternative diagnosis (*see Tables 4 and 5*) have a low probability of asthma.

- ✓ If there is a low probability of asthma and/or an alternative diagnosis is more likely, investigate for the alternative diagnosis, reconsidering asthma if the clinical picture changes or an alternative diagnosis is not confirmed. If reconsidering asthma, undertake or refer for further tests to investigate for a diagnosis of asthma.

Table 3: A monitored initiation of treatment in patients with suspected asthma

In patients with suspected asthma	
1.	Record the patient as having 'suspected asthma'.
2.	Proceed to a carefully monitored initiation of treatment. The initial choice of treatment will be based on an assessment of the degree of asthma severity. Typically this will be six weeks of inhaled steroids through a device the patient can use (<i>see sections 7.2, 8.1, 8.4</i>) but in more acute clinical circumstances a course of oral steroids may be appropriate (<i>see section 9.3.3</i>).
3.	Assess the baseline status using a validated questionnaire (eg Asthma Control Questionnaire or Asthma Control Test) (<i>see Table 8</i>) and/or lung function tests (spirometry or peak expiratory flow) (<i>see sections 3.2.2, 3.2.3</i>).
4.	Arrange a follow-up appointment in 6–8 weeks in order to assess response to treatment.
5.	At the follow-up appointment, symptomatic response may be assessed with a validated questionnaire (<i>see Table 8</i>). Lung function may be monitored with FEV ₁ at clinic visits or domiciliary serial peak flows.
If the objective response is good (ie a clinically important improvement in symptoms and/or substantial increase in lung function)	
6.	Confirm the diagnosis of asthma and record the basis on which the diagnosis was made.
7.	Adjust the treatment according to the response (eg, titrating down the dose of inhaled steroid) to the lowest dose that maintains the patient free of symptoms. Careful observation during a trial of withdrawing treatment will also identify patients whose improvement was due to spontaneous remission (this is particularly important in children).
8.	Provide self-management education and a personalised asthma action plan (<i>see section 5.2.2</i>) before arranging repeat prescribing so that the patient is aware of the action to take if their control deteriorates.
If the objective response is poor or equivocal	
9.	Discuss adherence and recheck inhaler technique as possible causes of treatment failure.
10.	Arrange further tests or consider alternative diagnoses (<i>see section 3.3.3</i>). It will usually be appropriate to withdraw the treatment.

Table 4: Clinical clues to alternative diagnoses in wheezy children

Clinical clue	Possible diagnosis
Perinatal and family history	
Symptoms present from birth or perinatal lung problem	Cystic fibrosis; chronic lung disease of prematurity; ciliary dyskinesia; developmental lung anomaly
Family history of unusual chest disease	Cystic fibrosis; neuromuscular disorder
Severe upper respiratory tract disease	Defect of host defence; ciliary dyskinesia
Symptoms and signs	
Persistent moist cough ⁷⁸	Cystic fibrosis; bronchiectasis; protracted bacterial bronchitis; recurrent aspiration; host defence disorder; ciliary dyskinesia
Excessive vomiting	Gastro-oesophageal reflux (with or without aspiration)
Paroxysmal coughing bouts leading to vomiting	Pertussis
Dysphagia	Swallowing problems (with or without aspiration)
Breathlessness with light-headedness and peripheral tingling	Dysfunctional breathing, panic attacks
Inspiratory stridor	Tracheal or laryngeal disorder
Abnormal voice or cry	Laryngeal problem
Focal signs in chest	Developmental anomaly; post-infective syndrome; bronchiectasis; tuberculosis
Finger clubbing	Cystic fibrosis; bronchiectasis
Failure to thrive	Cystic fibrosis; host defence disorder; gastro-oesophageal reflux
Investigations	
Focal or persistent radiological changes	Developmental lung anomaly; cystic fibrosis; postinfective disorder; recurrent aspiration; inhaled foreign body; bronchiectasis; tuberculosis

Table 5: Clinical clues to alternative diagnoses in adults

Clinical clue	Possible diagnosis
Without airflow obstruction	
Predominant cough without lung function abnormalities	Chronic cough syndromes; pertussis
Prominent dizziness, light-headedness, peripheral tingling	Dysfunctional breathing
Recurrent severe 'asthma attacks' without objective confirmatory evidence	Vocal cord dysfunction
Predominant nasal symptoms without lung function abnormalities	Rhinitis
Postural and food-related symptoms, predominant cough	Gastro-oesophageal reflux
Orthopnoea, paroxysmal nocturnal dyspnoea, peripheral oedema, pre-existing cardiac disease	Cardiac failure
Crackles on auscultation	Pulmonary fibrosis
With airflow obstruction	
Significant smoking history (ie, >30 pack-years), age of onset >35 years	COPD
Chronic productive cough in the absence of wheeze or breathlessness	Bronchiectasis*; inhaled foreign body*; obliterative bronchiolitis; large airway stenosis
New onset in smoker, systemic symptoms, weight loss, haemoptysis	Lung cancer*; sarcoidosis*

* may also be associated with non-obstructive spirometry

3.3.4 Intermediate probability of asthma based on initial structured clinical assessment

Adults and children who have some, but not all, of the typical features of asthma on an initial structured clinical assessment (see Table 2) or who do not respond well to a monitored initiation of treatment (see Table 3) have an intermediate probability of asthma.^{20, 22, 69} They require clinical assessment and investigation before a diagnosis can be made and, unless the clinical condition is acute, before treatment is commenced or continued. Particular care may be needed in conditions known to overlap with or mimic asthma, for example COPD (which may need to be distinguished from fixed airflow obstruction as a result of airway remodelling in chronic asthma), obesity, anxiety/panic, or dysfunctional breathing.

2+
3

Spirometry enables differentiation of obstructive and non-obstructive lung function, which determines the differential diagnosis (see Tables 4 and 5) and approach to investigation. Spirometry is useful for confirming the diagnosis of asthma but is not sufficiently specific to rule it out.^{17, 79}

D Spirometry, with bronchodilator reversibility as appropriate, is the preferred initial test for investigating intermediate probability of asthma in adults, and in children old enough to produce reliable results on testing.

Adults and children with airways obstruction

Asthma is the commonest cause of airways obstruction identified through spirometry in children. Obstruction due to other disorders is much more common in adults than in children. Patients may have more than one cause of airflow obstruction, which complicates the interpretation of any test. In particular, asthma and COPD commonly coexist in adults.

A bronchodilator reversibility test and/or a monitored initiation of treatment (typically six weeks of inhaled corticosteroids (ICS) can establish whether or not the airflow obstruction reverses to normal with treatment. Evidence of a symptomatic response, ideally using objective measures of asthma control and lung function, should be sought at a follow-up visit. If there is significant reversibility or improvement in symptom scores, confirm the diagnosis of asthma and record the basis on which the diagnosis was made. Continue to treat as asthma, but aim to find the minimum effective dose of therapy.

If the patient remains asymptomatic consider a trial of reduction or withdrawal of treatment. This is particularly important in children in whom natural resolution of symptoms is more common than in adults.

- ✓ In adults and children with an intermediate probability of asthma and airways obstruction identified through spirometry, undertake reversibility tests and/or a monitored initiation of treatment assessing the response to treatment by repeating lung function tests and objective measures of asthma control.

Adults and children without airways obstruction

In patients with normal spirometry results consider arranging challenge tests with methacholine, exercise or mannitol in order to test for airway hyper-responsiveness.^{17, 29, 51} Alternatively, a positive FeNO test indicates the presence of eosinophilic inflammation and increases the probability of asthma.^{17, 32, 79-82} Investigation of atopic status, serum-specific IgE and allergen skin-prick tests may be of value in selected patients; a normal result reduces the probability of asthma.¹⁷ Consider performing additional investigations such as full lung function tests and a chest X-ray in any patient presenting with atypical or additional symptoms or signs. A study in primary care in children aged six and under concluded that a chest X-ray, in the absence of a clinical indication, need not be part of the initial diagnostic work up but may be reserved for children with severe disease or clinical clues suggesting other conditions.⁸³

- ✓ In adults and children with an intermediate probability of asthma and normal spirometry results, undertake challenge tests and/or measurement of FeNO to identify eosinophilic inflammation.

Children unable to undertake spirometry

In some children, and particularly preschool children, there is insufficient evidence at the first consultation to make a firm diagnosis of asthma, but no features to suggest an alternative diagnosis. There are several possible approaches to reaching a diagnosis in this group. Which approach is taken will be influenced by the frequency and severity of the symptoms. These approaches include:


Watchful waiting with review. In children with mild intermittent wheeze and other respiratory symptoms that occur only with viral upper respiratory infections, it is often reasonable to give no maintenance treatment and to plan a review of the child after an interval agreed with the parents/carers.

Monitored initiation of treatment (see Table 3). Most children under five years of age and some older children cannot perform spirometry. In these children, offer a monitored initiation of treatment for a specific period. The choice of treatment (for example inhaled corticosteroids) depends on the severity and frequency of symptoms.

Monitor treatment for six to eight weeks and if there is clear evidence of clinical improvement, the treatment should be continued and they should be regarded as having asthma (it may be appropriate to consider a trial of withdrawal of treatment at a later stage). If the treatment trial is not beneficial, then consider tests for alternative conditions and referral for specialist assessment.

- ✓ In children with an intermediate probability of asthma who cannot perform spirometry:
- consider watchful waiting if the child is asymptomatic
 - offer a carefully monitored initiation of treatment if the child is symptomatic.

Figure 1: Diagnostic algorithm



¹ In children under 5 years and others unable to undertake spirometry in whom there is a high or intermediate probability of asthma, the options are monitored initiation of treatment or watchful waiting according to the assessed probability of asthma.

3.3.5 Diagnostic indications for referral

At any point in the diagnostic algorithm, there may be a need for referral for additional investigations and/or specialist advice. Some key indications for referral to specialist care are listed in Table 6.

Table 6: Diagnostic indications for specialist referral

Adults	Children
Referral for tests not available in primary care	
Diagnosis unclear	Diagnosis unclear
Suspected occupational asthma (symptoms that improve when patient is not at work, adult-onset asthma and workers in high-risk occupations) ⁸⁴	
Poor response to asthma treatment	Poor response to monitored initiation of asthma treatment
Severe/life-threatening asthma attack	Severe/life-threatening asthma attack
'Red flags' and indicators of other diagnoses	
Prominent systemic features (myalgia, fever, weight loss)	Failure to thrive
Unexpected clinical findings (eg crackles, clubbing, cyanosis, cardiac disease, monophonic wheeze or stridor)	Unexplained clinical findings (eg focal signs, abnormal voice or cry, dysphagia, inspiratory stridor)
Persistent non-variable breathlessness	Symptoms present from birth or perinatal lung problem
Chronic sputum production	Excessive vomiting or possetting
Unexplained restrictive spirometry	Severe upper respiratory tract infection
Chest X-ray shadowing	Persistent wet or productive cough
Marked blood eosinophilia	Family history of unusual chest disease
	Nasal polyps
Patient or parental anxiety or need for reassurance	

3.4 Organisation of diagnostic services

A structured clinical assessment and some diagnostic tests (for example spirometry with bronchodilator reversibility) are readily available in primary care, although specialist expertise may be needed in young children. Other tests, such as FeNO and skin-prick testing, are only available in some secondary care settings and a few primary care practices. Some tests (for example challenge tests) will require referral to a diagnostic centre.

In the future, this may require additional provision of specialist-led diagnostic services to support general practitioner (GP) assessment. For example a regional asthma-COPD diagnostic service in the Netherlands available to support GPs' assessment reported that the service agreed with the GPs' working diagnosis of asthma in 62% of cases, and was able to provide a diagnosis for 95% of the patients in whom GPs were uncertain.^{15, 71}

3

C Streamlined referral pathways should be developed for tests which are not routinely available in primary care.

3.5 Wheezing in preschool children and the future risk of developing persistent asthma

Several factors are associated with a risk of developing persisting wheezing or asthma through childhood.^{76, 85} The presence of these factors increases the probability that a child with respiratory symptoms will have asthma.

These factors include:

Age at presentation

The natural history of wheeze is dependent on age at first presentation. In general, the earlier the onset of wheeze, the better the prognosis. Cohort studies show a break point at around two years; most children who present before this age become asymptomatic by mid-childhood.⁸⁶⁻⁸⁹ Coexistent atopy is a risk factor for persistence of wheeze independent of age of presentation. **2++**

Sex

Male sex is a risk factor for asthma in prepubertal children. Female sex is a risk factor for the persistence of asthma in the transition from childhood to adulthood.^{90, 91} Boys with asthma are more likely to grow out of their asthma during adolescence than girls.^{62, 86, 90, 92-105}

Severity and frequency of previous wheezing episodes

Frequent or severe episodes of wheezing in childhood are associated with recurrent wheeze that persists into adolescence.^{74, 77, 86, 88, 94, 106-108}

Coexistence of atopic disease

A history of other atopic conditions such as eczema and rhinitis increases the probability of asthma. Positive tests for atopy in a wheezing child also increase the likelihood of asthma. A raised specific IgE to wheat, egg white, or inhalant allergens, such as house dust mite and cat dander, predicts later childhood asthma.^{109, 110} **2++**

Other markers of allergic disease at presentation, such as positive skin-prick tests and a raised blood eosinophil count, are related to the severity of current asthma and persistence through childhood.

Family history of atopy

A family history of atopy is the most clearly defined risk factor for atopy and asthma in children. The strongest association is with maternal atopy, which is an important risk factor for the childhood onset of asthma and for recurrent wheezing that persists throughout childhood.^{87, 102, 105, 111, 112} **2++**

Abnormal lung function

Persistent reductions in baseline airway function and increased airway responsiveness during childhood are associated with having asthma in adult life.⁹¹ **3**

4 Monitoring asthma

Regular review of people with asthma offers the opportunity to monitor current symptom control and the impact asthma is having on daily activities and quality of life, to assess future risk of asthma attacks, and to link these to management options.

Asthma is best monitored by routine clinical review on at least an annual basis by a healthcare professional with appropriate training in asthma management. The review can be undertaken in primary and/or secondary care according to clinical need and local service arrangements (*see section 14.3*). The suggested components of a review are listed in Table 7.

- ✓ The core components of an asthma review that should be assessed and recorded on at least an annual basis are current symptoms, future risk of attacks, management strategies, supported self management, and growth in children.

Table 7: Components of an asthma review

Parameters	Suggested assessment	Further information
Current symptom control	<ul style="list-style-type: none"> • Bronchodilator use • Validated symptom score • Time off work/school due to asthma 	Section 4.2, Table 8
Future risk of attacks	<ul style="list-style-type: none"> • Past history of asthma attacks • Oral corticosteroid use • Prescription data: frequent short-acting β_2 agonist (SABA) and infrequent ICS • Exposure to tobacco smoke 	Section 4.3 Section 7.5.3 Sections 7.1.1, 9.1.2 Section 6.2.3
Tests/ investigations	<ul style="list-style-type: none"> • Lung function (spirometry or by PEF) • Growth (height and weight centile) in children 	Section 4.4.1
Management	<ul style="list-style-type: none"> • Inhaler technique • Adherence (self report, prescription refill frequency) • Non-pharmacological management (trigger avoidance, breathing exercises) • Pharmacological management - consider multimorbidity and polypharmacy 	Section 8 Section 5.4 Section 6 Section 7 Sections 7.7.3–7.7.7
Supported self-management	<ul style="list-style-type: none"> • Education/discussion about self management • Provision/revision of a written personalised asthma action plan 	Sections 5.2.1, 5.3 Section 5.2.2

4.1 Targeting care

Identifying people with poor symptom control and at future risk of asthma attacks enables targeting of care:

- at the level of the individual patient, their families and carers (eg by increasing frequency of review, commencing/increasing preventer medication, personalisation of an asthma action plan, avoiding triggers such as smoking, shifting the necessity/concerns balance for ICS treatment)
- from a public health perspective (eg addressing any inequities of care due to deprivation)
- for policy makers (ensuring access to care).

Targeting care on people/populations identified as being poorly controlled or at greater risk of asthma attacks has the potential to improve the quality of life of people with asthma and their families.

Research is needed to understand the health service impact that stepping up care for a high-risk population has on the resources available for those at lower risk. Promotion of supported self management at a policy and organisational level (*see section 5.5*) is likely to be an important strategy for ensuring the safety of all people with asthma.

4.2 Monitoring current asthma symptom control

When assessing asthma symptoms, a general question, such as “how is your asthma today?”, is likely to yield a non-specific answer such as, “I am OK”. Using direct questions, such as “do you use your reliever (blue inhaler) every day/or do you use it more than two times per week”, or the Royal College of Physicians’ ‘3 Questions’,¹¹³ is more likely to yield useful information about current control.¹¹⁴ Symptomatic asthma control is best measured using validated questionnaires, for example the Asthma Control Questionnaire (ACQ) or Asthma Control Test (ACT) or Childhood Asthma Control Test (C-ACT) (*see Table 8*).

- ✓ When asking about asthma symptoms, use specific questions, such as the Royal College of Physicians’ ‘3 Questions’ or questions about reliever use, with positive responses prompting further assessment with a validated questionnaire to assess symptom control.
- ✓ Whenever practicable, children should be asked about their own symptoms; do not rely solely on parental report.

Table 8: Summary of tools that can be used to assess current asthma symptom control

Measurement	Methodology	Measurement characteristics	Comments
Royal College of Physicians '3 Questions' ¹¹³	Yes/no or graded response to three questions about the impact of asthma symptoms on sleep, daytime symptoms, and activities.	No to all questions consistent with controlled asthma. Any positive answers should prompt further assessment.	Direct questions for use in day-to-day clinical practice, ¹¹⁴ but this is not a validated control questionnaire.
Asthma Control Questionnaire (ACQ) ¹¹⁵⁻¹¹⁸	Five questions about symptoms over the preceding week plus optional reliever use and FEV ₁ ¹¹⁷	Well controlled ≤ 0.75 Poor control ≥ 1.5 Minimal important difference 0.5	Well validated in adults and children aged 5 and over. Available in many languages
Asthma Control Test (ACT) ^{119, 120} and Childhood ACT (C-ACT) for 4-11 year olds ¹²¹	ACT: five questions about symptoms, reliever use, and overall control over preceding four weeks. C-ACT: seven questions, four for the child, three for parent/carer	Poor control < 20 Minimal important difference 3 ¹²²	ACT well validated in adults and C-ACT in children 4-11 years
Mini Asthma Quality of Life Questionnaire (AQLQ) ^{116, 123, 124} Paediatric Asthma Quality of Life Questionnaire (PAQLQ) for 7-11 year olds ¹²⁵	15 questions in four domains (symptoms, activity limitations, emotional function and environmental stimuli) over preceding two weeks. The PAQLQ has 23 questions	Mean score (between one and seven) across the four domains; higher scores indicate better quality of life. Minimal important difference 0.5	Mini AQLQ well validated in adults and PAQLQ in children 7-11 years.

4.3 Predicting future risk of asthma attacks

Identifying future risk of asthma attacks is an important component in the delivery of personalised asthma care.¹²⁶

4.3.1 Adults

It is possible to identify an adult with asthma at increased risk of an asthma attack and to stratify the degree of risk associated with different markers. The factors associated with increased risk in adults are shown in Table 9.

2++
3

Table 9: Factors associated with increased risk of future asthma attacks in adults

Level of increased risk	Factor
Greatly increased risk	History of previous asthma attacks ¹²⁷⁻¹³⁷
Moderately increased risk	Poor control ^{128, 131-136, 138-142} (assess at every routine review using objective patient-reported control questionnaires, eg ACT ^{129, 135, 136, 141} or ACQ ^{133, 139}) Inappropriate or excessive SABA use ^{127, 132, 140, 142}
Slightly increased risk	Older age ^{127, 132, 137, 143} Female ^{127, 131, 132, 137, 143} Reduced lung function ^{127, 130, 133, 135, 142} Obesity ^{127, 130, 132, 143} Smoking ^{127, 132, 143} Depression ^{132, 143, 144}
Unclear (evidence limited or equivocal)	A history of anaphylaxis ^{127, 132} Comorbid gastro-oesophageal reflux ^{127, 132} COPD ¹⁴³ Raised FeNO at routine reviews ^{129, 138, 145, 146} Blood eosinophilia ^{127, 129, 132, 136, 146-148} Poor adherence ^{127, 136, 149, 150}

There is insufficient evidence in adults to say if the following factors are associated with an increased risk of future asthma attacks: concomitant prescription of paracetamol or non-steroidal anti-inflammatory medicines,¹³² deprivation,¹⁴³ being underweight,¹⁴³ frequent febrile upper respiratory tract infection,¹³⁷ diabetes,¹³² pollution,¹⁵¹ black or Indian ethnicity,^{130, 136} stepping down ICS treatment.¹³⁷

4.3.2 School-aged children

It is possible to identify a child (aged 5–12 years) with asthma who is at increased risk of an asthma attack and to stratify the degree of risk associated with different markers. The factors associated with an increased risk in school-aged children are shown in Table 10.¹⁵²

2++

Table 10: Factors associated with an increased risk of future asthma attacks in school-aged children¹⁵²

Level of increased risk	Factor
Greatly increased risk	A history of previous asthma attacks Persistent asthma symptoms
Moderately increased risk	Suboptimal drug regimen (the ratio of the number of prescriptions for controller medication to total number of prescriptions for asthma medication <0.5 ¹⁵³) Comorbid atopic/allergic disease Low-income family Vitamin D deficiency
Slightly increased risk	Younger age Exposure to environmental tobacco smoke Obesity Low parental education
No increased risk	Gender Urban residence
Unclear (evidence equivocal)	Reduced lung function Raised FeNO at routine reviews Positive skin-prick tests History of allergen exposure

There is insufficient evidence in children to say if the following factors are associated with an increased risk of future asthma attacks: serum total IgE, family history of atopy, age of onset of asthma, duration of asthma, comorbidities (gastro-oesophageal reflux disease, diabetes), special needs, parental health, parental marital status.

4.3.3 Preschool children

Evidence to support risk prediction in preschool children (under 5 years of age) is limited. One large UK-based study (n=424,326; 17,320 under 5) in primary care concluded that it was possible to identify a preschool child with asthma at increased risk of an asthma attack.¹⁴³ Young children treated with regular preventer medication had an incremental increase in risk of an asthma attack reflecting the severity of their asthma. Other markers of slight/moderate increased risk were comorbid atopic disease, younger age, low socioeconomic status, male gender, and being overweight.

3

A US study of 300 preschool children measured differences in response to various treatment strategies, and concluded that those with raised blood eosinophils (>200/ μ L) and sensitisation to at least one aeroallergen had the greatest increased risk of future asthma attack but also the most favourable response to daily ICS therapy overall.¹⁵⁴

3

>12 yrs 5-12 yrs <5 yrs

D	B	D	Assess risk of future asthma attacks at every asthma review by asking about history of previous attacks, objectively assessing current asthma control, and reviewing reliever use.
	B	D	In children, regard comorbid atopic conditions, younger age, obesity, and exposure to environmental tobacco smoke as markers of increased risk of future asthma attacks.
D			In adults, regard older age, female gender, reduced lung function, obesity, smoking, and depression as markers of a slightly increased risk of future asthma attacks.
✓	Clinicians should target care (including tailoring frequency of review, optimising pharmacological therapy, personalising supported self management) to reduce the patient's risk status.		
✓	Healthcare policy should target vulnerable groups, ensure equitable access to care, and promote reduction in environmental tobacco smoke.		

4.3.4 People with severe asthma

In children and adults with severe asthma (defined as more than two asthma attacks a year or persistent symptoms with SABA use more than twice a week despite specialist-level therapy; *see section 7.5*), evidence from observational studies shows that a history of asthma attacks, current level of symptom control and lung function provide valuable knowledge to evaluate risk of future asthma attacks. These patients will usually be under the care of a specialist asthma clinic. Predictors of future attacks were:

- previous asthma attack^{128, 129, 133, 142}
- very poor symptom control in adults^{128, 133, 140, 142, 155}
- greater SABA use^{140, 142}
- lower lung function (PEF or FEV₁ in adults; PEF or FEV₁/FVC ratio in children)^{133, 142}
- raised FeNO.¹²⁹

3

>12 yrs 5-12 yrs <5 yrs

D	D		In individuals with severe asthma, assess risk of future asthma attacks at each visit by asking structured questions about asthma control, reviewing history of previous attacks and measuring lung function.
---	---	--	---

4.4 Physiological measures

4.4.1 Spirometry and peak expiratory flow

Both spirometry and PEF are widely available (*see section 3.2*). Spirometry is reproducible, demonstrates airflow obstruction and can be used in children as young as five. The lower limit of normal should be used to define obstruction rather than a fixed FEV₁/FVC ratio.^{156, 157}

In adults, there is evidence that reduced lung function (PEF or FEV₁) is associated with an increased risk of acute attacks.^{127, 128, 130, 133, 135, 142} The evidence in children is less clear,¹⁵² although reduced lung function (PEF or FEV₁/FVC ratio) has been identified as a risk factor in children with severe asthma (*see section 4.3*).^{133, 142}

Some adult patients have an accelerated decline in lung function in terms of FEV₁. Risk factors and treatment strategies for these patients are poorly defined and further research in this area is a priority.

4.4.2 Fractional exhaled nitric oxide

The evidence for use of biomarkers largely relates to fractional exhaled nitric oxide (FeNO) testing. FeNO testing is non-invasive and potentially suitable for adults and schoolchildren and is less reliant on operator skill than, for example, spirometry. Results can, however, be affected by other factors (*see section 3.2.4*) and a normal result in a person with non-eosinophilic asthma may be falsely reassuring; results must be interpreted correctly within the clinical context of each individual.

There is some evidence from systematic reviews that the use of FeNO testing to guide the management of asthma in adults and children may reduce exacerbation rates or their severity, but not have an impact on symptom control.¹⁵⁸⁻¹⁶⁰ Different FeNO cut-offs and heterogeneous outcome measures and definitions amongst the included studies are reflected in inconsistent outcomes. For example, in one review, adults in the FeNO group had fewer asthma attacks than those in the control group but there was no effect on steroid courses or hospitalisations.¹⁵⁸ All the reviews concluded that, although there may be some benefit in those experiencing frequent exacerbations, there is insufficient evidence to recommend the routine use of FeNO testing for the monitoring of asthma in adults or children.

1++
1+

B Except in specialist asthma clinics, the routine use of FeNO testing to monitor asthma in adults or children is not recommended.

4.4.3 Eosinophils

A systematic review of six RCTs (five in adults) including 374 adults and 55 children with asthma, showed that the use of sputum eosinophil analysis to guide treatment can reduce asthma exacerbation rates in adults. None of the studies, however, showed a difference in ICS use between the groups. The included studies were heterogeneous in design with different outcomes, eosinophil percentages and definitions of, for example, exacerbation, making comparison difficult. Five of the six studies did not group patients by asthma severity. There was insufficient evidence in children on which to base conclusions.¹⁶¹

1++

Further research is needed to determine which patients are most likely to benefit from this approach. However, the limited availability and technical demands of undertaking sputum eosinophil analysis mean it is unlikely to be a useful approach in routine clinical practice (*see section 3.2.4*).

Blood eosinophil analysis may be a useful predictor of future risk of asthma attacks in adults (*see Table 9*) but no studies were found that evaluated the use of blood eosinophil analysis for the routine monitoring of asthma.

B The routine use of sputum eosinophilia to monitor asthma in adults or children is not recommended.

4.5 Other approaches

A number of other approaches have been explored, including lung sound analysis, an index of bronchial asthma control in adults,¹⁶² and exhaled breath temperature in children.¹⁶³ There is currently insufficient evidence on these approaches on which to base a recommendation.

3

5 Supported self management

Self management has been defined as the tasks that individuals must undertake to live with chronic conditions including, “having the confidence to deal with medical management, role management and emotional management of their conditions”.¹⁶⁴ In the context of asthma, self management has focused on the medical aspects of living with a variable condition and emphasised the importance of recognising and acting on symptoms and signs of deterioration. Personalised asthma action plans (PAAPs), however, need to be seen in the context of the broader challenges of living with asthma.¹⁶⁵

5.1 Effectiveness of supported self management

There is a substantial body of evidence to show that self-management education incorporating written PAAPs improves health outcomes for people with asthma. Twenty-two systematic reviews of 261 randomised controlled trials (RCTs) encompass evidence from a broad range of demographic, clinical and healthcare contexts.¹⁶⁶⁻¹⁸⁷ In addition, 35 RCTs provide further evidence about self management in preschool children,¹⁸⁸⁻¹⁹⁶ ethnic minorities,¹⁹⁷⁻²⁰⁸ and primary care-based populations.^{206, 209-217}

1++
1+
1-
2++
2+
2-

Self-management education delivered to adults or children with asthma (and/or their parents/carers):

- reduces emergency use of healthcare resources, including emergency department (ED) visits, hospital admissions and unscheduled consultations^{166, 168, 172-175, 177, 187}
- improves markers of asthma control, including reduced symptoms and days off work, and improves quality of life.^{166, 168, 169, 175, 177-179}

1++
1+
2++
2+

Patients with all severities of asthma were included in these systematic reviews, although some focused specifically on people who had attended EDs,¹⁸⁷ or with severe or difficult asthma.¹⁷² Most self-management education was delivered in healthcare settings, but some specifically evaluated school,¹⁸¹ home,¹⁸³ or community-based interventions.¹⁸⁴ Typically, education was delivered by healthcare professionals either in individual consultations or group settings, but some systematic reviews included technology-based interventions,^{170, 171} or were part of community-health interventions for deprived and/or ethnic minority groups.^{185, 186}

5.2 Components of a self-management programme

Successful programmes varied considerably, but core components included structured education, reinforced with written PAAPs, although the duration, intensity and format for delivery varied.

5.2.1 Patient education

Education is a core component of effective self-management programmes in adults^{166, 172, 187} and children.¹⁷³⁻¹⁷⁷ There is evidence that educational interventions that were supported by a written PAAP and regular professional review were more effective than less intensive regimes.^{166, 169, 174, 176, 177}

1++
1+
2++
2+

Information technology (IT)-based education has been shown to have potential,

but as yet there is no consistent evidence on which to base recommendations on format, target audiences or the context in which it should be delivered.¹⁷⁰

5.2.2 Personalised asthma action plans

Written PAAPs (for example, those for adults and children from Asthma UK, available at www.asthma.org.uk/advice/manage-your-asthma/action-plans) are crucial components of effective self-management education.^{91, 166, 168, 178-180, 187} One systematic review identified the features of PAAPs associated with beneficial outcomes (see Table 11).¹⁶⁸

These include:

- specific advice about recognising loss of asthma control, assessed by symptoms or peak flows or both.^{91, 168, 169} In children, symptom-based written plans are effective in reducing emergency consultations for asthma, although (in older children) plans based on peak flow may be as effective for other outcomes.^{178, 179} 1++
1+
- actions, summarised as two or three action points, to take if asthma deteriorates, including seeking emergency help, starting oral steroids (which may include provision of an emergency course of steroid tablets), restarting or temporarily increasing fourfold (as opposed to just doubling) ICS, as appropriate to clinical severity (see Table 11 for further advice).¹⁶⁸

A All people with asthma (and/or their parents or carers) should be offered self-management education, which should include a written personalised asthma action plan and be supported by regular professional review.

A In adults, written personalised asthma action plans may be based on symptoms and/or peak flows: symptom-based plans are generally preferable for children.

Every asthma consultation is an opportunity to review, reinforce and extend both the patient's knowledge and skills. This is true whether the patient is seen in primary care, the ED or the outpatient clinic. It is important to recognise that education is a process and not a single event.

- ✓ • A hospital admission represents a window of opportunity to review self-management skills. No patient should leave hospital without a written personalised asthma action plan.
- An acute consultation offers the opportunity to determine what action the patient has already taken to deal with the asthma attack. Their self-management strategy may be reinforced or refined and the need for consolidation at a routine follow up considered.
- A consultation for an upper respiratory tract infection or other known trigger is an opportunity to rehearse with the patient their self management in the event of their asthma deteriorating.
- Education should include personalised discussion of issues such as trigger avoidance and occupational exposure to support people and their families living with asthma.
- Every opportunity should be taken to remind patients and carers of the importance of achieving a smoke-free environment.
- Brief simple education linked to patient goals is most likely to be acceptable to patients.

The role of telehealthcare interventions in supporting self management is covered in section 14.4.

Table 11 Summary of the key components of a written personalised asthma action plan (adapted from Gibson et al)¹⁶⁸

Component of an action plan	Result	Practical considerations
<p><i>Format of action points:</i></p> <p>Symptom v peak-flow triggered</p> <p>Standard written instructions</p> <p>Traffic light configuration</p>	<p>Similar effect</p> <p>Consistently beneficial</p> <p>Not clearly better than standard instructions</p>	<p>Asthma UK personalised asthma action plans include both symptoms and peak-flow levels at which action should be taken (<i>see www.asthma.org.uk/advice/manage-your-asthma/action-plans</i>).</p>
<p><i>Number of action points:</i></p> <p>2–3 action points</p> <p>4 action points</p>	<p>Consistently beneficial</p> <p>Not clearly better than 2–3 points</p>	<p>Suggested action points in PAAPs are:</p> <p>PEF <80% best: quadruple inhaled corticosteroids (<i>see section 5.2.3</i>)</p> <p>PEF <60% best: commence oral steroids and seek same-day medical advice</p> <p>PEF <50% best: seek urgent medical advice</p>
<p><i>Peak expiratory flow (PEF) levels:</i></p> <p>Based on percentage personal best PEF</p> <p>Based on percentage predicted PEF</p>	<p>Consistently beneficial</p> <p>Not consistently better than usual care</p>	<p>Personal best should be assessed once treatment has been optimised and peak flows are stable. Best peak flow should be updated every few years in adults, and, if a peak flow is being used, more frequently in growing children.</p>
<p><i>Treatment instructions:</i></p> <p>Individualised using inhaled and oral steroids</p>	<p>Consistently beneficial</p>	<p>Adults may be advised to quadruple their inhaled corticosteroid dose (<i>see section 5.2.3</i>).</p> <p>For those on maintenance and reliever therapy (MART) regimes (<i>see section 7.3.5</i>).</p> <p>Patients who have stopped medication should be reminded to restart their inhaled corticosteroids.</p> <p>Patients may safely hold an emergency supply of prednisolone tablets for use if their symptoms continue to deteriorate and/or if their peak flow falls to 60% of their best.</p>

5.2.3 Increasing inhaled corticosteroids to abort an asthma attack

Regular, daily use of preventer medication is the best means of avoiding asthma attacks and the need for increased (rescue) doses of ICS. Trials of PAAPs have typically included advice to increase the dose of ICS at the start of an asthma attack in an attempt to abort the attack.

Overall, a Cochrane review including eight RCTs, five in adults (n=1,247) and three in children (n=422, age range 3–18), in people with mild to moderate asthma, showed no significant difference in the odds of requiring rescue oral corticosteroids or of having adverse events between those taking an increased dose of ICS for an exacerbation and those continuing with a stable dose, although the authors did not rule out the possibility of benefits in some patient groups.²¹⁸

However, the doses of ICS used during exacerbations, maintenance doses of ICS and concomitant medications varied between studies. Six of the eight studies recommended doubling ICS during attacks, four of which reported 'high' adherence (>80%) with maintenance regimes. One of the included studies reported that quadrupling ICS did not reduce the number of steroid courses, although it halved the risk of needing oral steroids in the subgroup (23%) who needed to start the increased dose. Another of the included studies reported that quintupling ICS significantly reduced the number of patients with attacks requiring steroids compared with steady low-dose ICS (18% v 38%).

1++

A further pragmatic RCT in adults aged 16 and over (n=1,871) with broad inclusion criteria evaluated quadrupling the patient's usual ICS dose for up to 14 days at the start of an asthma attack as part of a personalised self-management plan. They reported 19% fewer severe asthma exacerbations in those receiving the increased dose (adjusted hazard ratio 0.81, 95% confidence interval (CI) 0.71 to 0.92), although the absolute benefit was small (proportion requiring oral steroids was 45% v 52% in the intervention and control groups, respectively). The number of hospitalisations was significantly lower in the intervention group although admissions were infrequent (0.3% v 1.9%). Local candidiasis and oral dysphonia occurred more frequently in the intervention compared with the control group (7% v 2%).²¹⁹

1+

In contrast, an efficacy trial in children aged 5–11 years (n=254) with a reported adherence to daily ICS of 98%, evaluated quintupling a standardised dose of fluticasone for seven days at the early signs of loss of asthma control. There was no significant difference in severe attacks between those receiving the increased dose and those who continued on the lower dose. There was a non-significant trend to a lower growth rate in the intervention group.²²⁰

1+

It may be that highly adherent patients have little extra to gain by increasing ICS and that, in many patients, the beneficial effect arises because adherence to regular preventer therapy was poor. There is no evidence in 'real world' situations, where non-adherence is widespread, to inform recommendations in those under 16 years of age, although effective PAAPs in children have included this step.

The use of a single combination inhaler for maintenance and reliever therapy (MART) is an alternative approach to the introduction of a fixed-dose twice-daily combination inhaler which ensures an increased dosage of ICS as formoterol reliever treatment is increased (*see section 7.3.5* for a description of the MART regime). This approach might suit some individuals.

B In personalised asthma action plans for adults, consider advising quadrupling ICS at the onset of an asthma attack and for up to 14 days in order to reduce the risk of needing oral steroids.

- ✓ • Consider adherence before recommending increasing ICS as patients who are highly adherent (>90%) may have a ceiling effect and gain no additional benefit from increasing ICS at the onset of an attack.
- Weigh the benefit/risk ratio of recommending quadrupling ICS at the start of an asthma attack in people already on high dose ICS especially if they are experiencing frequent attacks and/or are still requiring oral steroids.

- ✓ For people on fixed-dose combination inhalers, increasing the dose of ICS may best be achieved by adding a single ICS inhaler.

5.3 Self management in specific patient groups

A range of different patient populations are included in the trials of self management. It cannot be assumed that a successful intervention in one setting will be feasible or appropriate in another.

5.3.1 Primary care

Studies of self-management interventions based in primary care have shown that they can:

- reduce emergency use of healthcare resources, including ED attendances, hospital admissions and unscheduled consultations^{206, 212} 1++
1+
- improve markers of asthma control.^{206, 209, 210, 212-215, 221}

Implementation of self-management interventions is challenging. The improved asthma control demonstrated in trials of interventions delivered by members of the research team^{206, 212} or in a centrally administered initiative^{213, 214} are reflected in some,^{209, 210, 215, 221} but not all,^{216, 217} trials in which members of the practice team are trained to deliver self-management education in routine clinical care. 1++
1+

One study showed no difference in outcomes when self-management education was delivered by lay people compared with practice asthma nurses.²¹¹ Studies based in the USA suggest that in deprived and/or ethnic communities the involvement of community health workers reduces ED attendance.¹⁸⁴ 2

A Self-management education, supported by a written personalised asthma action plan, should be offered to all patients on general practice 'active asthma' registers.

A Primary care practices should ensure that they have trained professionals and an environment conducive to providing supported self management.

- ✓ Implementation of self-management interventions is challenging in the non-specialist environment of primary care and needs to consider not only specific training in self-management skills, but also the logistics of when and how self-management education is incorporated into routine care. Strategies that have been used in effective interventions include:
- the use of proactive triggers to ensure routine reviews
 - structured protocols for asthma reviews
 - support from community pharmacists
 - routine mailing of educational resources
 - telephone calls to provide ongoing support and advice
 - IT-based education and monitoring
 - involvement of community workers to support clinical teams in deprived and/or ethnic minority communities.

5.3.2 Secondary care

There is good evidence that self-management education targeted at people who have a history of ED attendances¹⁸⁷ or hospital admissions^{207, 222} can reduce subsequent use of healthcare resources. Self-management education delivered prior to discharge can reduce readmissions and should be a core component of discharge planning (*see section 9.9.7*).²²³⁻²²⁵ **1++**

One wide-reaching review of the evidence for self management in people with severe or difficult asthma concluded that provision of psychoeducational interventions (especially those incorporating formal self management) may reduce hospital admissions and, in children, improve symptoms.¹⁷² **2+**

A Prior to discharge, inpatients should receive written personalised asthma action plans, given by healthcare professionals with expertise in providing asthma education.

5.3.3 Schoolchildren

School-based asthma education has been shown to:

- improve process outcomes (knowledge, self efficacy, self-management behaviours)¹⁸¹
- improve markers of asthma control (number of days and nights with asthma symptoms, school absences, asthma-related quality of life).^{181, 182}

The considerable heterogeneity in school-based interventions means it is not possible to specify which components should be included. Interventions incorporated combinations of classroom teaching for all pupils, peer support groups, individual education sessions with school nurses, interactive computer programmes, and involvement of parents.¹⁸¹ **1+ 1-**

A School health services should consider providing in-school asthma self-management education programmes provided by appropriately trained personnel.

5.3.4 Preschool children

There is a paucity of evidence about effective self-management strategies delivered to parents of preschool children. Trials recruiting only preschool children (aged five years or under) showed no impact on emergency use of healthcare resources, including ED visits, hospital admissions and unscheduled consultations,^{190, 195} and no¹⁹⁰ or limited¹⁹⁵ reduction in symptoms, despite increased ownership of PAAPs.¹⁹⁵

1+
1-

Other trials including preschool children and children up to the age of eight years showed only small and often transient effects of no apparent clinical significance.^{188, 189, 192-194}

5.3.5 Ethnic minority groups

Interventions specifically designed for ethnic minority groups, predominantly deprived African-American, Hispanic or Puerto Rican populations from inner cities in the USA,^{186, 197-205} can:

1++
1+
1-
2+

- reduce emergency use of healthcare resources, including ED attendances, hospital admissions and unscheduled consultations^{185, 186, 200, 201, 203}
- improve markers of asthma control^{185, 186, 197, 198, 201, 203}
- improve process outcomes (knowledge).^{185, 186, 202, 204}

In two UK-based RCTs, however, interventions which provided appropriate language materials and were delivered by bilingual professionals were reported as showing no or less benefit on healthcare outcomes in the South Asian population compared with the benefits seen in the white European population.^{206, 207}

1++
1+

There is insufficient evidence to identify all the aspects of cultural tailoring which may potentially contribute to effectiveness of self-management interventions, but addressing language barriers (for example with appropriate language materials and bilingual support) is not sufficient to enable an intervention to deliver equivalent outcomes in an ethnic minority group compared with a white European group.^{206, 207}

1++
1+

The strategies employed in ethnic minority groups are varied and include community-based neighbourhood projects,^{199, 203, 204} family-based education,²⁰⁰ nurse-led home visits,¹⁹⁸ IT-based programmes,^{197, 201, 202} and school-based educational interventions.^{201, 205} No single strategy stands out as being always effective, or always ineffective. Lack of engagement with programmes and high drop-out rates are major barriers to effectiveness of self-management interventions.^{198, 199, 203, 204} Reconfiguration of the supporting healthcare system appears to increase the impact.¹⁸⁶

1++
1+
1-

B

Culturally appropriate supported self-management education should be provided for people with asthma from ethnic minority groups. Addressing language barriers is insufficient.

- ✓ Consideration should be given to:
 - translation of materials into community languages with ethnically appropriate pictures
 - asthma educators fluent in community languages
 - identifying culturally appropriate support agencies within the local community
 - inclusion of culturally specific beliefs and practices
 - reference to culturally appropriate role models
 - involvement of a local community health worker to support clinical teams.

5.4 Adherence and concordance

The term adherence (or compliance) embodies a traditional model of prescriptive care which refers to the objectively measured usage of prescribed medication, or frequency of monitoring. The term concordance signifies a negotiated agreement between the professional and the patient. Non-concordance describes an inability of both parties to come to an understanding, not merely a failure of the patient to follow the healthcare professional's instructions.²²⁶ Sharing decision making and achieving concordance improves, though does not guarantee, adherence.²²⁷

5.4.1 Adherence to monitoring and treatment

Adherence to regular monitoring with peak-flow meters, even in clinical drug trials is poor, with recorded daily use as low as 6%.^{228, 229} The lack of evidence supporting long-term peak-flow monitoring,^{37, 230-232} however, does not negate the use of home peak-flow monitoring at critical times, for example at diagnosis and initial assessment, when assessing response to changes in treatment, and as part of a PAAP during asthma attacks.²³² Comparison should be with the patient's best (not predicted) peak flow.¹⁶⁸

It is estimated that between a third and a half of all medicines prescribed for long-term conditions are not taken as recommended.²³³ Evidence in people with asthma confirms that there is widespread non-adherence to regular preventer medication²³⁴⁻²³⁹ that increases over time.^{234, 236} Poor adherence should always be considered when there is a failure to control asthma symptoms.

1+
3

Non-adherence to medication use may be intentional and/or unintentional and may be understood as the result of the interaction of perceptual factors (for example beliefs about illness and treatment) and practical factors (forgetfulness, capacity, resources and opportunity).²³³

4

A widely recognised model for understanding patients' decisions about medication use is the Necessity-Concerns framework which describes the balance between the potential benefits and 'necessity' of taking prescribed treatment and the perceived disadvantages or 'concerns' about taking medication.²⁴⁰ The relative weight of these opposing arguments influences the decision to take medication (or not).^{241, 242}

5.4.2 Assessing medication adherence

In most clinical contexts, the key strategies for assessing adherence are self reporting and the prescribing record, although biochemical assays may have a role in asthma clinics for patients with severe asthma (*see section 10.2.1*). In a research context electronic dose monitoring is the gold standard; counting doses used is another approach that is frequently used.

Patient self reporting is simple, inexpensive and feasible in most clinical settings. Self reporting typically overestimates adherence by a third compared to electronic monitoring^{233, 236, 239} or dose counting.^{234, 235} This applies both in trial populations^{234, 236, 239} and clinical settings.²³⁵ Underuse is over-reported,^{234-236, 239} and overuse is under-reported,²³⁹ reflecting socially acceptable answers.²³³ Patients/caregivers who report missing doses or not taking medication are likely to be non-adherent,^{233, 235, 236} though their estimate of dosages taken may still be inaccurate.²³³ Being non-judgemental, and asking specific questions about use of a treatment over a short time period (for example, in the last week/month) can help elicit an accurate response.²³³ Questionnaires have been validated for use in research,²⁴⁰ but have not been validated as a tool in clinical use.²³³

1+
3
4

Computerised prescribing records

Computerised prescribing records, normally readily available in primary care consultations and/or pharmacy dispensing records, provide a useful indication of adherence to prescribed asthma regimens. At an individual level, prescribing data do not correlate with self-reported adherence and may be a useful strategy for opening a discussion about suspected poor adherence.²⁴¹ At a population level, formulae (such as 'proportion of days covered' by the prescription recorded over a defined period) have been devised to assess adherence from routine prescribing/dispensing databases.^{237, 238, 241, 243}

Biomarker testing

Biomarker testing with FeNO or biochemical urinary assays (for example for a metabolite of fluticasone propionate) may have a role in establishing (non-) adherence in people with severe/difficult asthma.^{244, 245} Suppression of FeNO after five days of directly observed inhaled steroid dosage has been shown to be an objective test to distinguish adherent from non-adherent patients with difficult asthma (*see section 10.2.1*).²⁴⁴

2+

Electronic monitoring

Electronic monitoring is the gold standard for assessing adherence in the research context, although not normally available in routine clinical practice.^{236, 239} Dose counting is also used as a comparator, although unlikely to be feasible in a clinical context.^{234, 235}

1+
3

D To assess adherence, ask specific questions about medication use and assess prescribing and any other data available. Explore attitudes to medication as well as practical barriers to adherence in a non-judgemental way.

- ✓ Questions about adherence should be open ended, acknowledge that poor adherence is the norm, and avoid use of potentially judgmental terminology. The questions are designed to stimulate an open discussion.
- Explore perceived benefits (“How do you think that the inhaler is helping you control your asthma?” “Are there times when you find that you don’t need your inhaler?”)
 - Ask about adverse reactions (“How much bother do you have from side effects?”)
 - Acknowledge general concerns about regular medication (“Some people worry about taking regular medication... what do you think?”)
 - Acknowledge practical difficulties with regular medication (“People sometimes find it difficult to remember to take regular treatment...”)
 - Ask about adherence over a specific time period (“How often did you use your preventer inhaler last week?”)

5.4.3 Interventions to improve medication adherence

Five systematic reviews were identified that evaluated interventions to improve adherence in people with asthma.²⁴⁶⁻²⁵⁰ Multifaceted interventions have demonstrated modest improvements in adherence,²⁴⁹⁻²⁵³ the clinical significance of these improvements is, however, unclear with limited, or sometimes no, observed effect on clinical outcomes.^{248, 250, 251, 253-255}

1++
1+
2+

The observed effect was greater if the intervention:

- included behavioural components (studies were in mixed populations with chronic conditions, including asthma)²⁵⁶⁻²⁵⁸
- included practical facilitators (such as simplified dosage regimes),²⁵³ strategies to aid integration into daily routines,²⁵⁹ and automated reminders^{249, 251, 252, 254, 255, 260}
- is monitored, delivered and sustained as part of a comprehensive programme of accessible proactive asthma care.²⁴⁹

1++
1+
2+
3

IT-based strategies to support adherence such as text message reminders in children and young people,^{251, 252, 260} feedback on adherence,²⁵⁵ and refill reminders,²⁵⁴ showed some promise, especially if they were interactive,^{247, 254, 255} although it is unclear how long improvements in adherence are sustained.

1++
1+

The value of electronic interventions to support adherence may be diminished in patient groups who are either unable to, or lack confidence in, accessing electronic formats, for instance some older adults and those with a learning disability or cognitive impairment. The financial implications of accessing applications on mobile devices, for example for low-income families, also need to be considered. Some studies included financial incentives.

School-based interventions depend on the child actually being at school and having someone to deliver the intervention at the school in a consistent manner. This has practical implications for implementation, for example the need for training.

Overall, interventions to improve medication adherence do not clearly improve clinical outcomes, and should therefore be considered as components of, as opposed to replacement for, ongoing supportive care (*see section 14.4.1*).²⁴⁹

D Adherence to long-term asthma treatment should be routinely and regularly addressed by all healthcare professionals within the context of a comprehensive programme of accessible proactive asthma care.

- ✓ Initiatives to promote adherence to regular treatment should consider:
- information requirements, for example individual and/or group sessions, written/electronic materials, ongoing access to information
 - practical facilitators, for example simple dosage regimes, dose counters, reminders
 - behavioural support, for example regular monitoring including assessment of medication use with feedback, counselling, psychological therapies
 - context – accessible proactive asthma care, for example Chronic Care Model²⁴⁹
 - consultation skills required to achieve shared decision making: adherence is more likely when the patient and the healthcare professional agree that the action is appropriate.

5.5 Implementation in practice

Despite the robust evidence base for self-management education, implementation in routine practice remains poor with only a third of people with asthma having a PAAP.^{261, 262} Implementation in routine clinical practice depends as much on the context in which it is delivered as the content of the intervention. Despite the diversity of healthcare systems, the evidence reviewed identified consistent messages that are suitable for adoption and adaptation in different healthcare settings.

A systematic review (including 14 RCTs, 2,438 patients, 107 doctors and 43 primary care teams) investigated the promotion of PAAP ownership and usage.²⁶³ In addition, 19 implementation studies from the USA,^{213, 216, 264-270} UK,^{8, 217, 271, 272} Scandinavia,²⁷³⁻²⁷⁵ Italy,²⁷⁶ and Brazil^{277, 278} were identified.

5.5.1 Types of intervention

The interventions in the implementation studies adopted four main strategies:

- primarily professional training^{216, 217}
- primarily organisational change^{8, 271, 273}
- primarily patient education^{213, 265-268, 276}
- a whole systems approach with components operating explicitly at patient, professional and organisational levels.^{264, 269, 270, 272, 274, 275, 277, 278}

Study designs varied and included five cluster randomised trials,^{216, 217, 266, 267, 271} a preference trial with randomised groups,²¹³ a controlled implementation study,⁸ eight studies based on longitudinal, often large, databases,^{264, 265, 268-270, 274, 275, 278} one control cohort,²⁷⁷ two uncontrolled before-and-after studies,^{272, 276} and one cross-sectional study.²⁷³

5.5.2 Implementation of interventions

Complex whole systems interventions in which motivated informed patients and trained professionals operate within an organisation with a culture of supported asthma self management were associated with:

- improved knowledge²⁷⁰ and action plan ownership^{263, 268, 272}
- reduced unscheduled care,^{268-270, 274, 277, 278} and improved markers of control.^{269, 270, 274, 275}

1++
1-

Implementing single components of the whole systems approach is insufficient to bring about consistent benefits. Improving professionals' knowledge is a core component of effective self-management programmes, but on its own does not improve clinical outcomes.^{216, 217} Organisational change to support self management improves process outcomes such as the proportion of patients with PAAPs or achieving a review,^{8, 271, 273} but improved asthma control in only one of the studies.²⁷³ Targeting the patient with educational material,²⁶⁸ support from pharmacists,²⁶⁵ school,^{266, 276} or telephone calls^{213, 266, 267} improved medication use,^{213, 267} knowledge²⁶⁶ and ownership of PAAPs,²⁶⁵ and had variable effects on clinical outcomes.

2++
2+

B Commissioners and providers of services for people with asthma should consider how they can develop an organisation which prioritises and actively supports self management. This should include strategies to proactively engage and empower patients and train and motivate professionals as well as providing an environment that promotes self management and monitors implementation.

6 Non-pharmacological management

There is a common perception amongst patients and carers that there are numerous environmental, dietary and other triggers of asthma and that avoiding these triggers will improve asthma and reduce the requirement for pharmacotherapy. Failure to address a patient, parent or carer's concern about environmental triggers may compromise concordance with recommended pharmacotherapy. Evidence that non-pharmacological management is effective can be difficult to obtain and more well-controlled intervention studies are required.

- This section distinguishes prevention activities as follows:
- primary prevention – interventions introduced before the onset of disease and designed to reduce its incidence.
- secondary prevention – interventions introduced after the onset of disease to reduce its impact.

6.1 Primary prevention

The evidence for primary interventional strategies is based predominantly on observational studies, although some interventions have been tested using experimental methods. Many are multifaceted and it can be difficult to disentangle the effects of one exposure or intervention from another.

6.1.1 Monofaceted and multifaceted allergen avoidance

Early life exposure to allergens (including aeroallergens and ingested food allergens) may lead to allergic sensitisation and so potentially increase the risk of subsequent asthma, particularly in children at high risk (that is, children with a family history of asthma or atopy, particularly a parental history). It is unclear whether the risk of developing asthma in children is reduced by interventions to reduce exposure to single allergens (monofaceted), or whether multifaceted interventions targeting the reduction of more than one type of allergen exposure simultaneously will lead to a better outcome or be more effective.

A Cochrane review of trials comparing single (six studies) or multiple (three studies) interventions with a no-intervention control, reported that in children who are at risk of developing childhood asthma there may be a role for multifaceted interventions which involve both dietary allergen reduction and environmental change to reduce exposure to inhaled allergens. Such interventions reduced the odds of a doctor diagnosing asthma later in childhood by half in those over five years of age, odds ratio (OR) 0.52, 95% CI 0.32 to 0.85.²⁷⁹ However, the effect of these multifaceted interventions on wheeze reported by parents was inconsistent and there was no beneficial effect on night-time coughing or breathlessness. These interventions can be costly, demanding and inconvenient to families, and the cost effectiveness is not established. Healthcare professionals can discuss and support this intervention in families who are motivated to follow the demanding programme. **1++**

In children at risk of developing asthma, there is no evidence that reducing in utero or early life exposure to single allergens (either to aeroallergens such as house dust mites or pets, or food allergens) is effective in reducing asthma and single (monofaceted) interventions were not significantly more effective than controls in the reduction of any outcomes.²⁸⁰ **1++**

A Measures to reduce in utero or early life exposure to single aeroallergens, such as house dust mites or pets, or single food allergens, are not recommended for the primary prevention of asthma.

A For children at risk of developing asthma, complex, multifaceted interventions targeting multiple allergens may be considered in families able to meet the costs, demands and inconvenience of such a demanding programme.

6.1.2 Aeroallergen avoidance

House dust mites

Exposure to high levels of house dust mite allergen in early life is associated with an increased likelihood of sensitisation to house dust mite by three to seven years of age.²⁸¹ Sensitisation to house dust mite is an important risk factor for the development of asthma,^{282, 283} and a few studies have suggested that exposure to high levels of house dust mites early in life increases the risks of subsequent asthma.^{284, 285} A UK study showed that low levels of house dust mite and cat allergen exposures in early life increased the risk of IgE sensitisation and asthma at five years, with some attenuation at high levels of exposure, but there were significant associations with family history and birth order.²⁸⁶

Outcomes from intervention studies attempting to reduce exposure to house dust mites are inconsistent. A multifaceted Canadian intervention study showed a reduced prevalence of doctor-diagnosed asthma but no impact on other allergic diseases, positive skin-prick tests or bronchial hyper-responsiveness;²⁸⁷ others have shown no effect on either allergic sensitisation or symptoms of allergic diseases.²⁸⁸ In one UK study, early results from environmental manipulation started in early pregnancy and focused mainly on house dust mite avoidance, showed reductions in some respiratory symptoms in the first year of life.²⁸⁹ Subsequent results showed a paradoxical effect with increased allergy but better lung function in the intervention group.²⁹⁰

1+

The considerable variation in the methodology used in these studies precludes the pooling of data or meta-analyses.

A Healthcare professionals should not recommend house dust mite aeroallergen avoidance for the primary prevention of asthma.

Pets in the home

A large number of birth cohort studies, longitudinal cohort studies and cross-sectional studies have addressed whether exposure to pets in the home in early life increases or reduces the subsequent risk of asthma and allergy, with contradictory results. Four systematic reviews, synthesising evidence from overlapping data sources, have provided conflicting results. One review concluded that exposure to cats in early life has a slight preventative effect on subsequent asthma, while exposure to dogs increases risk.²⁹¹ Another concluded, in contrast, that perinatal dog exposure protects against asthma, with no effect from cats.²⁹² Methodological factors, however, such as avoidance behaviour in at-risk families and other potential confounders, may have affected the analyses. Two further reviews concluded that exposure to cats and/or dogs in early childhood did not impact on asthma or wheeze in school-aged children.^{293,}

²⁹⁴ The most methodologically sound review pooled individual participant data

2++
2+
2-

from 11 European birth cohort studies and so was able to harmonise exposure, outcome and age-group definitions and use individual data rather than pooled risk estimates in heterogeneous groups, to minimise potential confounding.²⁹⁴ This review concluded that exposure to cats and/or dogs in infancy does not impact on a diagnosis of asthma or on wheezing symptoms in later life, although may influence allergic sensitisation, and that parents should not make choices on pet ownership based on the desire to prevent or reduce asthma symptoms. Several of the studies and reviews reported reduced allergic sensitisation in those with early exposure to pets, but the clinical significance of this is uncertain.

2++
2+
2-

B Healthcare professionals should not offer advice on pet ownership as a strategy for preventing childhood asthma.

6.1.3 Food allergen avoidance

Sensitisation to foods, particularly eggs, frequently precedes the development of aeroallergy and subsequent asthma.²⁹⁵ Food allergen avoidance in pregnancy and postnatally has not been shown to prevent the later development of asthma.²⁹⁶ Allergen avoidance during pregnancy may adversely affect maternal, and perhaps fetal, nutrition.²⁹⁷ High-dose food allergen exposure during pregnancy may reduce subsequent sensitisation rates by inducing tolerance.²⁹⁸

1+

B In the absence of any evidence of benefit and given the potential for adverse effects, maternal food allergen avoidance during pregnancy and lactation is not recommended as a strategy for preventing childhood asthma.

6.1.4 Breastfeeding

A systematic review of observational studies on the allergy preventive effects of breastfeeding indicates that it is effective for all infants irrespective of family history of allergy. The preventive effect is more pronounced in infants at high risk provided they are breastfed for at least four months.²⁹⁹ However, not all studies have demonstrated benefit and a large birth cohort study reported no protective effect against atopy and asthma.³⁰⁰

2+

Observational studies have the potential to be confounded by, for example higher rates of breastfeeding in atopic families, and taking this into account, the weight of evidence is in favour of breastfeeding as a preventive strategy.

C Breastfeeding should be encouraged for its many benefits, including a potential protective effect in relation to early asthma.

6.1.5 Modified infant milk formulae

Trials of modified milk formulae have not included sufficiently long follow up to establish whether there is any impact on asthma. A Cochrane review identified inconsistencies in findings and methodological concerns amongst studies, which mean that hydrolysed formulae cannot currently be recommended as part of an asthma prevention strategy.³⁰¹ A review of the use of soy formulae found no significant effect on asthma or any other allergic disease.³⁰²

1+

In the absence of any evidence of benefit from the use of modified infant milk formulae it is not possible to recommend it as a strategy for preventing childhood asthma.

6.1.6 Weaning

There are conflicting data on the association between early introduction of allergenic foods into the infant diet and the subsequent development of allergy and atopic eczema. No evidence was identified in relation to asthma.³⁰³ In one study late introduction of egg was associated with a non-significant increase in wheezing in preschool children.³⁰⁴

In the absence of evidence on outcomes in relation to asthma no recommendations on modified weaning can be made.

6.1.7 Nutritional supplementation

Fish Oils

Fish oils have a high level of omega-3 polyunsaturated fatty acids (n-3PUFAs). Western diets have a low intake of n-3PUFAs with a corresponding increase in intake of n-6PUFAs. This change has been associated with increasing rates of allergic disease and asthma.³⁰³ Two RCTs have investigated early-life fish oil dietary supplementation in relation to asthma outcomes in children at high risk of atopic disease (at least one parent or sibling had atopy with or without asthma). In a study, powered only to detect differences in cord blood, maternal dietary fish oil supplementation during pregnancy was associated with reduced cytokine release from allergen-stimulated cord blood mononuclear cells. However, effects on clinical outcomes at one year, in relation to atopic eczema, wheeze and cough, were marginal.³⁰⁵ In a second study, fish oil supplementation started in early infancy with or without additional house dust mite avoidance was associated with a significant reduction in wheeze at 18 months of age. By five years of age fish oil supplementation was not associated with effects on asthma or other atopic diseases.³⁰⁶

1+

In the absence of any evidence of benefit from the use of fish oil supplementation in pregnancy it is not possible to recommend it as a strategy for preventing childhood asthma.

Other nutrients

A number of observational studies have suggested an increased risk of subsequent asthma following reduced (maternal) intakes of selenium (based on umbilical cord levels),³⁰⁷ or vitamin E based on maternal pregnancy intake.³⁰⁸ No intervention studies in relation to selenium or vitamin E have yet been conducted and overall there is insufficient evidence to make any recommendations on maternal dietary supplementation as an asthma prevention strategy.³⁰³ Observational studies suggest that intervention trials are warranted.

6.1.8 Weight reduction in overweight and obese patients

There is consistent evidence that being overweight or obese increases the risk of a subsequent physician diagnosis of asthma by up to 50% in children and adults of both sexes.^{309, 310} A high birth weight is also associated with a higher risk of asthma.³⁰⁹ The quality of the evidence is, however, low as there was no adjustment for confounders. In addition, since obesity can have direct effects on respiratory symptoms and on lung mechanics, the mechanism of this relationship is unclear.

2-

Two systematic reviews looking at the association between being overweight or obese in childhood and the development of asthma concluded that high body mass index (BMI) increases the risk of incident asthma, with a dose-dependent relationship that was stronger in boys.^{311, 312} These reviews are, however, based on epidemiological studies and cannot confirm a causal link. 2+

A systematic review of the association between maternal obesity and gestational weight gain in pregnancy, and childhood asthma, concluded that maternal obesity was associated with an increased risk of diagnosed asthma and of ever-wheeze in children from these pregnancies, with each 1 kg/m² increase in maternal BMI associated with a 2–3% increase in odds of childhood asthma. High gestational weight gain was associated with higher odds of asthma or ever-wheeze in children (OR 1.16).³¹³ Prospective studies of weight-loss programmes during pregnancy for obese women and those with high gestational weight gain are needed to clarify the role of this intervention in the prevention of asthma in children resulting from these pregnancies. 2+

C Weight reduction is recommended in obese patients to promote general health and to reduce subsequent respiratory symptoms consistent with asthma.

C Obese and overweight children should be offered weight-loss programmes to reduce the likelihood of respiratory symptoms suggestive of asthma.

6.1.9 Microbial exposure

The 'hygiene hypothesis' suggested that early exposure to microbial products would switch off allergic responses thereby preventing allergic diseases such as asthma. The hypothesis is supported by some epidemiological studies comparing large populations who have or have not had such exposure.^{314, 315}

The concept is sometimes described as the 'microbial exposure hypothesis'. A double-blinded placebo-controlled trial of the probiotic *Lactobacillus rhamnosus GG* given to mothers resulted in a reduced incidence of atopic eczema in their children but had no effect on IgE antibody or allergic skin-test responses. The small sample size and short follow up in this study limit its interpretation.³¹⁶ There remains insufficient understanding of the ecology of gut flora in infancy in relation to outcomes. Bifidobacteria may be more important than lactobacilli in reducing susceptibility to allergic disease.³¹⁷

There is insufficient evidence to indicate that the use of dietary probiotics in pregnancy reduces the incidence of childhood asthma.

6.1.10 Avoidance of tobacco smoke and other air pollutants

No evidence was identified to support a link between exposure to environmental tobacco smoke (ETS) or other air pollutants and the induction of allergy.

There is an increased risk of infant wheezing associated with maternal smoking during pregnancy which adversely affects infant lung function.^{318–321} Evidence suggests that early-life ETS exposure is associated with later persistent asthma,^{322, 323} with a strong interaction with genetic polymorphisms which affect antioxidant activity.³²⁴ 2+

B Current and prospective parents should be advised of the many adverse effects which smoking has on their children including increased wheezing in infancy and increased risk of persistent asthma.

The limited data on antenatal or early-life exposure to other pollutants suggest similar effects to those for ETS, namely increased infant wheezing, enhanced by additional ETS exposure and antioxidant gene variations.³²⁵⁻³²⁷ There is one small study suggesting that vitamin C supplementation will modify the combined effects of genetic polymorphisms and pollution on lung function in children with asthma.³²⁸ Further research is required before recommendations for practice can be made.

3
4

6.1.11 Immunisation

In keeping with the microbial exposure hypothesis some studies have suggested an association between tuberculin responsiveness and subsequent reduced prevalence of allergy, implying a protective effect of Bacillus Calmette-Guérin (BCG). At present, it is not possible to determine whether poor tuberculin responsiveness represents an underlying defect which increases the risk of allergy and asthma or whether the immunisation itself has a protective effect.³²⁹

2+

Investigation of the effects of other childhood immunisation suggests that at worst there is no influence on subsequent allergic disease and may be some protective effect against the development of asthma.³³⁰

C All childhood immunisations should proceed normally as there is no evidence of an adverse effect on the incidence of asthma.

6.2 Secondary prevention

6.2.1 House dust mite avoidance

Allergic sensitisation to house dust mite-associated aeroallergens is common in people with asthma and exposure to house dust can act as a trigger in sensitised asthmatic individuals. Physical (for example mattress covers, vacuum cleaning, heating, ventilation, freezing, washing, air filtration and ionisers) and chemical (acaricides) measures to reduce house dust mite (HDM) aeroallergen levels and so reduce exposure have been advocated but there has been uncertainty as to whether the currently available physical and chemical measures, alone or in conjunction, can reduce the exposure levels sufficiently to allow a clinically relevant effect to be apparent.

A systematic review of 72 studies (64 RCTs and eight non-RCTs) included 37 studies evaluating single interventions (seven acaricides, nine air purification, one high-efficiency particulate air-filtration, 17 mattress covers, two pest-control measures, one pet removal) and 30 studies evaluating multicomponent strategies. The included studies enrolled adults, children or mixed populations. Taking a narrative approach, the review concluded that single component interventions are not effective at improving asthma control or reducing asthma attacks despite HDM levels being significantly reduced in many studies. Multicomponent interventions were found to have some clinical effects. However, the heterogeneity of interventions, how studies were combined and the small number of studies precluded definitive conclusions.³³¹ There is, therefore, continuing clinical uncertainty about which HDM avoidance measures may be clinically effective in asthma and further research is required.

1+

B Physical and chemical methods of reducing house dust mite levels in the home (including acaricides, mattress covers, vacuum cleaning, heating, ventilation, freezing, washing, air filtration and ionisers) should not be routinely recommended by healthcare professionals for the management of asthma.

6.2.2 Other allergens

Animal allergens, particularly from cats and dogs, are potent provokers of asthma symptoms. The reported effects of removal of pets from homes are paradoxical, with either no benefit for asthma^{332, 333} or a potential for continued high exposure to induce a degree of tolerance.³³⁴ In homes where there is no cat but still detectable cat allergen, there may be a benefit from introducing additional avoidance measures such as high-efficiency vacuum cleaners for patients allergic to cats, although there is insufficient evidence on which to base a recommendation.³³¹

Although fungal exposure has been strongly associated with hospitalisation and increased mortality in asthma, no controlled trials have addressed the efficacy of reducing fungal exposure in relation to control of asthma. Cockroach allergy is not a common problem in the UK and studies of attempts to avoid this allergen elsewhere have produced conflicting results.³³¹

Studies of individual aeroallergen avoidance strategies show that single interventions have limited or no benefit.³³¹ A multifaceted approach is more likely to be effective if it addresses all the indoor asthma triggers but there remains considerable uncertainty about which, if any, are the most effective strategies.³³¹ A strategy with a potential impact on mites, mould allergens and indoor pollutants is the use of a mechanical ventilation system to reduce humidity and increase indoor air exchange. A systematic review of this topic concluded that more research is required to determine whether this approach is effective.³³⁵

6.2.3 Smoking

Direct or passive exposure to cigarette smoke adversely affects quality of life, lung function, need for rescue medications for acute episodes of asthma and long-term control with ICS.³³⁶⁻³³⁹

In children with asthma, exposure to environmental tobacco smoke is associated with worsening asthma symptoms.³⁴⁰ Smoking cessation interventions aimed at families and carers have been shown to reduce childhood respiratory symptoms including those associated with asthma.³⁴¹ One study in adults with asthma suggested that smoking cessation improved asthma-specific quality of life, symptoms and drug requirements.³⁴²

1++
2+

Uptake of smoking in teenagers increases the risks of persisting asthma. One study showed a doubling of risk for the development of asthma over six years in children aged 14 who started to smoke (*see section 7.2.6 for the effect of smoking on treatment*).³⁴³

3

B People with asthma and parents/carers of children with asthma should be advised about the dangers of smoking and second-hand tobacco smoke exposure, and should be offered appropriate support to stop smoking.

6.2.4 Air pollution

Challenge studies demonstrate that various pollutants can enhance the response to allergen inhalation in patients with asthma.^{344, 345} Time-series and other observational studies suggest that air pollution may provoke acute asthma attacks or aggravate existing chronic asthma although the effects are very much less than in those with infection or allergen exposure.^{346, 347} Increased asthma symptoms in young children (mean age ≤ 9) have been linked, in observational studies, to exposure to air pollutants, including particulates, nitrogen dioxide, sulphur dioxide and ozone.³⁴⁰ Much less attention has been focused on indoor pollutants in relation to asthma and more work is required.^{348, 349}

Information on current levels of air pollution, recommended actions and health advice is available from The Daily Air Quality Index (available at www.uk-air.defra.gov.uk).

6.2.5 Electrolytes

Increasing dietary sodium has been implicated in the geographical variations in asthma mortality and high sodium intake is associated with increased bronchial hyper-responsiveness.³⁵⁰⁻³⁵² A systematic review of intervention studies reducing salt intake identified only minimal effects and concluded that dietary salt reduction could not be recommended in the management of asthma.³⁵³ Low magnesium intake has been associated with a higher prevalence of asthma with increasing intake resulting in reduced bronchial hyper-responsiveness and higher lung function.³⁵⁴ Magnesium plays a beneficial role in the treatment of asthma through bronchial smooth muscle relaxation, leading to the use of intravenous or inhaled preparations of magnesium sulphate for acute asthma attacks.³⁵⁵ Studies of oral supplementation are limited and more trials are required.³⁵⁶⁻³⁵⁸

6.2.6 Fish oils/lipids

In vitro studies suggest that supplementing the diet with n-3PUFAs, which are most commonly found in fish oils, might reduce the inflammation associated with asthma.^{359, 360} Results from observational studies are inconsistent and a Cochrane review of nine RCTs concluded that there was insufficient evidence to recommend fish oil supplementation for the treatment of asthma.³⁶¹

6.2.7 Antioxidants

Observational studies have reported that low intakes of vitamin C, vitamin E and selenium are associated with a higher prevalence of asthma.³⁰³ Intervention studies suggest that neither supplementation with vitamin C, vitamin E nor selenium is associated with clinical benefits in people with asthma.³⁶²⁻³⁶⁴ Observational studies in both adults and children have also consistently shown that a high intake of fresh fruit and vegetable is associated with less asthma and better pulmonary function.³⁶⁵⁻³⁷¹ No intervention studies evaluating the intake of fruit or vegetables and their effects on asthma have been reported.

6.2.8 Vitamin D

A systematic review of nine RCTs (adults, n=658, children, n=435) examined whether administration of vitamin D reduced severe asthma exacerbations (defined as those requiring oral corticosteroids) or improved asthma symptom control. In three of the nine included trials (n=680/1093), of predominantly adults with mild to moderate asthma on treatment with ICS, vitamin D reduced the risk

1++

of severe asthma exacerbation. The number of exacerbations in children was too low to evaluate this outcome.³⁷² 1++

A further subgroup analysis reported that it was not clear whether the reduction in risk of exacerbation was confined to people with lower baseline vitamin D status. Vitamin D dosage regimes varied between trials and no evidence was provided about the optimum dose of vitamin D or circulating 25-hydroxyvitamin D concentrations. Serious adverse effects did not vary between those receiving vitamin D or placebo.³⁷³ 1++

Further research is required on whether the effects of vitamin D supplementation are confined to people with lower baseline vitamin D status, and into the effects in children, and in people with frequent severe asthma attacks.

6.2.9 Weight reduction in overweight and obese patients with asthma

The current evidence base for weight reduction interventions to improve asthma control is inadequate in quantity and quality. A Cochrane review concluded that as the benefit of weight loss as an intervention for asthma control is uncertain, "...clinicians should be prepared to help patients to make a decision that is consistent with their own values...".³⁷⁴ 1++

Two RCTs in adults and one pilot RCT in children investigating the effects of interventions to reduce weight on asthma control and biomarkers of asthma severity, reported reductions in BMI but varying effects on asthma control and biomarkers.³⁷⁵⁻³⁷⁷ The pilot study in children (n=32) reported that a 10-week dietary intervention improved asthma control and lung function but had no effect on inflammation. This study was not, however, powered to determine clinical changes; baseline differences between control and intervention groups and in interactions with healthcare staff may have influenced the results.³⁷⁵ In adults, a trial (n=46) combining dietary (including two free meal replacements a day) and exercise (free gym membership and personal training sessions) components reported improved lung function, asthma symptoms and biomarkers of neutrophilic inflammation with a 5–10% weight loss.³⁷⁷ A larger trial (n=330), however, reported no significant differences in asthma outcomes between obese adults with asthma receiving a weight-loss intervention (combining dietary and exercise components) and those in the control group. Weight loss of more than 10% in either group was, however, associated with improvements in asthma symptom control compared with those with unchanged weight.³⁷⁶ 1+

Although evidence is limited, these studies show that dietary and weight-loss interventions are feasible in overweight or obese adults and children with asthma and that they may improve asthma control, lung function and inflammation, although weight loss of greater than 10% may be necessary to achieve benefit.

B Weight-loss interventions (including dietary and exercise-based programmes) should be considered for overweight and obese adults and children with asthma to improve asthma control.

6.2.10 Probiotics

Studies have suggested that an imbalance in gut flora is associated with a higher risk of development of allergy.³⁷⁸ Trials have investigated the use of probiotics in the treatment of established allergic disease with variable results.^{379, 380} Only one study focused on asthma, finding a decrease in eosinophilia but no effect on clinical parameters.³⁸¹ 1+
2+

In the absence of evidence of benefit, it is not possible to recommend the use of probiotics in the management of asthma.

6.2.11 Immunisation

A number of large studies have concluded that high vaccination coverage has no significant impact on any allergic outcome or asthma. There is a suggestion that the higher the vaccine coverage the greater the possibility that there is a degree of protection against the development of allergy in the first years of life.³⁸²⁻³⁸⁵

There is some discussion about whether BCG immunisation may confer protection against allergy and asthma. Research has focused on primary prophylaxis (*see section 6.1.11*), although there are some studies investigating the use of BCG, with or without allergen, as a means to switch off allergic immune responses. There are some observations suggesting that benefit might occur,³⁸⁶ but results of trials have been disappointing.^{387, 388} This is an area that requires further investigation.

There has been concern that influenza vaccination might aggravate respiratory symptoms, although any such effect would be outweighed by the benefits of the vaccination.³⁸⁹ Studies in children have suggested that immunisation with the vaccine does not exacerbate asthma,³⁹⁰ but has a small beneficial effect on quality of life in children with asthma.³⁹¹ The immune response to the immunisation may be adversely affected by high-dose ICS therapy and this requires further investigation.³⁹² A Cochrane review of pneumococcal vaccine found very limited evidence to support its use specifically in individuals with asthma.³⁹³

1++

B Immunisations should be administered independent of any considerations related to asthma. Responses to vaccines may be attenuated by high-dose inhaled corticosteroids.

6.2.12 Acupuncture

A Cochrane review of 21 trials highlighted many methodological problems with the studies reviewed. Only seven of the trials, involving 174 patients, employed randomisation to active or sham acupuncture points (with no recognised activity) for the treatment of persistent or chronic asthma. Blinding was a major problem in the assessment of the results and there were considerable inconsistencies in methodology. The review concluded that there was no evidence for a clinically valuable benefit from acupuncture and no significant benefits in relation to lung function.³⁹⁴ A later systematic review and meta-analysis of 11 RCTs found no evidence of an effect in reducing asthma severity but a suggestion that where bronchoconstriction was induced to establish efficacy of acupuncture there was a beneficial effect. Concern was expressed about potential publication bias in favour of positive outcome studies.³⁹⁵ Two other trials of acupuncture in relation to induced asthma were also negative.^{396, 397}

1+

6.2.13 Air ionisers

Ionisers have been widely promoted as being of benefit for patients with asthma. A Cochrane review of six studies, five using negative ion generators and one with a positive ion generator, found no evidence that air ionisers are of benefit in reducing symptoms in patients with asthma. One of the included studies in children (n=12, age range 3–11) showed that positively-ionised air was associated with bronchoconstriction, and another (n=20, age range 9–15) showed an increase in night-time cough, although this was not statistically significant.³⁹⁸

1++
1+

A Air ionisers are not recommended for the treatment of asthma.

6.2.14 Breathing exercises

Behavioural programmes centred on breathing exercises and dysfunctional breathing reduction techniques (including physiotherapist-delivered breathing programmes such as the Papworth method, and the Buteyko method) can lead to modest improvements in asthma symptoms and quality of life, and reduce bronchodilator requirement in adults with asthma, although have little effect on lung function or airway inflammation.³⁹⁹⁻⁴⁰² These techniques involve instruction by a trained therapist in exercises to reduce respiratory rate, minute volume and to promote nasal, diaphragmatic breathing. Trials that include more than five hours of intervention appeared more likely to be effective than shorter courses.⁴⁰⁰ They should ideally be provided as part of integrated medical care.

1++
1+

A systematic review of inspiratory muscle training for adults with asthma (n=113), including five RCTs, reported that evidence for its use was inconclusive.⁴⁰³

1++

One high-quality RCT, including 655 adults with asthma and impaired asthma-related quality of life, demonstrated that breathing retraining can be successfully delivered as a self-guided audiovisual programme, leading to equivalent quality-of-life benefits, measured by the AQLQ, and likely greater cost effectiveness compared with a programme delivered face-to-face by a physiotherapist.³⁹⁹ There were, however, clinically important improvements in AQLQ scores in a substantial proportion of the usual treatment group as well as in the two intervention groups.

1++

In a systematic review of yoga for asthma including 15 RCTs (13 in adults) and 1,048 participants (number of children not specified), meta-analysis of five of the eight studies that included quality of life as an outcome, suggested that yoga may improve quality of life,⁴⁰⁴ although improvements were mostly observed in trials which did not include a sham or placebo intervention in the control arm. Furthermore, the yoga interventions studied included breathing, postures, and meditation, and results were not presented for the effects of breathing exercises alone. Although current evidence does not support yoga as a routine intervention for people with asthma, it could be considered as an additional therapy or as an alternative to other forms of breathing exercises.⁴⁰⁵

1++

There is currently insufficient evidence on breathing exercises or yoga in children and adolescents aged 12 and under on which to base a recommendation.^{404, 406}

A Breathing exercise programmes (including face-to-face physiotherapist-taught methods and audiovisual programmes) can be offered to adults with asthma as an adjuvant to pharmacological treatment to improve quality of life and reduce symptoms.

6.2.15 Herbal and traditional chinese medicine

A Cochrane review identified 17 trials, nine of which reported some improvement in lung function but it was not clear that the results would be generalisable.⁴⁰⁷ A more recent double-blinded placebo-controlled trial of a Chinese herb decoction (Ding Chuan Tang) showed improvement in airway hyper-responsiveness in children with stable asthma.⁴⁰⁸ It is difficult to disentangle the effects of multiple ingredients; Ding Chuan Tang for example contains nine components. A second study, of 100 children with asthma, found that a five-herb mixture gave some benefits in relation to lung function and symptoms compared with placebo.⁴⁰⁹

1+

The conclusions of these trials of Chinese herbal therapy are not generalisable due to variations in the herbal mixtures and study designs. There are likely to be pharmacologically active ingredients in the mixtures and further investigations are warranted. There is a need for large appropriately powered controlled studies.

6.2.16 Homeopathy

A Cochrane review identified only three methodologically sound RCTs, two of which reported some positive effects. A criticism of the studies was that they did not employ individualised homeopathy, which is the essence of this approach to treatment.⁴¹⁰ A more recent trial of individualised homeopathy in childhood asthma, which was placebo controlled and appropriately powered, failed to show any evidence of benefit over conventional treatment in primary care.⁴¹¹

1++
1+

6.2.17 Hypnosis and relaxation therapies

A systematic review of relaxation therapies, including hypnotherapy, identified five controlled trials, two of which showed some benefits. Overall the methodology of the studies was poor and the review concluded that there was a lack of evidence of efficacy but that muscle relaxation could conceivably benefit lung function in patients with asthma.⁴¹²

1++

6.2.18 Manual therapy including massage and spinal manipulation

A Cochrane review identified four relevant RCTs.⁴¹³ The two trials of chiropractic suggest that there is no role for this modality of treatment in the management of asthma. No conclusions can be drawn on massage therapy.

6.2.19 Physical exercise training

A Cochrane review has shown no effect of physical training on PEF, FEV₁, FVC or ventilation at maximal exercise capacity (V_{Emax}).⁴¹⁴ However, oxygen consumption, maximum heart rate, and work capacity all increased significantly. Most studies discussed the potential problems of exercise-induced asthma, but none made any observations on this phenomenon. As physical training improves indices of cardiopulmonary efficiency, it should be seen as part of a general approach to improving lifestyle and rehabilitation in people with asthma, with appropriate precautions advised about exercise-induced asthma (*see section 7.7.2*).

6.2.20 Family therapy

A Cochrane review identified two trials (n=55) showing that family therapy may be a useful adjunct to medication in children with asthma.⁴¹⁵ Small study size limits the ability to form recommendations.



For those with difficult asthma in childhood, there may be a role for family therapy as an adjunct to pharmacotherapy.

7 Pharmacological management

The aim of asthma management is control of the disease. Complete control of asthma is defined as:

- no daytime symptoms
- no night-time awakening due to asthma
- no need for rescue medication
- no asthma attacks
- no limitations on activity including exercise
- normal lung function (in practical terms FEV₁ and/or PEF > 80% predicted or best)
- minimal side effects from medication.

✓ Lung function measurements cannot be used reliably to guide asthma management in children under five years of age.

Patients may have different goals and may wish to balance the aims of asthma management against the potential side effects or inconvenience of taking medication necessary to achieve perfect control.

A phased approach aims to abolish symptoms as soon as possible and to optimise peak flow by starting treatment at the level most likely to achieve this. Patients should start treatment at the level most appropriate to the initial severity of their asthma. The aim is to achieve early control and to maintain it by increasing treatment as necessary and decreasing treatment when control is good (*see Figures 2 and 3 for summaries of management in adults and children*).

✓ Before initiating a new drug therapy practitioners should check adherence with existing therapies (*see section 5.4*), check inhaler technique (*see section 8*), and eliminate trigger factors (*see section 6*).

Until May 2009 all doses of ICS in this section were referenced against beclometasone dipropionate (BDP) given via chlorofluorocarbon metered dose inhalers (CFC-MDIs). BDP-CFC is now unavailable. Differences in how the doses of ICS (as well as other drug classes) are expressed (metered dose or delivered dose, ie the dose that leaves the mouthpiece), mean it is increasingly difficult to cover all the possible doses in the text. The doses of ICS are therefore expressed as very low (generally paediatric doses), low (generally starting dose for adults), medium and high (*see Tables 12 and 13*). Adjustments to doses will have to be made for other inhaler devices and other corticosteroid molecules (*see section 8.2*).

Recommendations in sections 7 and 8 have been graded and the supporting evidence assessed for adults and adolescents over 12 years old, children aged 5–12 years, and children aged under 5 years. The evidence is less clear in children under two and the threshold for seeking an expert opinion should be lowest in these children.

1	2	3	1 Adults and adolescents aged over 12
			2 Children aged 5–12 years
			3 Children under 5 years

Recommendation does not apply to this age group.

7.1 Intermittent reliever therapy

Adults and children with a diagnosis of asthma should be prescribed a short-acting bronchodilator to relieve symptoms. For those with infrequent short-lived wheeze occasional use of reliever therapy may be the only treatment required. For exercise-induced asthma see section 7.7.2.

The following medicines act as short-acting bronchodilators:

- inhaled short-acting β_2 agonists⁴¹⁶
- inhaled ipratropium bromide⁴¹⁷
- theophyllines.⁴¹⁶

>12 years	5–12 years	<5 years
1++	1+	4
1+	1++	
1++		

Short-acting inhaled β_2 agonists work more quickly and/or with fewer side effects than the alternatives.⁴¹⁶

A	B	D	Prescribe an inhaled short-acting β_2 agonist as short-term reliever therapy for all patients with symptomatic asthma.

7.1.1 Frequency of dosing of inhaled short-acting β_2 agonists

Good asthma control is associated with little or no need for short-acting β_2 agonist. Short-acting β_2 agonists should only be used as required for the relief of symptoms.

✓	Anyone prescribed more than one short-acting bronchodilator inhaler device a month should be identified and have their asthma assessed urgently and measures taken to improve asthma control if this is poor.
---	---

7.2 Regular preventer therapy

Treatments have been judged on their ability to improve symptoms, improve lung function, and prevent asthma attacks, with an acceptable safety profile. Improvement of quality of life, while important, is the subject of too few studies to be used to make recommendations at present.

7.2.1 Inhaled corticosteroids

Inhaled corticosteroids are the most effective preventer drug for adults and older children for achieving overall treatment goals.^{418–421} There is a body of evidence demonstrating that, at recommended doses, they are also safe and effective in children under five with asthma.^{422–431}

>12 years	5–12 years	<5 years
1++	1++	1++

Many non-atopic children under five with recurrent episodes of viral-induced wheezing do not go on to have chronic atopic asthma. The majority do not require treatment with regular ICS (*see section 3.3*).

A	A	A	Inhaled corticosteroids are the recommended preventer drug for adults and children for achieving overall treatment goals.

Inhaled corticosteroids should be considered for adults, children aged 5–12 and children under the age of five with any of the following features: using inhaled β_2 agonists three times a week or more; symptomatic three times a week or more; or waking one night a week. In addition, ICS should be considered in adults and children aged 5–12 who have had an asthma attack requiring oral corticosteroids in the last two years.^{432–436}

>12 years	5–12 years	<5 years
1+	1+	1+

Inhaled corticosteroids should be considered for patients with any of the following asthma-related features:

B	C		• asthma attack in the last two years
B	B	B	• using inhaled β_2 agonists three times a week or more
B	B	B	• symptomatic three times a week or more
B	C	✓	• waking one night a week.

Alternative initial preventer therapies are available but are less effective than ICS (*see section 7.2.7*).

7.2.2 Starting dose of inhaled corticosteroids

In mild to moderate asthma, starting at high doses of ICS and stepping down confers no benefit.⁴³⁷

>12 years	5–12 years	<5 years
1+	1+	

- ✓ Start patients at a dose of inhaled corticosteroids appropriate to the severity of disease.
- ✓ A reasonable starting dose of inhaled corticosteroids will usually be low dose for adults (*see Table 12*) and very low dose for children (*see Table 13*).
- ✓ Titrate the dose of inhaled corticosteroid to the lowest dose at which effective control of asthma is maintained.

7.2.3 Frequency of dosing of inhaled corticosteroids

Most current ICS are slightly more effective when taken twice rather than once daily, but may be used once daily in some patients with milder disease and good or complete control of their asthma.^{416, 419, 434, 438, 439}

>12 years	5–12 years	<5 years
1+	1+	1+

There is little evidence of benefit for dosage frequency more than twice daily.⁴¹⁹

An RCT comparing daily ICS with intermittent (rescue) ICS in children aged 6–18 years with mild persistent asthma suggests that daily ICS are more effective at preventing asthma attacks.⁴⁴⁰

>12 years	5–12 years	<5 years
1++	1++	

A	A	A	Give inhaled corticosteroids initially twice daily (except ciclesonide which is given once daily).
A	A		Once-a-day inhaled corticosteroids at the same total daily dose can be considered if good control is established.

7.2.4 Comparison of inhaled corticosteroids

Many studies comparing different ICS are of inadequate design and have been omitted from further assessment. In view of the clear differences between normal volunteers and asthma patients in the absorption of ICS, data from normal volunteers have not been taken into account. Only studies in which more than one dose of at least one of the ICS or both safety and efficacy had been studied together in the same trial were evaluated.

Non-blinded studies also had to be considered because of the problems of obtaining competitors' delivery devices.

Beclometasone dipropionate and budesonide are approximately equivalent in clinical practice, although there may be variations with different delivery devices. There is limited evidence from two open studies of suboptimal design that budesonide via the Turbohaler® is more clinically effective.⁴⁴¹ However, at present a 1:1 ratio should be assumed when changing between BDP and budesonide.

Fluticasone propionate provides equal clinical activity to BDP and budesonide at half the dosage. The evidence that it causes fewer side effects at doses with equal clinical effect is limited. Mometasone appears to provide equal clinical activity to BDP and budesonide at half the dosage.⁴⁴² It is difficult to establish the exact equipotent dose of fluticasone furoate.^{443, 444}

7.2.5 Safety of inhaled corticosteroids

The safety of ICS is of crucial importance and a balance between benefits and risks for each individual needs to be assessed. Account should be taken of other topical steroid therapy when assessing systemic risk. It has been suggested that steroid warning cards (for example the *High Dose Inhaled Corticosteroid Safety Card* developed by the London Respiratory Network for NHS England⁴⁴⁵) should be issued to patients on higher dose ICS, but the benefits and possible disadvantages, particularly with regard to adherence, to such a policy remain to be established.

Adults

There is little evidence that low doses of ICS cause any short-term detrimental effects apart from the local side effects of dysphonia and oral candidiasis. However, the possibility of long-term effects on bone has been raised. One systematic review reported no effect on bone density at doses up to 1,000 micrograms BDP per day.⁴⁴⁶ The significance of small biochemical changes in adrenocortical function is unknown.



Titrate the dose of inhaled corticosteroid to the lowest dose at which effective control of asthma is maintained.

Children

Administration of medium- or high-dose ICS may be associated with systemic side effects.⁴⁴⁷ These may include growth failure and adrenal suppression. Isolated growth failure is not a reliable indicator of adrenal suppression and monitoring growth cannot be used as a screening test of adrenal function.^{438, 448}

Clinical adrenal insufficiency has been identified in a small number of children who have become acutely unwell at the time of intercurrent illness. Most of these children had been treated with high doses of ICS. The dose or duration of ICS treatment required to place a child at risk of clinical adrenal insufficiency is unknown but is likely to occur at ≥ 800 micrograms BDP per day or equivalent (medium dose ICS and above; see Table 12). The low-dose adrenocorticotrophic hormone test is considered to provide a physiological stimulation of adrenal responsiveness but it is not known how useful such a sensitive test is at predicting clinically relevant adrenal insufficiency.^{449, 450} In addition, it is unknown how frequently tests of adrenal function would need to be repeated if a child remained on high-dose inhaled corticosteroid. At higher doses, add-on agents, for example, long-acting β_2 agonists, should be actively considered.

While the use of ICS may be associated with adverse effects (including the potential to reduce bone mineral density) with careful ICS dose adjustment this risk is likely to be outweighed by their ability to reduce the need for multiple bursts of oral corticosteroids for acute asthma attacks.⁴⁵¹

- ✓ Monitor growth (height and weight centile) of children with asthma on an annual basis.
- ✓ The lowest dose of inhaled corticosteroids compatible with maintaining asthma control should be used.

For children treated with medium- or high-dose ICS:

- ✓ Specific written advice about steroid replacement in the event of a severe intercurrent illness or surgery should be part of the management plan.
- ✓ The child should be under the care of a specialist paediatrician for the duration of the treatment.

Adrenal insufficiency is a possibility in any child maintained on ICS presenting with shock or a decreased level of consciousness; serum biochemistry and blood glucose levels should be checked urgently. Intramuscular (IM) hydrocortisone may also be required.

7.2.6 Smoking and inhaled corticosteroids

Current and previous smoking reduces the effect of ICS, which may be overcome with increased doses.^{336, 452}

>12 years	5-12 years	<5 years
1+		

Patients should be advised that smoking reduces the effectiveness of therapy.



Clinicians should be aware that higher doses of inhaled corticosteroids may be needed in patients who are smokers or ex-smokers.

7.2.7 Other preventer therapies

Inhaled corticosteroids are the first choice preventer drug. There are alternative, less effective preventer therapies for patients taking short-acting β_2 agonists alone.

- Leukotriene receptor antagonists (LTRA) have some beneficial clinical effect^{419, 453, 454}
 - In children under five years who are unable to take ICS, leukotriene receptor antagonists may be used as an alternative preventer^{455, 456}
- Sodium cromoglicate and nedocromil sodium
 - Sodium cromoglicate is of some benefit in adults^{416, 457} and is effective in children aged 5-12⁴⁵⁸
 - Nedocromil sodium is of benefit in adults and children >5^{416, 459}
 - There is no clear evidence of benefit with sodium cromoglicate in children aged <5⁴⁶⁰
- Theophyllines have some beneficial effect^{416, 418}
- Antihistamines and ketotifen are ineffective.⁴⁶¹

>12 years	5-12 years	<5 years
1++	1++	1+
		1+
1+	1+	
1++	1+	
1++	1++	1++
1++	1++	1++

Table 12: Categorisation of inhaled corticosteroids by dose – adults* (see also Figure 2)

ICS	Dose		
	Low dose	Medium dose	High dose#
Pressurised metered dose inhalers (pMDI)			
Beclometasone dipropionate			
Non-proprietary	100 micrograms two puffs twice a day	200 micrograms two puffs twice a day	200 micrograms four puffs twice a day
Clenil Modulite pMDI	100 micrograms two puffs twice a day	200 micrograms two puffs twice a day	250 micrograms two puffs twice a day 250 micrograms four puffs twice a day
Kelhale pMDI (extrafine)	50 micrograms two puffs twice a day	100 micrograms two puffs twice a day	100 micrograms four puffs twice a day
Qvar pMDI (extrafine) Qvar Autohaler (extrafine) Qvar Easi-Breathe (extrafine)	50 micrograms two puffs twice a day	100 micrograms two puffs twice a day	100 micrograms four puffs twice a day
Soprobec pMDI	100 micrograms two puffs twice a day	200 micrograms two puffs twice a day	250 micrograms two puffs twice a day 250 micrograms four puffs twice a day
Ciclesonide			
Alvesco pMDI	80 micrograms two puffs once a day	160 micrograms two puffs once a day	160 micrograms two puffs twice a day
Fluticasone propionate			
Flixotide Evohaler	50 micrograms two puffs twice a day	125 micrograms two puffs twice a day	250 micrograms two puffs twice a day
Dry powder inhalers (DPI)			
Beclometasone			
Non-proprietary Easyhaler	200 micrograms one puff twice a day	200 micrograms two puffs twice a day	n/a
Budesonide			
Non-proprietary Easyhaler	100 micrograms two puffs twice a day	200 micrograms two puffs twice a day	400 micrograms two puffs twice a day
Budelin Novolizer	n/a	200 micrograms two puffs twice a day	200 micrograms four puffs twice a day
Pulmicort Turbohaler	100 micrograms two puffs twice a day 200 micrograms one puff twice a day	200 micrograms two puffs twice a day 400 micrograms one puff twice a day	400 micrograms two puffs twice a day
Fluticasone propionate			
Flixotide Accuhaler	100 micrograms one puff twice a day	250 micrograms one puff twice a day	500 micrograms one puff twice a day
Mometasone			
Asmanex Twisthaler	200 micrograms one puff twice a day	400 micrograms one puff twice a day	n/a

* Different products and doses are licensed for different age groups and some are not licensed for use in children. Prior to prescribing, the relevant summary of product characteristics (SPC) should be checked (www.medicines.org.uk/emc).

High doses (shaded boxes) should only be used after referring the patient to specialist care.

(Table 12 continues on next page)

Table 12 (continued): Categorisation of inhaled corticosteroids by dose – adults*
(see also Figure 2)

ICS	Dose		
	Low dose	Medium dose	High dose#
Combination inhalers			
Beclometasone dipropionate (extrafine) with formoterol			
Fostair (pMDI)	100/6 one puff twice a day	100/6 two puffs twice a day	200/6 two puffs twice a day
Fostair (NEXThaler)	100/6 one puff twice a day	100/6 two puffs twice a day	200/6 two puffs twice a day
Budesonide with formoterol			
DuoResp Spiromax	160/4.5 one puff twice a day	160/4.5 two puffs twice a day 320/9 one puff twice a day	320/9 two puffs twice a day
Symbicort Turbohaler	100/6 two puffs twice a day 200/6 one puff twice a day	200/6 two puffs twice a day 400/12 one puff twice a day	400/12 two puffs twice a day
Fobumix Easyhaler	80/4.5 two puffs twice a day 160/4.5 one puff twice a day	160/4.5 two puffs twice a day 320/9 one puff twice a day	320/9 two puffs twice a day
Fluticasone propionate with formoterol			
Flutiform MDI	50/5 two puffs twice a day	125/5 two puffs twice a day	250/10 two puffs twice a day
Flutiform K-haler	50/5 two puffs twice a day	125/5 two puffs twice a day	n/a
Fluticasone propionate with salmeterol			
Aerivio Spiromax	n/a	n/a	500/50 one puff twice a day
AirFluSal Forspiro	n/a	n/a	500/50 one puff twice a day
AirFluSal pMDI	n/a	125/25 two puffs twice a day	250/25 two puffs twice a day
Aloflute pMDI	n/a	125/25 two puffs twice a day	250/25 two puffs twice a day
Combisal pMDI	50/25 two puffs twice a day	125/25 two puffs twice a day	250/25 two puffs twice a day
Fusacomb Easyhaler	n/a	250/50 one puff twice a day	500/50 one puff twice a day
Sereflo pMDI	n/a	125/25 two puffs twice a day	250/25 two puffs twice a day
Seretide Accuhaler	100/50 one puff twice a day	250/50 one puff twice a day	500/50 one puff twice a day
Seretide Evohaler	50/25 two puffs twice a day	125/25 two puffs twice a day	250/25 two puffs twice a day
Sirdupla pMDI	n/a	125/25 two puffs twice a day	250/25 two puffs twice a day
Stalpex Orbicel	n/a	n/a	500/50 one puff twice a day
Fluticasone furoate with vilanterol			
Relvar Ellipta	n/a	92/22 one puff once a day	184/22 one puff once a day

* Different products and doses are licensed for different age groups and some are not licensed for use in children. Prior to prescribing, the relevant summary of product characteristics (SPC) should be checked (www.medicines.org.uk/emc).

High doses (shaded boxes) should only be used after referring the patient to specialist care.

Table 13: Categorisation of inhaled corticosteroids by dose – children*
(see also Figure 3)

ICS	Dose		
	Very low dose	Low dose	Medium dose#
Pressurised metered dose inhalers (pMDI) with spacer			
Beclometasone dipropionate			
Non-proprietary	50 micrograms two puffs twice a day	100 micrograms two puffs twice a day	200 micrograms two puffs twice a day
Clenil Modulite	50 micrograms two puffs twice a day	100 micrograms two puffs twice a day	200 micrograms two puffs twice a day
Qvar (extrafine) Qvar autohaler Qvar Easi-breathe	n/a	50 micrograms two puffs twice a day	100 micrograms two puffs twice a day
Soprobec	50 micrograms two puffs twice a day	100 micrograms two puffs twice a day	200 micrograms two puffs twice a day
Ciclesonide			
Alvesco Aerosol inhaler	n/a	80 micrograms two puffs once a day	160 micrograms two puffs once a day
Fluticasone propionate			
Flixotide Evohaler	50 micrograms one puff twice a day	50 micrograms two puffs twice a day	125 micrograms two puffs twice a day
Dry powder inhalers (DPI)			
Budesonide			
Non-proprietary Easyhaler	n/a	100 micrograms two puffs twice a day	200 micrograms two puffs twice a day
Pulmicort Turbohaler	100 micrograms one puff twice a day	100 micrograms two puffs twice a day 200 micrograms one puff twice a day	200 micrograms two puffs twice a day 400 micrograms one puff twice a day
Fluticasone propionate			
Flixotide Accuhaler	50 micrograms one puff twice a day	100 micrograms one puff twice a day	250 micrograms one puff twice a day
Mometasone			
Asmanex Twisthaler	n/a	200 micrograms one puff twice a day	n/a
Combination inhalers			
Budesonide with formoterol			
Symbicort Turbohaler	100/6 one puff twice a day	100/6 two puffs twice a day 200/6 one puff twice a day	n/a
Fluticasone propionate with salmeterol			
Combisal MDI	n/a	50/25 two puffs twice a day	n/a
Seretide Accuhaler	n/a	100/50 one puff twice a day	n/a
Seretide Evohaler	n/a	50/25 two puffs twice a day	n/a

* Different products and doses are licensed for different age groups and some are not licensed for use in children. Prior to prescribing, the relevant summary of product characteristics (SPC) should be checked (www.medicines.org.uk/emc).

Medium doses (shaded boxes) should only be used after referring the patient to specialist care.

7.3 Initial add-on therapy

A proportion of patients with asthma may not be adequately controlled with low-dose ICS alone. Before initiating a new drug therapy practitioners should recheck adherence (*see section 5.4*), inhaler technique and eliminate trigger factors. The duration of a trial of add-on therapy will depend on the desired outcome. For instance, preventing nocturnal awakening may require a relatively short trial of treatment (days or weeks), whereas preventing asthma attacks or decreasing steroid tablet use may require a longer trial of therapy (weeks or months). If there is no response to treatment the drug should be discontinued.

7.3.1 Criteria for introduction of add-on therapy

No exact dose of ICS can be deemed the correct dose at which to add another therapy. The addition of other treatment options to ICS has been investigated at doses from 200–1,000 micrograms BDP in adults and up to 400 micrograms BDP in children.^{462–465} Many patients will benefit more from add-on therapy than from increasing ICS above doses as low as 200 micrograms BDP/day. At doses of ICS above 800 micrograms BDP/day side effects become more frequent. An absolute threshold for introduction of add-on therapy in all patients cannot be defined.

>12 years	5–12 years	<5 years
1++	1+	

7.3.2 Inhaled long-acting β_2 agonist

The addition of an inhaled long-acting β_2 agonist (LABA) to ICS alone improves lung function and symptoms, and decreases asthma attacks in adults and children.^{462, 466–472}


Long-acting inhaled β_2 agonists should not be used without ICS.⁴⁷³

>12 years	5–12 years	<5 years
1++	1++	

Evidence to guide the choice of initial add-on therapy is stronger in adults than in children. On the basis of current evidence, LABA is the first choice initial add-on therapy in adults.

In children, options for initial add-on therapy are limited to LABA and LTRA, with evidence to support both individually, but insufficient evidence to support use of one over the other (*see section 7.4.2*).⁴⁷² LABA are not licensed for use in children under four years of age and evidence for use of LTRA in this age group is limited to studies comparing LTRA with ICS or placebo (*see section 7.2.7*).

1++	1++	



The first choice as add-on therapy to inhaled corticosteroids in adults is an inhaled long-acting β_2 agonist, which should be considered before increasing the dose of inhaled corticosteroid.

In children aged five and over, an inhaled long-acting β_2 agonist or a leukotriene receptor antagonist can be considered as initial add-on therapy.

7.3.3 Safety of long-acting β_2 agonist

Following a review in 2007 of LABA in the treatment of adults, adolescents, and children with asthma, the Medicines and Healthcare products Regulatory Agency (MHRA) further reviewed the use of LABA, specifically in children younger than 12 years of age and concluded that the benefits of these medicines used in conjunction with ICS in the control of asthma symptoms outweigh any apparent risks.⁴⁷⁴



Long-acting inhaled β_2 agonists should only be started in patients who are already on inhaled corticosteroids, and the inhaled corticosteroid should be continued.

7.3.4 Combination inhaled corticosteroid/long-acting β_2 agonist inhalers

In efficacy studies, where there is generally good adherence, there is no difference in efficacy in giving ICS and a LABA in combination or in separate inhalers.⁴⁷⁵

>12 years	5-12 years	<5 years
1++	1++	

In clinical practice it is generally considered that combination inhalers aid adherence and also have the advantage of guaranteeing that the LABA is not taken without the ICS.



Combination inhalers are recommended to:

- guarantee that the long-acting β_2 agonist is not taken without inhaled corticosteroid
- improve inhaler adherence.

7.3.5 Single combination inhaler for maintenance and reliever therapy

The use of a single combination inhaler for maintenance and reliever therapy (MART) is an alternative approach to the introduction of a fixed-dose twice-daily combination inhaler which might suit some individuals. It relies on the rapid onset of reliever effect with formoterol and by including a dose of inhaled corticosteroid ensures that, as the need for a reliever increases, the dose of preventer medication is also increased. This underpins the self-management plan which must be provided with a MART regime. *See section 5.2.3* for a description of increasing inhaled corticosteroids at the onset of an attack. Maintenance and reliever therapy may also lower the overall dose of ICS needed to prevent asthma attacks.

A systematic review comparing a combined ICS/LABA inhaler as MART with ICS alone or with current best practice (ICS with or without LABA) showed that maintenance and reliever therapy can reduce the risk of asthma attacks requiring oral steroids in patients who are not well controlled on ICS alone and who have a history of asthma attacks.⁴⁷⁶ The review reported more withdrawals due to adverse events in the maintenance and reliever therapy group (possibly because patients did not adjust well to the change in inhaler) compared with the current best practice group, but no significant difference between the groups in serious adverse events. All trials were funded by the manufacturer.

>12 years	5-12 years	<5 years
1++		

In a subsequent systematic review including 16 RCTs and 22,748 patients, a meta-analysis of five of the 16 studies concluded that, in patients aged 12 and over, use of MART was associated with a reduced risk of asthma attacks compared with standard ICS/LABA treatment including the same dose of ICS as the MART group (in five of the nine studies), or to standard ICS/LABA treatment including a higher dose of ICS than the MART group (in two of 16 studies). Among children aged 4-11, a subgroup analysis from a single RCT reported that MART was associated with a lower risk of asthma attacks compared with standard ICS/LABA treatment including the same dose of ICS as the MART group. There was no difference in impact on quality of life (QoL), asthma control, lung function and asthma medication use between MART and regular fixed-dose treatment regimens.⁴⁷⁷ Fifteen of the 16 studies used a combination of budesonide and formoterol in a dry powder inhaler.

1++		
-----	--	--

If this management option is introduced the total regular dose of daily ICS should not be decreased. Patients taking rescue doses of their combination inhaler once a day or more on a regular basis should have their treatment reviewed. Careful education of patients about the specific issues around this management strategy is required.

At present, maintenance and reliever therapy is only licensed for use with budesonide/formoterol or beclomethasone/formoterol. The summaries of product characteristics should be consulted for age-appropriate prescribing and maximum dosing regimens. Not all combination products are licensed for maintenance and reliever therapy. The appropriate combination inhaler should be prescribed by brand name.



Consider the option of combined maintenance and reliever therapy in adult patients who have a history of asthma attacks on medium dose ICS or ICS/LABA.

7.4 Additional controller therapies

If control remains poor on low-dose (adults) or very low-dose (children aged five and over) ICS plus a LABA, recheck the diagnosis, assess adherence to existing medication and check inhaler technique before increasing therapy. If more intense treatment is appropriate, then the following options can be considered.

7.4.1 Increased dose of inhaled corticosteroids

If there is an improvement when LABA is added, but control remains suboptimal, continue with the LABA and increase the dose of ICS to medium (adults) or low dose (children 5–12 years). If there is no improvement when a LABA is added, consider stopping the LABA before increasing the dose of ICS.⁴⁷⁵

>12 years	5–12 years	<5 years
4	4	

7.4.2 Leukotriene receptor antagonists

Evidence to support the use of LTRA as an add-on therapy to ICS plus LABA is lacking and evidence for their use is largely based on extrapolation from trials of LTRA as add-on therapy to ICS alone. The addition of LTRA to ICS may provide improvement in lung function, a decrease in asthma attacks, and an improvement in symptoms in adults and children over five years of age, although reported benefits differ between studies and evidence is limited in children.^{454, 478, 479}

>12 years	5–12 years	<5 years
1++	1++	

A systematic review of studies comparing the addition of LTRA to ICS with the addition of LABA to ICS showed that the addition of LABA to ICS was more effective at reducing asthma attacks (the primary outcome) and improving secondary outcomes including SABA use, symptoms and quality of life in adults, although differences were generally small. There was insufficient evidence on which to base conclusions regarding which add-on therapy is more effective in children.⁴⁷²

>12 years	5–12 years	<5 years
1++	1++	

In adults, the addition of LTRA to ICS is superior to ICS alone and has a similar effect on asthma control to high-dose ICS. High-dose ICS, however, appears superior to ICS-LTRA for some pulmonary function indices, although further studies to investigate this are required.⁴⁸⁰

>12 years	5–12 years	<5 years
1+		

In adults, if there is no improvement following addition of a LABA, consider stopping the LABA and initiating a trial of LTRA.



If asthma control remains suboptimal after the addition of an inhaled long-acting β_2 agonist then:

- increase the dose of inhaled corticosteroids from low dose to medium dose in adults or from very low dose to low dose in children (5–12 years), if not already on these doses.

or

- consider adding a leukotriene receptor antagonist.

7.5 Specialist therapies

In a small proportion of patients asthma is not adequately controlled on the recommended initial or additional controller therapies (see sections 7.3 and 7.4). There are very few clinical trials in this specific patient group to guide management. For this reason, these patients should be referred for specialist care.



All patients whose asthma is not adequately controlled on recommended initial or additional controller therapies should be referred for specialist care.

7.5.1 Tiotropium bromide

A review of RCTs in adults taking tiotropium bromide, a long-acting muscarinic antagonist (LAMA), in addition to ICS plus LABA compared with ICS plus LABA, reported fewer asthma exacerbations (although results were inconclusive), improved lung function and some benefits relating to asthma control in those taking tiotropium, but no improvement in quality of life. Evidence relating to serious adverse effects was inconclusive but fewer non-serious adverse events were reported in those taking tiotropium. In two of the three trials included in the review patients were taking high-dose ICS.⁴⁸¹ The addition of tiotropium to high-dose ICS plus LABA may confer some additional benefit although results are currently inconclusive. Further research is needed to confirm possible benefits or harms of tiotropium in combination with different doses of ICS/LABA.⁴⁸¹

>12 years	5–12 years	<5 years
1++		

There is insufficient evidence to suggest that addition of tiotropium to ICS in patients inadequately controlled on ICS alone has any benefit over addition of LABA to ICS.⁴⁸² The addition of LABA to ICS remains the first choice for add-on treatment in adults. In adults with asthma who do not respond to ICS plus LABA, the addition of tiotropium to ICS is a possible, although 'off-label' alternative.^{483, 484}

>12 years	5–12 years	<5 years
1++ 1+		

A review comparing the addition of tiotropium to ICS with increased dose of ICS in adults found only one study suitable for inclusion and insufficient evidence to determine if adding tiotropium to ICS ('off-label' use) is safer or more effective than increasing the dose of ICS.⁴⁸⁵

>12 years	5–12 years	<5 years
1+		

7.5.2 Other approaches

Theophyllines may improve lung function and symptoms, but are associated with an increase in adverse events.⁴⁶³

>12 years	5–12 years	<5 years
1+	1-	

Addition of short-acting anticholinergics is generally of no value.^{464, 486} Addition of nedocromil to ICS is of marginal benefit.^{457, 465}

>12 years	5–12 years	<5 years
1+		



If control remains inadequate after stopping a LABA and increasing the dose of inhaled corticosteroid, consider sequential trials of add-on therapy, ie leukotriene receptor antagonists or theophyllines.

The following recommendations are largely based on extrapolation from trials of add-on therapy to ICS alone (*see sections 7.3 and 7.4*).

D	D	<p>If asthma control remains inadequate on medium-dose (adults) or low-dose (children) of inhaled corticosteroid plus a long-acting β_2 agonist or a leukotriene receptor antagonist, the following interventions can be considered:</p> <ul style="list-style-type: none"> • increase the inhaled corticosteroids to high dose (adults)/ medium dose (children 5–12 years)* <i>or</i> • add a leukotriene receptor antagonist (if not already trialed) <i>or</i> • add tiotropium (adults) <i>or</i> • add a theophylline.
----------	----------	---

*at high doses of inhaled corticosteroid via a pMDI, a spacer should be used.

There are no controlled trials indicating which of these is the best option, although the potential for side effects is greater with theophyllines.

- ✓ If a trial of an add-on treatment is ineffective, stop the drug (or in the case of increased dose of inhaled corticosteroid, reduce to the original dose).
- ✓ Although there are no controlled trials, children (all ages) who are under specialist care may benefit from a trial of higher doses ICS (greater than 800 micrograms/day) before moving to use of oral steroids.

7.5.3 Continuous or frequent use of oral steroids

The aim of treatment is to control asthma using the lowest possible doses of medication.

Some patients with very severe asthma not controlled with high-dose ICS, and who have also been tried on or are still taking LABA, LTRA, tiotropium (adults only) or theophyllines, may require regular long-term steroid tablets. These patients should already be under the care of a specialist asthma service.

- ✓ For the small number of patients not controlled on high-dose therapies, use daily steroid tablets in the lowest dose providing adequate control.
- ✓ Patients requiring frequent or continuous use of oral corticosteroids should be under the care of a specialist asthma service.

Patients on long-term steroid tablets (for example, longer than three months) or requiring frequent courses of steroid tablets (for example three to four per year) will be at risk of systemic side effects.⁴⁴⁹ To prevent and treat steroid tablet-induced side effects:

- blood pressure should be monitored
- urine or blood sugar and cholesterol should be checked: diabetes mellitus and hyperlipidaemia may occur
- bone mineral density should be monitored in adults. When a significant reduction occurs, treatment with a long-acting bisphosphonate should be offered. See also, SIGN 142 Management of osteoporosis and the prevention of fragility fractures.⁴⁸⁷

- bone mineral density should be monitored in children >5 ⁴⁸⁸
- growth (height and weight centile) should be monitored in children
- cataracts and glaucoma may be screened for through community optometric services.

Prednisolone is the most widely used steroid for maintenance therapy in patients with chronic asthma. There is no evidence that other steroids offer an advantage.

Although popular in paediatric practice, there are no studies to show whether alternate day steroids produce fewer side effects than daily steroids. No evidence was identified to guide timing of dose or dose splitting.

7.5.4 Monoclonal antibody

Anti-IgE monoclonal antibody

Omalizumab given by subcutaneous injection can reduce the steroid burden for the patient without increasing the risk of adverse events.⁴⁸⁹⁻⁴⁹¹ Three systematic reviews reported reductions in asthma exacerbations in patients with moderate or severe allergic asthma receiving omalizumab compared with placebo in addition to oral corticosteroids or ICS.⁴⁸⁹⁻⁴⁹¹ These studies all reported that more patients on omalizumab compared with placebo withdrew steroids.

>12 years	5-12 years	<5 years
1++ 2++	1++ 2++	

Omalizumab is given as a subcutaneous injection every two or four weeks depending on the patient's IgE level and weight. Local skin reactions may occur. Anaphylaxis, presenting as bronchospasm, hypotension, syncope, urticaria, and/or angioedema of the throat or tongue have been reported after administration of omalizumab occurring as early as the first dose, and as late as one year. Due to concerns about anaphylaxis, the first three doses of omalizumab should only be administered to patients in a healthcare setting under direct medical supervision.

Guidance on when to consider treatment can be found in NICE technology appraisal guidance TA278.⁴⁸⁹

Anti-IL-5 monoclonal antibody

A systematic review of anti-interleukin-5 (IL-5) monoclonal antibody therapies including trials of mepolizumab (four trials; two intravenous, one subcutaneous, one mixed), reslizumab (four trials intravenous) and benralizumab (five trials subcutaneous), and 6,000 patients aged 12 years and over, most of whom had severe eosinophilic asthma, reported reduced asthma exacerbation rates and emergency department/unscheduled care visits with mepolizumab and benralizumab, and reduced asthma exacerbation rates with reslizumab compared with placebo. No serious excess adverse events were reported although significantly more patients receiving benralizumab than placebo discontinued treatment due to adverse events and this requires further investigation.⁴⁹² The review did not look at the potential steroid-sparing effect of anti-IL5 therapies. Use of intravenous mepolizumab is not currently licensed.

>12 years	5-12 years	<5 years
1++		

An RCT of 135 patients with severe eosinophilic asthma receiving 100 mg of mepolizumab subcutaneously or placebo every four weeks, reported a significant glucocorticoid-sparing effect with mepolizumab (28% v 11%, respectively), improved secondary outcomes including fewer exacerbations and improved ACQ-5 scores, and a similar safety profile.⁴⁹³

>12 years	5-12 years	<5 years
1++		

No studies were found that directly compared omalizumab with mepolizumab. A systematic review and meta-analysis, however, concluded that mepolizumab was of equivalent benefit compared to omalizumab in patients eligible for both treatments.⁴⁹⁴ A network meta-analysis comparing omalizumab with mepolizumab showed similar adverse event rates for omalizumab and intravenous mepolizumab (not a licensed route of administration) and a reduction in adverse events compared with placebo and/or baseline therapy (mean annualised asthma exacerbation rate 1.22 v 2.29 omalizumab; 1.28 v 2.56 mepolizumab).⁴⁹⁵

>12 years	5-12 years	<5 years
1+		

Head-to-head trials comparing omalizumab with mepolizumab and other IL-5 therapies and of different IL-5 therapies are needed to confirm the relative clinical and cost effectiveness of each approach.

Guidance on use of mepolizumab, reslizumab and benralizumab differs in England/Wales and Scotland and the relevant NICE or SMC advice should, therefore, be checked prior to considering these treatment approaches.

B	B		Omalizumab given by subcutaneous injection may be considered in eligible patients with a high oral corticosteroid burden.
A			Mepolizumab (subcutaneous), reslizumab (intravenous) and benralizumab (subcutaneous) may be considered in eligible patients with a high oral corticosteroid burden.
✓			<ul style="list-style-type: none"> Patients being considered for monoclonal antibody treatment should be assessed to confirm the diagnosis of asthma, that uncontrolled asthma is the cause of their ongoing symptoms, and that they are adherent with current treatment. An asthma specialist with expertise in monoclonal antibody treatment should assess patients prior to undergoing treatment, and treatment should take place in a specialist centre with the appropriate resources and training, including access to an intensive care unit. Patients undergoing monoclonal antibody treatment should have their details entered onto the UK Severe Asthma Registry.

7.5.5 Other agents

Immunosuppressants (methotrexate, ciclosporin and oral gold) decrease long-term steroid tablet requirements, but all have significant side effects. There is no evidence of persisting beneficial effect after stopping them; and there is marked variability in response.⁴⁹⁶

>12 years	5-12 years	<5 years
1++	3	

✓ Immunosuppressants (methotrexate, ciclosporin and oral gold) may be given as a three-month trial, once other drug treatments have proved unsuccessful. Their risks and benefits should be discussed with the patient and their treatment effects carefully monitored. Treatment should be in a centre with experience of using these medicines.

Colchicine and intravenous immunoglobulin have not been shown to have any beneficial effect in adults.⁴⁹⁶

1+		
----	--	--

Continuous subcutaneous terbutaline infusion has been reported to be beneficial in patients with severe asthma but efficacy and safety have not been assessed in RCTs.⁴⁹⁷⁻⁴⁹⁹

4	3	3
---	---	---

Anti-tumour necrosis factor alpha (anti-TNF alpha) therapy has been investigated in patients with severe asthma but these studies are too small and too short term to allow recommendation of anti-TNF alpha therapy outside the context of a controlled clinical trial.^{500, 501}

7.5.6 Immunotherapy for asthma

Studies using both subcutaneous and sublingual allergen immunotherapy (SCIT and SLIT) have shown some benefit in reducing asthma symptoms and bronchial hyper-reactivity (BHR) in children and adults currently on a range of other preventative strategies including ICS. There are, however, few studies comparing immunotherapy with ICS or of adding immunotherapy to ICS so there is difficulty precisely defining where in asthma management this approach should sit.

Subcutaneous immunotherapy

Trials of allergen-specific immunotherapy by subcutaneous injection of increasing doses of allergen extracts have consistently demonstrated beneficial effects compared with placebo in the management of allergic asthma. Allergens included house dust mite, grass pollen, tree pollen, cat and dog allergen and moulds. Cochrane reviews have concluded that immunotherapy reduces asthma symptoms, the use of asthma medications and improves BHR; the most recent of these included 42 trials with house dust mite, 27 with pollen, 10 with animal allergens, two with *Cladosporium* mould, two with latex and six with multiple allergens.⁵⁰²

The effect of immunotherapy is difficult to quantify due to the use of different symptom scores and variation in the way outcomes are reported. Reductions in asthma medication use and a small symptomatic benefit have been reported but there are significant side effects including 1 in 16 patients reporting a local adverse reaction and 11% reporting a systemic adverse reaction defined as anaphylaxis, asthma, rhinitis, urticaria or a combination of these.⁵⁰² Immunotherapy is not licensed for the treatment of asthma; the current licence is for allergic rhinitis induced by grass pollen.

One study directly compared allergen immunotherapy with ICS and found that symptoms and lung function improved more rapidly in the group on ICS.⁵⁰³

Immunotherapy for allergic rhinitis has been shown to have a carry-over effect after therapy has stopped.⁵⁰⁴

>12 years	5-12 years	<5 years
1++		
1++	1++	
2+		
3		



The use of subcutaneous immunotherapy is not recommended for the treatment of asthma in adults or children.

Sublingual immunotherapy

There has been increasing interest in the use of sublingual immunotherapy (SLIT), which is associated with fewer adverse reactions than subcutaneous immunotherapy.

A systematic review including 52 studies of SLIT in adults and children (n=5,077), most of whom had intermittent or mild asthma symptoms, showed no clear benefit of SLIT. Asthma symptoms scores were the most commonly reported outcome measure (in 42/52 studies) and although overall results were inconclusive, there was some evidence of improvements in asthma symptom scores (in nine studies) and/or medication use (in five studies). Symptom scores and medication use were, however, mostly assessed using unvalidated scales and meta-analysis was not possible. There was no evidence of improvement in lung function,

>12 years	5-12 years	<5 years
1++	1++	

	>12 years	5-12 years	<5 years
quality of life or asthma attacks, although data on these outcomes was limited. Adverse events were significantly more common in those receiving SLIT (absolute increase 327/1,000 SLIT v 222/1,000 control) although these were mostly mild or transient. Serious adverse events were rare with only five of 22 studies reporting any events and no difference between groups in the rate of events (1.3%); all events were thought to be unrelated to treatment. ⁵⁰⁵	1++	1++	

Despite the large volume of evidence evaluating the safety and clinical effectiveness of SLIT in adults and children, heterogeneity in studies (including in doses, allergens, treatment duration, other asthma medication and presence of asthma symptoms), together with the lack of data on its long-term effectiveness and concerns about study quality, mean there is currently insufficient evidence to recommend use of SLIT in adults or children with asthma.

Sublingual immunotherapy is not licensed for use in the treatment of asthma.

B	B	Sublingual immunotherapy is not recommended for the treatment of asthma in children or adults.
----------	----------	---

7.5.7 Bronchial thermoplasty

The aim of bronchial thermoplasty is to reduce bronchial smooth muscle mass, thus reducing the capacity for bronchoconstriction. Currently only a few UK centres offer this treatment which has considerable cost and resource implications.

	>12 years	5-12 years	<5 years
A systematic review of three RCTs (n=429) looking at the use of bronchial thermoplasty for moderate or severe persistent asthma in adults (aged 18 and over) showed a significantly lower rate of severe asthma exacerbations at 12 months in those treated with bronchial thermoplasty in one trial that included a sham intervention in the control group. A second trial, with no sham intervention in the control group, showed a decrease in severe exacerbations in both the intervention and control groups. There were no significant differences in asthma control, lung function, changes in doses of regular medication or use of rescue medication between the intervention and control groups. A small, but statistically significant improvement in quality-of-life scores (measured using AQLQ) with bronchial thermoplasty compared with control groups was seen only in the two studies without a sham intervention. In the study with a sham intervention, QoL scores improved in both groups. ⁵⁰⁶	1++		

Bronchial thermoplasty is an invasive procedure and is associated with an increased rate of adverse respiratory events in the short term. Significantly more patients receiving bronchial thermoplasty than controls were admitted to hospital because of respiratory adverse events within the first 12 weeks following treatment (8 per 100 v 2 per 100; risk ratio (RR) 3.5, 95% CI 1.26 to 9.68). By 12 months following treatment, there was no difference between groups. ⁵⁰⁶	1++		
--	-----	--	--

A systematic review looking at the long-term efficacy and safety of bronchial thermoplasty, including the same three RCTs, reported a significant reduction in respiratory adverse events in patients after five years compared to one year following treatment, although these results were not compared to a control group who had not received bronchial thermoplasty. There was no difference in the number of ED visits or hospitalisations for respiratory adverse events between one and five years of follow up in those treated with bronchial thermoplasty. The longer-term effects of bronchial thermoplasty, beyond five years following treatment, are not known. ⁵⁰⁷	1+		
---	----	--	--

Further research is needed to identify which patients with asthma might benefit from bronchial thermoplasty. However, it is likely that patients who remain uncontrolled despite optimal medical treatment and who have been considered for biological treatments and are either unsuitable for or fail a trial of such a treatment may be an appropriate group, as other treatment options for these patients are elusive. There are no trials comparing the efficacy of bronchial thermoplasty with biological treatments for people with asthma.

B

Bronchial thermoplasty may be considered for the treatment of adult patients (aged 18 and over) with severe asthma who have poorly-controlled asthma despite optimal medical therapy.



- Patients being considered for bronchial thermoplasty should be assessed to confirm the diagnosis of asthma, that uncontrolled asthma is the cause of their ongoing symptoms, and that they are adherent with current treatment.
- An asthma specialist with expertise in bronchial thermoplasty should assess patients prior to undergoing treatment, and treatment should take place in a specialist centre with the appropriate resources and training, including access to an intensive care unit.
- Patients undergoing bronchial thermoplasty should have their details entered onto the UK Severe Asthma Registry.

Figure 2: Summary of management in adults





Figure 3: Summary of management in children



7.6 Decreasing treatment

Decreasing therapy once asthma is controlled is recommended, but often not implemented leaving some patients overtreated. There are few studies that have investigated the most appropriate way to decrease treatment. A study in adults on high-dose ICS has shown that for patients who are stable it is reasonable to attempt to halve the dose of ICS every three months.⁵⁰¹

Some children with milder asthma and a clear seasonal pattern to their symptoms may have a more rapid dose reduction during their 'good' season.

- ✓ Regular review of patients as treatment is decreased is important. When deciding which drug to decrease first and at what rate, the severity of asthma, the side effects of the treatment, time on current dose, the beneficial effect achieved, and the patient's preference should all be taken into account.
- ✓ Patients should be maintained at the lowest possible dose of inhaled corticosteroid. Reduction in inhaled corticosteroid dose should be slow as patients deteriorate at different rates. Reductions should be considered every three months, decreasing the dose by approximately 25–50% each time.

7.7 Specific management issues

7.7.1 Asthma attacks

There is some limited evidence that LTRA may be used intermittently in children with episodic asthma. Treatment is started at the onset of either asthma symptoms or of coryzal symptoms and continued for seven days.⁵⁰⁸

>12 years	5–12 years	<5 years
	1+	1+

7.7.2 Exercise-induced asthma

The following medicines have been shown to give protection against exercise-induced asthma:

- inhaled corticosteroids^{420, 509, 510}
- short-acting β_2 agonists^{416, 511}
- long-acting β_2 agonists⁵¹²
- theophyllines^{513, 514}
- leukotriene receptor antagonists⁵¹⁵
- sodium cromoglicate or nedocromil sodium.⁵¹⁶

>12 years	5–12 years	<5 years
1++	1++	
1++	1++	
1++	1++	
1 ⁻	2 ⁺	
1++	2 ⁺	
1++	2 ⁺	

The following medicines do not give protection against exercise-induced asthma at normal doses:

- anticholinergics⁵¹⁷
- ketotifen⁵¹⁸
- antihistamine.⁵¹⁹

1+	1+
1+	1+
1++	1++

Long-acting β_2 agonists and LTRA provide more prolonged protection than SABA, but a degree of tolerance develops with LABA particularly with respect to duration of action. No tolerance has been demonstrated with LTRA.^{514, 515, 520}

1++	1++
-----	-----

- ✓ For most patients, exercise-induced asthma is an expression of poorly-controlled asthma and regular treatment including inhaled corticosteroids should be reviewed.

If exercise is a specific problem in patients taking inhaled corticosteroids who are otherwise well controlled, consider adding one of the following therapies:

A	C		• leukotriene receptor antagonists
A	A		• long-acting β_2 agonists
C	C		• sodium cromoglicate or nedocromil sodium
C	C		• theophyllines.

Immediately prior to exercise, inhaled SABA are the drug of choice.^{416, 511}

>12 years	5-12 years	<5 years
1++	1++	

A	A		Immediately prior to exercise, inhaled short-acting β_2 agonists are the drug of choice.
---	---	--	--

7.7.3 Comorbid rhinitis

Patients with asthma often have rhinitis. The most effective therapy for rhinitis is intranasal steroids.^{521, 522} Treatment of allergic rhinitis with intranasal steroids has not been shown, in double-blinded placebo-controlled trials, to improve asthma control.

>12 years	5-12 years	<5 years
1+	1+	

7.7.4 Allergic bronchopulmonary aspergillosis

In adult patients with allergic bronchopulmonary aspergillosis, itraconazole may decrease steroid tablet dose and improve asthma control.⁵²³

>12 years	5-12 years	<5 years
2+		

C			In adult patients with allergic bronchopulmonary aspergillosis, a four-month trial of itraconazole should be considered.
---	--	--	---

- ✓ Careful monitoring for side effects, particularly hepatic, is recommended.

7.7.5 Aspirin-intolerant asthma

There are theoretical reasons to suggest that LTRA might be of particular value in the treatment of aspirin-intolerant asthma. However, there is little evidence to justify managing patients with aspirin-intolerant asthma in a different manner to other patients with asthma, apart from the rigorous avoidance of non-steroidal anti-inflammatory medications.^{524, 525}

7.7.6 Comorbid gastro-oesophageal reflux

A Cochrane review of twelve double-blinded controlled trials found that treatment of gastro-oesophageal reflux disease (GORD) had no benefit on asthma symptoms or lung function when both conditions were present. Reduction in dry cough was observed although this was probably not due to improved asthma control.^{526, 527}

A systematic review identified a single RCT which found that proton-pump inhibitors did not improve asthma symptoms in children with GORD.⁵²⁸

	>12 years	5-12 years	<5 years
A further systematic review, including 11 trials and 2,524 patients who had received at least four weeks of daily therapy with proton-pump inhibitors found a small but statistically significant improvement in morning peak expiratory flow (8.86 L/min, 95% CI 2.35 to 15.02) in study participants compared with controls, but no differences in asthma symptom score, Asthma Quality of Life Questionnaire score, evening PEF, FEV ₁ and adverse events. The review concluded that there was insufficient evidence to support the routine use of proton-pump inhibitors in the treatment of asthma. ⁵²⁹	1++		

7.7.7 Beta blockers

Beta blockers, including eye drops, are contraindicated in patients with asthma. Current guidance can be found in the British National Formulary.⁴

8 Inhaler devices

8.1 Technique and training

Studies of technique and the effects of training have used arbitrary non-standardised scores making comparison difficult. Although technique will have some bearing, it does not necessarily relate to clinical effectiveness.

The proportion of patients making no mistakes with an inhaler in one well-conducted study was 23–43% for pMDI, 53–59% for DPI and 55–57% for pMDI + spacer. When technique was assessed as number of steps correct out of the total number of steps, pMDI + spacer was slightly better than DPI.⁵³⁰

Teaching technique improved the correct usage score from a mean of 60% to 79%. Figures for no mistakes after teaching were 63% for pMDI, 65% for DPI, and 75% for breath-actuated MDI (the latter figure based on one study of 2,467 patients).⁵³⁰

>12 years	5–12 years	<5 years
1++		



Prescribe inhalers only after patients have received training in the use of the device and have demonstrated satisfactory technique.

8.2 β_2 agonist delivery

8.2.1 Acute asthma

A pMDI + spacer is at least as good as a nebuliser at treating mild and moderate asthma attacks in children and adults.^{531–534}

>12 years	5–12 years	<5 years
1++	1++	



Children and adults with mild and moderate asthma attacks should be treated with a pMDI + spacer with doses titrated according to clinical response.

There are insufficient data on which to make recommendations in acute severe or life-threatening asthma.

8.2.2 Stable asthma

For children aged 0–5, there is no evidence comparing nebulisers with other inhalers and the data are insufficiently extensive or robust to draw conclusions for pMDI compared with DPI.

In children aged 5–12 there is no significant difference between a pMDI + spacer and a DPI. In adults there is no significant difference between a pMDI ± spacer and a DPI. The lower pulse rate with a pMDI compared with a Turbohaler is the only difference with regard to side effects. Patients have been shown to prefer a Turbohaler to a pMDI.^{530, 535, 536}

>12 years	5–12 years	<5 years
1++	1++	



In children aged 5–12, a pMDI + spacer is as effective as any other hand-held inhaler.

In adults, a pMDI ± spacer is as effective as any other hand-held inhaler, but patients may prefer some types of DPI.

There are no data to make recommendations for children under five years old.

- ✓ Choice of reliever inhaler for stable asthma should be based on patient preference and assessment of correct use. Many patients will not be prepared to carry a spacer.

8.3 Inhaled corticosteroids for stable asthma

No comparative data on ICS for stable asthma in children under five years old were identified.

For the delivery of ICS in children aged 5–12 years with stable asthma, a pMDI is as effective as a Clickhaler.^{537, 538} No significant clinical difference was found between a pMDI and a Turbohaler at half the dose for the same drug (budesonide).^{530, 539} This comparison cannot necessarily be made against other ICS/device combinations.

In adults, there is no clinical difference in effectiveness of a pMDI ± spacer compared to a DPI. A breath-actuated MDI is as effective as a pMDI. More recent DPIs are as effective as older DPIs. Nebulisers have not been shown to be superior to pMDIs + spacer for delivery of ICS in patients with chronic asthma. A specialised specific nebuliser may provide improved lung function and reduced rescue therapy use, but at high prescribed doses. Higher doses (>2 mg) are generally only licensed for use from a nebuliser.^{530, 539}

>12 years	5–12 years	<5 years
1++	1++	



In children aged 5–12 years, a pMDI + spacer is as effective as any DPI.



In adults, a pMDI ± spacer is as effective as any DPI.

No recommendation can be given for nebulised therapy in children aged 5–12 years and there is no evidence relating to children aged under 5 years old.

8.4 Prescribing devices

There is no evidence to dictate an order in which devices should be tested for those patients who cannot use a pMDI. In the absence of evidence, the most important points to consider are patient preference and local cost.

- ✓
 - The choice of device may be determined by the choice of drug.
 - If the patient is unable to use a device satisfactorily an alternative should be found.
 - The patient should have their ability to use the prescribed inhaler device (particularly for any change in device) assessed by a competent healthcare professional (*see section 8.1*).
 - The medication needs to be titrated against clinical response to ensure optimum efficacy.
 - Reassess inhaler technique as part of the structured clinical review (*see section 14.3*).
- ✓ Generic prescribing of inhalers should be avoided as this might lead to people with asthma being given an unfamiliar inhaler device which they are not able to use properly.

- ✓ In young children, a pMDI and spacer is the preferred method of delivery of β_2 agonists and inhaled corticosteroids. A face mask is required until the child can breathe reproducibly using the spacer mouthpiece. Where this is ineffective a nebuliser may be required.

No prospective controlled trials were found that compared use of different devices for preventer and reliever treatments with use of the same device for both treatments. Two cross-sectional studies found an association between increased errors in the use of inhalers when different types of inhaler were used (see section 7.3.4).^{540, 541}

>12 years	5-12 years	<5 years
3		

- ✓ Prescribing mixed inhaler types may cause confusion and lead to increased errors in use. Using the same type of device to deliver preventer and reliever treatments may improve outcomes.

8.5 Use and care of spacers

- ✓
 - The spacer should be compatible with the pMDI being used. A change in spacer may alter effective dose delivered.
 - The drug should be administered by repeated single actuations of the metered dose inhaler into the spacer, each followed by inhalation.
 - There should be minimal delay between pMDI actuation and inhalation.
 - Tidal breathing is as effective as single breaths.
 - Spacers should be cleaned monthly rather than weekly as per manufacturer's recommendations or performance is adversely affected. They should be washed in detergent and allowed to dry in air. The mouthpiece should be wiped clean of detergent before use.
 - Drug delivery via a spacer may vary significantly due to static charge. Metal and other antistatic spacers are not affected in this way.
 - Plastic spacers should be replaced at least every 12 months but some may need changing at 6 months.

8.6 Environmental impact of metered dose inhalers

Metered dose inhalers contain propellants which are liquefied, compressed gases used as a driving force and an energy source for atomisation of the drug. Chlorofluorocarbons (CFCs), which were used originally, are potent greenhouse gases and ozone-depleting substances, and were phased out under the Montreal Protocol. They have been replaced by two hydrofluoroalkane (HFA) propellants: 1,1,1,2-tetrafluoroethane (HFA134a) and 1,1,1,2,3,3,3-heptafluoropropane (HFA227ea), identified as having a high global-warming potential.⁵⁴² As a result of this change, MDIs currently contribute an estimated 3.5% of the carbon footprint of the NHS in the UK.⁵⁴³ The UK has a high proportion of MDI use (70%) compared with the rest of Europe (< 50%) and Scandinavia (10–30%).⁵⁴⁴

- ✓ Prescribers, pharmacists and patients should be aware that there are significant differences in the global-warming potential of different MDIs and that inhalers with low global-warming potential should be used when they are likely to be equally effective. Where there is no alternative to MDIs, lower volume HFA134a inhalers should be used in preference to large volume or HFA227ea inhalers.
- ✓ Patients should be encouraged to ask the pharmacy they use if they can recycle their used inhalers.

9 Management of acute asthma

9.1 Lessons from asthma deaths and near-fatal asthma

Confidential enquires into over 200 asthma deaths in the UK conclude there are factors associated with the disease, the medical management and the patient's behaviour or psychosocial status which contribute to death. Most deaths occurred before admission to hospital.⁵⁴⁵⁻⁵⁴⁹ The report of the UK-wide National Review of Asthma Deaths (NRAD) in 2014 reiterates many of the findings from earlier studies.⁵⁵⁰

9.1.1 Disease factors

Most patients who died of asthma had chronically severe asthma. In a minority the fatal attack occurred suddenly in a patient with mild or moderately severe background disease.^{545-549, 551} **2++**

9.1.2 Medical management

Many of the deaths occurred in patients who had received inadequate treatment with ICS or steroid tablets and/or inadequate objective monitoring of their asthma. Follow up was inadequate in some and others should have been referred earlier for specialist advice. There was widespread underuse of written management plans. Heavy or increasing use of SABA therapy was associated with asthma death.^{545-549, 552, 553} The NRAD report recommended that prescription of more than 12 SABA inhalers a year should prompt review of a patient's management.⁵⁵⁰ **2++**

Deaths continue to be reported following inappropriate prescription of beta blockers and non-steroidal anti-inflammatory drugs; all asthma patients should be asked about past reactions to these agents (*see sections 7.7.7 and 7.7.5*).

Patients with an acute asthma attack should not be sedated unless this is to allow anaesthetic or intensive care procedures (*see section 9.3.12*).⁵⁵¹

The NRAD report highlighted that there is an increased risk of death within one month of discharge from hospital following an acute attack and that follow up in primary care is therefore essential (*see section 9.6*).⁵⁵⁰

9.1.3 Adverse psychosocial and behavioural factors

Behavioural and adverse psychosocial factors were recorded in the majority of patients who died of asthma.⁵⁴⁵⁻⁵⁴⁹ The most important of these are shown in Table 14.

Compared with control patients admitted to hospital with asthma, those who died were significantly more likely to have learning difficulties, psychosis or prescribed antipsychotic drugs, financial or employment problems, repeatedly failed to attend appointments or discharged themselves from hospital, drug or alcohol abuse, obesity or a previous near-fatal attack.^{554, 555} **2++**

Compared with control patients with asthma in the community, patients who died had more severe disease, more likelihood of a hospital admission or visit to the ED for their asthma in the previous year, more likelihood of a previous near-fatal attack, poor medical management, failure to measure pulmonary function, and non-adherence.

B

Healthcare professionals must be aware that patients with severe asthma and one or more adverse psychosocial factors are at risk of death.

Table 14: Patients at risk of developing near-fatal or fatal asthma^{545-549, 552, 553}

A combination of severe asthma recognised by one or more of:
<ul style="list-style-type: none"> • previous near-fatal asthma, eg previous ventilation or respiratory acidosis • previous admission for asthma, especially if in the last year • requiring three or more classes of asthma medication • heavy use of β_2 agonist • repeated attendances at ED for asthma care, especially if in the last year.
AND adverse behavioural or psychosocial features recognised by one or more of:
<ul style="list-style-type: none"> • non-adherence with treatment or monitoring • failure to attend appointments • fewer GP contacts • frequent home visits • self discharge from hospital • psychosis, depression, other psychiatric illness or deliberate self harm • current or recent major tranquilliser use • denial • alcohol or drug abuse • obesity • learning difficulties • employment problems • income problems • social isolation • childhood abuse • severe domestic, marital or legal stress.

Studies comparing near-fatal asthma with deaths from asthma have concluded that patients with near-fatal asthma have identical adverse factors to those described in Table 14, and that these contribute to the near-fatal asthma attack.⁵⁵⁶⁻⁵⁵⁸ Compared with patients who die, those with near-fatal asthma are significantly younger, are significantly more likely to have had a previous near-fatal asthma attack, are more likely to have ready access to acute medical care, and are less likely to have concurrent medical conditions or to experience delay in receiving medical care. 2+

With near-fatal asthma it is advisable to involve a close relative when discussing future management.

Patients with difficult asthma should also be identified (*see section 10.1*).



Keep patients who have had a near-fatal asthma attack under specialist supervision indefinitely.

9.1.4 Seasonal factors

In the UK there is a peak of asthma deaths in July and August in people aged up to 44 years and in December and January in older people.^{556, 559} 2++

9.1.5 Prediction and prevention of a severe asthma attack

Most attacks of asthma severe enough to require hospital admission develop relatively slowly over a period of six hours or more. In one study, over 80% developed over more than 48 hours.⁵⁶⁰⁻⁵⁶⁵ There is therefore time for effective action to reduce the number of attacks requiring hospitalisation. There are many similarities between patients who die from asthma, patients with near-fatal asthma and control patients with asthma who are admitted to hospital. **2++**



A respiratory specialist should follow up patients admitted with a severe asthma attack for at least one year after the admission.

9.2 Acute asthma in adults

Annexes 3-5 contain algorithms summarising the recommended treatment for patients presenting with moderate, acute severe or life-threatening asthma in general practice (*see Annex 3*), the ED (*see Annex 4*), and hospital (*see Annex 5*).

9.2.1 Recognition of acute asthma

Definitions of increasing levels of severity of acute asthma attacks are provided in Table 15.⁵⁶⁶⁻⁵⁷¹ Predicted PEF values should be used only if the recent best PEF (within two years) is unknown.⁵⁷² **2+ 4**

9.2.2 Self treatment by patients developing acute or uncontrolled asthma

Patients with asthma, and all patients with severe asthma, should have an agreed written PAAP and their own peak-flow meter, with regular checks of inhaler technique and adherence. They should know when and how to increase their medication and when to seek medical assistance. Written PAAPs can decrease hospitalisation for,¹⁶⁶ and deaths from asthma (*see section 5.3.2*).⁵⁷³

9.2.3 Initial assessment

All possible initial contact personnel, for example practice receptionists, ambulance call takers, NHS 111 (England and Wales), NHS 24 (Scotland), and out-of-hours providers, should be aware that asthma patients complaining of respiratory symptoms are at risk of becoming seriously unwell very quickly. Such patients should have immediate access to a healthcare professional trained in the emergency treatment of asthma. The assessments required to determine whether the patient is suffering from an acute attack of asthma, the severity of the attack and the nature of treatment required are detailed in Tables 15 and 16. It may be helpful to use a systematic recording process. Proformas have proved useful in the ED setting.⁵⁷⁴

Table 15: Levels of severity of acute asthma attacks in adults⁵⁶⁶⁻⁵⁷¹

Moderate acute asthma	Increasing symptoms PEF >50–75% best or predicted No features of acute severe asthma	
Acute severe asthma	Any one of: - PEF 33–50% best or predicted - respiratory rate $\geq 25/\text{min}$ - heart rate $\geq 110/\text{min}$ - inability to complete sentences in one breath	
Life-threatening asthma	Any one of the following in a patient with severe asthma:	
	Clinical signs	Measurements
	Altered conscious level	PEF <33% best or predicted
	Exhaustion	SpO ₂ <92%
	Arrhythmia	PaO ₂ <8 kPa
	Hypotension	'normal' PaCO ₂ (4.6–6.0 kPa)
	Cyanosis	
	Silent chest	
	Poor respiratory effort	
Near-fatal asthma	Raised PaCO ₂ and/or requiring mechanical ventilation with raised inflation pressures ⁵⁵⁵⁻⁵⁵⁷	

SpO₂: oxygen saturation measured by a pulse oximeter

PaO₂: partial arterial pressure of oxygen

kPa: kilopascals

PaCO₂: partial arterial pressure of carbon dioxide

9.2.4 Prevention of acute deterioration

A register of patients at risk may help healthcare professionals in primary care to identify patients who are more likely to die from their asthma. A system should be in place to ensure that these patients are contacted if they fail to attend for follow up.

9.2.5 Criteria for referral

D Refer to hospital any patients with features of acute severe or life-threatening asthma.

Other factors, such as failure to respond to treatment, social circumstances or concomitant disease, may warrant hospital referral.

Table 16: Initial assessment of symptoms, signs and measurements

Clinical features	<p>Clinical features can identify some patients with severe asthma, eg severe breathlessness (including too breathless to complete sentences in one breath), tachypnoea, tachycardia, silent chest, cyanosis, accessory muscle use, altered consciousness or collapse.^{566-571, 575}</p> <p>None of these singly or together is specific. Their absence does not exclude a severe attack.</p>	2+
PEF or FEV₁	<p>Measurements of airway calibre improve recognition of the degree of severity, the appropriateness or intensity of therapy, and decisions about management in hospital or at home.^{576, 577}</p> <p>PEF or FEV₁ are useful and valid measures of airway calibre. PEF is more convenient in the acute situation. PEF expressed as a percentage of the patient's previous best value is most useful clinically. PEF as a percentage of predicted gives a rough guide in the absence of a known previous best value. Different peak-flow meters give different readings. Where possible the same or similar type of peak-flow meter should be used.</p>	2+
Pulse oximetry	<p>Measure oxygen saturation (SpO₂) with a pulse oximeter to determine the adequacy of oxygen therapy and the need for arterial blood gas measurement. The aim of oxygen therapy is to maintain SpO₂ 94-98%.⁵⁷⁸</p>	
Blood gases	<p>Patients with SpO₂ <92% (irrespective of whether the patient is on air or oxygen) or other features of life-threatening asthma require arterial blood gas measurement.^{566-569, 571, 579} SpO₂ <92% is associated with a risk of hypercapnia. Hypercapnia is not detected by pulse oximetry.⁵⁷⁹ In contrast, the risk of hypercapnia with SpO₂ >92% is much less.⁵⁷⁸</p>	2+ 4
Chest X-ray	<p>Chest X-ray is not routinely recommended in the absence of:</p> <ul style="list-style-type: none"> - suspected pneumomediastinum or pneumothorax - suspected consolidation - life-threatening asthma - failure to respond to treatment satisfactorily - requirement for ventilation. 	4
Systolic paradox	<p>Systolic paradox (<i>pulsus paradoxus</i>) is an inadequate indicator of the severity of an attack and should not be used.^{566-571, 580}</p>	2+

9.2.6 Criteria for admission

Adult patients with any feature of a life-threatening or near-fatal asthma attack or a severe asthma attack that does not resolve after initial treatment should be admitted to hospital. Admission may also be appropriate when peak flow has improved to greater than 75% best or predicted one hour after initial treatment but concerns remain about symptoms, previous history or psychosocial issues (see sections 9.1 and 9.2).^{556, 558, 566-571}

2++
2+

B Admit patients with any feature of a life-threatening or near-fatal asthma attack.

B Admit patients with any feature of a severe asthma attack persisting after initial treatment.

C Patients whose peak flow is greater than 75% best or predicted one hour after initial treatment may be discharged from ED unless they meet any of the following criteria, when admission may be appropriate:

- still have significant symptoms
- concerns about adherence
- living alone/socially isolated
- psychological problems
- physical disability or learning difficulties
- previous near-fatal asthma attack
- asthma attack despite adequate dose of oral corticosteroid prior to presentation
- presentation at night
- pregnancy.

9.3 Treatment of acute asthma in adults

9.3.1 Oxygen

Many patients with acute severe asthma are hypoxaemic.⁵⁸¹⁻⁵⁸⁴ Supplementary oxygen should be given urgently to hypoxaemic patients, using a face mask, Venturi mask or nasal cannulae with flow rates adjusted as necessary to maintain SpO₂ of 94–98%,⁵⁷⁸ taking care to avoid overoxygenation which may be detrimental.⁵⁸⁵

1+
2+
4

Emergency oxygen should be available in hospitals, ambulances and primary care.

Hypercapnia indicates the development of near-fatal asthma and the need for emergency specialist/anaesthetic intervention. In this situation care should be taken to avoid hypoxia as well as overoxygenation.

C Give controlled supplementary oxygen to all hypoxaemic patients with acute severe asthma titrated to maintain an SpO₂ level of 94–98%. Do not delay oxygen administration in the absence of pulse oximetry but commence monitoring of SpO₂ as soon as it becomes available.

9.3.2 β_2 agonist bronchodilators

In most cases inhaled β_2 agonists given in high doses act quickly to relieve bronchospasm with few side effects.⁵⁸⁶⁻⁵⁸⁸ There is no evidence for any difference in efficacy between salbutamol and terbutaline. Nebulised adrenaline (epinephrine), a non-selective β_2 agonist, does not have significant benefit over salbutamol or terbutaline.⁵⁸⁹

1++
1+

In patients with mild to moderate asthma attacks, β_2 agonists can be administered by repeated activations of a pMDI via an appropriate large volume spacer. There are insufficient data on which to make a recommendation about the use of metered dose inhalers with spacers in acute-severe or life-threatening asthma. In such patients, β_2 agonists should be administered by wet nebulisation driven by oxygen, if available.⁵³¹ Inhaled β_2 agonists are as efficacious and preferable to intravenous β_2 agonists (meta-analysis has excluded subcutaneous trials) in adult acute asthma in the majority of cases.⁵⁹⁰ If intravenous β_2 agonists are used, consider monitoring serum lactate.⁵⁹¹

1++
3

A Use high-dose inhaled β_2 agonists as first-line agents in patients with acute asthma and administer as early as possible. Reserve intravenous β_2 agonists for those patients in whom inhaled therapy cannot be used reliably.

✓ If intravenous β_2 agonists are used, consider monitoring serum lactate to monitor for toxicity.

Oxygen-driven nebulisers are preferred for nebulising β_2 agonist bronchodilators because of the risk of oxygen desaturation while using air-driven compressors.^{531, 566, 592}

1++

A flow rate of 6 L/min is required to drive most nebulisers. Where oxygen cylinders are used, a high-flow regulator must be fitted.⁵⁷⁸

4

The absence of supplemental oxygen should not prevent nebulised therapy from being administered when appropriate.⁵⁹³

4

A In hospital, ambulance and primary care, nebulisers for giving β_2 agonist bronchodilators should preferably be driven by oxygen.

✓ In patients with acute asthma with acute-severe or life-threatening features the nebulised route (oxygen-driven) is recommended.

Parenteral β_2 agonists, in addition to inhaled β_2 agonists, may have a role in ventilated patients or those in extremis, however there is limited evidence to support this.

Most acute asthma attacks will respond adequately to bolus nebulisation of β_2 agonists. Continuous nebulisation of β_2 agonists with an appropriate nebuliser may be more effective than bolus nebulisation in relieving acute asthma for patients with a poor response to initial therapy.⁵⁹⁴⁻⁵⁹⁷

1+

A In patients with severe asthma that is poorly responsive to an initial bolus dose of β_2 agonist, consider continuous nebulisation with an appropriate nebuliser.

✓ In patients with acute asthma with acute-severe or life-threatening features the nebulised route (oxygen-driven) is recommended.

Repeat doses of β_2 agonists at 15–30 minute intervals or give continuous nebulisation of salbutamol at 5–10 mg/hour (requires the appropriate nebuliser) if there is an inadequate response to initial treatment. Higher bolus doses, for example 10 mg of salbutamol, are unlikely to be more effective.

9.3.3 Steroid therapy

Steroids reduce mortality, relapses, subsequent hospital admission and requirement for β_2 agonist therapy. The earlier they are given in the acute attack the better the outcome.^{598, 599} **1++**

A Give steroids in adequate doses to all patients with an acute asthma attack.

Steroid tablets are as effective as injected steroids, provided they can be swallowed and retained.⁵⁹⁸ Prednisolone 40–50 mg daily or parenteral hydrocortisone 400 mg daily (100 mg six hourly) are as effective as higher doses.⁶⁰⁰ Where necessary soluble prednisolone (sodium phosphate) 5 mg tablets are available. In cases where oral treatment may be a problem consider intramuscular methylprednisolone (160 mg) as an alternative to a course of oral prednisolone.⁶⁰¹ **1++**

✓ Continue prednisolone (40–50 mg daily) until recovery (minimum 5 days).

Following recovery from the acute asthma attack steroids can be stopped abruptly. Doses do not need tapering provided the patient receives ICS (apart from patients on maintenance steroid treatment or rare instances where steroids are required for three or more weeks).^{602, 603} **1+**

It is not known if ICS provide further benefit in addition to systemic steroids.^{604, 605} **1+**

✓ Do not stop inhaled corticosteroids during prescription of oral corticosteroids.

9.3.4 Ipratropium bromide

Combining nebulised ipratropium bromide with a nebulised β_2 agonist produces significantly greater bronchodilation than β_2 agonist alone, leading to faster recovery and shorter duration of admission. Anticholinergic treatment is not necessary and may not be beneficial in milder asthma attacks or after stabilisation.^{606–608} **1++**

B Add nebulised ipratropium bromide (0.5 mg 4–6 hourly) to β_2 agonist treatment for patients with acute severe or life-threatening asthma or those with a poor initial response to β_2 agonist therapy.

9.3.5 Magnesium sulphate

A systematic review of 25 RCTs (13 including adults) involving 2,907 patients with asthma showed that nebulised magnesium sulphate when used in addition to nebulised β_2 agonist (with or without nebulised ipratropium) provided no benefit in terms of lung function or need for hospital admission.⁶⁰⁹ Subgroup analysis of the most severe patients was not possible due to heterogeneity in studies and the use of multiple different end-points. Some smaller studies noted modest improvements in lung function with nebulised magnesium in the most severe subgroup (presenting $FEV_1 < 50\%$), but the results were not significant. **1++**

A double-blinded, placebo-controlled study of 1,109 patients aged over 16 years presenting with an acute asthma attack to 34 emergency departments across the UK randomised patients to intravenous or nebulised magnesium or to placebo.⁶¹⁰ Many of these patients had $PEF > 50\%$ at presentation and the study failed to show improvement in either rate of hospital admission or breathlessness as judged by a visual analogue score. A single dose of intravenous magnesium sulphate is safe **1++**
1+

and may improve lung function and reduce intubation rates in patients with acute severe asthma.^{355, 611-613} Intravenous magnesium sulphate may also reduce hospital admissions in adults with acute asthma who have had little or no response to standard treatment. However, the heterogeneous nature of the studies included in this review and lack of information on the severity of the asthma attack or when intravenous magnesium was given in relation to standard treatment limit the conclusions that can be drawn.⁶¹³

1++
1+

The safety and efficacy of repeated intravenous (IV) doses of magnesium sulphate have not been assessed. Repeated doses could cause hypermagnesaemia with muscle weakness and respiratory fatigue.

A Nebulised magnesium sulphate is not recommended for treatment in adults with acute asthma.

B Consider giving a single dose of intravenous magnesium sulphate to patients with acute severe asthma (PEF <50% best or predicted) who have not had a good initial response to inhaled bronchodilator therapy.

✓ Magnesium sulphate (1.2–2 g IV infusion over 20 minutes) should only be used following consultation with senior medical staff.

9.3.6 Intravenous aminophylline

In an acute asthma attack, IV aminophylline is not likely to result in any additional bronchodilation compared with standard care with inhaled bronchodilators and steroids. Side effects such as arrhythmias and vomiting are increased if IV aminophylline is used.⁶¹⁴

1++

✓ Use IV aminophylline only after consultation with senior medical staff.

Some patients with near-fatal asthma or life-threatening asthma with a poor response to initial therapy may gain additional benefit from IV aminophylline (5 mg/kg loading dose over 20 minutes unless on maintenance oral therapy, then infusion of 0.5–0.7 mg/kg/hr). Such patients are probably rare and were not identified in a meta-analysis of trials.⁶¹⁴ If IV aminophylline is given to patients already taking oral aminophylline or theophylline, blood levels should be checked on admission. Levels should be checked daily for all patients on aminophylline infusions.

9.3.7 Leukotriene receptor antagonists

Current evidence on oral leukotriene receptor antagonists does not support their use in patients with acute asthma.⁶¹⁵ Further studies are required to assess whether IV treatment is effective and safe.

1++

9.3.8 Antibiotics

When an infection precipitates an asthma attack it is likely to be viral. The role of bacterial infection has been overestimated.⁶¹⁶ Decision making regarding the use of antibiotics in patients with acute asthma should be guided by objective measures including procalcitonin where available.^{617, 618}

1++
1+

B Routine prescription of antibiotics is not indicated for patients with acute asthma.

9.3.9 Heliox

The use of heliox, (helium/oxygen mixture in a ratio of 80:20 or 70:30), either as a driving gas for nebulisers, as a breathing gas, or for artificial ventilation in adults with acute asthma is not supported.^{619, 620} A systematic review of ten trials, including 544 patients with acute asthma, found no improvement in pulmonary function or other outcomes in adults treated with heliox, although the possibility of benefit in patients with more severe obstruction exists.^{621, 622} Heliox requires the use of specifically designed or modified breathing circuits and ventilators.

1++
1+

B Heliox is not recommended for use in patients with acute asthma outside a clinical trial setting.

9.3.10 Intravenous fluids

There are no controlled trials, observational or cohort studies of differing fluid regimes in patients with acute asthma. Some patients require rehydration and correction of electrolyte imbalance. Hypokalaemia can be caused or exacerbated by β_2 agonist and/or steroid treatment and must be corrected.

9.3.11 Nebulised furosemide

Although theoretically furosemide may produce bronchodilation, a review of three small trials failed to show any significant benefit of treatment with nebulised furosemide compared to β_2 agonists.⁶²³

1+

9.3.12 Critical care settings

In adults with acute asthma and a poor response to standard therapy (inhaled bronchodilators, steroids, oxygen and intravenous bronchodilators) other therapies may be considered in the appropriate critical care setting with the appropriate available expertise. There is little high-quality evidence to guide treatment at this stage of an acute asthma attack and it is important to involve a clinician with the appropriate skills in airway management and critical care support as early as possible.

Indications for admission to intensive care or high-dependency units include patients requiring ventilatory support and those with acute severe or life-threatening asthma who are failing to respond to therapy, as evidenced by:^{566, 567}

- deteriorating PEF
- persisting or worsening hypoxia
- hypercapnia
- arterial blood gas analysis showing fall in pH or rising hydrogen concentration
- exhaustion, feeble respiration
- drowsiness, confusion, altered conscious state
- respiratory arrest.

Ketamine

A review (including 12 case reports, three RCTs and five other observational studies) of ketamine use in adults and children in *status asthmaticus* reported that ketamine is a potential bronchodilator but that prospective trials are needed before conclusions about effectiveness can be drawn.⁶²⁴

2-

Extracorporeal membrane oxygenation

Extracorporeal membrane oxygenation (ECMO) is thought to help to provide adequate gas exchange whilst helping to prevent the barotraumas caused by aggressive mechanical ventilation. Currently, there are five centres in the UK with ECMO facilities for adults (Glenfield Hospital, Leicester; Papworth Hospital, Cambridge; Wythenshawe Hospital, Manchester; Guy's & St Thomas' Hospital, London; Royal Brompton & Harefield Hospital, London).

An international retrospective registry study of 272 adult patients with near-fatal asthma most of whom were put on venovenous ECMO showed a survival rate to hospital discharge of 83.5%. The rate of in-hospital complications was high (65.1%), the most common of which was haemorrhage (28.3%), most commonly at a manageable cannulation site (13.1%); only 1.5% died as a result of the haemorrhage. Other complications were renal (26.8%), cardiovascular (26.1%), mechanical (24.6%), metabolic (22.4%), infection (16.5%), neurologic (4.8%), and limb ischemia (2.6%). The most common cause of death was organ failure (37.8%, 17/45 complications). Long-term complications of ECMO were not considered.⁶²⁵

3

Although it is unclear which patients would benefit the most from venovenous ECMO, survivors were younger (34.7 ν 43.4, $p=0.001$), had a lower mean pH (7.1 ν 7.2, $p=0.045$), higher oxygen saturation (92.3 ν 85.2, $p=0.03$) and lower positive end-expiratory pressure (7.8 ν 11.5, $p=0.002$) than those who died.

Limitations of the registry include the lack of selection criteria for inclusion, and consequent lack of clarity about whether patients were on optimal or even similar ventilator settings, and the voluntary nature of reporting of cases which may lead to reporting bias. Despite these limitations, the use of ECMO provides a potential rescue therapy in patients with near-fatal asthma refractory to conventional ventilator treatment.

D Where available, extracorporeal membrane oxygenation may be considered in adults with near-fatal asthma refractory to conventional ventilator treatment.

Recombinant human deoxyribonuclease

A pilot RCT of the use of recombinant human deoxyribonuclease in severely ill, non-intubated adults with asthma refractory to bronchodilators reported no benefit from its use in this patient group.⁶²⁶

1+

✓ Adults with asthma not responding to standard treatment should be evaluated by a specialist with the appropriate experience and skills to use and assess medication encountered in critical care settings.

✓ In patients with acute severe or life-threatening asthma, anaesthetists and intensivists should be notified as soon as possible if there is no improvement in or deterioration of asthma.

Not all patients admitted to the intensive care unit (ICU) need ventilation, but those with worsening hypoxia or hypercapnia, drowsiness or unconsciousness and those who have had a respiratory arrest require intermittent positive pressure ventilation. Intubation in such patients is very difficult and should be performed by an anaesthetist or ICU consultant.^{566, 567}

2+

C All patients transferred to intensive care units should be accompanied by a doctor suitably equipped and skilled to intubate if necessary.

9.3.13 Non-invasive ventilation

Non-invasive ventilation (NIV) is well established in the management of ventilatory failure caused by extrapulmonary restrictive conditions and exacerbations of COPD. Hypercapnic respiratory failure developing during an acute asthmatic attack is an indication for urgent ICU admission. It is unlikely that NIV would replace intubation in these very unstable patients but it has been suggested that this treatment can be used safely and effectively.⁶²⁷

4

Evidence to support the use of NIV in adults is limited and inconclusive. A Cochrane review found only one trial of NIV, with 30 patients, which showed improvement in hospitalisation rates, discharge from emergency departments and lung function.⁶²⁸ Two further small studies suggest that NIV may be safe and feasible in treating patients with severe asthma exacerbations but provide little evidence of benefit compared with standard care.^{629, 630}

1++
1+
2-

Larger RCTs are needed to determine the role of NIV in treating patients with acute asthma.⁶²⁸ Future trials should include measurable clinical outcomes such as respiratory parameters, physiological variables and blood gases.



NIV should only be considered in an ICU or equivalent clinical setting.

9.4 Further investigation and monitoring



- Measure and record PEF 15–30 minutes after starting treatment, and thereafter according to the response. Measure and record PEF before and after nebulised or inhaled β_2 agonist.
- Record oxygen saturation by oximetry and maintain arterial SpO_2 at 94–98%.
- Repeat measurements of blood gas tensions within one hour of starting treatment if:
 - the initial PaO_2 is <8 kPa unless SpO_2 is $>92\%$; or
 - the initial PaCO_2 is normal or raised; or
 - the patient's condition deteriorates.



- Measure them again if the patient's condition has not improved by 4–6 hours.
- Measure and record the heart rate.
- Measure serum potassium and blood glucose concentrations.
- Measure the serum theophylline concentration if aminophylline is continued for more than 24 hours (aim for a concentration of 10–20 mg/L or 55–110 mol/L).

9.5 Asthma management protocols and proformas

The use of structured proformas facilitates improvements in the process of care in emergency departments and hospital wards and improves patient outcomes. The use of this type of documentation can assist data collection aimed at determining the quality of care and outcomes.^{574, 631, 632}

2++

9.6 Hospital discharge and follow up

Annex 5 summarises management of acute asthma in hospital.

An asthma care bundle developed by the BTS is also available from the BTS website (www.brit-thoracic.org.uk).

9.6.1 Timing of discharge

No single physiological parameter defines absolutely the timing of discharge from an admission with acute asthma. Patients should have clinical signs compatible with home management, be on reducing amounts of β_2 agonist (preferably no more than four hourly) and be on medical therapy they can continue safely at home.

Although diurnal variability of PEF is not always present during an asthma attack, evidence suggests that patients discharged with PEF <75% best or predicted and with diurnal variability >25% are at greater risk of early relapse and readmission.^{633, 634}

2+

9.6.2 Patient education

Following discharge from hospital or emergency departments, a proportion of patients reattend with more than 15% reattending within two weeks. Some repeat attenders need emergency care, but many delay seeking help, and are undertreated and/or undermonitored.⁶³⁵

2+

Prior to discharge, trained staff should give asthma education. This should include education on inhaler technique and PEF record keeping, with a written PEF and symptom-based PAAP being provided allowing the patient to adjust their therapy within recommendations. These measures have been shown to reduce morbidity after the asthma attack and reduce relapse rates.⁶³⁶

1++

Some patients may use emergency departments rather than primary care services for their asthma care. Education has been shown to reduce subsequent hospital admission and improve scheduled appointments and self-management techniques but does not improve reattendance at emergency departments.¹⁸⁷

1++

For the above groups there is a role for a trained asthma liaison nurse based in, or associated with, the emergency department.¹⁸⁷

Patient education is covered in section 5.2.1

9.6.3 Follow up

A careful history should elicit the reasons for the asthma attack and explore possible actions the patient should take to prevent future emergency presentations.

Medication should be altered depending upon the assessment and the patient provided with an asthma action plan aimed at preventing relapse, optimising treatment and preventing delay in seeking assistance in the future.

Prior to discharge, follow up should be arranged with the patient's general practitioner or asthma nurse within two working days and with a hospital specialist asthma nurse or respiratory physician at about one month after admission.

In a small RCT, follow-up care by a nurse specialist was as effective and safe as that given by a respiratory doctor.⁶³⁷ 1+

Assisting patients in making appointments while being treated for an acute asthma attack in emergency departments may improve subsequent attendance at primary care centres.⁶³⁸ 1+



It is essential that the patient's primary care practice is informed within 24 hours of discharge from the emergency department or hospital following an asthma attack. Ideally this communication should be directly with a named individual responsible for asthma care within the practice.

9.7 Acute asthma in children

The assessment of acute asthma in children under five can be difficult. Intermittent wheezing attacks are usually triggered by viral infection and the response to asthma medication may be inconsistent. Prematurity and low birth weight are risk factors for recurrent wheezing. The differential diagnosis of symptoms includes aspiration pneumonitis, pneumonia, bronchiolitis, tracheomalacia, and complications of underlying conditions such as congenital anomalies and cystic fibrosis. This guideline is intended for children who are thought to have acute wheeze related to underlying asthma and should be used with caution in younger children who do yet have a considered diagnosis of asthma, particularly those under two years of age. The guideline is not intended for children under one year of age unless directed by a respiratory paediatrician. The guideline should not be used to treat acute bronchiolitis.

9.7.1 Clinical assessment

Table 17 details criteria for assessment of severity of acute asthma attacks in children.

Annexes 6–9 contain algorithms summarising the recommended treatments for children presenting with acute or uncontrolled asthma in general practice (*see Annex 6*), the ED (*see Annex 7*), and hospital (*see Annexes 8 and 9*).

Table 17: Levels of severity of acute asthma attacks in children⁶³⁹

Moderate acute asthma	Able to talk in sentences $\text{SpO}_2 \geq 92\%$ $\text{PEF} \geq 50\%$ best or predicted Heart rate $\leq 140/\text{min}$ in children aged 1–5 years $\leq 125/\text{min}$ in children >5 years Respiratory rate $\leq 40/\text{min}$ in children aged 1–5 years $\leq 30/\text{min}$ in children >5 years	
Acute severe asthma	Can't complete sentences in one breath or too breathless to talk or feed $\text{SpO}_2 < 92\%$ PEF 33–50% best or predicted Heart rate $> 140/\text{min}$ in children aged 1–5 years $> 125/\text{min}$ in children >5 years Respiratory rate $> 40/\text{min}$ in children aged 1–5 years $> 30/\text{min}$ in children >5 years	
Life-threatening asthma	Any one of the following in a child with severe asthma:	
	Clinical signs	Measurements
	Exhaustion	$\text{PEF} < 33\%$ best or predicted
	Hypotension	$\text{SpO}_2 < 92\%$
	Cyanosis	
	Silent chest	
	Poor respiratory effort	
	Confusion	

Before children can receive appropriate treatment for an acute asthma attack in any setting, it is essential to assess accurately the severity of their symptoms. The following clinical signs should be recorded:

- Pulse rate
 - increasing tachycardia generally denotes worsening asthma; a fall in heart rate in life-threatening asthma is a preterminal event
- Respiratory rate and degree of breathlessness
 - ie too breathless to complete sentences in one breath or to feed
- Use of accessory muscles of respiration
 - best noted by palpation of neck muscles
- Amount of wheezing
 - which might become biphasic or less apparent with increasing airways obstruction
- Degree of agitation and conscious level
 - always give calm reassurance.

Clinical signs correlate poorly with the severity of airways obstruction.⁶⁴⁰⁻⁶⁴³ Some children with acute severe asthma do not appear distressed. 2++

- ✓ Decisions about admission should be made by trained clinicians after repeated assessment of the response to bronchodilator treatment.

9.7.2 Pulse oximetry

Accurate measurements of oxygen saturation are essential in the assessment of all children with acute wheezing. Oxygen saturation monitors should be available for use by all healthcare professionals assessing acute asthma in both primary and secondary care settings.

Low oxygen saturations after initial bronchodilator treatment selects a group of patients with more severe asthma.^{640, 643} 2++

- B Consider intensive inpatient treatment of children with SpO₂ <92% in air after initial bronchodilator treatment.**

9.7.3 Peak expiratory flow

PEF measurements can be of benefit in assessing children who are familiar with the use of such devices. The best of three PEF measurements, ideally expressed as a percentage of personal best, can be useful in assessing the response to treatment.

A measurement of <50% predicted PEF or FEV₁ with poor improvement after initial bronchodilator treatment is predictive of a more prolonged asthma attack.

9.7.4 Chest X-ray

Chest X-rays rarely provide additional useful information and are not routinely indicated.^{644, 645}

- ✓ A chest X-ray should be performed if there is subcutaneous emphysema, persisting unilateral signs suggesting pneumothorax, lobar collapse or consolidation and/or life-threatening asthma not responding to treatment.

9.7.5 Blood gases

Blood gas measurements should be considered if there are life-threatening features not responding to treatment. Arteriolised ear lobe blood gases can be used to obtain an accurate measure of pH and PaCO₂.⁵⁷⁸ If ear lobe sampling is not practicable a finger prick sample can be an alternative. Normal or raised PaCO₂ levels are indicative of worsening asthma. A more easily obtained free flowing venous blood PaCO₂ measurement of <6 kPa (45 millimetres of mercury) excludes hypercapnia.⁵⁷⁸ 4

9.8 Initial treatment of acute asthma in children

There is good evidence supporting recommendations for the initial treatment of children with acute asthma presenting to primary and secondary healthcare centres. There is less evidence to guide the use of second-line therapies to treat the small number of severe cases of acute asthma poorly responsive to first-line measures. Despite this, the risks of death and other adverse outcomes after admission to hospital are extremely low irrespective of the treatment options chosen.

Emergency departments attending to children with acute asthma should have a nurse trained in paediatrics available on duty at all times and staff familiar with the specific needs of children. Using a proforma can increase the accuracy of severity assessment.

The use of an assessment-driven algorithm and an integrated care pathway has been shown to reduce hospital stay without substantial increases in treatment costs.⁶⁴⁶

4

D The use of structured care protocols detailing bronchodilator usage, clinical assessment, and specific criteria for safe discharge is recommended.

9.8.1 Oxygen

✓ Children with life-threatening asthma or SpO₂ <94% should receive high-flow oxygen via a tight-fitting face mask or nasal cannula at sufficient flow rates to achieve normal saturations of 94–98%.

9.8.2 Inhaled short-acting β_2 agonists

Inhaled β_2 agonists are the first-line treatment for acute asthma in children aged two years and over.⁶⁴⁷⁻⁶⁵⁰ Assessment of response should be based on accurately recorded clinical observations and repeat measurements of oxygenation (SpO₂) (see Table 17). Children receiving β_2 agonists via a pMDI + spacer are less likely to have tachycardia and hypoxia than when the same drug is given via a nebuliser.⁵³¹ In children under two who have a poor initial response to β_2 agonists administered with adequate technique, consider an alternative diagnosis and other treatment options.

1+

A Inhaled β_2 agonists are the first-line treatment for acute asthma in children.

✓ Discontinue long-acting β_2 agonists when short-acting β_2 agonists are required more often than four hourly.

A A pMDI + spacer is the preferred option for children with mild to moderate asthma.

Children under three years of age are likely to require a face mask connected to the mouthpiece of a spacer for successful drug delivery. Inhalers should be actuated into the spacer in individual puffs and inhaled immediately by tidal breathing (for five breaths).

Frequent doses of β_2 agonists are safe for the treatment of acute asthma,⁶⁴⁷⁻⁶⁴⁹ although children with mild symptoms benefit from lower doses.⁶⁵⁰

1+

B Individualise drug dosing according to severity and adjust according to the patient's response.

Two to four puffs of salbutamol (100 micrograms via a pMDI + spacer) might be sufficient for mild asthma attacks, although up to 10 puffs might be needed for more severe attacks. Single puffs should be given one at a time and inhaled separately with five tidal breaths. Relief from symptoms should last 3–4 hours. If symptoms return within this time a further or larger dose (maximum 10 puffs) should be given and the parents/carer should seek urgent medical advice.

Children with severe or life-threatening asthma ($\text{SpO}_2 < 92\%$) should receive frequent doses of nebulised bronchodilators driven by oxygen (2.5–5 mg salbutamol). If there is poor response to the initial dose of β_2 agonists, subsequent doses should be given in combination with nebulised ipratropium bromide (see section 9.8.3). Doses of nebulised bronchodilator can be repeated every 20–30 minutes. Continuous nebulised β_2 agonists are of no greater benefit than the use of frequent intermittent doses in the same total hourly dosage.^{651, 652} Once improving on two- to four-hourly salbutamol, patients should be switched to a pMDI and spacer treatment as tolerated.

Schools can hold a generic reliever inhaler enabling them to treat an acutely wheezy child whilst awaiting medical advice. This is safe and potentially life saving.

- ✓ Increase β_2 agonist dose by giving one puff every 30–60 seconds, according to response, up to a maximum of ten puffs.
- ✓ Parents/carers of children with an acute asthma attack at home, and symptoms not controlled by up to 10 puffs of salbutamol via a pMDI and spacer, should seek urgent medical attention.
- ✓ If symptoms are severe additional doses of bronchodilator should be given as needed whilst awaiting medical attention.
- ✓ Paramedics attending to children with an acute asthma attack should administer nebulised salbutamol, using a nebuliser driven by oxygen if symptoms are severe, whilst transferring the child to the emergency department.
- ✓ Children with severe or life-threatening asthma should be transferred to hospital urgently.

9.8.3 Ipratropium bromide

There is good evidence for the safety and efficacy of frequent doses of ipratropium bromide (every 20–30 minutes) used in addition to β_2 agonists for the first two hours of a severe asthma attack. Benefits are more apparent in the most severe patients.⁶⁵³

1+

A If symptoms are refractory to initial β_2 agonist treatment, add ipratropium bromide (250 micrograms/dose mixed with the nebulised β_2 agonist solution).

Frequent doses up to every 20–30 minutes (250 micrograms/dose mixed with 5 mg of salbutamol solution in the same nebuliser) should be used for the first few hours of admission. Salbutamol dose should be tapered to one to two hourly thereafter according to clinical response. The ipratropium dose should be tapered to four to six hourly or discontinued.

- ✓ Repeated doses of ipratropium bromide should be given early to treat children who are poorly responsive to β_2 agonists.

9.8.4 Steroid therapy

The early use of steroids in emergency departments and assessment units can reduce the need for hospital admission and prevent a relapse in symptoms after initial presentation.^{598, 599} Benefits can be apparent within three to four hours. In head-to-head comparisons there is insufficient evidence to suggest that dexamethasone offers an advantage over prednisolone for the management of mild to moderate acute asthma in children. Further studies may indicate whether a single dose of dexamethasone may offer clinical benefit over multiple doses of prednisolone.⁶⁵⁴⁻⁶⁵⁶

1+
1-

A large UK study of preschool children with mild to moderate wheeze associated with viral infection showed no reduction in hospital stay (or other outcomes) following treatment with oral steroids. In the acute situation, it is often difficult to determine whether a preschool child has asthma or episodic viral wheeze. Children with severe symptoms requiring hospital admission should still receive oral steroids. In children who present with moderate to severe wheeze without a previous diagnosis of asthma it is still advisable to give oral steroids. For children with frequent episodes of wheeze associated with viruses caution should be taken in prescribing multiple courses of oral steroids.⁶⁵⁷

1++

A Give oral steroids early in the treatment of acute asthma attacks in children.

B Oral prednisolone is the steroid of choice for asthma attacks in children unless the patient is unable to tolerate the dose.

Use a dose of 10 mg of prednisolone for children under two years of age, a dose of 20 mg for children aged 2–5 years and a dose of 30–40 mg for children older than five years.

Oral and intravenous steroids are of similar efficacy.^{600, 658, 659} Intravenous hydrocortisone (4 mg/kg repeated four hourly) should be reserved for severely affected children who are unable to retain oral medication.

1+

Larger doses do not appear to offer a therapeutic advantage for the majority of children.⁶⁶⁰ There is no need to taper the dose of steroid tablets at the end of treatment.^{602, 603}

2+



- Use a dose of 10 mg prednisolone for children under two years of age, 20 mg for children aged 2–5 years and 30–40 mg for children older than five years. Those already receiving maintenance steroid tablets should receive 2 mg/kg prednisolone up to a maximum dose of 60 mg.
- Repeat the dose of prednisolone in children who vomit and consider intravenous steroids in those who are unable to retain orally ingested medication.
- Treatment for up to three days is usually sufficient, but the length of course should be tailored to the number of days necessary to bring about recovery. Tapering is unnecessary unless the course of steroids exceeds 14 days.

Inhaled corticosteroids

There is insufficient evidence to support the use of ICS as alternative or additional treatment to steroid tablets for children with acute asthma.^{604, 661-668}

1++
1+
1-

A Do not use inhaled corticosteroids in place of oral steroids to treat children with an acute asthma attack.

Children with chronic asthma not receiving regular preventative treatment will benefit from starting ICS as part of their long-term management. There is no evidence that increasing the dose of ICS is effective in treating acute symptoms, but it is good practice for children already receiving ICS to continue with their usual maintenance doses.

✓ It is good practice for children already receiving inhaled corticosteroids to continue with their usual maintenance dose during an asthma attack whilst receiving additional treatment.

9.8.5 Antibiotics

There is insufficient evidence to support or refute the role of antibiotics in acute asthma, but the majority of acute asthma attacks are triggered by viral infection.⁴⁵⁸

✓ Do not give antibiotics routinely in the management of children with acute asthma.

9.8.6 Leukotriene receptor antagonists

Initiating oral montelukast in primary care settings, early after the onset of an acute asthma attack, can result in decreased asthma symptoms and the need for subsequent healthcare attendances in those with mild asthma attacks.^{508, 669} Current evidence shows no benefit for the addition of leukotriene receptor antagonists to standard asthma treatment for moderate to severe asthma attacks.⁶¹⁵

1++
1+

9.8.7 Nebulised magnesium sulphate

There is no evidence to support the use of nebulised magnesium sulphate, either in place of or in conjunction with inhaled β_2 agonists, in children with mild to moderate asthma.⁶⁰⁹ A subgroup analysis from a large RCT suggests a possible role in children with more severe asthma attacks ($\text{SpO}_2 < 92\%$) or with short duration of deterioration. Further studies are required to evaluate which clinical groups would benefit the most from this intervention.⁶⁷⁰

1++

A Nebulised magnesium sulphate is not recommended for children with mild to moderate asthma attacks.

C Consider adding 150 mg magnesium sulphate to each nebulised salbutamol and ipratropium in the first hour in children with a short duration of acute severe asthma symptoms presenting with an $\text{SpO}_2 < 92\%$.

9.9 Second-line treatment of acute asthma in children

Children with continuing severe asthma despite optimal first-line treatments, frequent nebulised β_2 agonists and ipratropium bromide plus oral steroids, and those with life-threatening features, need urgent review by a specialist with a view to management in an appropriate high-dependency area or transfer to a paediatric intensive care unit to receive second-line intravenous therapies.

Three options, IV magnesium sulphate, IV β_2 agonist or IV aminophylline can be considered. In one RCT comparing all three agents in 100 children, a bolus of magnesium sulphate was shown to reduce clinical symptoms faster than the other treatments. There were no significant side effects documented in the magnesium sulphate group.⁶⁷¹ A systematic review of four paediatric trials comparing IV salbutamol with IV aminophylline demonstrated equivalence. One study found a shorter length of stay in the aminophylline group although these patients received a bolus followed by an infusion, compared to a single bolus of IV salbutamol. Both IV salbutamol and IV aminophylline can cause side effects and should be administered with appropriate monitoring.⁶⁷²

1++
1+

9.9.1 Intravenous salbutamol

The role of intravenous β_2 agonists in addition to nebulised treatment remains unclear.⁵⁹⁰ One study has shown that an IV bolus of salbutamol given in addition to near-maximal doses of nebulised salbutamol results in clinically significant benefits for those with moderate to severe asthma.⁵⁹⁰

1+

B Consider early addition of a single bolus dose of intravenous salbutamol (15 micrograms/kg over 10 minutes) in a severe asthma attack where the child has not responded to initial inhaled therapy.

A continuous intravenous infusion of salbutamol should be considered when there is uncertainty about reliable inhalation or for severe refractory asthma. This should be given in a high dependency unit with continuous electrocardiogram (ECG) monitoring and twice daily electrolyte monitoring. Doses above 1–2 micrograms/kg/min (200 micrograms/ml solution) should be given in a paediatric intensive care unit setting (up to 5 micrograms/kg/min). Nebulised bronchodilators should be continued while the patient is receiving intravenous bronchodilators. Once the patient is improving the intravenous infusion should be reduced before reducing the frequency of nebulised bronchodilators.

✓ When inserting an IV cannula take a blood sample to measure serum electrolytes. Serum potassium levels are often low after multiple doses of β_2 agonists and should be replaced.

✓ If intravenous β_2 agonist infusions are used, consider monitoring serum lactate to monitor for toxicity.

9.9.2 Intravenous aminophylline

There is no evidence that aminophylline is of benefit for mild to moderate asthma and side effects are common and troublesome.^{612, 614, 673, 674} One well-conducted study has shown evidence of benefit in children with acute severe asthma unresponsive to multiple doses of β_2 agonists and steroids, although the loading dose used was double that currently recommended in the UK and a third of patients were withdrawn from active medication because of vomiting.⁶⁷⁵

1+
2+

A Aminophylline is not recommended in children with mild to moderate acute asthma.

B Consider aminophylline for children with severe or life-threatening asthma unresponsive to maximal doses of bronchodilators and steroids.

A 5 mg/kg loading dose should be given over 20 minutes (omit in those receiving maintenance oral theophyllines) with ECG monitoring followed by a continuous infusion at 1 mg/kg/hour. Measure serum theophylline levels in patients already receiving oral treatment and in those receiving prolonged treatment.

9.9.3 Intravenous magnesium sulphate

Intravenous magnesium sulphate is a safe treatment for acute asthma in children not responding to first-line treatment.⁶⁷⁶ Doses of up to 75 mg/kg/day (maximum 2 g) have been used. One additional trial (n=34 receiving magnesium sulphate) reported that the potential side effect of hypotension with a single dose of IV magnesium sulphate is rare.⁶⁷¹

1++
1+

B In children who respond poorly to first-line treatments, consider the addition of intravenous magnesium sulphate as first-line intravenous treatment (40 mg/kg/day).

9.9.4 Other therapies

Heliox

There is no evidence to support the use of heliox for the treatment of acute asthma in childhood.

9.9.5 Critical care settings

In children with acute asthma and a poor response to standard therapy (inhaled bronchodilators, steroids, oxygen and intravenous bronchodilators) other therapies may be considered in the appropriate critical care setting with the appropriate available expertise. There is little high-quality evidence to guide treatment at this stage of an acute asthma attack and it is important to involve a clinician with the appropriate skills in airway management and critical care support as early as possible.

Ketamine

A systematic review of the use of ketamine for the management of acute asthma attacks in children found only one small study (n=68), among non-intubated children, suitable for inclusion. No benefit from ketamine compared with placebo in terms of respiratory rate, oxygen saturation, hospital admission rate, need for mechanical ventilation, or need for other adjuvant therapy was found.⁶⁷⁷

1++
2-

Sevoflurane

A small (n=7) non-comparative study of sevoflurane in children with life-threatening asthma reported that sevoflurane inhalation corrects high levels of PaCO₂ and provides clinical improvement in mechanically ventilated children.⁶⁷⁸ Use of this agent is, however, limited to areas with appropriate scavenging facilities to extract gas in order to protect healthcare staff.

3

Extracorporeal membrane oxygenation

There is no good quality evidence on the use of ECMO in children, probably reflecting, in part, the low number of children who would be suitable for this approach. Extracorporeal membrane oxygenation has, however, been used successfully in other forms of critical respiratory failure in children for a number of years and there are four paediatric ECMO centres in the UK that would consider treating children with near-fatal asthma who are not responding to conventional treatment (Glenfield Hospital, Leicester; The Freeman Hospital, Newcastle; The Royal Hospital for Children, Glasgow; and Great Ormond Street Hospital, London).

Recombinant human deoxyribonuclease

There is no evidence to support the use of recombinant human deoxyribonuclease in acute asthma in children.



Children with asthma not responding to standard treatment should be evaluated by a specialist with the appropriate experience and skills to use and assess medication familiar to those in critical care settings.

9.9.6 Non-invasive ventilation

A systematic review of NIV for acute asthma in children included two RCTs (n=40) comparing NIV as add-on therapy to usual care versus usual care in children under 18 years of age hospitalised for an acute asthma attack. Both included studies used bilevel positive airway pressure only. Both included studies reported improvements in asthma symptom scores. This finding is, however, based on a small number of participants and on trials assessed as having a high risk of bias.⁶⁷⁹

1++

A further, observational, study reported that NIV is feasible in children with severe asthma within the paediatric intensive care unit setting, but did not include a control group for comparison of clinical outcomes.⁶⁸⁰

3

Although there is some evidence that NIV is safe and feasible for use in this population, there is little evidence of its effectiveness and insufficient evidence on which to base a recommendation.

Future trials, including measurable clinical outcomes such as respiratory parameters, physiological variables and blood gases, are needed to assess the role of NIV in treating children with *status asthmaticus*.

9.9.7 Discharge planning

Children can be discharged when stable on 3–4 hourly inhaled bronchodilators that can be continued at home.⁶⁸¹ Peak expiratory flow and/or FEV₁ should be >75% of best or predicted and SpO₂ >94%. An asthma care bundle developed by BTS is also available from the BTS website (www.brit-thoracic.org.uk). Adult studies show that optimal care comprising self monitoring, regular review and a written PAAP can improve outcomes.¹⁶⁶ Acute asthma attacks should be considered a failure of preventive therapy and thought should be given about how to help families avoid further severe episodes.

Discharge plans should address the following:

- the diagnosis – clearly document the criteria used to diagnose asthma
- check inhaler technique
- consider the need for preventer treatment or optimising/adjusting previously prescribed preventer treatments
- provide a written PAAP for subsequent asthma attacks with clear instructions about the use of bronchodilators and the need to seek urgent medical attention in the event of worsening symptoms not controlled by up to 10 puffs of salbutamol 4 hourly
- assess exposure to environmental tobacco smoke or actual smoking in older children and refer to suitable agencies where appropriate
- identify the trigger of the acute attack and discuss future management plans for exposure
- arrange follow up by primary care services within two working days
- arrange follow up in a paediatric asthma clinic at about one month after admission
- arrange referral to a paediatric respiratory specialist if there have been life-threatening features.

Many children with recurrent episodes of wheeze triggered by viruses do not go on to develop atopic asthma. The need for regular preventer treatment may depend on the severity and frequency of episodes. Many may not require inhaled corticosteroids.



It is essential that the patient's primary care practice is informed within 24 hours of discharge from the emergency department or hospital following an asthma attack. Ideally this communication should be directly with a named individual responsible for asthma care within the practice.

10 Difficult asthma

10.1 Defining and assessing difficult asthma

The term difficult asthma generally refers to a clinical situation where a prior diagnosis of asthma exists, and asthma-like symptoms and asthma attacks persist despite prescription of high-dose asthma therapy. There is no definition of difficult asthma in children or adults that is universally agreed, and specifically at what level of treatment prescription or asthma attack frequency the term difficult asthma should apply. Consequently there are no precise data on the prevalence of this clinical problem. Previous consensus studies have suggested failure to achieve symptom control despite prescribed high-dose ICS as a minimum requirement, or have stipulated a treatment level equivalent to at least high-dose ICS (adults) or medium-dose ICS (children) plus a LABA or LTRA before labelling as 'difficult'.^{682, 683}

In this guideline difficult asthma is defined as persistent symptoms and/or frequent asthma attacks despite treatment with high-dose ICS (adults) or medium-dose ICS (children) plus a LABA (age 5 and over) or LTRA; or medium-dose ICS (adults) or low-dose ICS (children) plus a LABA (age 5 and over) or LTRA and an appropriate additional therapy (*see section 7.5.2*); or continuous or frequent use of oral steroids (*see section 7.5.3*).

Observational uncontrolled studies in participants with difficult asthma, using multidisciplinary assessment models have identified high rates of alternative or coexistent diagnoses and psychological comorbidity.^{97, 684-686} These uncontrolled studies, using systematic multidisciplinary assessment and management, have suggested improved outcomes in adults and children, but controlled clinical trials are required. Within this broadly defined group of participants with difficult asthma, a proportion will have refractory disease, which is relatively resistant to currently available therapies. This group can only be identified after detailed evaluation, including exclusion of alternative causes of persistent symptoms, management of other comorbidities and confirmation of adherence with therapy.

D Patients with difficult asthma should be systematically evaluated, including:

- **confirmation of the diagnosis of asthma, and**
- **identification of the mechanism of persisting symptoms and assessment of adherence to therapy.**

D This assessment should be facilitated through a dedicated multidisciplinary difficult asthma service, by a team experienced in the assessment and management of difficult asthma.

10.2 Factors contributing to difficult asthma

10.2.1 Poor adherence

Poor adherence with asthma medication is associated with poor asthma outcome in adults and children (*see section 5.4*). Two UK studies in adults attending specialist difficult asthma services documented high levels of poor adherence identified by low prescription filling. A study of 182 patients in the Northern Ireland Regional Difficult Asthma Service found that 63 patients (35%) filled 50% or fewer inhaled

3

2+
3

LABA/ICS prescriptions and 88% admitted poor adherence with inhaled therapy after initial denial; 23 of the 51 patients (45%) prescribed oral steroids were found to be non-adherent using serum prednisolone/cortisol testing.⁶⁸⁷ In another study, 75 of 115 (65%) patients filled prescriptions for <80% of ICS medication and had significantly worse lung function, higher sputum eosinophil counts and prior ventilation compared to adherent patients.⁶⁸⁸ A study of 71 school-aged children with persistent symptoms, despite high-dose treatment or continuous or frequent use of oral steroids, attending one hospital in London, found that 56 (79%) had potentially modifiable risk factors, the two most common of which were psychosocial factors (59%) and medication issues including adherence (48%). In 39 children (55%) the factors identified and the interventions recommended meant that further escalation of treatment was avoided.⁶⁸⁹ In a paediatric case-control series, poor adherence based on prescription records was identified in 22% of children with difficult to control asthma, although adherence was not reported in the stable controls.⁶⁹⁰ In a descriptive study of 100 adult participants with a physician diagnosis of 'severe asthma', 28 patients were on >15 mg prednisolone and of these nine (32%) were found to be non-adherent with prednisolone.⁶⁸⁵

2+
3

There is a need to identify patients who have poor control solely as a result of poor adherence to simple therapies that are currently available. In theory, improving adherence through monitoring and intervention could potentially reduce asthma attacks, target resources for genuine therapy-resistant cases and reduce overall health costs by minimising asthma attacks, hospitalisation and health resource use.

Monitoring adherence is likely to be beneficial to asthma control and there is some evidence that it can improve lung function and quality of life.⁶⁹¹ Adherence monitoring based on self assessment is unlikely to be accurate and objective measures are therefore needed. An ancillary study to an RCT showed that there was very poor agreement between objective (doses remaining in Turbohaler device) and subjective (self-reported) measurements of adherence in children aged 5–12 years with mild or moderate asthma and airway hyper-responsiveness to methacholine, and that self reporting failed to detect poor adherence.²³⁴ Objective measurement of non-adherence based on FeNO suppression in adults with difficult asthma was demonstrated in one study although further validation of this test is required.²⁴⁴ Some other objective measures such as prescription filling are problematical because patients may fill prescriptions but not take the medication.

1+

C Healthcare professionals should always consider poor adherence to maintenance therapy before escalating treatment in patients with difficult asthma.

10.2.2 Psychosocial factors

Fatal and near-fatal asthma have been associated with adverse psychosocial factors (*see section 9.1.3*). Most observational studies^{97, 685, 692–695} and a case-control study⁶⁹⁶ in patients with difficult asthma have also suggested a high level of psychological morbidity, though this observation has not been universal.^{697, 698}

2+
3

A meta-analysis of behavioural adjustment in children suggested increasing asthma severity, defined on the basis of treatment requirements, was associated with greater behavioural difficulties.⁶⁹⁹ The core issue of cause and effect remains unclear; specifically the extent to which persistent asthma symptoms, despite aggressive treatment, results in psychological morbidity or whether pre-existing psychological morbidity makes asthma difficult to control. 2++

There is a lack of evidence that interventions specifically targeting psychological morbidity in patients with difficult asthma are of benefit. A small proof of concept study targeting treatment of depression demonstrated a reduction in oral steroid use,⁷⁰⁰ and an observational study in high-risk children with asthma suggested potential benefit from joint consultation with a child psychiatrist, with an improvement in symptom scores and adherence to therapy.⁷⁰¹ However, a non-blinded randomised intervention study in adults with difficult asthma showed no benefit from a six-month nurse-delivered psychoeducational programme.⁷⁰² A meta-analysis of psychoeducational interventions in patients with difficult asthma concluded that many of the studies were of poor quality, although there was some evidence of a positive effect from psychosocial educational interventions on hospital admissions in adults and children and on symptoms in children. There was not enough evidence to warrant significant changes in clinical practice and little information available on cost effectiveness.⁷⁰³ 1+
3

C Healthcare professionals should be aware that difficult asthma is commonly associated with coexistent psychological morbidity.

D Assessment of coexistent psychological morbidity should be performed as part of a difficult asthma assessment. In children this may include a psychosocial assessment of the family.

10.2.3 Dysfunctional breathing

Observational uncontrolled studies in patients with difficult asthma have identified high rates of dysfunctional breathing as an alternative or concomitant diagnosis to asthma. The dysfunctional breathing may cause symptoms that mimic asthma or coexist with asthma, worsening symptoms.^{97, 685} It remains unclear what is the best mechanism for identifying and managing this problem. 3

D Dysfunctional breathing should be considered as part of the assessment of patients with difficult asthma.

10.2.4 Allergy

Acute asthma has been associated with IgE-dependent sensitisation to indoor allergens.⁷⁰⁴ In case-control studies, mould sensitisation has been associated with recurrent admission to hospital and oral steroid use^{705, 706} and with intensive care unit admissions and respiratory arrest.^{707, 708} There is no published evidence of any intervention study in this patient group. Research in this area is required. 2++
3

C In patients with difficult asthma and recurrent hospital admission, allergen testing to moulds should be performed.

10.2.5 Monitoring airway response

Two blinded RCTs and one open randomised study have supported the use of titrating steroid treatment against sputum eosinophilia in adults with moderate to severe asthma, with greatest benefit seen in patients receiving higher doses of ICS therapy.⁷⁰⁹⁻⁷¹¹ In the study with the largest number of patients receiving high-dose ICS treatment, patients who were considered to be poorly adherent with treatment, or had inadequately controlled aggravating factors, such as rhinitis or gastro-oesophageal reflux were specifically excluded.⁷⁰⁹ Case series have suggested that sputum induction is safe in patients with difficult to control asthma.^{67, 712-715}

1+
1-
3

Controlled studies using FeNO to target treatment have not specifically targeted adults or children with difficult asthma.^{716, 717}

1+

B In patients with difficult asthma, consider monitoring induced sputum eosinophil counts to guide steroid treatment.

11 Asthma in adolescents

11.1 Definitions

Adolescence is the transitional period of growth and development between puberty and adulthood, defined by the WHO as between 10 and 19 years of age.³

There is international agreement on best practice for working with adolescents with health problems outlined in consensus publications.⁷¹⁸⁻⁷²⁰ Key elements of working effectively with adolescents in the transition to adulthood include:⁷²¹

- seeing them on their own, separate from their parents, for part of the consultation, and
- discussing confidentiality and its limitations.

For diagnosing and managing asthma in adolescents, the evidence base is limited. Much recent research has focused on the prevalence of asthma and ecological risk associations rather than on diagnosis and management of asthma in adolescents.

11.2 Prevalence of asthma in adolescence

Asthma is common in adolescence with a prevalence of wheeze in 13–14 year olds in Western Europe in the past 12 months (current wheeze) of 14.3%.⁷²² For more severe asthma (defined as ≥ 4 attacks of wheeze or ≥ 1 night per week sleep disturbance from wheeze or wheeze affecting speech in the past 12 months) the prevalence was 6.2%.

There is evidence of underdiagnosis of asthma in adolescents, with estimates of 20–30% of all asthma present in this age group being undiagnosed.⁷²²⁻⁷²⁵ This has been attributed to under reporting of symptoms. A number of risk factors have been independently associated with underdiagnosis including: female gender, smoking (both current smoking and passive exposure), low socioeconomic status, family problems, low physical activity and high body mass, and race/ethnicity.⁷²⁵ Children with undiagnosed frequent wheezing do not receive adequate healthcare for their illness⁷²⁵ and the health consequences of not being diagnosed with asthma are substantial.^{726, 727}

Although feasible, there is insufficient evidence to support screening for asthma in adolescents.^{728, 729}



Clinicians seeing adolescents with any cardiorespiratory symptoms should ask about symptoms of asthma.

11.3 Diagnosis and assessment

11.3.1 Exercise-related symptoms

Exercise-related wheezing and breathlessness are common asthma symptoms in adolescents. However, these symptoms are poor predictors of exercise-induced asthma. Only a minority of adolescents referred for assessment of exercise-induced respiratory symptoms show objective evidence of exercise-induced bronchospasm.⁷³⁰ Other diagnoses producing reproducible symptoms on exercise

include normal physiological exercise limitation, with and without poor physical fitness, vocal cord dysfunction, dysfunctional breathing, habit cough, and supraventricular tachycardia.⁵⁴

Most exercise-related wheezing in adolescents can be diagnosed and managed by careful clinical assessment.⁷³¹ The absence of other features of asthma and an absent response to pretreatment with β_2 agonist make exercise-induced asthma unlikely. Exercise testing with cardiac and respiratory monitoring that reproduces the symptoms may be helpful in identifying the specific cause.⁵⁴

11.3.2 Use of questionnaires

When using questionnaires, the prevalence of current symptoms is higher when the adolescent completes the questions rather than the parents, while questions about the last 12 months give similar results between the parents and the adolescent.⁷³²

In one study in adolescents, internet and written questionnaires about asthma provided equivalent results.⁷³³ The ACQ and the ACT have been validated in adolescents with asthma (see Table 8).¹¹⁸

11.3.3 Quality of life measures

Quality of life scales (such as AQLQ12+) can be used in adolescents.^{734, 735}

11.3.4 Lung function

In adolescents with asthma, tests of airflow obstruction and airway responsiveness may provide support for a diagnosis of asthma. However, most adolescents with asthma have normal lung function despite having symptoms.

11.3.5 Bronchial hyper-reactivity

Although many children with asthma go into long-lasting clinical remission at adolescence, BHR may persist. Whether persisting BHR reflects ongoing airway inflammation is debated.⁷³⁶

A negative response to an exercise test is helpful in excluding asthma in children with exercise-related breathlessness.⁵⁴

11.4 Risk factors

There is a body of evidence from cohort studies highlighting risk factors for asthma in adolescents.

11.4.1 Atopy

Studies confirm that atopic dermatitis and atopic rhinitis are amongst the factors most strongly associated with asthma persisting into teenage years.⁷³⁷⁻⁷⁴⁰

11.4.2 Prematurity and early life wheezing

Adolescents who were very low birth weight due to prematurity (as opposed to intrauterine growth retardation) were more prone to chronic cough, wheezing and asthma and showed medium and small airway obstruction compared with matched controls.⁷⁴¹

Frequent or severe episodes of wheezing in childhood are associated with recurrent wheeze that persists into adolescence.^{74, 77, 86, 88, 94, 106-108, 740}

11.4.3 Gender

During adolescence there is a reversal of the gender association of asthma with the disease being more prevalent in females than males from 13–14 years onwards.⁷⁴² The same change is seen with asthma attacks, with risk of an asthma admission in females becoming double that observed in males from around 13–14 years.⁷⁴³ This phenomenon has been attributed to a greater incidence of asthma among teenage girls.⁷⁴⁴

11.4.4 Chlorinated swimming pools

Exposure to chlorinated swimming pools has been associated with an increased risk of asthma, airway inflammation and some respiratory allergies.⁷⁴⁵ Such associations were not found among adolescents without atopy or in those who attended copper-silver sanitised pools.⁷⁴⁶

11.5 Comorbidities and modifiable behaviours

11.5.1 Anxiety and depressive disorders

Asthma in adolescence is associated with an increased likelihood of major depression, panic attacks and anxiety disorder. This may reflect effects of common factors associated with anxiety and depressive disorders rather than a direct causal link with asthma.⁷⁴⁷ In young people with asthma, the presence of an anxiety or depressive disorder is highly associated with increased asthma symptom burden.⁷⁴⁸ Depressive symptoms were one risk factor identified in children and adolescents who died of asthma. Assessment of anxiety may help identify individuals who are at risk for poorer asthma-specific quality of life.⁷⁴⁹

Clinical conditions associated with anxiety may be mistaken for, or overlap with asthma. These include dysfunctional breathing (hyperventilation syndrome and sighing dyspnoea), vocal cord dysfunction, and psychogenic cough. These conditions can present acutely and may often be frightening to the young person. This may lead to a cycle of bronchodilator overuse, which then further exacerbates the symptoms. Detailed medical assessment with careful attention to the adolescent's personal perceptions and experiences of their symptoms is required to make an accurate diagnosis.⁷⁵⁰

Brief screening questionnaires for anxiety and depression suitable for use in adolescents are available and may help identify those with significant anxiety and depression.⁷⁵¹

11.5.2 Obesity

The evidence on whether asthma is more common in overweight and obese adolescents with asthma is conflicting.^{737, 752-754} While weight reduction in obese adults with asthma improves lung function, symptoms, morbidity and health status, this has not yet been established in adolescents with asthma.

11.5.3 Gastro-oesophageal reflux and gastro-oesophageal reflux disease

Gastro-oesophageal reflux and GORD is common in patients with asthma, including adolescents.⁷⁵⁵ A systematic review confirmed an association between GORD and asthma in children and adolescents in secondary and tertiary referral settings. The nature of the association, however, is unclear.⁷⁵⁶ There is no evidence that treatment for GORD improves asthma symptoms in children and adolescents with GORD and asthma.^{527, 528}

11.6 Asthma attacks and the risk of hospital admission

Clinical characteristics and markers of severity, including frequent respiratory symptoms, airway hyper-responsiveness, atopy, and low lung function, identify those at high risk of hospitalisation for asthma, particularly with respect to multiple admissions.⁷⁵⁷

11.7 Long-term outlook and entry into the workplace

A long-term follow-up study of vocational and working careers found that adolescents and young adults (10–22 years) with relatively mild asthma had slightly more limitations in vocational and professional careers than those without asthma. They had a small increased risk of limitations in daily activity attributable to respiratory health and of absence from work. In the majority, however, the differences amounted to only a few days per year.⁷⁵⁸ Young adults with asthma had a low awareness of occupations that might worsen asthma (for example exposure to dusts, fumes, sprays, exertion and temperature changes) and did not generally discuss career plans with their GP. Further details about occupational asthma can be found in section 13.



Clinicians should discuss future career choices with adolescents with asthma and highlight occupations that might increase susceptibility to work-related asthma symptoms.

11.8 Non-pharmacological management

11.8.1 Tobacco smoking and environmental exposure to tobacco smoke

Exposure to passive smoking remains a significant health risk.

One study of asthma morbidity among urban young adolescents (mean approximately 11 years of age) found at baseline that 28% of caregivers reported exposure to environmental tobacco smoke (ETS) in the home and 19% reported exposure outside the primary household. Children who received a 20-minute educational intervention about ETS exposure and whose ETS exposure had decreased at follow up had fewer hospitalisations ($p=0.034$) and emergency department visits ($p\leq 0.001$) reported in the next 12 months, as well as fewer episodes of poor asthma control ($p=0.042$).⁷⁵⁹

In a national survey in Denmark, 37.7% of adolescents with asthma smoked currently, 16.5% daily. Smoking was more common in girls. More of those with asthma smoked daily, smoked more cigarettes and had tried to quit smoking.⁷⁶⁰

2+
3
4

Among adolescents, smoking is a risk factor for asthma.^{738, 761-763} A longitudinal study of asthma and allergic disease in school children in Sweden found that both passive and active smoking were significantly related to asthma and wheeze in adolescents. Maternal ETS exposure was associated with lifetime symptoms, but daily smoking among the adolescents was more strongly related to current symptoms.⁷⁶⁴

2+
3
4

Young people aged 12–17 years who have a strong commitment to quit smoking should be offered advice on how to stop and encouraged to use local NHS smoking cessation services by providing details of when, where and how to access them.

- ✓ Adolescents with asthma (and their parents and carers) should be encouraged to avoid exposure to environmental tobacco smoke, and should be informed about the risks and urged not to start smoking.
- ✓ Adolescents with asthma should be asked if they smoke personally. If they do and wish to stop, they should be offered advice on how to stop and encouraged to use local NHS smoking cessation services.

11.8.2 Complementary and alternative medicine

In a small study of Italian teenagers, 16% of those included had used complementary and alternative medicine (CAM) (homeopathy, acupuncture, herbal medicines).⁷⁶⁵ In a study in the USA, 80% of urban adolescents (aged 13–18 years) with asthma reported that they had used CAM, most commonly rubs, herbal teas, prayer and massage.⁷⁶⁶ While most adolescents used CAM with conventional asthma therapy, 27% reported they used it instead of prescribed therapy⁷⁶⁶ suggesting that CAM use may be a marker of non-adherence to prescribed asthma treatment.

2-

- ✓ Healthcare professionals should be aware that complementary and alternative medicine use is common in adolescents and should ask about its use.

11.9 Pharmacological management

Specific evidence about the pharmacological management of adolescents with asthma is limited and is usually extrapolated from paediatric and adult studies. Recommendations for pharmacological management of asthma in children and adults can be found in section 7.

11.10 Inhaler devices

Specific evidence about inhaler device use and choice in adolescents is limited. Inhaler devices are covered in section 8.

Two small studies comparing two different types of inhalers in adolescents found that both DPIs and pMDIs plus spacer are of value in adolescent asthma.^{767, 768} There were no differences between the two inhaler devices in terms of symptoms or lung function but patients preferred the DPI.

2+
3

Although adolescents with asthma may be competent at using their inhaler devices, their actual adherence to treatment may be affected by other factors such as preference. In particular, many adolescents prescribed a pMDI with spacer do not use the spacers as they are felt to be too inconvenient.^{769, 770}

- ✓ Adolescent preference for inhaler device should be taken into consideration as a factor in improving adherence to treatment.
- ✓ As well as checking inhaler technique it is important to enquire about factors that may affect inhaler device use in real life settings, such as school.
- ✓ Consider prescribing a more portable device (as an alternative to a pMDI with spacer) for delivering bronchodilators when away from home.

11.11 Organisation and delivery of care

11.11.1 Healthcare setting

Very little evidence was identified to determine the best healthcare setting to encourage attendance amongst adolescents with asthma.

A two-year follow-up study found that a multidisciplinary day programme improved asthma control in a group of adolescents with very severe asthma. This study involved a highly selected group of patients and a wide range of interventions and is not generalisable to most adolescents with asthma.⁶⁸⁴

3

11.11.2 Schools as a setting for healthcare delivery and asthma education

Some innovative approaches have used schools as a setting for asthma education and review (*see section 14.5*). One focus has been on healthcare delivery, such as school-based clinics. Evidence from a single cluster RCT suggests that school-based, nurse-led asthma clinics increase the uptake of asthma reviews in adolescents from 51% in practice care to 91%.⁷⁷¹ Knowledge of asthma, inhaler techniques and positive attitudes increased and a majority of the adolescents preferred the setting, but there was no improvement in clinical outcomes. This may be because the nurses were not able to change or prescribe treatment (which relied on a separate visit to a doctor).

Other approaches have used schools as a setting for asthma education including peer-led education. In a single well-conducted RCT, peer-led education in schools improved quality of life, asthma control and days off school for adolescents with asthma.⁷⁷² In a study in the USA, a randomised trial of a web-based tailored asthma management programme delivered using school computers found that, after 12 months, students reported fewer symptoms, school days missed, restricted-activity days, and hospitalisations for asthma than control students. The programme was inexpensive to deliver.²⁰¹

1+

A number of countries, particularly Australia and New Zealand, have developed national programmes to ensure that schools can deliver appropriate first aid and emergency response to students with asthma as well as encouraging participation in sporting activities.⁷⁷³

B School-based clinics may be considered for adolescents with asthma to improve attendance.

B Peer-led interventions for adolescents in the school setting should be considered.

- ✓ Integration of school-based clinics with primary care services is essential.

11.11.3 Transition to adult services

Transition to adult services is important for all adolescents with asthma, irrespective of the asthma severity. No studies on transition of adolescents with asthma to adult services were identified although there are many studies looking at transition of adolescents with chronic illness. Few studies compare different approaches and many recommendations come from consensus statements rather than randomised controlled trials.⁷¹⁸⁻⁷²⁰

It is important that the process of transition is co-ordinated and that a healthcare professional be identified to oversee transition and either link with a counterpart in adult services or remain involved until the young person is settled within adult services.^{774, 775}

- ✓ In the initial period after transition to adult services in secondary care, adolescents are best seen by one consultant to build their confidence and encourage attendance.

11.11.4 Preparation for transition

Transition should be seen as a process and not just the event of transfer to adult services.⁷⁷⁴ It should begin early, be planned, involve the young person, and be both age and developmentally appropriate.⁷⁷⁴

- Young people should be given the opportunity to be seen without their parents/carers.
- Transition services must address the needs of parents/carers whose role in their child's life is evolving at this time.
- Transition services must be multidisciplinary and multiagency. Optimal care requires a co-operative working relationship between adult and paediatric services, particularly where the young person has complex needs with multiple specialty involvement.
- Co-ordination of transitional care is critical. There should be an identified co-ordinator who supports the young person until he or she is settled within the adult system.
- Young people should be encouraged to take part in transition/support programmes and/or put in contact with other appropriate youth support groups.
- The involvement of adult physicians prior to transfer supports attendance and adherence to treatment.
- Transition services must undergo continued evaluation.

11.12 Patient education and self management

11.12.1 Education in self management

Section 5 covers supported self management, including the components of a self-management programme.

Effective transition care involves preparing adolescents with asthma to take independent responsibility for their own asthma management and enabling them to negotiate the health system effectively. Clinicians need to educate and empower adolescents to manage as much of their asthma care as they are

capable of doing while supporting parents to gradually hand over responsibility for management to their child. The specific knowledge, attitudes and skills that underpin independent self-management practices in adolescents with asthma are that they can:⁷⁷⁶

- name and explain their condition
- list their medications, treatments or other management practices (eg special diet)
- explain why each medication or management practice is necessary
- remember to take their medications most of the time
- answer questions asked of them by doctors or other healthcare professionals
- ask questions of their doctor or other healthcare professional
- arrange (and cancel) appointments
- consult with a doctor or other healthcare professional without a parent/carer
- remember to order more medication before it runs out
- have prescriptions filled at the pharmacy
- develop the desire for their healthcare to be independent of their parents/carers
- prioritise their health over (some) other desires.

For adolescents with asthma, the available evidence about self management is mainly qualitative and provides insight about the concerns adolescents have about their asthma and its management. Adolescents with asthma report embarrassment over using inhalers in front of others, sadness over not being able to take part in normal activities, frustration and anger at the way they are treated by their families (for example being limited in what they are allowed to do, being fussed over by parents). They also report specific anxieties around fear of dying and feeling guilty over the effect their illness has on the rest of the family. They are concerned about needing to rely on someone else when they have a bad asthma attack and that teachers do not know what to do. They stress the importance of support from friends at school, especially those with asthma.^{777, 778}

Studies of adolescents with chronic illness (including asthma) have highlighted factors that adolescents feel are important in delivering education about self management to them.⁷⁷⁹ These included:

- education must be adapted to meet individual needs and repeated and developed as understanding and experience increases and should include emotional support for coping with feelings
- education should be delivered by educators that respect, engage, encourage and motivate the adolescents
- accompanying information, both written and oral, should be personalised rather than general and use non-medical language that adolescents can understand
- education should be delivered in an appropriate and uninterrupted setting and make appropriate use of information technology.

D

Design of individual or group education sessions delivered by healthcare professionals should address the needs of adolescents with asthma.

11.12.2 Adherence

Adherence with asthma treatment, and with asthma trigger avoidance, is often poor in adolescents. The evidence for poor adherence comes mainly from questionnaire-based and qualitative studies rather than objective electronic monitoring.⁷⁸⁰

When directly asked, most adolescents admit they do not always follow their treatment plans. Reasons for not adhering include both unintentional reasons (confusion about medications and forgetfulness) and intentional reasons (inhalers being ineffective/hard to use, treatment plan too complicated, more important things to do, concern about adverse effects, denial, can't be bothered and embarrassment).^{770, 781} Background factors, such as younger age, family size, exercise and not smoking or drinking alcohol as well as disease-related factors such as sense of normality, energy and will power, support from the parents, physicians and nurses, and a positive attitude towards the disease and treatment were related to good reported adherence.⁷⁸²

Non-adherence to medication regimens in adolescents has been linked to other health-risk behaviours including tobacco, alcohol and drug use and also to depression.⁷⁸³ Not only are specific behaviours such as smoking, poor adherence to medication regimens or medical review appointments detrimental to asthma control, they also have been highlighted as potential beacons of distress in adolescents.⁷⁸⁴ Clinical tools such as the Home, Education/Employment, Activities, Drugs, Sexuality, Suicide/depression adolescent health screen provide practitioners with an easily usable psychosocial screen.⁷⁸⁵

Strategies to improve adherence in adolescents emphasise the importance of focusing on the individual and their lifestyle and using individualised asthma planning and personal goal setting.⁷⁸⁶ One study found that once-daily supervised asthma preventer therapy at school improved asthma control and quality of life.⁷⁸⁷

3

12 Asthma in pregnancy

12.1 Natural history and management of stable asthma

The majority of women with asthma have normal pregnancies and the risk of complications is small in those with well-controlled asthma. Several physiological changes occur during pregnancy that could worsen or improve asthma, but it is not clear which, if any, are important in determining the course of asthma during pregnancy. Pregnancy can affect the course of asthma and asthma and its treatment can affect pregnancy outcomes.

12.1.1 Course of asthma in pregnancy

The natural history of asthma during pregnancy is extremely variable. In a prospective cohort study of 366 pregnancies in 330 women with asthma, the asthma worsened during pregnancy in 35%.⁷⁸⁸ A prospective cohort study of 1,739 pregnant women showed an overall improvement in 23% and deterioration in 30.3%.⁷⁸⁹ The conclusions of a meta-analysis of 14 studies is in agreement with the commonly quoted generalisation that during pregnancy about one third of asthma patients experience an improvement in their asthma, one third experience a worsening of symptoms, and one third remain the same.⁷⁹⁰ There is also some evidence that the course of asthma is similar in successive pregnancies.^{788, 791} A systematic review showed no effect of pregnancy or stage of pregnancy on FEV₁.⁷⁹²

1+
2+

Studies suggest that 11–18% of pregnant women with asthma will have at least one emergency department visit for acute asthma and of these 62% will require hospitalisation.^{793, 794} Severe asthma is more likely to worsen during pregnancy than mild asthma,⁷⁸⁸ but some patients with very severe asthma may experience improvement, whilst symptoms may deteriorate in some patients with mild asthma. In a large study in the USA, the rates of asthma attack were 13%, 26% and 52% in those with mild, moderate and severe asthma respectively.⁷⁸⁹ The corresponding rates of hospitalisation were 2%, 7% and 27%.

2+
2-

A systematic review concluded that if symptoms do worsen this is most likely in the second and third trimesters, with the peak in the sixth month.⁷⁹¹ In a large cohort study, the most severe symptoms were experienced by patients between the 24th and 36th week of pregnancy. Thereafter symptoms decreased significantly in the last four weeks and 90% of patients had no asthma symptoms during labour or delivery. Of those who did, only two patients required anything more than inhaled bronchodilators.⁷⁸⁸ A further study has confirmed that patients are least likely to have an asthma attack in the last month of pregnancy.⁷⁹⁵

2+

12.1.2 Effect of asthma in pregnancy

A systematic review has shown that baseline asthma severity does determine what happens to the course of asthma in pregnancy and asthma may affect the risk of adverse outcomes.⁷⁹⁶ A cohort study comparing 198 pregnant women with asthma to 198 women without asthma reported that non-atopic patients with asthma tend to have more severe asthma. Pre-eclampsia was also more common in this group. However with adequate surveillance and treatment, pregnancy and delivery complications can be avoided.⁷⁹⁷

2++
2+

Uncontrolled asthma is associated with many maternal and fetal complications, including hyperemesis, hypertension, pre-eclampsia, vaginal haemorrhage, complicated labour, fetal growth restriction, preterm birth, increased perinatal mortality, and neonatal hypoxia.^{789, 798-801} A large Swedish population-based study using record linkage data demonstrated increased risks for preterm birth, low birth weight, perinatal mortality and pre-eclampsia in women with asthma. The risks for preterm delivery and low birth weight were higher in women with more severe asthma necessitating admission.⁸⁰²

2+

A large prospective cohort study comparing women with moderate and severe asthma with women without asthma found the only significant difference was an increased Caesarean section rate (OR 1.4, 95% CI 1.1 to 1.8).⁷⁸⁹ Logistic regression analysis of the severe group showed an increased risk of gestational diabetes (adjusted odds ratio (AOR) 3.0, 95% CI 1.2 to 7.8) and preterm delivery <37 weeks (AOR 2.2, 95% CI 1.2 to 4.2) but this could have been an effect of corticosteroids. In the Yale asthma study no effect of asthma symptoms or severity was seen on preterm delivery but oral steroids increased the rate of preterm delivery and reduced gestation by 2.2 weeks (AOR 1.05, 95% CI 1.01 to 1.09).⁸⁰³ Daily asthma symptoms were associated with an increased risk of fetal growth restriction (AOR 2.25, 95% CI 1.25 to 4.06) and there was a 24% increase with each increased symptom step. This is supported by a systematic review of four studies that concluded asthma exacerbation in pregnancy increases the risk of low birth weight.⁸⁰⁴ The RR was 2.54 (95% CI 1.52 to 4.25) compared with women without asthma. In a large cohort study of 2,123 women with asthma, there was an association of both mean FEV₁ and mean FEV₁ <80% predicted with gestational hypertension, preterm delivery <37 weeks and <32 weeks, and low birth weight.⁸⁰⁵

2+

2++

In contrast, if asthma is well controlled throughout pregnancy there is little or no increased risk of adverse maternal or fetal complications.^{788, 793} Pregnancy should therefore be an indication to optimise therapy and maximise lung function in order to reduce the risk of acute asthma attacks.

2+

C Monitor pregnant women with moderate/severe asthma closely to keep their asthma well controlled.

B Women should be advised of the importance of maintaining good control of their asthma during pregnancy to avoid problems for both mother and baby.

✓ Advise women who smoke about the dangers for themselves and their babies and give appropriate support to stop smoking.

12.2 Management of acute asthma in pregnancy

The management of acute asthma in pregnancy may be affected by concerns about harmful effects of medication on the fetus. In a prospective controlled study of 51 pregnant women and 500 non-pregnant women presenting with acute asthma to an emergency department in Boston, USA, pregnant patients with asthma were less likely to receive appropriate treatment with steroids and, as a result, were more likely to experience ongoing asthma attacks at two weeks.⁸⁰⁶ Available studies give little cause for concern regarding treatment side effects and the maternal and fetal risks of uncontrolled asthma are much greater than the risks from using conventional asthma medications for management of acute asthma. In five confidential enquiries into maternal deaths in the UK (covering

2+

1994–2008) there were 22 deaths from asthma.^{807–811} A report from the Intensive Care National Audit and Research Centre on female admissions to adult critical care units in England, Wales and Northern Ireland between 2009 and 2012 found that of 1,188 currently pregnant women, 94 (8%) were admitted with acute asthma and of 5,605 postpartum women, 32 (0.6%) were admitted with acute asthma.⁸¹²

2+

Oxygen should be delivered to maintain saturation 94–98% in order to prevent maternal and fetal hypoxia.⁵⁷⁸ When interpreting arterial blood gases in pregnancy it should be remembered that the progesterone-driven increase in minute ventilation may lead to relative hypocapnia and a respiratory alkalosis, and higher PaO₂^{813, 814} but oxygen saturations are unaltered.⁸¹⁵ Acidosis is poorly tolerated by the fetus.

4

Drug therapy should be given as for a non-pregnant patient with acute asthma, including nebulised β_2 agonists and early administration of steroid tablets (*see section 9*).^{788, 794, 795, 798, 799} In severe cases, intravenous β_2 agonists, aminophylline or intravenous bolus magnesium sulphate can be used as indicated.⁸¹⁶

2+

Continuous fetal monitoring should be performed when asthma is uncontrolled or severe, or when fetal assessment on admission is not reassuring. Consideration should be given to early referral to critical care services as impaired ventilatory mechanics in late pregnancy can lower functional residual capacity and may result in earlier oxygen desaturation.⁸¹⁷ Pregnant women may be more difficult to intubate due to anatomical changes especially if they have pre-eclampsia.⁸¹⁸

C In pregnant patients, give drug therapy for acute asthma as for non-pregnant patients including systemic steroids and magnesium sulphate.

D In pregnant patients with acute asthma, deliver high-flow oxygen immediately to maintain saturation 94–98%.

D Acute severe asthma in pregnancy is an emergency and should be treated vigorously in hospital.

✓ Continuous fetal monitoring is recommended for pregnant women with acute severe asthma.

✓ For women whose asthma is poorly controlled during pregnancy there should be close liaison between the respiratory physician and obstetrician, with early referral to critical care physicians for women with acute severe asthma.

12.3 Drug therapy in pregnancy

In general, the medicines used to treat asthma are safe in pregnancy.^{819, 820} A large UK population-based case-control study found no increased risk of major congenital malformations in children of women receiving asthma treatment in the year before or during pregnancy.⁸²¹ The risk of harm to the fetus from severe or chronically undertreated asthma outweighs any small risk from the medications used to control asthma.

2+

C Counsel women with asthma regarding the importance and safety of continuing their asthma medications during pregnancy to ensure good asthma control.

12.3.1 β_2 agonists

No significant association has been demonstrated between major congenital malformations or adverse perinatal outcome and exposure to short-acting β_2 agonists.⁸¹⁹⁻⁸²³ A prospective study of 259 pregnant patients with asthma who were using bronchodilators compared with 101 pregnant patients with asthma who were not, and 295 control participants, found no differences in perinatal mortality, congenital abnormalities, prematurity, mean birth weight, Apgar scores or labour/delivery complications.⁸²⁴ A case-control study including 2,460 infants exposed to short-acting β_2 agonists found no increased risk of congenital malformations in exposed infants.⁷⁸⁹

2++
2+
3

With regard to LABAs, evidence from prescription event monitoring suggests that salmeterol is safe in pregnancy and although there are some data on formoterol, numbers are small.^{825, 826} A systematic review of studies including 190 exposures to LABA demonstrated no increased risk of congenital malformations, preterm delivery or pre-eclampsia.⁸²⁷ A case-control study including 156 infants exposed to LABA found no increased risk of major congenital malformations.⁸²¹ As in other settings, LABAs should be used with an ICS, ideally as a combination product.⁸²⁸

2+

Data on the use of combination products in pregnancy are limited although there are no theoretical reasons why these would be more harmful than the same agents given separately. There are some safety data for seretide (salmeterol/fluticasone propionate) but with small numbers.⁸²⁹

C Use short-acting β_2 agonists as normal during pregnancy.

C Use long-acting β_2 agonists as normal during pregnancy.

12.3.2 Inhaled corticosteroids

No significant association has been demonstrated between major congenital malformations or adverse perinatal outcome and exposure to ICS.^{819, 821, 827, 830-838} A meta-analysis of four studies of ICS use in pregnancy showed no increase in the rate of major malformations, preterm delivery, low birth weight or pregnancy-induced hypertension.⁸³⁹ The UK case-control study included 1,429 infants exposed to ICSs and found no increased risk of major congenital malformations.⁸²¹

2++
2+
2-

Inhaled anti-inflammatory treatment has been shown to decrease the risk of an acute attack of asthma in pregnancy and the risk of readmission following an asthma attack.^{794, 795} A randomised placebo-controlled trial of inhaled beclometasone compared with oral theophylline in moderate asthma in pregnancy showed no difference in the primary outcome of one or more asthma attacks resulting in medical intervention, but inhaled beclometasone was better tolerated.⁷⁸⁹

2+

B Use inhaled corticosteroids as normal during pregnancy.

12.3.3 Theophyllines

No significant association has been demonstrated between major congenital malformations or adverse perinatal outcome and exposure to methylxanthines.^{819, 840}

2+

For women requiring theophylline to maintain asthma control, measurement of theophylline levels is recommended. Since protein binding decreases in pregnancy, resulting in increased free drug levels, a lower therapeutic range is probably appropriate.⁸⁴¹

4

C Use oral and intravenous theophyllines as normal during pregnancy.

D Check blood levels of theophylline in pregnant women with acute severe asthma and in those critically dependent on therapeutic theophylline levels.

12.3.4 Steroid tablets

There is much published literature showing that steroid tablets are not teratogenic,⁸¹⁹ but there is a slight concern that they may be associated with oral clefts. Data from several studies have failed to demonstrate this association with first trimester exposure to steroid tablets.^{842, 843} One case-control study, however, found a significant association, although this increase is not significant if only paired controls are considered.⁸⁴⁴ Although one meta-analysis reported an increased risk, a prospective study by the same group found no difference in the rate of major birth defects in prednisolone-exposed and control babies.⁸⁴⁵ A further population-based case-control study revealed a crude odds ratio of corticosteroid exposure from four weeks before through to 12 weeks after conception of 1.7 (95% CI, 1.1 to 2.6) for cleft lip.⁸⁴⁶ Another case-control study including 262 exposed infants found no such association, although this was not limited to first trimester exposure.⁸²¹

2+

2-

The association is therefore not definite and even if it is real, the benefit to the mother and the fetus of steroids for treating a life-threatening disease justify the use of steroids in pregnancy.^{800, 813} Moreover, the various studies of steroid exposure include many patients with conditions other than asthma, and the pattern of steroid use was generally as a regular daily dose rather than as short courses, which is how asthma patients would typically receive oral steroids.

2+

Prednisolone is extensively metabolised by placental enzymes so only 10% reaches the fetus, making this the oral steroid of choice to treat maternal asthma in pregnancy. Pregnant women with acute asthma attacks are less likely to be treated with steroid tablets than non-pregnant women.⁸⁰⁶ Failure to administer steroid tablets when indicated increases the risk of ongoing asthma attacks and therefore the risk to the mother and her fetus.

2+

Some studies have found an association between steroid tablet use and pregnancy-induced hypertension or pre-eclampsia, preterm labour⁷⁹⁷ and fetal growth but severe asthma may be a confounding variable.⁸⁴⁷

2+

C Use steroid tablets as normal when indicated during pregnancy for women with severe asthma. Steroid tablets should never be withheld because of pregnancy. Women should be advised that the benefits of treatment with oral steroids outweigh the risks.

12.3.5 Leukotriene receptor antagonists

Data regarding the safety of LTRAs in pregnancy are limited. A case-control study with 96 cases exposed to LTRAs found no increased risk of major malformations between women with asthma exposed to LTRA and women with asthma taking only β_2 agonists.⁸⁴⁷ A systematic review found no increased risk of malformations or preterm delivery in nine exposed women.^{803, 827} Three studies looking at infant outcomes in women exposed to LTRAs (two in women taking montelukast) showed no increased risk of congenital malformations.⁸⁴⁸⁻⁸⁵⁰

2++
2+
2-
4

C If leukotriene receptor antagonists are required to achieve adequate control of asthma then they should not be withheld during pregnancy.

12.3.6 Sodium cromoglicate and nedocromil sodium

No significant association has been demonstrated between major congenital malformations or adverse perinatal outcome and exposure to sodium cromoglicate and nedocromil sodium.^{819, 827, 847}

2+

C Use sodium cromoglicate and nedocromil sodium as normal during pregnancy.

12.3.7 Immunomodulation therapy

There are as yet no clinical data on the use of omalizumab for moderate-severe allergic asthma in pregnancy.

12.4 Management during labour

Acute attacks of asthma are very rare in labour, perhaps due to endogenous steroid production. In women receiving steroid tablets there is a theoretical risk of maternal hypothalamic-pituitary-adrenal axis suppression. Women with asthma may safely use all forms of usual labour analgesia.

In some studies there is an association between asthma and an increased Caesarean section rate,^{797, 851, 852} but this may be due to planned Caesarean sections or inductions of labour rather than due to any direct effect of asthma on intrapartum indications.⁷⁹⁵ A large prospective cohort study comparing women with moderate and severe asthma with women without asthma found the only significant difference was an increased Caesarean section rate (OR 1.4, 95% CI 1.1 to 1.8).⁷⁸⁹

2+

Data suggest that the risk of postpartum asthma attacks is increased in women having Caesarean sections.⁸⁵¹ This may relate to the severity of their asthma rather than to the Caesarean section, or to factors such as postoperative pain with diaphragmatic splinting, hypoventilation and atelectasis. Prostaglandin E2 may safely be used for labour inductions.⁸⁴¹ Prostaglandin F2 α (carboprost/hemobate®) used to treat postpartum haemorrhage due to uterine atony may cause bronchospasm.⁸⁴¹ Although ergometrine may cause bronchospasm particularly in association with general anaesthesia,⁸⁴¹ this is not a problem encountered when syntometrine (syntocinon/ergometrine) is used for postpartum haemorrhage prophylaxis.

2-
3

Although suppression of the fetal hypothalamic-pituitary-adrenal axis is a theoretical possibility with maternal systemic steroid therapy, there is no evidence from clinical practice or the literature to support this.⁸⁵³

- ✓ Advise women that an acute asthma attack is rare in labour.
- ✓ Advise women to continue their usual asthma medications in labour.
- ✓ In the absence of an acute severe asthma attack, reserve Caesarean section for the usual obstetric indications.
- ✓ If anaesthesia is required, regional blockade is preferable to general anaesthesia in women with asthma due to the potential risk of bronchospasm with certain inhaled anaesthetic agents.
- ✓ Women receiving steroid tablets at a dose exceeding prednisolone 7.5 mg per day for more than two weeks prior to delivery should receive parenteral hydrocortisone 100 mg 6–8 hourly during labour.
- D Use prostaglandin F2α with extreme caution in women with asthma because of the risk of inducing bronchoconstriction.**

12.5 Drug therapy for breastfeeding mothers

The medicines used to treat asthma, including steroid tablets, have been shown in early studies to be safe to use in breastfeeding mothers.⁸⁵⁴ There is less experience with newer agents. Less than 1% of the maternal dose of theophylline is excreted into breast milk.⁸⁵⁴ 2+

Prednisolone is secreted in breast milk, but milk concentrations of prednisolone are only 5–25% of those in serum.⁵²⁷ The proportion of an oral or intravenous dose of prednisolone recovered in breast milk is less than 0.1%.^{855–857} For maternal doses of at least 20 mg once or twice daily the nursing infant is exposed to minimal amounts of steroid with no clinically significant risk.^{855–857} 2+
3

- C Encourage women with asthma to breastfeed.**
- C Use asthma medications as normal during lactation, in line with manufacturers' recommendations.**

13 Occupational asthma

13.1 Incidence

The true frequency of occupational asthma is not known, but underreporting is likely. Published reports, which come from surveillance schemes, compensation registries or epidemiological studies, estimate that occupational asthma may account for about 9–15% of adult onset asthma.⁸⁵⁸⁻⁸⁶⁰ It is now the commonest industrial lung disease in the developed world with over 400 reported causes.⁸⁶¹⁻⁸⁶³ 2++

The diagnosis should be suspected in all adults with symptoms of airflow limitation, and positively searched for in those with high-risk occupations or exposures. Patients with pre-existing asthma aggravated non-specifically by dust and fumes at work (work-aggravated asthma) should be distinguished from those with pre-existing asthma who become additionally sensitised to an occupational agent.

B In patients with adult onset, or reappearance of childhood asthma, healthcare professionals should consider that there may be an occupational cause.

13.2 At-risk populations

Several hundred agents have been reported to cause occupational asthma and new causes are reported regularly in the medical literature.

The most frequently reported causative agents include isocyanates, flour and grain dust, colophony and fluxes, latex, animals, aldehydes and wood dust.⁸⁶⁴⁻⁸⁷² 2++

The workers most commonly reported to occupational asthma surveillance schemes include paint sprayers, bakers and pastry makers, nurses, chemical workers, animal handlers, welders, food processing workers and timber workers.^{864, 865, 867, 869-875} 2++

Workers reported to be at increased risk of developing asthma include bakers, food processors, forestry workers, chemical workers, plastics and rubber workers, metal workers, welders, textile workers, electrical and electronic production workers, storage workers, farm workers, waiters, cleaners, painters, dental workers and laboratory technicians.⁸⁷⁶⁻⁸⁷⁹ 2+

13.3 Diagnosis

Occupational asthma should be considered in all workers with symptoms of airflow limitation (*see Annex 10*). The best screening question to ask is whether symptoms improve on days away from work. This is more sensitive than asking whether symptoms are worse at work, as many symptoms deteriorate in the hours after work or during sleep. The use of non-leading questions is advocated.⁸⁸⁰ Asthma symptoms reported by the use of a questionnaire to be better on days away from work have been shown to have a sensitivity of 58–100% for subsequently validated occupational asthma and specificities of 45–100%, with wheeze and shortness of breath the symptoms most commonly reported.⁸⁸¹ There is also some evidence that free histories taken by experts may have a higher sensitivity than patient questionnaires administered by experts, but their specificity may be lower for a diagnosis of occupational asthma.⁸⁸¹ 4

One study notes a relatively low positive predictive value of work related symptoms.⁸⁸² 3

- ✓ Adults with suspected asthma or unexplained airways obstruction should be asked:
- Are you the same, better, or worse on days away from work?
 - Are you the same, better, or worse on holiday?
- Those with positive answers should be investigated for occupational asthma.

Occupational asthma can be present when tests of lung function are normal, limiting their use as a screening tool. Asthmatic symptoms improving away from work can produce false negative diagnoses, so further validation is needed.

Serial measurement of peak respiratory flow is the most readily available initial investigation, and the sensitivity and specificity of serial peak-flow measurement in the diagnosis of occupational asthma are high.⁸⁸³⁻⁸⁹⁰ 3

Although skin-prick tests or blood tests for specific IgE are available, there are few standardised allergens commercially available which limits their use. A positive test denotes sensitisation, which can occur with or without disease. The diagnosis of occupational asthma can usually be made without specific bronchial provocation testing, considered to be the gold standard diagnostic test. The availability of centres with expertise and facilities for specific provocation testing is very limited in the UK and the test itself is time consuming.

As a general observation, the history is more useful in excluding occupational asthma than in confirming it. A significant proportion of workers with symptoms that improve on days away from work or on holiday have been shown by objective tests not to have occupational asthma.⁸⁹¹ 3

D In suspected work-related asthma, the diagnosis of asthma should be confirmed using standard objective criteria.

13.3.1 Sensitivity and specificity of serial peak-flow measurements

In a meta-analysis of 31 studies in which a variety of reference standards were used, the pooled sensitivity and specificity of serial PEF measurements were 75% and 79% respectively. Higher values (82% and 88%) were obtained from pooling studies where more complete series of measurements had been made, achieved by 61% of the analysed population. Visual analysis was more sensitive (78% v 71%) but less specific (69% v 91%) than computer-based methods.⁸⁹⁰ 2+

There are several validated methods for interpreting serial PEF records for a diagnosis of occupational asthma which differ in their minimal data requirements. The original discriminant analysis method requires:

- at least three days in each consecutive work period
- at least four evenly spaced readings per day
- at least three series of consecutive days at work with three periods away from work (usually about three weeks).⁸⁹²

Shorter records without the requirement for three consecutive days at work can be analysed using the area between curves score. This requires at least eight readings a day on eight work days and three rest days.⁸⁹³ A statistical method using the addition of timepoint analysis requires the waking time to be similar on rest and work days.⁸⁹⁴ 3

The analysis is best done with the aid of a criterion-based expert system. Suitable record forms and support are available from www.occupationalasthma.com

D Objective diagnosis of occupational asthma should be made using serial peak-flow measurements, with at least four readings per day.

13.3.2 Diagnosis of validated cases of occupational asthma using IgE testing

A review by the British Occupational Health Research Foundation states that, "...the respective sensitivities and specificities of the ability of skin-prick or serological tests to detect specific IgE vary between allergens and depend on the setting of positive cut-offs".⁸⁸¹ The sensitivities and specificities of serum-specific IgE antibodies to low molecular weight agents depends on whether the antibodies have been properly characterised and the availability of appropriate hapten conjugates. The presence of specific IgE confirms sensitisation but alone does not confirm the presence of occupational asthma, nor necessarily its cause.⁸⁸¹ The review concluded that skin-prick testing or tests for specific IgE should be used in the investigation of occupational asthma caused by high molecular weight agents but are less sensitive for detecting specific IgE and occupational asthma caused by low molecular weight agents. In neither case are the tests specific for diagnosing asthma.⁸⁸¹

D Skin-prick testing or tests for specific IgE should be used in the investigation of occupational asthma caused by high molecular weight agents.

D Skin-prick testing or tests for specific IgE should not be used in the investigation of occupational asthma caused by low molecular weight agents.

13.3.3 Non-specific reactivity

Studies of non-specific reactivity are confounded by the different methods used, different cut-offs for normality and the interval between last occupational exposure and the performance of the test (an increase in time interval may allow recovery of initial hyper-reactors). A single measurement of non-specific reactivity has been shown to have only moderate specificity and sensitivity for the validation of occupational asthma and changes in non-specific reactivity at and away from work alone have only moderate sensitivity and specificity for diagnosis.^{881, 895}

D A single measurement of non-specific reactivity should not be used for the validation of occupational asthma.

13.3.4 Specific bronchial provocation testing

Specific inhalation challenges (SIC) with occupational agents should only be carried out in hospitals with expertise in using occupational agents, and should always include: a control challenge on a separate day; a gradual increase of exposure to the suspected occupational agent; close monitoring of airway calibre during the challenge and for at least six hours after the end of the exposure.⁸⁹⁶ When carrying out specific challenge testing, an increased duration of allergen exposure may increase the overall diagnostic sensitivity of the tests.⁸⁹⁷

A positive SIC is one in which the FEV₁ falls by $\geq 15\%$ from baseline; either within the first hour after exposure (an immediate reaction) or later (a late reaction) or both. Alternatively for late reactions, two measurements below the 95% CI for three days away from exposure have been validated as a positive test.⁸⁹⁸ Equivocal reactions can sometimes be clarified by finding changes in non-specific bronchial responsiveness, sputum eosinophils or exhaled nitric oxide. Specific inhalation challenge is generally a safe procedure; excessive reactions are rare with <3% of patients needing repeated doses of a bronchodilator and steroid treatment.

The sensitivity and specificity of SIC are high but not easily quantified as the method is usually used as the reference standard for the diagnosis of occupational asthma. False negative tests also occur, and SIC testing may be of less value where complex workplace exposures cannot be replicated in the laboratory. SIC remains the gold standard for making a diagnosis of occupational asthma.

13.3.5 Sputum eosinophilia

Eosinophilic bronchial inflammation can be assessed by cell counts in fresh sputum, induced by inhaling hypertonic saline.^{881, 895, 899} Studies have shown that induced sputum eosinophilia is not sufficiently sensitive or specific to help in the diagnosis of occupational asthma although it may help in the interpretation of equivocal SIC reactions.⁸⁸¹ In the clinical setting the absence of sputum eosinophilia does not exclude a diagnosis of occupational asthma.⁸⁸¹

2+
3
4

13.3.6 Exhaled nitric oxide

The 2010 review by the British Occupational Health Research Foundation states that, "...the measurement of exhaled nitric oxide produced by inflammatory and epithelial cells in the respiratory tract is non-invasive and has been studied extensively in non-occupational asthma, although it has not been fully validated as an effective diagnostic test for occupational asthma". The review concluded that the role of exhaled nitric oxide measurements in the diagnosis of occupational asthma in this setting is not established.⁸⁸¹

4

13.3.7 Exhaled breath condensate

Exhaled breath condensate may offer assistance in those undergoing diagnostic testing for occupational asthma. Its definitive utility is not yet understood.^{900, 901}

3

13.4 Management of occupational asthma

The aim of management is to identify the cause, remove the worker from exposure, and for the worker to have worthwhile employment.

Complete avoidance of exposure may or may not improve symptoms and bronchial hyper-responsiveness. Both the duration of continued exposure following the onset of symptoms and the severity of asthma at diagnosis may be important determinants of outcome. Early diagnosis and early avoidance of further exposure, either by relocation of the worker or substitution of the hazard offer the best chance of complete recovery. Workers who remain in the same job and continue to be exposed to the same causative agent after diagnosis are unlikely to improve and symptoms may worsen. The likelihood of improvement or resolution of symptoms or of preventing deterioration is greater in workers who have no further exposure to the causative agent.^{885, 902-910}

2++

Several studies have shown that the prognosis for workers with occupational asthma is worse for those who remain exposed for more than one year after symptoms develop, compared with those removed earlier.⁹¹¹⁻⁹¹³

D Relocation away from exposure should occur as soon as diagnosis is confirmed, and ideally within 12 months of the first work-related symptoms of asthma.

There is consistent evidence from clinical and workforce case series that about one third of workers with occupational asthma are unemployed after diagnosis. It is unclear whether this risk is higher than that for other adults with asthma.⁹¹⁴⁻⁹¹⁶ The risk of unemployment may fall with increasing time after diagnosis.⁹¹⁷ There is consistent evidence that loss of employment following a diagnosis of occupational asthma is associated with loss of income. Adults with occupational asthma may find employment more difficult than adults with non-occupational asthma.^{915, 916} Approximately one third of workers with occupational asthma have been shown to be unemployed up to six years after diagnosis.⁹¹³⁻⁹²¹

2-

14 Organisation and delivery of care

14.1 Care pathways

Clinical care pathways are “...structured multidisciplinary care plans used by health services to detail essential steps in the care of patients with a specific clinical problem. They aim to link evidence to practice and optimise clinical outcomes whilst maximising clinical efficiency.”⁹²²

There is little high-quality evidence from randomised trials addressing the impact of care pathways for asthma. Pathways have usually been implemented through a training session or programme. Two interventions, one to establish pathways for the management of people with high-risk asthma in UK primary care, the other to establish pathways for children with acute and chronic asthma in New Zealand primary care, led to non-significant reductions in ED attendance and hospitalisation.^{923, 924} Pathways for inpatient care can improve processes of care, such as prescription of oral prednisolone and use of written asthma action plans in children,⁹²⁵ and can reduce length of stay for children,^{646, 926} but have not improved follow up in general practice after discharge.⁹²⁷

1+
1-

Further well-conducted studies are needed to define the benefits of care pathways for asthma. These should include large studies suitably powered to clarify the impact of pathways promoting systematic management of people with high-risk asthma in UK primary care, and pathways integrating asthma care across the primary/secondary care interface.

14.2 Educating clinicians

There is strong evidence that educating clinicians can improve health outcomes for patients. Two large Cochrane systematic reviews (covering all clinical conditions, not just asthma) found that:

- educational outreach visits (for example training visits to general practices) lead to small to moderate improvements in outcomes⁹²⁸
- mixed interactive and didactic education is more effective than either alone.⁹²⁹

1++

Several models of clinician education specifically for asthma have been tested in randomised trials and these broadly support the conclusions of the two Cochrane reviews. The most consistently effective of these for asthma comprises educational outreach visits which deliver multifaceted training, based on theoretical models of behaviour change, including training in consultation styles and delivery of key messages. Several studies have tested the American-developed Physician Asthma Care Education (PACE) paediatric asthma programme,^{189, 930} or adaptations of it for Australian and UK practice,^{217, 931} and have shown reductions in ED visits,⁹³⁰ improved symptom control,²¹⁷ and increased use of written asthma action plans.⁹³¹ The PACE intervention has not been tested for adult populations and there is little experience of its use in the UK.

1++
1+

In the USA, peer education comprising intensive training of a ‘practice asthma champion’ who in turn trained and supported colleagues, led to fewer asthma attacks in children.⁹³² Practice asthma champions were trained in pharmacotherapy and physician behaviour change techniques, and received ongoing support for their role as a ‘change agent’. They received guideline summaries, key targets

1+

for their physician colleagues and feedback on their colleagues' performance along with monthly support from a nurse co-ordinator. When this peer education programme was combined with intensively trained outreach nurses implementing patient reviews (the Planned Care Model), children experienced fewer asthma symptoms and fewer asthma attacks.

1+

These interventions illustrate that, to effect change, interventions need to be of sufficient intensity to engage with, and change, the way practices are organised.

Less intensive educational interventions, such as brief outreach visits comprising simple group education are less effective, showing no impact on symptoms, quality of life, or healthcare use.⁹³³⁻⁹³⁶

1+
1-

Remote IT educational interventions, such as remote spirometry training,⁹³⁷ may be effective but have not been widely tested.

Further large-scale studies, carried out in the UK, are needed to test the impact of intensive educational interventions, such as adapted PACE and peer education programmes.

B Training for primary care clinicians should include educational outreach visits using multifaceted programmes that include consultation training including goal setting.

14.3 Asthma clinics

14.3.1 Structured review

Proactive clinical review of people with asthma improves clinical outcomes. Evidence for benefit is strongest when reviews include discussion and use of a written PAAP.¹⁶⁶ Benefits include reduced school or work absence, reduced asthma attack rate, improved symptom control and reduced attendance at the emergency department.^{938, 939} Proactive structured review, as opposed to opportunistic or unscheduled review, is associated with reduced rates of asthma attack and days lost from normal activity.⁹⁴⁰⁻⁹⁴² It is difficult to be prescriptive about the frequency of review as need will vary with the severity of the disease. Outcome is probably similar whether a practice nurse, or a general practitioner conducts the review. Clinicians trained in asthma management achieve better outcomes for their patients.^{941, 943, 944}

1+
2+
3
4

A In primary care, people with asthma should be reviewed regularly by a nurse or doctor with appropriate training in asthma management. Review should incorporate a written action plan.

✓ It is good practice to audit the percentage of patients reviewed annually. Consider focusing on particular groups such as those overusing bronchodilators, patients on higher dose therapies, those with asthma attacks or from groups with more complex needs.

14.3.2 Primary care asthma clinics

Primary care asthma clinics can be defined as a "...proactive system of care sited in primary care (for example GP clinic) which occupies a defined and often regular clinical session for the routine review of patients with asthma".⁹⁴⁵

Within primary care, structured reviews may be delivered as appointments in routine surgeries, or within dedicated asthma clinics.

One systematic review which included three small studies of the asthma clinic model, showed no evidence of improvement in important outcomes such as hospitalisation, ED attendances, or quality of life, although there was a reduction in night-time waking, and no evidence that clinics were cost effective.⁹⁴⁵ The poor quality of the included studies led the review to conclude that there was a lack of evidence to inform the best way to organise structured asthma care in practice. **1++**

There is, however, no evidence that these clinics do harm. Asthma reviews in primary care may best be carried out, however, during routine surgeries rather than a dedicated asthma clinic.

14.3.3 Specialist asthma clinics

The evidence for whether specialist asthma clinics improve outcomes for people with severe or difficult asthma was limited to one systematic review, including 17 studies, many of poor quality and underpowered.¹⁷² The review focused on psychoeducational interventions mostly for adults and adolescents (age 16 or older) with difficult or severe asthma, so provided incomplete evidence on the ideal content of such clinics. The review found that these interventions reduced hospitalisations (but not ED attendances) in adults and children, and improved symptoms in children. The authors concluded that the strength of evidence was insufficient to change practice. **2+**

Further trials testing the impact of clinics run by specialists in asthma care are needed.

C Consider including psychoeducational interventions in clinics for adults and children with difficult asthma.

14.4 Telehealthcare

Telehealthcare is evolving rapidly and terminology is changing and is used inconsistently in the literature and in practice. In this guideline, 'telehealthcare' is used as an overarching term for all technology-enabled healthcare. Within this, telemonitoring implies collection and transfer of patient data; teleconsultation is the use of technology to enable remote consultation between a patient and a clinician; and telemedicine is interprofessional consultation.

14.4.1 Supporting self management

Telehealthcare embraces a range of functionalities which target different aspects of self-management behaviour including automated medication reminders to improve adherence,⁹⁴⁶ educational games to improve knowledge^{170, 181, 266, 947} or effect behavioural change,^{201, 948, 949} and telemonitoring with various levels of professional oversight to support self management.^{212, 950-955} These functions may use different IT modalities (text messaging,^{171, 956} automated telephone calls,²⁶⁷ 'apps',⁹⁵³ computer games,^{170, 181, 266} cloud-based electronic health records,⁹⁵³⁻⁹⁵⁵) and may be delivered in different contexts (primary/community care,^{212, 267, 952, 953} hospital outpatients,⁹⁴⁷ school based^{181, 201, 266, 949}) which may influence their impact. In the fast moving context of telehealthcare, the aim of the intervention and the theoretical underpinning is likely to be more important to interpreting the evidence than details of the mode of delivery. **1++**
1+
1-
2++
2+

Automated reminders to improve adherence

In the short term, and in the context of a clinical trial, automated reminders (delivered by text messaging, alarms, or automated telephone calls) can improve adherence to medication, but do not have an impact on clinical outcomes.⁹⁴⁶ As part of more complex telehealthcare interventions, reminders may contribute to improved adherence to monitoring or medication use.^{171, 954-956}

1+
1-
2++

Computer-based educational games to improve knowledge or affect behaviour

Educational games improved asthma knowledge in most, but not all participants in school-based interventions,^{170, 181, 266} and children attending a UK outpatients clinic.⁹⁴⁷ The latter study showed reduced school absenteeism and the number of steroid courses,⁹⁴⁷ but overall there is an inconsistent effect on clinical outcomes,^{170, 181, 266} and no impact on use of healthcare resources.^{181, 266, 947}

1+
1-
2+

Games based on behavioural change theories have resulted in some improvement in self-management skills, although impact on symptoms and use of healthcare resources is variable. A generic health behaviour game which targeted teenagers with specific behavioural traits (such as rebelliousness, poor emotional support or low self esteem), improved asthma control, reduced absenteeism, and reduced admissions, but did not reduce ED attendances.^{201, 949}

1+

Telemonitoring to support self management

Telemonitoring, the transmission of monitoring data from a patient to an electronic health record which can be shared with (or monitored by) healthcare professionals, is promoted as having the potential to improve outcomes.

Some studies have demonstrated improvement in at least one clinical outcome, such as measures of asthma control,^{952, 955} lung function,²¹² quality of life, reduced risk of activity limitation,²¹² and school absenteeism, exacerbations, and use of unscheduled care.²¹² Other trials, however, have shown no impact on asthma control or use of healthcare resources.^{951, 953}

1++
1+

These interventions are heterogeneous, and the impact of the telemonitoring is likely to be strongly influenced by the demographic context (deprivation status and cultural background^{212, 952}), and the level of professional support provided (frequency of monitoring,^{212, 954} personalisation of feedback,⁹⁵⁴ access to case-management support⁹⁵²). People with poorly-controlled asthma have the potential to gain more by engaging with telemonitoring than those whose control is already optimal.⁹⁵⁵ Telehealthcare-supported self management offered no clinical benefits over care delivered in traditional ways that was already guideline standard.⁹⁵³

1++
1+

Despite the heterogeneous interventions, the overarching findings from the systematic reviews are consistent and show that telehealthcare:

- can improve process outcomes, such as knowledge,^{170, 181, 957} adherence to monitoring,¹⁷¹ self-efficacy/self-management skills,^{181, 948, 957} and increased use of preventer medication,^{946, 956, 957} at least in the short term⁹⁴⁶
- has an inconsistent effect on clinical outcomes, such as symptoms,^{170, 171, 181, 948, 950, 951, 956, 957} SABA use,¹⁷⁰ lung function,^{170, 171, 950, 956, 957} school absenteeism,^{181, 957} activity limitation,^{950, 957} quality of life,^{181, 950, 951, 957} and oral steroid courses⁹⁴⁸
- generally has no effect on unscheduled use of healthcare resources (such as hospitalisations and ED attendances),^{170, 951, 956, 957} out-of-hours consultations,⁹⁵¹ and GP consultations^{951, 957}

1++
1-
2++
2+
2-

- has cost implications relating to providing and supporting telehealthcare services^{171, 951} | 1++
1-
- has no identified harms and whilst the telehealthcare intervention was often no better than usual care, there were no instances in which it was less effective. | 2++
2+
2-

Telehealthcare is a means of delivering care, not a panacea. Overall, clinical outcomes with telehealthcare are at least as good as, though not consistently superior to, traditionally delivered care. Information technology-based approaches may, therefore, be considered where organisational/clinical/social circumstances or clinician and patient preferences or convenience suggest they may be appropriate.

C

Telehealthcare may be considered as an option for supporting self management.

14.4.2 Remote consulting

Remote consulting can be either asynchronous, with information exchanged sequentially, for example via email, text or web, or synchronous, with information exchange by, for example, telephone.

Evidence to support either approach in patients with asthma is very limited. Two systematic reviews of asynchronous remote consulting covering 15 RCTs and 52 other studies, most of them observational, included only four studies addressing asthma, two of them RCTs, one of which was of poor quality.^{958, 959} Although both reviews suggest that asynchronous telehealthcare led to significant reductions in healthcare use and some improvement in disease status (for example HbA1c in diabetes), the evidence relating to asthma is limited and of low quality and no conclusions can be drawn about its effectiveness in this patient group. | 1+

Evidence to support synchronous consulting in patients with asthma is also limited and, in general, did not address major outcomes of importance. Of four RCTs identified,^{213, 267, 960, 961} two were considered to be of low methodological quality.^{213, 267} There is some evidence to suggest that synchronous consulting can lead to improvements in parental QoL,⁹⁶⁰ and equivalent health status to people reviewed in 'traditional' face-to-face consultations.⁹⁶¹ | 1++
1-

14.4.3 Computerised decision support systems

Computerised decision support systems (CDSS) can broadly be divided into systems targeted at healthcare professionals and integrated within the electronic health record, and web-based systems that are used by patients (and their healthcare professionals) to support self management.

A systematic review of eight RCTs considering the impact on asthma control of CDSS used by healthcare practitioners found little effect on patient outcomes because the healthcare practitioners rarely used the CDSS being evaluated and when used, rarely followed the advice given. Future CDSS need to align better with professional workflows so that pertinent and timely advice is easily accessible within the consultation. The authors concluded that integration of CDSS into electronic health records is cumbersome and a major factor in their ineffectiveness.⁹⁶² | 1+

A second review of 19 RCTs concluded that CDSS can improve chronic disease processes and outcomes. This conclusion, however, reflects the inclusion of four | 1++

trials of systems used by patients to promote self management, three of which reported improved asthma control or QoL, although one, with a high risk of bias, improved symptoms and QoL but led to increased unscheduled care.⁹⁶³ 1++

A Computerised decision support systems for patient use can be considered as an approach to supporting self management.

14.5 School-based interventions

Most school-based asthma interventions focus on education delivered by adults (usually healthcare professionals) to school children.¹⁸¹ Other approaches include peer education, whereby students are trained and then, in turn, train their peers,^{772, 964} web-based programmes,²⁰¹ or directly-observed therapy with ICS medication,⁷⁸⁷ which may additionally include education of parents (*see section 11.11.2*).⁹⁶⁵ One study tested a multifaceted intervention combining education of schoolchildren with additional training of their doctor, including provision of self-management plans.⁹⁶⁶ Most evaluations have been based in the USA, often involving minority ethnic groups not directly applicable to the UK.

Education for children in schools generally led to improvements in symptom control and quality of life, but had no impact on healthcare use.¹⁸¹ Peer education was effective for adolescents⁷⁷² but not preteens.⁹⁶⁴ In two studies, directly observed therapy improved symptom control.^{787, 965} Of all the school-based interventions tested, Bruzzese's multifaceted programme had the most impact, improving symptoms, quality of life, emergency department use and hospitalisation.⁹⁶⁶ 1+
1-
2+

B Consider a multifaceted approach to school-based asthma education programmes targeting children's healthcare professionals as well as the children themselves.

14.6 Ethnicity/culture-based interventions

The majority of studies examining ethnicity and culture-based interventions that tailor asthma education for people from minority ethnic groups have been carried out in the USA. Further details on the aspects of tailoring can be found in section 5.3.5.

A review of system-level interventions concluded that the most effective at reducing further healthcare use were those targeted at people who had attended emergency care or had been hospitalised.¹⁸⁶ Interventions were usually intensive, multisession clinic-based programmes. They were nurse led or used experts including pharmacists or allergy specialists.¹⁸⁶ These findings mirror the little work published in the UK, which showed that a clinic based in primary care was ineffective,²⁰⁶ while a specialist nurse-led intervention targeted at those attending emergency care reduced further unscheduled care, albeit less in people from ethnic minority groups than in those from white populations.²⁰⁷ 1+

Further studies examining the impact of interventions on people from minority ethnic groups in the UK are needed. 1+

C Establish intensive clinic-based programmes for people from ethnic minority groups who have attended emergency care.

14.7 Lay-led interventions

Educational interventions led by lay, rather than healthcare professionals, have become popular in the last decade. The NHS Expert Patient Programme, a six-week group education programme, is an example. Programmes are usually generic; people attending may have a range of conditions, not specifically asthma.

A systematic review including 17 RCTs of lay-led self-management education programmes was identified.⁹⁶⁷ Only two of the included trials specifically addressed people with asthma, and these found no improvements in breathlessness, health-related quality of life, healthcare use, days/nights spent in hospital, and no change in disease-specific knowledge. Overall, lay-led self-management interventions may lead to small, short-term improvements in participants' self efficacy, self-rated health, cognitive symptom management, and frequency of aerobic exercise. There is, however, currently no evidence to suggest that these interventions alter healthcare use or are cost effective.

1+

A

Lay-led self-management programmes for people with asthma are not recommended.

14.8 Pharmacist-led interventions

Pharmacists have opportunities to provide education for people with asthma, and furthermore, may be able to identify those with poor asthma control. The body of evidence for pharmacy-based interventions is, however, methodologically weak or of limited relevance. Two systematic reviews were assessed. One review addressed interventions in low and middle income countries, while another addressed pharmacy interventions more generally.^{968, 969}

Interventions generally involved educating community pharmacists to, in turn, educate patients.⁹⁷⁰⁻⁹⁷² Other models or elements included follow-up reviews for newly prescribed medication,⁹⁷³ identifying those with poor control by using questionnaires such as the Asthma Control Test,⁹⁷² searching prescribing databases for patients using large numbers of reliever inhalers,⁹⁷⁴ and targeting reviews or referral to general practitioners.

Overall, the most consistent improvements in outcomes were seen in inhaler technique,⁹⁷⁰⁻⁹⁷² with a few studies showing improvements in reduced dispensing of, or need for, reliever inhalers.^{972, 974} There was no convincing evidence of reduction in healthcare use.

1+

1-

Further high-quality randomised trials testing pharmacist-led interventions to improve asthma outcomes are needed.

✓

Consider training pharmacists to provide education for people with asthma.

15 Provision of information

The provision of accurate information to patients and carers is of great importance in order to achieve good adherence to treatment and improved patient outcomes. Specific recommendations and good practice points relating to provision of information by healthcare professionals to patients and carers are found throughout this guideline. In addition, supported self management is covered in detail in section 5, including sections on personalised asthma action plans (*see section 5.2.2 and Table 11*) and adherence and concordance (*see section 5.4*).

Patient versions of this guideline, in booklet form, covering the management of asthma in adults (for patients and their families and carers) and the management of asthma in children (for parents and carers) are available on the SIGN website www.sign.ac.uk, (*see section 15.2*) or directly from SIGN and could be a useful addition to the patient's PAAP. Healthcare professionals are encouraged to inform patients and carers that these booklets are available. The patient versions are reviewed and updated in line with the clinical guideline. In addition to information on care and treatment, the booklets include contact details for, and brief information about, a number of organisations that provide information for patients (*see section 15.3*).

15.1 Checklist of information for patients and carers

This section gives examples of the information patients/carers may find helpful at the key stages of the patient journey. The checklist was designed by members of the guideline development group based on their experience and their understanding of the evidence base. In developing the checklist, consideration was given to what patients and carers valued. The checklist is neither exhaustive nor exclusive.

Assessment and diagnosis

- Fully explain symptoms and triggers, giving examples to help. Ask the patient questions to ensure they understand.
- Explain to the patient that diagnosing and managing asthma is not straightforward, and that they might be trying a few different tests and medicines. Explain to the patient that the results of tests or medicine trials may mean more tests and trying different medicines.
- Explain the different tests using clear, concise, jargon-free language. Show the materials that will be used in these tests, for example a spirometer.
- Ensure patients are kept informed about which tests will be performed, when they are likely to be carried out, and what the results mean.
- Explain and show equipment (inhalers and spacers), how it is used, how often the patient should use it, where they will get these from.
- Encourage patients and their families to discuss their questions and concerns during appointments and reviews. This will help patients to get the most from their appointments or reviews.

Ongoing care (monitoring)

- Advise patients and their families of the need to work in partnership with them to allow a holistic approach to managing their asthma.
- Ensure the patient is aware that they do not have to wait until their regular review if they have concerns that they need to discuss with their healthcare professional sooner.
- Encourage people to take a notebook to appointments to allow them to record key information.
- Offer a summary of discussions at the end of every appointment and check the patient's understanding.
- Ensure appropriate information is given to patients to encourage them to take responsibility for their asthma, for example making sure that they are familiar with personal asthma action plans and filling these in with them if they do not have one.
- Be sensitive to and aware of how culture and beliefs affect a patient's asthma and lifestyle. For example offer action plans in different language as appropriate.
- Listen carefully to the needs and priorities of patients and carers.
- Explain what happens if the patient reaches a crisis point of an asthma attack.

Medicines (pharmacological management)

- Inform patients of side effects from medication when prescribing and reviewing medication and reassure them that these are normal. Listen to any concerns.
- Explain in clear, jargon-free language, any new medicines, and reasons for changing medicines.
- Check and optimise inhaler technique.

Non-pharmacological management

- Remain open minded and open to discussing things that may help manage symptoms alongside medicines. Different things might help different people.

Self management

- Ask patients to think about asthma triggers, for example perfumes, cleaning products, smoke, etc.
- Ask patients what they do to help them to manage their asthma, for example do they keep a diary, notebook, use an app, peak-flow meter, etc.
- Provide and explain a personal asthma action plan (PAAP).

Asthma attacks

- Introduce yourself.
- Discuss with the patient their personal asthma action plan before they leave hospital.
- Discuss with the patient and their family or carer what happens after they leave the hospital, for example explain that they need to make an appointment with their doctor or asthma nurse.

Asthma in pregnancy
<ul style="list-style-type: none"> • Communicate with the labour team to ensure they are aware of any at-risk patients. • Discuss with the patient any changes to their asthma action plan and make sure the patient understands any changes.
Asthma in young people
<ul style="list-style-type: none"> • Involve children and young people from the start and encourage them to take responsibility for managing their asthma. Listen to and address their needs fully and ask children and young people the following questions. <ul style="list-style-type: none"> - Have you had any asthma attacks? What were you doing at the time? - Have you been breathless? - Have you been taking your medication? If not, what were the reasons for this?
Work-related asthma
<ul style="list-style-type: none"> • Explain that people may find they have issues at work, for example triggers may be present. • Discuss what can be done to help at work.

15.2 Publications from SIGN

SIGN patient versions of guidelines are documents that 'translate' guideline recommendations and their rationales, originally developed for healthcare professionals, into a form that is more easily understood and used by patients and the public. They are intended to:

- help patients and carers understand what the latest evidence supports around diagnosis, treatment and self care
- empower patients to participate fully in decisions around management of their condition in discussion with healthcare professionals
- highlight for patients where there are areas of uncertainty.

The following SIGN patient versions for asthma can be accessed at:
www.sign.ac.uk/patient-publications.html

- Managing asthma in adults: a booklet for adults, partners, friends, family members and carers. (SIGN 2019)
- Managing asthma in children: a booklet for parents, carers and family members. (SIGN 2019)
- Asthma in pregnancy: a booklet for women, partners, friends, family members and carers. (SIGN 2019)

15.3 Sources of further information

15.3.1 National organisations for people who have asthma

Asthma UK

18 Mansell Street, London, E11 8AA

Asthma UK's Helpline: 0300 222 5800 (9am-5pm, Mon-Fri) – nurses provide advice for people with asthma and for healthcare professionals.

General enquiries: info@asthma.org.uk

Tel: 0300 222 5800

www.asthma.org.uk

Asthma UK is a charity dedicated to improving the health and wellbeing of people who are affected by asthma. The charity provides a wide range of information and resources on their website, including downloadable asthma action plans. Printed information booklets and other resources are available on request and bulk copies are available for purchase by healthcare professionals.

British Lung Foundation

73–75 Goswell Road, London, EC1V 7ER

Helpline: 08458 50 50 20

Tel: 020 7688 5555

www.blf.org.uk

The British Lung Foundation aims to help people understand and live with lung disease. They run the Breathe Easy support network which offers information, support and friendship to anyone affected by lung disease.

15.3.2 Other organisations

Allergy UK

Planwell House, Lefa Business Park, Edginton Way, Sidcup, Kent, DA14 5BH

Helpline: 01322 619898

www.allergyuk.org

Allergy UK is a charity which aims to increase people's understanding and awareness of allergies, and helps people manage their allergies.

ASH (Action on Smoking and Health)

First Floor, 144–145 Shoreditch High Street, London, E1 6JE

Tel: 020 7739 4732

www.ash.org.uk

ASH is the leading voluntary organisation campaigning for effective tobacco-control legislation and providing an expert information service.

NHS 111

Freephone: 111

This is a 24-hour helpline for people in England and Wales. It is led by nurses who provide confidential healthcare advice and information 24 hours, 365 days a year.

NHS 24

Freephone: 111
www.nhs24.scot

This is a 24-hour helpline for people in Scotland. It is led by nurses who provide confidential healthcare advice and information 24 hours, 365 days a year.

Department of Work and Pensions (DWP)

www.dwp.gov.uk

The website gives details of state benefits patients may be entitled to.

16 The evidence base

16.1 Systematic literature review

The evidence base for this guideline was synthesised in accordance with SIGN methodology. A systematic review of the literature was carried out using an explicit search strategy devised by a SIGN Evidence and Information Scientist. Databases searched include Medline, Embase, Cinahl, PsycINFO and the Cochrane Library. Internet searches were carried out on various websites. The main searches were supplemented by material identified by individual members of the development group. Each of the selected papers was evaluated by two members of the group using standard SIGN methodological checklists before conclusions were considered as evidence.

The evidence base builds on the reviews carried out for the original (2003) version of the guideline and subsequent updates. Annex 2 provides details of the time period covered for each topic. A copy of the search narrative, including listings of strategies, is available on the SIGN website as part of the supporting material for this guideline.

16.2 Recommendations for research

The guideline development group was not able to identify sufficient evidence to answer all of the key questions asked in this update of the guideline (*see Annex 1*). The following areas for further research have been identified:

- Clinical prediction models for quantifying risk need to be developed and prospectively validated in adults, children aged 5–12 and children under five years of age. Does risk assessment based on these factors improve outcomes when used prospectively in routine clinical practice?
- In monitoring asthma, what level of risk is associated with factors where evidence is currently limited or equivocal (*see Tables 9 and 10*)?
- Does incorporation of assessment of risk of future asthma attacks (potentially using a risk score) into routine care improve outcomes?
- What is the utility of FeNO measurement in guiding asthma treatment to improve asthma outcomes, such as reduced asthma attacks or increased asthma control, in different patient groups?
- What is the impact of poverty, urban/rural living and ethnicity on asthma outcomes in the UK setting?
- In children under five years of age, what factors are associated with increased risk of acute asthma/wheezing attacks? Do risk factors in this age group differ from those in older children?
- What features of available apps lead to improvements in adherence to medication and which have any impact on clinical outcomes?
- Which approaches to improving medication adherence are most effective and sustainable in patients with asthma?
- How effective are house dust mite and other allergen reduction measures in asthma? A systematic review/meta-analysis is required including only high-quality trials that i) use interventions that are documented to reduce allergen exposure, ii) follow up participants for a sufficient time for important clinical

outcomes to become apparent, iii) provides separate analyses for children and adults, and iv) accounts for any changes in asthma medication over the course of the trial.

- What are the potential beneficial effects of vitamin D supplementation in people with asthma, particularly children and people with frequent severe asthma exacerbations, with different baseline vitamin D levels?
- How effective are breathing exercises in children with asthma?
- What components of individualised multicomponent allergen reduction strategies are effective at improving asthma control and reducing exacerbations?
- Do strategies to reduce environmental allergens improve asthma control and reduce exacerbations in specific subgroups of people with asthma, eg children?
- How effective is montelukast in patients without allergic rhinitis and/or atopic dermatitis?
- Development of an agreed universal definition of 'asthma exacerbation' to allow comparison of this outcome between studies.
- Classification of asthma-related and non-asthma related adverse events to allow comparison of adverse events between studies.
- Which, if any, subgroups of children benefit most from addition of LTRA as compared with LABA as additional add-on therapy to ICS alone?
- What are the short- and long-term steroid-sparing effects of monoclonal antibody therapies in adults and children on different treatment regimens?
- Does the effectiveness of treatment with monoclonal antibodies decrease over time and/or does clinically relevant antibody sensitisation occur, and if so, at what point does/do these occur?
- What markers of response are there to enable targeting of monoclonal antibody therapy?
- Does suppression of IgE or IL-5 have any long-term effects on the recipient's immune function?
- What is the short- and long-term effectiveness and safety of subcutaneous and sublingual immunotherapy in asthma in studies with optimal design and patient-centric endpoints, such as asthma control and exacerbations? Does effectiveness differ between different products or between patients with different characteristics?
- Which patients with asthma might benefit most from bronchial thermoplasty and what are the long-term outcomes and safety of this treatment?
- What is the place of bronchial thermoplasty in the management of severe asthma compared with other options such as biological treatments?
- What is the relative clinical effectiveness and safety of bronchial thermoplasty compared with monoclonal antibody treatments?
- What is the role of non-invasive ventilation and high-flow oxygen therapy in treating children with severe exacerbations of asthma, and what is their effect on measurable outcomes including respiratory parameters, physiological variables and blood gases?
- In considering treatment with extracorporeal membrane oxygenation (ECMO) what is the definition of life-threatening or standard care?
- What is the clinical effectiveness and safety of ECMO treatment in patients with asthma taking anticoagulants?

17 Development of the guideline

17.1 Introduction

SIGN is a collaborative network of clinicians, other healthcare professionals and patient organisations and is part of Healthcare Improvement Scotland. SIGN guidelines are developed by multidisciplinary groups of practising clinicians using a standard methodology based on a systematic review of the evidence. Further details about SIGN and the guideline development methodology are contained in 'SIGN 50: A guideline developer's handbook', available at www.sign.ac.uk. This guideline was developed according to the 2011 edition of SIGN 50.

SIGN and BTS have worked in partnership since 2001 to produce the British Guideline on the Management of Asthma. Governance arrangements including a Memorandum of Understanding between SIGN and BTS approved by Healthcare Improvement Scotland, SIGN Council and the BTS Board of Trustees, are in place. These arrangements cover production of each update and appointment of members to the overall Guideline Development Group.

17.2 Guideline development group

Dr James Paton (Co-chair)	Clinical Reader and Honorary Consultant Paediatrician, Royal Hospital for Children, Glasgow
Dr John White (Co-chair)	Consultant Respiratory Physician, York District Hospital
Mr Joe Annandale	Respiratory Nurse Specialist, Prince Philip Hospital, Llanelli
Dr Anne Boyter	Senior Lecturer, Strathclyde Institute of Pharmacy and Biomedical Sciences, Glasgow
Ms Juliet Brown	Evidence and Information Scientist, Healthcare Improvement Scotland
Ms Beatrice Cant	Programme Manager, SIGN Executive
Dr Toby Capstick	Consultant Pharmacist, St James' University Hospital, Leeds
Dr Richard Chavasse	Consultant in Respiratory Paediatrics, St George's Hospital, London
Dr Luke Daines	Academic Clinical Fellow, Asthma UK Centre for Applied Research, Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, and General Practitioner, Craiglockhart Medical Group, Edinburgh
Dr Andrew Deacon	ST4 Adult Respiratory Medicine, Manchester Royal Infirmary
Dr Rebecca Devaney	ST6 Paediatric Respiratory Medicine, Queen's Medical Centre, Nottingham
Dr Sinan Eccles	Consultant Adult Respiratory Medicine, Royal Glamorgan Hospital, Pontyclun

Professor David Fishwick	Consultant Respiratory Physician, Brearley Chest Clinic, Northern General Hospital, Sheffield
Ms Karen Gibson	Asthma Nurse Specialist, Norfolk and Norwich University Hospital, Norwich
Mrs Toni Gibson	Lay Representative
Professor Chris Griffiths	Professor of Primary Care, Centre for Primary Care and Public Health, London
Dr Nicola Littlewood	Consultant in Emergency Medicine, Queen Elizabeth University Hospital, Glasgow
Dr David Lo	ST8 Paediatric Respiratory Medicine, Leicester Royal Infirmary
Dr Kenneth MacLeod	Consultant in Paediatric Respiratory Medicine, Royal Hospital for Sick Children, Edinburgh
Dr Alexander Mathioudakis	NIHR Academic Clinical Fellow and Honorary Lecturer in Respiratory Medicine, University of Manchester
Ms Tina Morrow	Lay representative
Dr Rob Niven	Senior Lecturer in Respiratory Medicine, Withenshaw Hospital, Manchester
Dr Rebecca Normansell	Joint Co-ordinating Editor Cochrane Airways, Population Health Research Institute, St George's, University of London
Professor Hilary Pinnock	Professor of Primary Care Respiratory Medicine, Asthma UK Centre for Applied Research, Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, and General Practitioner, Whitstable Medical Practice, Kent
Professor Graham Roberts	Professor and Honorary Consultant Paediatrician, University of Southampton
Dr Stephen Scott	Consultant in Respiratory Medicine, Countess of Chester Hospital, Chester
Dr Diana Slim	Respiratory Registrar, Bristol Royal Infirmary
Mrs Lynne Smith	Evidence and Information Scientist, Healthcare Improvement Scotland
Dr Fatimazahra Tharoo	General Practitioner, Birmingham
Professor Steve Turner	Professor and Honorary Consultant Paediatrician, Department of Child Health, University of Aberdeen
Ms Sally Welham	Deputy Chief Executive, British Thoracic Society
Dr Sarah Winfield	Consultant Obstetrician, Leeds Teaching Hospitals NHS Trust
Mr Alex Woodward	Respiratory Physiotherapist, Leicestershire Partnership NHS Trust

The membership of the guideline development group was confirmed following consultation with the member organisations of SIGN. All members of the guideline development group made declarations of interest. A register of interests is available in the supporting material section for this guideline at www.sign.ac.uk

Guideline development and literature review expertise, support and facilitation were provided by the SIGN Executive.

All members of the SIGN Executive make yearly declarations of interest. A register of interests is available on the contacts page of the SIGN website www.sign.ac.uk

Euan Bremner	Project Officer
Karen Graham	Patient Involvement Advisor
Kirsty Allan	Administration Officer
Domenico Romano	Publications Designer
Gaynor Rattray	Guideline Co-ordinator

17.3 Acknowledgements

SIGN would like to acknowledge the guideline development group responsible for the development of SIGN 153: British guideline on the management of asthma, on which this guideline is based.

SIGN would like to acknowledge the PRISMS group who kindly provided the searches, quality assessment and data extraction for the implementation studies in asthma self-management (*see section 5.5*) based on their systematic review of self-management support interventions for people with long-term conditions conducted as part of a project funded by the National Institute for Health Research Health Services and Delivery Research programme (project number 11/1014/04). (Taylor SJC, Pinnock H, Epiphaniou E, et al. A rapid synthesis of the evidence on interventions supporting self-management for people with long-term conditions. (PRISMS Practical Systematic Review of Self-Management Support for long-term conditions). *Health Serv Deliv Res* 2014;2:54). The considered judgement and recommendations (in section 5.5) were developed by the self-management Evidence Review Group in accordance with SIGN methodology. The views and opinions expressed therein are those of the SIGN/BTS guideline development group and do not necessarily reflect those of the PRISMS authors, NIHR, NHS or the Department of Health.

17.4 Consultation and peer review

17.4.1 Consultation

Selected changes to this guideline were presented for discussion in draft form at the Winter Meeting of the British Thoracic Society in December 2018. All questions and comments raised at the meeting were addressed on the day and were also summarised and considered separately by the guideline development group. The draft guideline was also available on the SIGN and BTS websites for five weeks to allow all interested parties to comment. A total of eighteen organisations and seven individuals submitted formal responses as part of the open consultation. A report of the consultation and peer review comments and responses is available in the supporting material section for this guideline on the SIGN website. All expert referees and other contributors made declarations of interest and further details of these are available on request from the SIGN Executive.

17.4.2 Specialist review

This guideline was also reviewed in draft form by the following independent expert referees, who were asked to comment primarily on the comprehensiveness and accuracy of interpretation of the evidence base supporting the recommendations in the guideline. The guideline group addresses every comment made by an external reviewer, and must justify any disagreement with the reviewers' comments. All expert referees made declarations of interest and further details of these are available on request from the SIGN Executive.

SIGN and BTS are very grateful to these experts for their contribution to the guideline.

Dr Bernard Higgins	Consultant Respiratory Physician, Newcastle Upon Tyne Hospitals NHS Trust
Professor Richard Beasley	Director, Medical Research Institute of New Zealand and Adjunct Professor, University of Otago and Physician, Capital and Coast District Health Board, Wellington

17.4.3 Editorial group

As a final quality-control check, the guideline is reviewed by an editorial group comprising the relevant specialty representatives on SIGN Council and members of the Governance Committee for the BTS/SIGN British guideline on the management of asthma to ensure that the specialist reviewers' comments have been addressed adequately and that any risk of bias in the guideline development process as a whole has been minimised. The editorial group for this guideline was as follows. All members of the editorial group made declarations of interest and further details of these are available on request from the SIGN executive.

Mrs Sheila Edwards	Chief Executive, British Thoracic Society
Dr Luke Howard	Chair, BTS Standards of Care Committee
Dr Roberta James	Programme Lead, SIGN
Dr Karen Ritchie	Head of Knowledge and Information, Healthcare Improvement Scotland
Ms Sally Welham	Deputy Chief Executive, British Thoracic Society

Abbreviations

ACT	Asthma Control Test
ACQ	Asthma Control Questionnaire
AOR	adjusted odds ratio
Apgar score	A number expressing the physical condition of a newborn infant (a score of ten representing the best possible condition).
AQLQ	Asthma Quality of Life Questionnaire
BCG	Bacillus Calmette-Guérin
BDP	beclometasone dipropionate
BHR	bronchial hyper-reactivity
BMI	body mass index
BNF	British National Formulary
BTS	British Thoracic Society
CAM	complementary and alternative medicine
CDSS	computerised decision support systems
CFC	chlorofluorocarbon
C-ACT	Childhood Asthma Control Test
CI	confidence interval
COPD	chronic obstructive pulmonary disease
DPI	dry powder inhaler
ECG	electrocardiogram
ECMO	extracorporeal membrane oxygenation
ED	emergency department
ETS	environmental tobacco smoke
FeNO	fractional exhaled nitric oxide
FEV₁	forced expiratory volume in one second
FVC	forced vital capacity
GMC	General Medical Council
GORD	gastro-oesophageal reflux disease
GP	general practitioner
HbA1c	glycated haemoglobin
HDM	house dust mite
HFA	hydrofluoroalkane
ICS	inhaled corticosteroids
ICU	intensive care unit
IgE	immunoglobulin E
IL5	interleukin-5
IM	intramuscular
IT	information technology
IU	international unit

IV	intravenous
kU/L	kilounits of antibody per litre
kPa	kilopascals
LABA	long-acting β_2 agonist
LAMA	long-acting muscarinic antagonist
LTRA	leukotriene receptor antagonists
MA	marketing authorisation
MART	maintenance and reliever therapy
MDI	metered dose inhaler
MHRA	Medicines and Healthcare products Regulatory Agency
n-3PUFA	omega-3 polyunsaturated fatty acid
NICE	National Institute for Health and Care Excellence
NIV	non-invasive ventilation
NPV	negative predictive value
NRAD	National Review of Asthma Deaths
OR	odds ratio
PAAP	personalised asthma action plan
PACE	Physician Asthma Care Education
PaCO₂	partial arterial pressure of carbon dioxide
PaO₂	partial arterial pressure of oxygen
PAQLQ	Paediatric Asthma Quality of Life Questionnaire
PC₂₀	the provocative concentration of bronchoconstrictor (eg methacholine) required to cause a 20% fall in FEV ₁
PD₂₀	the provocative dose of bronchoconstrictor (eg methacholine) required to cause a 20% fall in FEV ₁
PEF	peak expiratory flow
pMDI	pressurised metered dose inhaler
ppb	parts per billion
PPV	positive predictive value
QoL	quality of life
RCT	randomised controlled trial
RR	risk ratio
SABA	short-acting β_2 agonist
SCIT	subcutaneous immunotherapy
SIC	specific inhalation challenge
SIGN	Scottish Intercollegiate Guidelines Network
SLIT	sublingual immunotherapy
SMC	Scottish Medicines Consortium
SpO₂	oxygen saturation measured by a pulse oximeter
TNF	tumour necrosis factor
V_Emax	ventilation at maximal exercise capacity

Annex 1

Key questions addressed in this update

This guideline is based on a series of structured key questions that define the target population, the intervention, diagnostic test, or exposure under investigation, the comparison(s) used and the outcomes used to measure efficacy, effectiveness, or risk. These questions form the basis of the systematic literature search.

Section	Key question
4.2, 4.4	<p>1. In people with asthma (<5, 5–12, >12) which marker/s is/are most effective for monitoring current asthma control?</p> <p>Population: people with asthma</p> <p>Interventions: symptom scores, physiological measurements (lung function tests, bronchial reactivity/airway challenge), other markers: exhaled air (eg FeNO), sputum, blood (eg eosinophil count), periostin</p> <p>Comparisons: as for interventions</p> <p>Outcomes: current asthma control</p>
4.3	<p>2. In people with asthma (<5, 5–12, >12), which individual, or combination of, characteristic/s effectively predict/s future loss of control and/or future risk of attacks?</p> <p>Population: people with asthma</p> <p>Interventions: symptom pattern, asthma control, asthma severity, previous history of attacks, atopy (including sensitisation, comorbid allergic conditions, family history), treatment adherence, behaviours (including smoking), social deprivation, biomarkers, polypharmacy</p> <p>Comparisons: none</p> <p>Outcomes: number of asthma attacks, frequency of asthma attacks</p>
5.2.3	<p>3. In people with asthma, at the onset of an asthma attack and as part of a self-management plan, does increasing the dose of ICS or adding an LTRA, compared to usual care, reduce the severity of or abort the asthma attack or improve symptom scores?</p> <p>Population: people with asthma</p> <p>Interventions: increasing ICS above current dose, adding LTRA</p> <p>Comparisons: usual care</p> <p>Outcomes: change in asthma control, aborting asthma attack, change in severity of asthma attack, avoidance of need for oral steroids</p>

6.2	4. What interventions (avoidance or reduction of exposure to environmental factors) in the home/school/outdoor environment improve asthma control and prevent or reduce severity of asthma attacks?
	<p>Population: people with asthma</p> <p>Interventions: avoidance of exposure to environmental factors, reduction of exposure to environmental factors, eg use of mattress covers for house dust mites</p> <p>Comparisons: no intervention to reduce exposure to environmental factors.</p> <p>Outcomes: asthma symptom control, number of asthma attacks, severity of asthma attacks</p>
6.2.14	5. In people aged 12 and over with asthma, is breathing training in addition to usual care effective at reducing asthma attacks, improving symptoms, reducing side effects, improving treatment adherence or improving lung function?
	<p>Population: people with asthma aged 12 and over</p> <p>Interventions: breathing training</p> <p>Comparisons: no breathing training (ie usual care)</p> <p>Outcomes: asthma attacks, asthma symptom control, adverse side effects, treatment adherence, lung function</p>
7.3	6. In people with asthma whose symptoms are not adequately controlled (poor control and/or frequent attacks) by low-dose (>12 years) or very low-dose (<5, 5–12 years) ICS, which initial add-on therapy is most effective at reducing asthma attacks, improving symptoms, reducing side effects, improving treatment adherence or improving lung function?
	<p>Population: people with asthma taking low-dose (>12 years) or very low-dose (<5, 5–12 years) ICS</p> <p>Interventions: increasing ICS above low/very-low dose, LABA, LTRA, LAMA, theophylline, slow-release β_2 agonist tablets</p> <p>Comparisons: increasing ICS above low/very-low dose, LABA, LTRA, LAMA, theophylline, slow-release β_2 agonist tablets</p> <p>Outcomes: asthma attacks, asthma symptom control, adverse side effects, treatment adherence, lung function</p>
7.4, 7.5	7. In people with asthma whose symptoms are not adequately controlled by low-dose (>12 years) or very low-dose (<5, 5–12 years) ICS plus a LABA, is adding an LTRA, LAMA, theophylline or slow-release β_2 agonist tablets, more effective than increasing the dose of ICS at reducing asthma attacks, improving symptoms, reducing side effects, improving treatment adherence or improving pulmonary/lung function?
	<p>Population: people with asthma taking low-dose (>12 years) or very low-dose (<5, 5–12 years) ICS plus a LABA</p> <p>Interventions: LTRA, LAMA, theophylline, slow-release β_2 agonist tablets</p> <p>Comparisons: increasing ICS dose above low-dose (>12 years) or very low-dose (<5, 5–12 years)</p> <p>Outcomes: asthma attacks, asthma symptom control, adverse side-effects, treatment adherence, pulmonary/lung function</p>

7.5.4	<p>8. In people with asthma who are not adequately controlled on high-dose ICS plus LABA or on oral corticosteroids, does addition of monoclonal antibodies (eg omalizumab, mepolizumab, reslizumab) reduce use of oral steroids, unscheduled care, side effects, or improve symptoms, treatment adherence or lung function?</p>
	<p>Population: people with asthma inadequately controlled on high-dose ICS plus a LABA or on oral corticosteroids</p> <p>Interventions: monoclonal antibodies (including omalizumab, mepolizumab, reslizumab, etc)</p> <p>Comparisons: no use of monoclonal antibodies</p> <p>Outcomes: reduction in unscheduled care (reduced asthma attacks, reduced steroid courses, reduced unplanned hospital/GP visits), adverse side effects, asthma symptom control, lung function, treatment adherence, reductions in treatment (ICS reduction, oral steroid reduction)</p>
7.5.7	<p>9. In people with asthma who are not adequately controlled on high-dose ICS plus a LABA or oral corticosteroids, does addition of bronchial thermoplasty reduce use of oral steroids, unscheduled care, side effects, or improve symptoms, treatment adherence or lung function?</p>
	<p>Population: people with asthma inadequately controlled on high-dose ICS plus a LABA or on oral corticosteroids</p> <p>Interventions: bronchial thermoplasty</p> <p>Comparisons: no bronchial thermoplasty</p> <p>Outcomes: reduction in unscheduled care (reduced asthma attacks, reduced steroid courses, reduced unplanned hospital/GP visits), adverse side effects, asthma symptom control, lung function, treatment adherence, reductions in treatment (ICS reduction, oral steroid reduction)</p>
7.5.6	<p>10. In people with asthma who are poly- or mono-sensitised, is sublingual immunotherapy compared to standard therapy effective at reducing asthma attacks, improving asthma control, improving treatment adherence or improving lung function?</p>
	<p>Population: people with asthma mono- or poly-sensitised</p> <p>Interventions: sublingual immunotherapy (SLIT)</p> <p>Comparisons: standard therapy</p> <p>Outcomes: asthma attacks, asthma symptom control, treatment adherence, lung function</p>
5.4.3	<p>11. What interventions in the home or workplace/school improve adherence with asthma treatments?</p>
	<p>Population: people with asthma</p> <p>Interventions: inhaler timers, inhaler alarms, directly observed therapy (by video, at school, in primary/secondary care settings)</p> <p>Comparisons: usual care</p> <p>Outcomes: asthma symptom control, number of asthma attacks, severity of asthma attacks</p>

9.3.12, 9.9.5	12. In the immediate treatment of people with life-threatening or near-fatal asthma, does extracorporeal membrane oxygenation (ECMO) or other potentially life-saving therapies, compared to usual care, improve patient survival or other outcomes?
	<p>Population: people experiencing a life-threatening or near-fatal asthma attack</p> <p>Interventions: extracorporeal membrane oxygenation (ECMO), ketamine, other rescue therapies</p> <p>Comparisons: usual care</p> <p>Outcomes: survival, morbidity</p>
6.2.8	13. In people with asthma, is supplementation with vitamin D compared to placebo effective at reducing asthma attacks, reducing side effects or improving lung function?
	<p>Population: people with asthma</p> <p>Interventions: vitamin D supplementation</p> <p>Comparisons: usual care</p> <p>Outcomes: asthma attacks, side effects, lung function</p>

Annex 2

Summary of search histories by section

This guideline is based on a series of structured key questions that define the target population, the intervention, diagnostic test, or exposure under investigation, the comparison(s) used and the outcomes used to measure efficacy, effectiveness, or risk. These questions form the basis of the systematic literature search.

Literature searches to support the various sections of this guideline are conducted on a rolling basis. This summary indicates the currency of the searches supporting each section. Searches in all databases began with the earliest year available at that time, which varied from database to database; for example, searches in Embase extended back to 1980 and in CINAHL to 1982. Specific date coverage is provided for Medline.

The 2019 revision saw updating of multiple sections of the guideline identified as priority areas by the guideline development group. Literature searches were conducted in Medline, Embase, CINAHL and the Cochrane Library for all topics to identify systematic reviews published between 2012 and March 2018. Additional literature search coverage for the specific topics considered in this update is described below.

Detailed search strategies are available on the SIGN website in the supplementary material section.

Section 4 Monitoring asthma

Monitoring current asthma symptom control

A broad search was carried out in May 2018 covering 2014–2018. No study design filter was applied.

Predicting future risk of asthma attacks

A broad search was carried out in May 2018 with no date limit. No study design filter was applied.

Section 5 Supported self management

Components of a self-management programme

A broad search was carried out in April 2018 with no date limit to identify studies which looked at people with asthma increasing the dose of ICS or adding an LTRA, compared to usual care, at the onset of an asthma attack and as part of a self-management plan. No study design filter was applied.

Section 6 Non-pharmacological management

Secondary non-pharmacological prevention

A broad search was carried out to identify studies which looked at what interventions (avoidance or reduction of exposure to environmental factors) in the home/school/outdoor environment improve asthma control and prevent or reduce severity of asthma attacks. The search covered 2014–2018 on Medline. No study design filter was applied.

A search was conducted in April 2018 to identify studies on breathing training. The search covered 2013–2018 in Medline, Embase, the Cochrane Library and CINAHL. An RCT filter was applied.

A search was conducted in May 2018 to identify studies on vitamin D supplementation. The search covered 2016–2018 in Medline, Embase, the Cochrane Library and CINAHL. An RCT filter was applied.

Section 7 Pharmacological management


The 2019 revision updated searches for inhaled steroids, long-acting β_2 agonists, theophyllines, leukotriene receptor antagonists, frequency and dose of inhaled steroids, monoclonal antibodies, sublingual immunotherapy and bronchial thermoplasty.

The Cochrane Library, Medline and Embase were searched from 2012–2018. SIGN systematic review and RCT filters were applied.


Section 9 Management of acute asthma

Broad searches were carried out in May/June 2018 with no date limit to identify studies which looked at extracorporeal membrane oxygenation (ECMO) or other potentially life-saving therapies for people with life-threatening or near-fatal asthma. No study design filter was applied.

Annex 3



Annex 4



Annex 5

Management of acute asthma in adults in hospital

Features of acute severe asthma

- Peak expiratory flow (PEF) 33–50% of best (use % predicted if recent best unknown)
- Can't complete sentences in one breath
- Respiration ≥ 25 breaths/min
- Pulse ≥ 110 beats/min

Life-threatening features

- PEF $< 33\%$ of best or predicted
- SpO₂ $< 92\%$
- Silent chest, cyanosis, or poor respiratory effort
- Arrhythmia or hypotension
- Exhaustion, altered consciousness

If a patient has any life-threatening feature, measure arterial blood gases. No other investigations are needed for immediate management.

Blood gas markers of a life-threatening attack:


- 'Normal' (4.6–6 kPa, 35–45 mmHg) PaCO₂
- Severe hypoxia: PaO₂ < 8 kPa (60 mmHg) irrespective of treatment with oxygen
- A low pH (or high H⁺)

Caution: Patients with severe or life-threatening attacks may not be distressed and may not have all these abnormalities. The presence of any should alert the doctor.

Near-fatal asthma

- Raised PaCO₂
- Requiring mechanical ventilation with raised inflation pressures

Peak Expiratory Flow Rate - Normal Values



IMMEDIATE TREATMENT

- Oxygen to maintain SpO₂ 94–98%
- β_2 bronchodilator (salbutamol 5 mg) via an oxygen-driven nebuliser
- Ipratropium bromide 0.5 mg via an oxygen-driven nebuliser
- Prednisolone tablets 40–50 mg or IV hydrocortisone 100 mg
- No sedatives of any kind
- Chest X-ray if pneumothorax or consolidation are suspected or patient requires mechanical ventilation

IF LIFE-THREATENING FEATURES ARE PRESENT:

- Discuss with senior clinician and ICU team
- Consider IV magnesium sulphate 1.2–2 g infusion over 20 minutes (unless already given)
- Give nebulised β_2 bronchodilator more frequently eg salbutamol 5 mg up to every 15–30 minutes or 10 mg per hour via continuous nebulisation (requires special nebuliser)

SUBSEQUENT MANAGEMENT

IF PATIENT IS IMPROVING continue:

- Oxygen to maintain SpO₂ 94–98%
- Prednisolone 40–50 mg daily or IV hydrocortisone 100 mg 6 hourly
- Nebulised β_2 bronchodilator with ipratropium 4–6 hourly

IF PATIENT NOT IMPROVING AFTER 15–30 MINUTES:

- Continue oxygen and steroids
- Use continuous nebulisation of salbutamol at 5–10 mg/hour if an appropriate nebuliser is available. Otherwise give nebulised salbutamol 5 mg every 15–30 minutes
- Continue ipratropium 0.5 mg 4–6 hourly until patient is improving

IF PATIENT IS STILL NOT IMPROVING:

- Discuss patient with senior clinician and ICU team
- Consider IV magnesium sulphate 1.2–2 g over 20 minutes (unless already given)
- Senior clinician may consider use of IV β_2 bronchodilator or IV aminophylline or progression to mechanical ventilation

MONITORING

- Repeat measurement of PEF 15–30 minutes after starting treatment
- Oximetry: maintain SpO₂ > 94 –98%
- Repeat blood gas measurements within 1 hour of starting treatment if:
 - initial PaO₂ < 8 kPa (60 mmHg) unless subsequent SpO₂ $> 92\%$ or
 - PaCO₂ normal or raised or
 - patient deteriorates
- Chart PEF before and after giving β_2 bronchodilator and at least 4 times daily throughout hospital stay

Transfer to ICU accompanied by a doctor prepared to intubate if:

- Deteriorating PEF, worsening or persisting hypoxia, or hypercapnia
- Exhaustion, altered consciousness
- Poor respiratory effort or respiratory arrest

DISCHARGE

When discharged from hospital, patients should have:

- Been on discharge medication for 12–24 hours and have had inhaler technique checked and recorded
- PEF $> 75\%$ of best or predicted and PEF diurnal variability $< 25\%$ unless discharge is agreed with respiratory physician
- Treatment with oral steroids (prednisolone 40–50 mg until recovery - minimum 5 days) and inhaled steroids in addition to bronchodilators
- Own PEF meter and written asthma action plan
- GP follow up arranged within 2 working days
- Follow-up appointment in respiratory clinic within 4 weeks

Patients with severe asthma (indicated by need for admission) and adverse behavioural or psychosocial features are at risk of further severe or fatal attacks.

- Determine reason(s) for exacerbation and admission
- Send details of admission, discharge and potential best PEF to GP

Annex 6

Management of acute asthma in children in general practice

Age 2–5 years

ASSESS AND RECORD ASTHMA SEVERITY

Moderate asthma

- SpO₂ ≥92%
- Able to talk
- Heart rate ≤140/min
- Respiratory rate ≤40/min

Acute severe asthma

- SpO₂ <92%
- Too breathless to talk
- Heart rate >140/min
- Respiratory rate >40/min
- Use of accessory neck muscles

Life-threatening asthma

- SpO₂ <92% plus any of:
- Silent chest
- Poor respiratory effort
- Agitation
- Confusion
- Cyanosis

- β₂ bronchodilator:
- via spacer ± facemask*
- Consider oral prednisolone 20 mg

- Oxygen via facemask to maintain SpO₂ 94–98% if available

- β₂ bronchodilator
- via nebuliser (preferably oxygen-driven), salbutamol 2.5 mg
- or, if nebuliser not available, via spacer*
- Oral prednisolone 20 mg

- β₂ bronchodilator with ipratropium:
- via nebuliser (preferably oxygen-driven), salbutamol 2.5 mg and ipratropium 0.25 mg every 20 minutes
- or, if nebuliser and ipratropium not available, β₂ bronchodilator via spacer*
- Oral prednisolone 20 mg
or IV hydrocortisone 50 mg if vomiting

Assess response to treatment 15 mins after β₂ bronchodilator

IF POOR RESPONSE ARRANGE ADMISSION

IF POOR RESPONSE REPEAT β₂ BRONCHODILATOR AND ARRANGE ADMISSION

REPEAT β₂ BRONCHODILATOR VIA OXYGEN-DRIVEN NEBULISER WHILST ARRANGING IMMEDIATE HOSPITAL ADMISSION

GOOD RESPONSE

- Continue β₂ bronchodilator via spacer or nebuliser, as needed but not exceeding 4 hourly
- **If symptoms are not controlled repeat β₂ bronchodilator and refer to hospital**
- Continue prednisolone until recovery (minimum 3–5 days)
- Arrange follow-up clinic visit within 48 hours
- Consider referral to secondary care asthma clinic if 2nd attack within 12 months.

POOR RESPONSE

- Stay with patient until ambulance arrives
- Send written assessment and referral details
- Repeat β₂ bronchodilator via oxygen-driven nebuliser in ambulance

LOWER THRESHOLD FOR ADMISSION IF:

- Attack in late afternoon or at night
- Recent hospital admission or previous severe attack
- Concern over social circumstances or ability to cope at home

NB: If a patient has signs and symptoms across categories, always treat according to their most severe features

* β₂ bronchodilator via spacer given one puff at a time, inhaled separately using tidal breathing, according to response, give another puff every 60 seconds up to a maximum of 10 puffs

Age >5 years

ASSESS AND RECORD ASTHMA SEVERITY

Moderate asthma

- SpO₂ ≥92%
- Able to talk
- Heart rate ≤125/min
- Respiratory rate ≤30/min
- PEF ≥50% best or predicted

Acute severe asthma

- SpO₂ <92%
- Too breathless to talk
- Heart rate >125/min
- Respiratory rate >30/min
- Use of accessory neck muscles
- PEF 33–50% best or predicted

Life-threatening asthma

- SpO₂ <92% plus any of:
- Silent chest
- Poor respiratory effort
- Agitation
- Confusion
- Cyanosis
- PEF <33% best or predicted

- β₂ bronchodilator:
- via spacer*
- Consider oral prednisolone 30–40 mg

- Oxygen via facemask to maintain SpO₂ 94–98% if available

- β₂ bronchodilator
- via nebuliser (preferably oxygen-driven), salbutamol 5 mg
- or, if nebuliser not available, via spacer*
- Oral prednisolone 30–40 mg

- β₂ bronchodilator with ipratropium:
- via nebuliser (preferably oxygen-driven), salbutamol 5 mg and ipratropium 0.25 mg every 20 minutes
- or, if nebuliser and ipratropium not available, β₂ bronchodilator via spacer*
- Oral prednisolone 30–40 mg or IV hydrocortisone 100 mg if vomiting

Assess response to treatment 15 mins after β₂ bronchodilator

IF POOR RESPONSE ARRANGE ADMISSION

IF POOR RESPONSE REPEAT β₂ BRONCHODILATOR AND ARRANGE ADMISSION

REPEAT β₂ BRONCHODILATOR VIA OXYGEN-DRIVEN NEBULISER WHILST ARRANGING IMMEDIATE HOSPITAL ADMISSION

GOOD RESPONSE

- Continue β₂ bronchodilator via spacer or nebuliser, as needed but not exceeding 4 hourly
- **If symptoms are not controlled repeat β₂ bronchodilator and refer to hospital**
- Continue prednisolone until recovery (minimum 3–5 days)
- Arrange follow-up clinic visit within 48 hours
- Consider referral to secondary care asthma clinic if 2nd attack within 12 months.

POOR RESPONSE

- Stay with patient until ambulance arrives
- Send written assessment and referral details
- Repeat β₂ bronchodilator via oxygen-driven nebuliser in ambulance

LOWER THRESHOLD FOR ADMISSION IF:

- Attack in late afternoon or at night
- Recent hospital admission or previous severe attack
- Concern over social circumstances or ability to cope at home

NB: If a patient has signs and symptoms across categories, always treat according to their most severe features

Annex 7

Management of acute asthma in children in emergency department

Age 2–5 years

ASSESS AND RECORD ASTHMA SEVERITY

Moderate asthma

- SpO₂ ≥92%
- No clinical features of severe asthma

NB: if a patient has signs and symptoms across categories, always treat according to their most severe features

Acute severe asthma

- SpO₂ <92%
- Too breathless to talk or eat
- Heart rate >140/min
- Respiratory rate >40/min
- Use of accessory neck muscles

Life-threatening asthma

SpO₂ <92% plus any of:

- Silent chest
- Poor respiratory effort
- Agitation
- Confusion
- Cyanosis

First line treatments

Oxygen via face mask/nasal prongs to achieve SpO₂ 94–98%

- β₂ bronchodilator:
 - via spacer ± facemask*
- Consider oral prednisolone 20 mg

β₂ bronchodilator

- via nebuliser (preferably oxygen-driven), salbutamol 2.5 mg
- or, if nebuliser not available, via spacer*
- Oral prednisolone 20 mg or IV hydrocortisone 4 mg/kg if vomiting
- If poor response add 0.25 mg nebulised ipratropium bromide to every nebulised β₂ bronchodilator and repeat every 20 minutes for 2 hours according to response

Reassess within 1 hour

First line treatments

Oxygen via face mask/nasal prongs to achieve SpO₂ 94–98%

- β₂ bronchodilator with ipratropium:
 - via nebuliser (preferably oxygen-driven), salbutamol 2.5 mg and ipratropium 0.25 mg
 - Repeat bronchodilators every 20–30 minutes
 - Oral prednisolone 20 mg or IV hydrocortisone 4 mg/kg if vomiting

Discuss with senior clinician, PICU team or paediatrician

Reassess within 1 hour

Age >5 years

ASSESS AND RECORD ASTHMA SEVERITY

Moderate asthma

- SpO₂ ≥92%
- PEF ≥50% best or predicted
- No clinical features of severe asthma

NB: if a patient has signs and symptoms across categories, always treat according to their most severe features

Acute severe asthma

- SpO₂ <92%
- PEF 33–50% best or predicted
- Heart rate >125/min
- Respiratory rate >30/min
- Use of accessory neck muscles

Life-threatening asthma

SpO₂ <92% plus any of:

- PEF <33% best or predicted
- Silent chest
- Poor respiratory effort
- Altered consciousness
- Cyanosis

First line treatments

Oxygen via face mask/nasal prongs to achieve SpO₂ 94–98%

- β₂ bronchodilator:
 - via spacer*
- Oral prednisolone 30–40 mg

β₂ bronchodilator

- via nebuliser (preferably oxygen-driven), salbutamol 5 mg
- or, if nebuliser not available, via spacer*
- Oral prednisolone 30–40 mg or IV hydrocortisone 4 mg/kg if vomiting
- If poor response add 0.25 mg nebulised ipratropium bromide to every nebulised β₂ bronchodilator and repeat every 20 minutes for 2 hours according to response

Reassess within 1 hour

First line treatments

Oxygen via face mask/nasal prongs to achieve SpO₂ 94–98%

- β₂ bronchodilator with ipratropium:
 - via nebuliser (preferably oxygen-driven), salbutamol 5 mg and ipratropium 0.25 mg
 - Repeat bronchodilators every 20–30 minutes
 - Oral prednisolone 30–40 mg or IV hydrocortisone 4 mg/kg if vomiting

Discuss with senior clinician, PICU team or paediatrician

Second line treatments

- Consider 2nd line treatments – see Annex 8
- Admit all cases if features of severe attack persist after initial treatment
- Arrange transfer to PICU/HDU if poor response to treatment as per local guidelines

DISCHARGE PLAN

- Continue β₂ bronchodilator 4 hourly as necessary
- Continue prednisolone 20 mg daily until recovery (minimum 3–5 days)
- Advise to contact GP if not controlled on above treatment
- Provide a written asthma action plan
- Review regular treatment
- Check inhaler technique
- Arrange GP follow up within 48 hours
- Arrange hospital asthma clinic follow up in 4–6 weeks if 2nd or subsequent attack in past 12 months.

Second line treatments

DISCHARGE PLAN

- Continue β₂ bronchodilator 4 hourly as necessary
- Continue prednisolone 30–40 mg daily until recovery (minimum 3–5 days)
- Advise to contact GP if not controlled on above treatment
- Provide a written asthma action plan
- Review regular treatment
- Check inhaler technique
- Arrange GP follow up within 48 hours
- Arrange hospital asthma clinic follow up in 4–6 weeks if 2nd or subsequent attack in past 12 months.

- Consider 2nd line treatments – see Annex 8
- Admit all cases if features of severe attack persist after initial treatment
- Arrange transfer to PICU/HDU if poor response to treatment as per local guidelines

* β₂ bronchodilator via spacer given one puff at a time, inhaled separately using tidal breathing; according to response, give another puff every 60 seconds up to a maximum of 10 puffs

Annex 8

Management of acute asthma in children in hospital

Age 2–5 years

Age >5 years

ASSESS AND RECORD ASTHMA SEVERITY

ASSESS AND RECORD ASTHMA SEVERITY

Moderate asthma

- $SpO_2 \geq 92\%$
- No clinical features of severe asthma

NB: If a patient has signs and symptoms across categories, always treat according to their most severe features

Acute severe asthma

- $SpO_2 < 92\%$
- Too breathless to talk or eat
- Heart rate $> 140/\text{min}$
- Respiratory rate $> 40/\text{min}$
- Use of accessory neck muscles

Life-threatening asthma

$SpO_2 < 92\%$ plus any of:

- Silent chest
- Poor respiratory effort
- Agitation
- Confusion
- Cyanosis

Moderate asthma

- $SpO_2 \geq 92\%$
- $PEF > 50\%$ best or predicted
- No clinical features of severe asthma

NB: If a patient has signs and symptoms across categories, always treat according to their most severe features

Acute severe asthma

- $SpO_2 < 92\%$
- $PEF 33\text{--}50\%$ best or predicted
- Heart rate $> 125/\text{min}$
- Respiratory rate $> 30/\text{min}$
- Use of accessory neck muscles

Life-threatening asthma

$SpO_2 < 92\%$ plus any of:

- $PEF < 33\%$ best or predicted
- Silent chest
- Poor respiratory effort
- Confusion
- Cyanosis

First-line treatments

First-line treatments

Oxygen via face mask/nasal prongs to achieve SpO_2 94–98%

Oxygen via face mask/nasal prongs to achieve SpO_2 94–98%

• β_2 bronchodilator:
- via spacer \pm facemask*

• Consider oral prednisolone 20 mg

• β_2 bronchodilator
- via nebuliser (preferably oxygen-driven), salbutamol 2.5 mg
- or, if nebuliser not available, via spacer*

• Oral prednisolone 20 mg or IV hydrocortisone 4 mg/kg if vomiting

• Repeat β_2 bronchodilator up to every 20–30 minutes according to response

• **If poor response** add 0.25 mg nebulised ipratropium bromide to every nebulised β_2 bronchodilator every 20 minutes for 1–2 hours

• β_2 bronchodilator with ipratropium:
- via nebuliser (preferably oxygen-driven), salbutamol 2.5 mg and ipratropium 0.25 mg

• Repeat bronchodilators every 20–30 minutes

• Oral prednisolone 20mg or IV hydrocortisone 4mg/kg if vomiting

• Consider adding 150 mg magnesium sulphate to each β_2 bronchodilator/ipratropium nebuliser in first hour

Discuss with senior clinician, PICU team or paediatrician

• β_2 bronchodilator:
- via spacer*

• Oral prednisolone 30–40 mg

• β_2 bronchodilator
- via nebuliser (preferably oxygen-driven), salbutamol 5 mg
- or, if nebuliser not available, via spacer*

• Oral prednisolone 30–40 mg or IV hydrocortisone 4 mg/kg if vomiting

• Repeat β_2 bronchodilator up to every 20–30 minutes according to response

• **If poor response** add 0.25 mg nebulised ipratropium bromide to every nebulised β_2 bronchodilator every 20 minutes for 1–2 hours

• β_2 bronchodilator with ipratropium:
- via nebuliser (preferably oxygen-driven), salbutamol 5 mg and ipratropium 0.25 mg

• Repeat bronchodilators every 20–30 minutes

• Oral prednisolone 30–40 mg or IV hydrocortisone 4mg/kg if vomiting

• Consider adding 150 mg magnesium sulphate to each β_2 bronchodilator/ipratropium nebuliser in first hour

Discuss with senior clinician, PICU team or paediatrician

Reassess within 1 hour

Reassess within 1 hour

Record respiratory rate, heart rate and oxygen saturation every 1–4 hours

Record respiratory rate, heart rate, oxygen saturation and PEF/FEV every 1–4 hours

Second-line treatments

Second-line treatments

RESPONDING

- Continue bronchodilators 1–4 hours as necessary
- Discharge when stable on 4-hourly treatment
- Continue prednisolone 20 mg daily until recovery (minimum 3–5 days)

At discharge

- Ensure stable on 4-hourly inhaled treatment
- Review the need for regular treatment and the use of inhaled steroids
- Review inhaler technique
- Provide a written asthma action plan for treating future attacks
- Arrange GP follow up within 48 hours
- Arrange hospital asthma clinic follow up in 4–6 weeks

NOT RESPONDING

- Continue 20–30 minute nebulisers
- Consider chest X-ray and blood gases
- Discuss with senior clinician, paediatrician or PICU
- Consider admission to HDU/PICU

Consider risks and benefits of:

- **Bolus IV infusion of magnesium sulphate** 40 mg/kg (max 2 g) over 20 minutes
- **Bolus IV salbutamol** 15 micrograms/kg if not already given
- Continuous IV salbutamol infusion 1–5 micrograms/kg/min (200 micrograms/ml solution)
- **IV aminophylline** 5 mg/kg loading dose over 20 minutes (omit in those receiving oral theophyllines)

followed by continuous infusion 1mg/kg/hour

Assess response before initiating each new treatment

RESPONDING

- Continue bronchodilators 1–4 hours as necessary
- Discharge when stable on 4-hourly treatment
- Continue prednisolone 30–40 mg daily until recovery (minimum 3–5 days)

At discharge

- Ensure stable on 4-hourly inhaled treatment
- Review the need for regular treatment and the use of inhaled steroids
- Review inhaler technique
- Provide a written asthma action plan for treating future attacks
- Arrange GP follow up within 48 hours
- Arrange hospital asthma clinic follow up in 4–6 weeks

NOT RESPONDING

- Continue 20–30 minute nebulisers
- Consider chest X-ray and blood gases
- Discuss with senior clinician, paediatrician or PICU
- Consider admission to HDU/PICU

Consider risks and benefits of:

- **Bolus IV infusion of magnesium sulphate** 40 mg/kg (max 2 g) over 20 minutes
- **Bolus IV salbutamol** 15 micrograms/kg if not already given
- Continuous IV salbutamol infusion 1–5 micrograms/kg/min (200 micrograms/ml solution)
- **IV aminophylline** 5 mg/kg loading dose over 20 minutes (omit in those receiving oral theophyllines)

followed by continuous infusion 1mg/kg/hour

Assess response before initiating each new treatment

RESPONDING

- Continue bronchodilators 1–4 hours as necessary
- Discharge when stable on 4-hourly treatment
- Continue prednisolone 30–40 mg daily until recovery (minimum 3–5 days)

At discharge

- Ensure stable on 4-hourly inhaled treatment
- Review the need for regular treatment and the use of inhaled steroids
- Review inhaler technique
- Provide a written asthma action plan for treating future attacks
- Arrange GP follow up within 48 hours
- Arrange hospital asthma clinic follow up in 4–6 weeks

NOT RESPONDING

- Continue 20–30 minute nebulisers
- Consider chest X-ray and blood gases
- Discuss with senior clinician, paediatrician or PICU
- Consider admission to HDU/PICU

Consider risks and benefits of:


- **Bolus IV infusion of magnesium sulphate** 40 mg/kg (max 2 g) over 20 minutes
- **Bolus IV salbutamol** 15 micrograms/kg if not already given
- Continuous IV salbutamol infusion 1–5 micrograms/kg/min (200 micrograms/ml solution)
- **IV aminophylline** 5 mg/kg loading dose over 20 minutes (omit in those receiving oral theophyllines)

followed by continuous infusion 1mg/kg/hour

Assess response before initiating each new treatment

* β_2 bronchodilator via spacer given one puff at a time, inhaled separately using tidal breathing; according to response, give another puff every 60 seconds up to a maximum of 10 puffs


Annex 9



¹ Management of acute asthma in children under 1 year should be under the direction of a respiratory paediatrician.

Annex 10

WORK-RELATED ASTHMA AND RHINITIS: CASE FINDING AND MANAGEMENT IN PRIMARY CARE

Guidelines for the identification, Management and Prevention of Occupational Asthma | www.bohrf.org.uk/projects/asthma.html

References

1. Scottish Intercollegiate Guidelines Network (SIGN). SIGN 50: A guideline developers' handbook. Edinburgh: SIGN; 2008. (SIGN publication no. 50). [cited 1st Jul 2014] Available from <http://www.sign.ac.uk/guidelines/fulltext/50/index.html>
2. British Thoracic Society, Scottish Intercollegiate Guidelines Network. British guideline on the management of asthma. *Thorax* 2003;58 Suppl 1:i1-94.
3. World Health Organisation. Health topics: adolescent health. 2010. Available from http://www.who.int/topics/adolescent_health/en: [Accessed. 1 Jul 2014].
4. Committee. JF. British National Formulary (online). London: BMJ Group and Pharmaceutical Press; Available from <http://www.medicinescomplete.com>: [Accessed 17 July 2019].
5. Medicines and Healthcare products Regulatory Agency. Off-label use or unlicensed medicines: prescribers' responsibilities. *Drug Safety Update* 2009;2(9):6-7.
6. Global Initiative for Asthma. Global strategy for asthma management and prevention. 2018. Available from <https://ginasthma.org/gina-reports/>: [Accessed. 03 May 2019].
7. British Medical Association (BMA) NE, NHS England. 2015/16 General Medical Services (GMS) contract Quality and Outcomes Framework (QOF): guidance for GMS contract 2014/15. London: NHS Employers; 2014; Available from <https://www.nhsemployers.org/-/media/Employers/Documents/Primary-care-contracts/QOF/2014-15/2014-15-General-Medical-Services-contract---Quality-and-Outcomes-Framework.pdf?la=en&hash=69CF3870EB19476ECDFF3A5CBAA35F08C34E00DC>: [Accessed 17 July 2019].
8. Pinnock H, Adlem L, Gaskin S, Harris J, Snellgrove C, Sheikh A. Accessibility, clinical effectiveness, and practice costs of providing a telephone option for routine asthma reviews: phase IV controlled implementation study. *Br J Gen Pract* 2007;57(542):714-22.
9. Schneider A, Faderl B, Schwarzbach J, Welker L, Karsch-Volk M, Jorres RA. Prognostic value of bronchial provocation and FENO measurement for asthma diagnosis--results of a delayed type of diagnostic study. *Respiratory Medicine* 2014;108(1):34-40.
10. Waring N. Use of new asthma BTM steps in one general practice--should asthmatics no longer on treatment be followed-up? *Primary Care Respiratory Journal* 1997;5:28-.
11. Greiver M, Lang C, Hunchuck J, Rothschild K, Lang C, Hunchuck J, et al. Improving the diagnosis of asthma in a primary care practice. *Can Fam Physician* 2012;58(7):773-4, e382-4.
12. Melbye H, Drivenes E, Dalbak LG, Leinan T, Hoegh-Henrichsen S, Ostrem A, et al. Asthma, chronic obstructive pulmonary disease, or both? Diagnostic labeling and spirometry in primary care patients aged 40 years or more. *Int J Chron Obstruct Pulmon Dis* 2011;6:597-603.
13. Ringsberg KC, Bjarneman P, Larsson R, Wallstrom E, Lowhagen O. Diagnosis of asthma in primary health care: a pilot study. *J Allergy (Cairo)* 2014;2014:898965.
14. Miravittles M, Andreu I, Romero Y, Sitjar S, Altes A, Anton E, et al. Difficulties in differential diagnosis of COPD and asthma in primary care. *Br J Gen Pract* 2012;62(595):e68-75.
15. Metting EI, Riemersma RA, Kocks JH, Piersma-Wichers MG, Sanderma R, van der Molen T. Feasibility and effectiveness of an Asthma/COPD service for primary care: a cross-sectional baseline description and longitudinal results. *NPJ Prim Care Respir Med* 2015;25:14101.
16. Prieto Centurion V, Huang F, Naureckas ET, Camargo CA, Jr., Charbeneau J, Joo MJ, et al. Confirmatory spirometry for adults hospitalized with a diagnosis of asthma or chronic obstructive pulmonary disease exacerbation. *BMC Pulmonary Medicine* 2012;12:73.
17. (NICE). NifHaCE. Asthma: diagnosis and monitoring of asthma in adults, children and young people - appendices A-R. London: NICE; 2017. Available from <https://www.nice.org.uk/guidance/ng80/evidence/appendices-ar-pdf-4656178048>: [Accessed 17 July 2019].
18. Galant SP, Crawford LJ, Morphew T, Jones CA, Bassin S. Predictive value of a cross-cultural asthma case-detection tool in an elementary school population. *Pediatrics* 2004;114(3):e307-16.
19. Jones CA, Morphew T, Clement LT, Kimia T, Dyer M, Li M, et al. A school-based case identification process for identifying inner city children with asthma: the Breathmobile program. *Chest* 2004;125(3):924-34.
20. Yu IT, Wong TW, Li W. Using child reported respiratory symptoms to diagnose asthma in the community. *Arch Dis Child* 2004;89(6):544-8.
21. Schneider A, Ay M, Faderl B, Linde K, Wagenpfeil S. Diagnostic accuracy of clinical symptoms in obstructive airway diseases varied within different health care sectors. *J Clin Epidemiol* 2012;65(8):846-54.
22. Tomita K, Sano H, Chiba Y, Sato R, Sano A, Nishiyama O, et al. A scoring algorithm for predicting the presence of adult asthma: a prospective derivation study. *Primary Care Respiratory Journal* 2013;22(1):51-8.
23. Cane RS, Ranganathan SC, McKenzie SA. What do parents of wheezy children understand by "wheeze"? *Archives of Disease in Childhood* 2000;82(4):327-32.
24. Schneider A, Gindner L, Tilemann L, Schermer T, Dinant GJ, Meyer FJ, et al. Diagnostic accuracy of spirometry in primary care. *BMC Pulm Med* 2009;9:31.

25. Bacharier LB, Strunk RC, Mauger D, White D, Lemanske RF Jr, Sorkness CA. Classifying asthma severity in children: Mismatch between symptoms, medication use, and lung function. *Am J Respir Crit Care Med* 2004;170(4):426-32.
26. Brouwer AFJ, Roorda RJ, Brand PLP. Home spirometry and asthma severity in children. *European Respiratory Journal* 2006;28(6):1131-7.
27. Verini M, Peroni DG, Rossi N, Nicodemo A, De Stradis R, Spagnolo C, et al. Functional assessment of allergic asthmatic children while asymptomatic. *Allergy Asthma Proc* 2006;27(4):359-64.
28. Dweik RA, Boggs PB, Erzurum SC, Irvin CG, Leigh MW, Lundberg JO, et al. An official ATS clinical practice guideline: interpretation of exhaled nitric oxide levels (FENO) for clinical applications. *Am J Respir Crit Care Med* 2011;184(5):602-15.
29. Joseph-Bowen J, de Klerk NH, Firth MJ, Kendall GE, Holt PG, Sly PD. Lung function, bronchial responsiveness, and asthma in a community cohort of 6-year-old children. *Am J Respir Crit Care Med* 2004;169(7):850-4.
30. Pellegrino R, Viegi G, Brusasco V, Crapo RO, Burgos F, Casaburi R, et al. Interpretative strategies for lung function tests. *Eur Respir J* 2005;26(5):948-68.
31. Quanjer PH, Stanojevic S, Cole TJ, Baur X, Hall GL, Culver BH, et al. Multi-ethnic reference values for spirometry for the 3-95-yr age range: the global lung function 2012 equations. *Eur Respir J* 2012;40(6):1324-43.
32. (NICE). NifHaCE. Asthma: diagnosis and monitoring of asthma in adults, children and young people. London: NICE; 2017. Available from <http://www.nice.org.uk/guidance/ng80>: [Accessed 17 July 2019].
33. (NICE) NifHaCE, (NICE). Chronic obstructive pulmonary disease in over 16s: diagnosis and management. London: NICE; 2010. Available from <http://www.nice.org.uk/guidance/cg101/resources/chronic-obstructivepulmonary-disease-in-over-16s-diagnosis-andmanagement-35109323931589>: [Accessed 17 July 2019].
34. Quanjer PH, Lebowitz MD, Gregg I, Miller MR, Pedersen OF. Peak expiratory flow: conclusion and recommendations of a working party of the European Respiratory Society. *Eur Respir J Suppl* 1997;24:2S-8S.
35. D'Alonzo GE, Steinijans VW, Keller A. Measurements of morning and evening airflow grossly underestimate the circadian variability of FEV₁ and peak expiratory flow rate in asthma. *Am J Respir Crit Care Med* 1995;152(3):1097-9.
36. Chowienczyk PJ, Parkin DH, Lawson CP, Cochrane GM. Do asthmatic patients correctly record home spirometry measurements? *BMJ* 1994;309(6969):1618.
37. Kamps AW, Roorda RJ, Brand PL. Peak flow diaries in childhood asthma are unreliable. *Thorax* 2001;56(3):180-2.
38. Higgins BG, Britton JR, Chinn S, Cooper S, Burney PG, Tattersfield AE. Comparison of bronchial reactivity and peak expiratory flow variability measurements for epidemiologic studies. *Am Rev Respir Dis* 1992;145(3):588-93.
39. Higgins BG, Britton JR, Chinn S, Jones TD, Jenkinson D, Burney PG, et al. The distribution of peak flow variability in a population sample. *Am Rev Respir Dis* 1989;140(5):1368-72.
40. Quackenboss JJ, Lebowitz MD, Krzyzanowski M. The normal range of diurnal changes in peak expiratory flow rates. Relationship to symptoms and respiratory disease. *Am Rev Respir Dis* 1991;143(2):323-30.
41. Thiadens HA, De Bock GH, Dekker FW, Huysman JA, Van Houwelingen JC, Springer MP, et al. Value of measuring diurnal peak flow variability in the recognition of asthma: a study in general practice. *Eur Respir J* 1998;12(4):842-7.
42. Lebowitz MD, Krzyzanowski M, Quackenboss JJ, O'Rourke MK. Diurnal variation of PEF and its use in epidemiological studies. *Eur Respir J Suppl* 1997(24):49s-56s.
43. Boezen HM, Schouten JP, Postma DS, Rijcken B. Distribution of peak expiratory flow variability by age, gender and smoking habits in a random population sample aged 20-70 yrs. *Eur Respir J* 1994;7(10):1814-20.
44. Siersted HC, Hansen HS, Hansen NC, Hyldebrandt N, Mostgaard G, Oxhøj H. Evaluation of peak expiratory flow variability in an adolescent population sample. The Odense Schoolchild Study. *Am J Respir Crit Care Med* 1994;149(3 Pt 1):598-603.
45. Gannon PF, Newton DT, Belcher J, Pantin CF, Burge PS. Development of OASYS-2: a system for the analysis of serial measurement of peak expiratory flow in workers with suspected occupational asthma. *Thorax* 1996;51(5):484-9.
46. Crapo RO, Casaburi R, Coates AL, Enright PL, Hankinson JL, Irvin CG, et al. Guidelines for methacholine and exercise challenge testing-1999. This official statement of the American Thoracic Society was adopted by the ATS Board of Directors, July 1999. *Am J Respir Crit Care Med* 2000;161(1):309-29.
47. Cockcroft DW, Killian DN, Mellon JJ, Hargreave FE. Bronchial reactivity to inhaled histamine: a method and clinical survey. *Clin Allergy* 1977;7(3):235-43.
48. Cockcroft DW, Murdock KY, Berscheid BA, Gore BP. Sensitivity and specificity of histamine PC20 determination in a random selection of young college students. *J Allergy Clin Immunol* 1992;89(1 Pt 1):23-30.
49. Brand PL, Postma DS, Kerstjens HA, Koeter GH. Relationship of airway hyperresponsiveness to respiratory symptoms and diurnal peak flow variation in patients with obstructive lung disease. The Dutch CNSLD Study Group. *Am Rev Respir Dis* 1991;143(5 Pt 1):916-21.
50. Ramsdale EH, Morris MM, Roberts RS, Hargreave FE. Bronchial responsiveness to methacholine in chronic bronchitis: relationship to airflow obstruction and cold air responsiveness. *Thorax* 1984;39(12):912-8.

51. Remes ST, Pekkanen J, Remes K, Salonen RO, Korppi M. In search of childhood asthma: questionnaire, tests of bronchial hyperresponsiveness, and clinical evaluation. *Thorax* 2002;57(2):120-6.
52. Joos GF, O'Connor B, Anderson SD, Chung F, Cockcroft DW, Dahlen B, et al. Indirect airway challenges. *Eur Respir J* 2003;21(6):1050-68.
53. Anderton RC, Cuff MT, Frith PA, Cockcroft DW, Morse JL, Jones NL, et al. Bronchial responsiveness to inhaled histamine and exercise. *J Allergy Clin Immunol* 1979;63(5):315-20.
54. Abu-Hasan M, Tannous B, Weinberger M. Exercise-induced dyspnea in children and adolescents: if not asthma then what? *Ann Allergy Asthma Immunol* 2005;94(3):366-71.
55. American Thoracic Society, European Respiratory Society. ATS/ERS recommendations for standardized procedures for the online and offline measurement of exhaled lower respiratory nitric oxide and nasal nitric oxide, 2005. *Am J Respir Crit Care Med* 2005;171(8):912-30.
56. Alving K, Malinowski A. Basic aspects of exhaled nitric oxide. *Exhaled biomarkers* 2010;49:1-31.
57. Bjerner L, Alving K, Diamant Z, Magnussen H, Pavord I, Piacentini G, et al. Current evidence and future research needs for FeNO measurement in respiratory diseases. *Respir Med* 2014;108(6):830-41.
58. Malmberg LP, Turpeinen H, Ryttilä P, Sarna S, Haahtela T. Determinants of increased exhaled nitric oxide in patients with suspected asthma. *Allergy* 2005;60(4):464-8.
59. Barreto M, Villa MP, Monti F, Bohmerova Z, Martella S, Montesano M, et al. Additive effect of eosinophilia and atopy on exhaled nitric oxide levels in children with or without a history of respiratory symptoms. *Pediatr Allergy Immunol* 2005;16(1):52-8.
60. Malmberg LP, Petays T, Haahtela T, Laatikainen T, Jousilahti P, Vartiainen E, et al. Exhaled nitric oxide in healthy nonatopic school-age children: determinants and height-adjusted reference values. *Pediatr Pulmonol* 2006;41(7):635-42.
61. Chan EY, Dundas I, Bridge PD, Healy MJ, McKenzie SA. Skin-prick testing as a diagnostic aid for childhood asthma. *Pediatr Pulmonol* 2005;39(6):558-62.
62. Castro-Rodriguez JA, Holberg CJ, Wright AL, Martinez FD. A clinical index to define risk of asthma in young children with recurrent wheezing. *Am J Respir Crit Care Med* 2000;162(4 Pt 1):1403-6.
63. Eysink PE, ter Riet G, Aalberse RC, van Aalderen WM, Roos CM, van der Zee JS, et al. Accuracy of specific IgE in the prediction of asthma: development of a scoring formula for general practice. *Br J Gen Pract* 2005;55(511):125-31.
64. Simpson A, Soderstrom L, Ahlstedt S, Murray CS, Woodcock A, Custovic A. IgE antibody quantification and the probability of wheeze in preschool children. *J Allergy Clin Immunol* 2005;116(4):744-9.
65. Kurukulaaratchy RJ, Fenn M, Matthews S, Arshad SH. Characterisation of atopic and non-atopic wheeze in 10 year old children. *Thorax* 2004;59(7):563-8.
66. Pavord ID, Pizzichini MM, Pizzichini E, Hargreave FE. The use of induced sputum to investigate airway inflammation. *Thorax* 1997;52(6):498-501.
67. Lex C, Payne DN, Zacharasiewicz A, Li AM, Wilson NM, Hansel TT, et al. Sputum induction in children with difficult asthma: safety, feasibility, and inflammatory cell pattern. *Pediatr Pulmonol* 2005;39(4):318-24.
68. Ryttilä P, Pelkonen AS, Metso T, Nikander K, Haahtela T, Turpeinen M. Induced sputum in children with newly diagnosed mild asthma: The effect of 6 months of treatment with budesonide or disodium cromoglycate. *Allergy* 2004;59(8):839-44.
69. Dundas I, Chan EY, Bridge PD, McKenzie SA. Diagnostic accuracy of bronchodilator responsiveness in wheezy children. *Thorax* 2005;60(1):13-6.
70. Anderson SD, Charlton B, Weiler JM, Nichols S, Spector SL, Pearlman DS. Comparison of mannitol and methacholine to predict exercise-induced bronchoconstriction and a clinical diagnosis of asthma. *Respir Res* 2009;10:4.
71. Lucas AE, Smeenk FJ, Smeele IJ, van Schayck OP, Smeenk FJ, Smeele IJ, et al. Diagnostic accuracy of primary care asthma/COPD working hypotheses, a real life study. *Respiratory Medicine* 2012;106(8):1158-63.
72. Buffels J, Degryse J, Liistro G, Decramer M, Degryse J, Liistro G, et al. Differential diagnosis in a primary care population with presumed airway obstruction: a real-life study. *Respiration* 2012;84(1):44-54.
73. Gerald LB, Grad R, Turner-Henson A, Hains C, Tang S, Feinstein R, et al. Validation of a multistage asthma case-detection procedure for elementary school children. *Pediatrics* 2004;114(4):e459-68.
74. Ly NP, Gold DR, Weiss ST, Celedon JC. Recurrent wheeze in early childhood and asthma among children at risk for atopy. *Pediatrics* 2006;117(6):e1132-8.
75. Marchant JM, Masters IB, Taylor SM, Cox NC, Seymour GJ, Chang AB. Evaluation and outcome of young children with chronic cough. *Chest* 2006;129(5):1132-41.
76. Kurukulaaratchy RJ, Matthews S, Holgate ST, Arshad SH. Predicting persistent disease among children who wheeze during early life. *European Respiratory Journal* 2003;22(5):767-71.
77. Schonberger H, van Schayck O, Muris J, Bor H, van den Hoogen H, Knottnerus A, et al. Towards improving the accuracy of diagnosing asthma in early childhood. *Eur J Gen Pract* 2004;10(4):138-45.
78. Marchant JM, Masters IB, Taylor SM, Chang AB. Utility of signs and symptoms of chronic cough in predicting specific cause in children. *Thorax* 2006;61(8):694-8.
79. Smith AD, Cowan JO, Filsell S, McLachlan C, Monti-Sheehan G, Jackson P, et al. Diagnosing asthma: comparisons between exhaled nitric oxide measurements and conventional tests. *Am J Respir Crit Care Med* 2004;169(4):473-8. (34 ref).

-
80. Goldstein MF, Veza BA, Dunskey EH, Dvorin DJ, Belecanech GA, Haralabatos IC. Comparisons of peak diurnal expiratory flow variation, postbronchodilator FEV(1) responses, and methacholine inhalation challenges in the evaluation of suspected asthma. *Chest* 2001;119(4):1001-10.
81. Hunter CJ, Brightling CE, Woltmann G, Wardlaw AJ, Pavord ID. A comparison of the validity of different diagnostic tests in adults with asthma. *Chest* 2002;121(4):1051-7.
82. Smith AD, Cowan JO, Brassett KP, Filsell S, McLachlan C, Monti-Sheehan G, et al. Exhaled nitric oxide: a predictor of steroid response. *Am J Respir Crit Care Med* 2005;172(4):453-9.
83. Hederos CA, Janson S, Andersson H, Hedlin G. Chest X-ray investigation in newly discovered asthma. *Pediatr Allergy Immunol* 2004;15(2):163-5.
84. Fishwick D, Barber CM, Bradshaw LM, Ayres JG, Barraclough R, Burge S, et al. Standards of care for occupational asthma: an update. *Thorax* 2012;67(3):278-80.
85. Kurukulaaratchy RJ, Matthews S, Arshad SH. Relationship between childhood atopy and wheeze: what mediates wheezing in atopic phenotypes? *Ann Allergy Asthma Immunol* 2006;97(1):84-91.
86. Jenkins MA, Hopper JL, Bowes G, Carlin JB, Flander LB, Giles GG. Factors in childhood as predictors of asthma in adult life. *British Medical Journal* 1994;309(6947):90-3.
87. Martinez FD, Wright AL, Taussig LM, Holberg CJ, Halonen M, Morgan WJ. Asthma and wheezing in the first six years of life. The Group Health Medical Associates. *N Engl J Med* 1995;332(3):133-8.
88. Sporik R, Holgate ST, Cogswell JJ. Natural history of asthma in childhood - a birth cohort study. *Archives of Disease in Childhood* 1991;66(9):1050-3.
89. Strachan DP, Butland BK, Anderson HR. Incidence and prognosis of asthma and wheezing illness from early childhood to age 33 in a national British cohort. *British Medical Journal* 1996;312(7040):1195-9.
90. Aberg N, Engstrom I. Natural history of allergic diseases in children. *Acta Paediatr Scand* 1990;79(2):206-11.
91. Toelle BG, Xuan W, Peat JK, Marks GB. Childhood factors that predict asthma in young adulthood. *European Respiratory Journal* 2004;23(1):66-70.
92. Anderson HR, Pottier AC, Strachan DP. Asthma from birth to age 23: incidence and relation to prior and concurrent atopic disease. *Thorax* 1992;47(7):537-42.
93. Barbee RA, Murphy S. The natural history of asthma. *J Allergy Clin Immunol* 1998;102(4 Pt 2):S65-72.
94. Blair H. Natural history of childhood asthma. 20-year follow-up. *Archives of Disease in Childhood* 1977;52(8):613-9.
95. Johnstone DE. A study of the natural history of bronchial asthma in children. *Am J Dis Child* 1968;115(2):213-6.
96. Laor A, Cohen L, Danon YL. Effects of time, sex, ethnic origin, and area of residence on prevalence of asthma in Israeli adolescents. *BMJ* 1993;307(6908):841-4.
97. Heaney LG, Conway E, Kelly C, Johnston BT, English C, Stevenson M, et al. Predictors of therapy resistant asthma: outcome of a systematic evaluation protocol. *Thorax* 2003;58(7):561-6.
98. Luyt DK, Burton PR, Simpson H. Epidemiological study of wheeze, doctor diagnosed asthma, and cough in preschool children in Leicestershire. *BMJ* 1993;306(6889):1386-90.
99. Martin AJ, McLennan LA, Landau LI, Phelan PD. The natural history of childhood asthma to adult life. *Br Med J* 1980;280(6229):1397-400.
100. Robertson CF, Heycock E, Bishop J, Nolan T, Olinsky A, Phelan PD. Prevalence of asthma in Melbourne schoolchildren: changes over 26 years. *British Medical Journal* 1991;302(6785):1116-8.
101. Roorda RJ. Prognostic factors for the outcome of childhood asthma in adolescence. *Thorax* 1996;51(Suppl 1):S7-12.
102. Sears MR, Holdaway MD, Flannery EM, Herbison GP, Silva PA. Parental and neonatal risk factors for atopy, airway hyper-responsiveness, and asthma. *Archives of Disease in Childhood* 1996;75(5):392-8.
103. Sherman CB, Tosteson TD, Tager IB, Speizer FE, Weiss ST. Early childhood predictors of asthma. *Am J Epidemiol* 1990;132(1):83-95.
104. Sigurs N, Bjarnason R, Sigurbergsson F, Kjellman B, Bjorksten B. Asthma and immunoglobulin E antibodies after respiratory syncytial virus bronchiolitis: a prospective cohort study with matched controls. *Pediatrics* 1995;95(4):500-5.
105. Tariq SM, Matthews SM, Hakim EA, Stevens M, Arshad SH, Hide DW. The prevalence of and risk factors for atopy in early childhood: a whole population birth cohort study. *J Allergy Clin Immunol* 1998;101(5):587-93.
106. Clough JB, Keeping KA, Edwards LC, Freeman WM, Warner JA, Warner JO. Can we predict which wheezy infants will continue to wheeze? *Am J Respir Crit Care Med* 1999;160(5 Pt 1):1473-80.
107. Dodge R, Martinez FD, Cline MG, Lebowitz MD, Burrows B. Early childhood respiratory symptoms and the subsequent diagnosis of asthma. *J Allergy Clin Immunol* 1996;98(1):48-54.
108. Reijonen TM, Kotaniemi-Syrjanen A, Korhonen K, Korppi M. Predictors of asthma three years after hospital admission for wheezing in infancy. *Pediatrics* 2000;106(6):1406-12.
109. Kotaniemi-Syrjanen A, Reijonen TM, Romppanen J, Korhonen K, Savolainen K, Korppi M. Allergen-specific immunoglobulin E antibodies in wheezing infants: the risk for asthma in later childhood. *Pediatrics* 2003;111(3):e255-61.
110. Sears MR, Herbison GP, Holdaway MD, Hewitt CJ, Flannery EM, Silva PA. The relative risks of sensitivity to grass pollen, house dust mite and cat dander in the development of childhood asthma. *Clin Exp Allergy* 1989;19(4):419-24.
111. Rona RJ, Duran-Tauleria E, Chinn S. Family size, atopic disorders in parents, asthma in children, and ethnicity. *J Allergy Clin Immunol* 1997;99(4):454-60.
-

112. Rusconi F, Galassi C, Corbo GM, Forastiere F, Biggeri A, Ciccone G, et al. Risk factors for early, persistent, and late-onset wheezing in young children. *SIDRIA Collaborative Group. Am J Respir Crit Care Med* 1999;160(5 Pt 1):1617-22.
113. Pearson MG, Bucknall CE, editors. *Measuring clinical outcome in asthma : a patient-focused approach* London: Royal College of Physicians of London; 1999.
114. Pinnock H, Burton C, Campbell S, Gruffydd-Jones K, Hannon K, Hoskins G, et al. Clinical implications of the Royal College of Physicians three questions in routine asthma care: a real-life validation study. *Prim Care Respir J* 2012;21(3):288-94.
115. Juniper EF, Bousquet J, Abetz L, Bateman ED, GOAL Committee. Identifying 'well-controlled' and 'not well-controlled' asthma using the Asthma Control Questionnaire. *Respir Med* 2006;100(4):616-21.
116. Juniper EF, O'Byrne PM, Guyatt GH, Ferrie PJ, King DR. Development and validation of a questionnaire to measure asthma control. *Eur Respir J* 1999;14(4):902-7.
117. Juniper EF, Svensson K, Mork AC, Ståhl E. Measurement properties and interpretation of three shortened versions of the asthma control questionnaire. *Respir Med* 2005;99(5):553-8.
118. Juniper EF, Gruffydd-Jones K, Ward S, Svensson K. Asthma Control Questionnaire in children: validation, measurement properties, interpretation. *Eur Respir J* 2010;36(6):1410-6.
119. Nathan RA, Sorkness CA, Kosinski M, Schatz M, Li JT, Marcus P, et al. Development of the asthma control test: a survey for assessing asthma control. *J Allergy Clin Immunol* 2004;113(1):59-65.
120. Schatz M, Sorkness CA, Li JT, Marcus P, Murray JJ, Nathan RA, et al. Asthma Control Test: reliability, validity, and responsiveness in patients not previously followed by asthma specialists. *J Allergy Clin Immunol* 2006;117(3):549-56.
121. Liu AH, Zeiger R, Sorkness C, Mahr T, Ostrom N, Burgess S, et al. Development and cross-sectional validation of the Childhood Asthma Control Test. *J Allergy Clin Immunol* 2007;119(4):817-25.
122. Schatz M, Kosinski M, Yarlas AS, Hanlon J, Watson ME, Jhingran P. The minimally important difference of the Asthma Control Test. *J Allergy Clin Immunol* 2009;124(4):719-23.
123. Juniper EF, Guyatt GH, Ferrie PJ, Griffith LE. Measuring quality of life in asthma. *Am Rev Respir Dis* 1993;147(4):832-8.
124. Bacharier LB, Guilbert TW, Zeiger RS, Strunk RC, Morgan WJ, Lemanske RF Jr, et al. Patient characteristics associated with improved outcomes with use of an inhaled corticosteroid in preschool children at risk for asthma. *J Allergy Clin Immunol* 2009;123(5):1077-82.
125. Juniper EF, Guyatt GH, Feeny DH, Ferrie PJ, Griffith LE, Townsend M. Measuring quality of life in children with asthma. *Qual Life Res* 1996;5(1):35-46.
126. Pavord ID, Beasley R, Agusti A, Anderson GP, Bel E, Brusselle G, et al. After asthma: redefining airways diseases. *Lancet* 2018;391(10118):350-400.
127. Blakey JD, Price DB, Pizzichini E, Popov TA, Dimitrov BD, Postma DS, et al. Identifying Risk of Future Asthma Attacks Using UK Medical Record Data: A Respiratory Effectiveness Group Initiative. *J Allergy Clin Immunol Pract* 2017;5(4):1015-24.e8.
128. Calhoun WJ, Haselkorn T, Mink DR, Miller DP, Dorenbaum A, Zeiger RS. Clinical burden and predictors of asthma exacerbations in patients on guideline-based steps 4-6 asthma therapy in the TENOR cohort. *J Allergy Clin Immunol Pract* 2014;2(2):193-200.
129. Kimura H, Konno S, Makita H, Taniguchi N, Shimizu K, Suzuki M, et al. Prospective predictors of exacerbation status in severe asthma over a 3-year follow-up. *Clin Exp Allergy* 2018;21:21.
130. Miller MK, Lee JH, Miller DP, Wenzel SE, Group TS. Recent asthma exacerbations: a key predictor of future exacerbations. *Respir Med* 2007;101(3):481-9.
131. O'Connor RD, Bleecker ER, Long A, Tashkin D, Peters S, Klingman D, et al. Subacute lack of asthma control and acute asthma exacerbation history as predictors of subsequent acute asthma exacerbations: evidence from managed care data. *J Asthma* 2010;47(4):422-8.
132. Price D, Wilson AM, Chisholm A, Rigazio A, Burden A, Thomas M, et al. Predicting frequent asthma exacerbations using blood eosinophil count and other patient data routinely available in clinical practice. *J Asthma Allergy* 2016;9:1-12.
133. Quezada W, Kwak ES, Reibman J, Rogers L, Mastrorade J, Teague WG, et al. Predictors of asthma exacerbation among patients with poorly controlled asthma despite inhaled corticosteroid treatment. *Ann Allergy Asthma Immunol* 2016;116(2):112-7.
134. Schatz M, Zeiger RS, Yang SJ, Chen W, Crawford W, Sajjan S, et al. The relationship of asthma impairment determined by psychometric tools to future asthma exacerbations. *Chest* 2012;141(1):66-72.
135. Tanaka A, Uno T, Sato H, Jinno M, Hirai K, Miyata Y, et al. Predicting future risk of exacerbations in Japanese patients with adult asthma: A prospective 1-year follow up study. *Allergol Int* 2017;66(4):568-73.
136. Tay TR, Wong HS, Tee A. Predictors of future exacerbations in a multi-ethnic Asian population with asthma. *J Asthma* 2018;1-8.
137. Tomita K, Sano H, Iwanaga T, Ishihara K, Ichinose M, Kawase I, et al. Association between episodes of upper respiratory infection and exacerbations in adult patients with asthma. *J Asthma* 2012;49(3):253-9.
138. Ko FWS, Hui DSC, Leung T-F, Chu H-Y, Wong GWK, Tung AHM, et al. Evaluation of the asthma control test: a reliable determinant of disease stability and a predictor of future exacerbations. *Respirology* 2012;17(2):370-8.
139. Meltzer EO, Busse WW, Wenzel SE, Belcheroff V, Weng HH, Feng J, et al. Use of the Asthma Control Questionnaire to predict future risk of asthma exacerbation. *J Allergy Clin Immunol* 2011;127(1):167-72.

140. Stephenson JJ, Quimbo RA, Gutierrez B. Subacute lack of asthma control as a predictor of subsequent acute asthma exacerbation in a managed care population. *Am J Manag Care* 2010;16(2):108-14.
141. Wei H-H, Zhou T, Wang L, Zhang H-P, Fu J-J, Wang L, et al. Current asthma control predicts future risk of asthma exacerbation: a 12-month prospective cohort study. *Chin Med J* 2012;125(17):2986-93.
142. Zeiger RS, Yegin A, Simons FER, Haselkorn T, Rasouliyan L, Szeffler SJ, et al. Evaluation of the National Heart, Lung, and Blood Institute guidelines impairment domain for classifying asthma control and predicting asthma exacerbations. *Ann Allergy Asthma Immunol* 2012;108(2):81-7.
143. Bloom CI, Nissen F, Douglas IJ, Smeeth L, Cullinan P, Quint JK. Exacerbation risk and characterisation of the UK's asthma population from infants to old age. *Thorax* 2018;73(4):313-20.
144. Zhang L, Zhang X, Zheng J, Wang L, Zhang H-P, Wang L, et al. Co-morbid psychological dysfunction is associated with a higher risk of asthma exacerbations: a systematic review and meta-analysis. *J Thorac Dis* 2016;8(6):1257-68.
145. Lehtimäki L, Csonka P, Mäkinen E, Isojärvi J, Hovi SL, Ahovuo-Saloranta A. Predictive value of exhaled nitric oxide in the management of asthma: a systematic review. *Eur Respir J* 2016;48(3):706-14.
146. Semprini R, Williams M, Semprini A, McDouall A, Fingleton J, Holweg C, et al. Type 2 Biomarkers and Prediction of Future Exacerbations and Lung Function Decline in Adult Asthma. *J Allergy Clin Immunol Pract* 2018;30:30.
147. Price DB, Rigazio A, Campbell JD, Bleecker ER, Corrigan CJ, Thomas M, et al. Blood eosinophil count and prospective annual asthma disease burden: a UK cohort study. *Lancet Respir Med* 2015;3(11):849-58.
148. Westerhof GA, de Groot JC, Amelink M, de Nijs SB, Ten Brinke A, Weersink EJ, et al. Predictors of frequent exacerbations in (ex)smoking and never smoking adults with severe asthma. *Respir Med* 2016;118:122-7.
149. Engelkes M, Janssens HM, de Jongste JC, Sturkenboom MCJM, Verhamme KMC. Medication adherence and the risk of severe asthma exacerbations: a systematic review. *Eur Respir J* 2015;45(2):396-407.
150. Killane I, Sulaiman I, MacHale E, Breathnach A, Taylor TE, Holmes MS, et al. Predicting asthma exacerbations employing remotely monitored adherence. *Healthc Technol Lett* 2016;3(1):51-5.
151. Lindgren P, Johnson J, Williams A, Yawn B, Pratt GC. Asthma exacerbations and traffic: examining relationships using link-based traffic metrics and a comprehensive patient database. *Environ Health* 2016;15(1):102.
152. Buelo A, McLean S, Julious S, Flores-Kim J, Bush A, Henderson J, et al. At-risk children with asthma (ARC): a systematic review. *Thorax* 2018;73(9):813-24.
153. National Committee for Quality Assurance. Use of appropriate medications for people with asthma and medication management for people with asthma. Available from <https://www.ncqa.org/hedis/measures/medication-management-for-people-with-asthma-and-asthma-medication-ratio/>; [Accessed. 03 May 2019].
154. Fitzpatrick AM, Jackson DJ, Mauger DT, Boehmer SJ, Phipatanakul W, Sheehan WJ, et al. Individualized therapy for persistent asthma in young children. *J Allergy Clin Immunol* 2016;138(6):1608-18.
155. Haselkorn T, Fish JE, Zeiger RS, Szeffler SJ, Miller DP, Chipps BE, et al. Consistently very poorly controlled asthma, as defined by the impairment domain of the Expert Panel Report 3 guidelines, increases risk for future severe asthma exacerbations in The Epidemiology and Natural History of Asthma: Outcomes and Treatment Regimens (TENOR) study. *J Allergy Clin Immunol* 2009;124(5):895-902.e1-4.
156. Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, et al. Standardisation of spirometry. *Eur Respir J* 2005;26(2):319-38.
157. Tweeddale PM, Alexander F, McHardy GJ. Short term variability in FEV₁ and bronchodilator responsiveness in patients with obstructive ventilatory defects. *Thorax* 1987;42(7):487-90.
158. Petsky Helen L, Kew Kayleigh M, Turner C, Chang Anne B. Exhaled nitric oxide levels to guide treatment for adults with asthma. *Cochrane Database of Systematic Reviews* 2016: Issue 9.
159. Petsky Helen L, Kew Kayleigh M, Chang Anne B. Exhaled nitric oxide levels to guide treatment for children with asthma. *Cochrane Database of Systematic Reviews* 2016: Issue 11.
160. Essat M, Harnan S, Gomersall T, Tappenden P, Wong R, Pavord I, et al. Fractional exhaled nitric oxide for the management of asthma in adults: a systematic review. *Eur Respir J* 2016;47(3):751-68.
161. Petsky Helen L, Li A, Chang Anne B. Tailored interventions based on sputum eosinophils versus clinical symptoms for asthma in children and adults. *Cochrane Database of Systematic Reviews* 2017: Issue 8.
162. Shimoda T, Obase Y, Nagasaka Y, Nakano H, Kishikawa R, Iwanaga T. Lung sound analysis can be an index of the control of bronchial asthma. *Allergol Int* 2017;66(1):64-9.
163. Hamill L, Ferris K, Kapande K, McConaghy L, Douglas I, McGovern V, et al. Exhaled breath temperature measurement and asthma control in children prescribed inhaled corticosteroids: A cross sectional study. *Pediatr Pulmonol* 2016;51(1):13-21.
164. Institute of Medicine of the National Academies. The 1st Annual Crossing the Quality Chasm Summit: a focus on communities. Washington D.C.: The National Academic Press; 2004.
165. Ring N, Jepson R, Hoskins G, Wilson C, Pinnock H, Sheikh A, et al. Understanding what helps or hinders asthma action plan use: a systematic review and synthesis of the qualitative literature. *Patient Educ Couns* 2011;85(2):e131-43.

166. Gibson PG, Powell H, Coughlan J, Wilson AJ, Abramson M, Haywood P, et al. Self-management education and regular practitioner review for adults with asthma (Cochrane Review). In: The Cochrane Library, 2003.
167. Lefevre F, Piper M, Weiss K, Mark D, Clark N, Aronson N. Do written action plans improve patient outcomes in asthma? An evidence-based analysis. *J Fam Pract* 2002;51(10):842-48.
168. Gibson PG, Powell H. Written action plans for asthma: an evidence-based review of the key components. *Thorax* 2004;59(2):94-9.
169. Powell H, Gibson PG. Options for self-management education for adults with asthma (Cochrane Review). In: The Cochrane Library, (1) 2003.
170. Bussey-Smith KL, Rossen RD. A systematic review of randomized control trials evaluating the effectiveness of interactive computerized asthma patient education programs. *Ann Allergy Asthma Immunol* 2007;98(6):507-16.
171. de Jongh T, Gurol-Urganci I, Vodopivec-Jamsek V, Car J, Atun R. Mobile phone messaging for facilitating self-management of long-term illnesses (Cochrane Review). In: The Cochrane Library, 2012.
172. Smith JR, Mugford M, Holland R, Noble MJ, Harrison BD. Psycho-educational interventions for adults with severe or difficult asthma: a systematic review. *J Asthma* 2007;44(3):219-41.
173. Boyd M, Lasserson TJ, McKean MC, Gibson PG, Ducharme FM, Haby M. Interventions for educating children who are at risk of asthma-related emergency department attendance (Cochrane Review). In: The Cochrane Library, 2009.
174. Coffman JM, Cabana MD, Halpin HA, Yelin EH. Effects of asthma education on children's use of acute care services: a meta-analysis. *Pediatrics* 2008;121(3):575-86.
175. Wolf FM, Guevara JP, Grum CM, Clark NM, Cates CJ. Educational interventions for asthma in children (Cochrane Review). In: The Cochrane Library, 2010.
176. Clarke SA, Calam R. The effectiveness of psychosocial interventions designed to improve health-related quality of life (HRQOL) amongst asthmatic children and their families: a systematic review. *Qual Life Res* 2012;21(5):747-64.
177. Bravata DM, Gienger AL, Holty JE, Sundaram V, Khazeni N, Wise PH, et al. Quality improvement strategies for children with asthma: a systematic review. *Archives of Pediatrics & Adolescent Medicine* 2009;163(6):572-81.
178. Bhogal S, Zemek R, Ducharme F. Written action plans for asthma in children (Cochrane Review). In: The Cochrane Library, (1) 2006.
179. Zemek RL, Bhogal SK, Ducharme FM. Systematic review of randomized controlled trials examining written action plans in children: what is the plan? *Arch Pediatr Adolesc Med* 2008;162(2):157-63.
180. Kessler KR. Relationship between the use of asthma action plans and asthma exacerbations in children with asthma: a systematic review. *J Asthma Allergy Educ* 2011;2(1):11-21.
181. Coffman JM, Cabana MD, Yelin EH. Do school-based asthma education programs improve self-management and health outcomes? *Pediatrics* 2009;124(2):729-42.
182. Ahmad E, Grimes DE. The effects of self-management education for school-age children on asthma morbidity: a systematic review. *J Sch Nurs* 2011;27(4):282-92.
183. Welsh EJ, Hasan M, Li L. Home-based educational interventions for children with asthma (Cochrane Review). In: The Cochrane Library, 2011.
184. Viswanathan M, Kraschnewski J, Nishikawa B, Morgan LC, Thieda P, Honeycutt A, et al. Outcomes of community health worker interventions. *Evid Rep Technol Assess (Full Rep)* 2009;181:1-144.
185. Bailey EJ, Cates CJ, Kruske SG, Morris PS, Brown N, Chang AB. Culture-specific programs for children and adults from minority groups who have asthma (Cochrane Review). In: The Cochrane Library, (2) 2009.
186. Press VG, Pappalardo AA, Conwell WD, Pincavage AT, Prochaska MH, Arora VM. Interventions to improve outcomes for minority adults with asthma: a systematic review. *J Gen Intern Med* 2012;27(8):1001-15.
187. Tapp S, Lasserson T, Rowe B. Education interventions for adults who attend the emergency room for acute asthma (Cochrane Review). In: The Cochrane Library, 2010.
188. Wilson SR, Latini D, Starr NJ, Fish L, Loes LM, Page A, et al. Education of parents of infants and very young children with asthma: a developmental evaluation of the Wee Wheezers program. *J Asthma* 1996;33(4):239-54.
189. Clark NM, Gong M, Schork MA, Evans D, Roloff D, Hurwitz M, et al. Impact of education for physicians on patient outcomes. *Pediatrics* 1998;101(5):831-6.
190. Stevens CA, Wesseldine LJ, Couriel JM, Dyer AJ, Osman LM, Silverman M. Parental education and guided self-management of asthma and wheezing in the pre-school child: a randomised controlled trial. *Thorax* 2002;57(1):39-44.
191. Butz AM, Malveaux FJ, Eggleston PA, Thompson L, K H, Rand CS. A review of community-based asthma interventions for inner-city children. *Pediatr Asthma Allergy Immunol* 1994;8(3):149-56.
192. Eakin MN, Rand CS, Bilderback A, Bollinger ME, Butz A, Kandasamy V, et al. Asthma in Head Start children: effects of the Breathmobile program and family communication on asthma outcomes. *J Allergy Clin Immunol* 2012;129(3):664-70.
193. Hederos CA, Janson S, Hedlin G. Group discussions with parents have long-term positive effects on the management of asthma with good cost-benefit. *Acta Paediatr* 2005;94(5):602-8.
194. Hederos CA, Janson S, Hedlin G. Six-year follow-up of an intervention to improve the management of preschool children with asthma. *Acta Paediatr* 2009;98(12):1939-44.

195. Szczepanski R, Jaeschke R, Spindler T, Ihorst G, Forster J, Group AS. Preschoolers' and parents' asthma education trial (P2AET)--a randomized controlled study. *Eur J Pediatr* 2010;169(9):1051-60.
196. Warschburger P, von Schwerin A-D, Buchholz HT, Petermann F. An educational program for parents of asthmatic preschool children: short- and medium-term effects. *Patient Educ Couns* 2003;51(1):83-91.
197. Bartholomew LK, Gold RS, Parcel GS, Czyzewski DI, Sockrider MM, Fernandez M, et al. Watch, Discover, Think, and Act: evaluation of computer-assisted instruction to improve asthma self-management in inner-city children. *Patient Educ Couns* 2000;39(2-3):269-80.
198. Brown JV, Bakeman R, Celano MP, Demi AS, Kobrynski L, Wilson SR. Home-based asthma education of young low-income children and their families. *J Pediatr Psychol* 2002;27(8):677-88.
199. Fisher EB, Strunk RC, Sussman LK, Sykes RK, Walker MS. Community organization to reduce the need for acute care for asthma among African American children in low-income neighborhoods: the Neighborhood Asthma Coalition. *Pediatrics* 2004;114(1):116-23.
200. La Roche MJ, Koinis-Mitchell D, Gualdrón L. A culturally competent asthma management intervention: a randomized controlled pilot study. *Ann Allergy Asthma Immunol* 2006;96(1):80-5.
201. Joseph CL, Peterson E, Havstad S, Johnson CC, Hoerauf S, Stringer S, et al. A web-based, tailored asthma management program for urban African-American high school students. *Am J Respir Crit Care Med* 2007;175(9):888-95.
202. Mosnaim GS, Cohen MS, Rhoads CH, Rittner SS, Powell LH. Use of MP3 players to increase asthma knowledge in inner-city African-American adolescents. *Int J Behav Med* 2008;15(4):341-6.
203. Flores G, Bridon C, Torres S, Perez R, Walter T, Brotanek J, et al. Improving asthma outcomes in minority children: a randomized, controlled trial of parent mentors. *Pediatrics* 2009;124(6):1522-32.
204. Martin M, Catrambone C, Kee R, Evans A, Sharp L, Lyttle C, et al. Improving asthma self-efficacy: developing and testing a pilot community-based asthma intervention for African American adults. *J Allergy Clin Immunol* 2009;123(1):153-9.
205. Velsor-Friedrich B, Militello LK, Richards MH, Harrison PR, Gross IM, Romero E, et al. Effects of coping-skills training in low-income urban African-American adolescents with asthma. *J Asthma* 2012;49(4):372-9.
206. Moudgil H, Marshall T, Honeybourne D. Asthma education and quality of life in the community: a randomised controlled study to evaluate the impact on white European and Indian subcontinent ethnic groups from socioeconomically deprived areas in Birmingham, UK. *Thorax* 2000;55(3):177-83.
207. Griffiths C, Foster G, Barnes N, Eldridge S, Tate H, Begum S, et al. Specialist nurse intervention to reduce unscheduled asthma care in a deprived multiethnic area: the east London randomised controlled trial for high risk asthma (ELECTRA). *British Medical Journal* 2004;328(7432):144.
208. Poureslami I, Nimmon L, Doyle-Waters M, Rootman I, Schulzer M, Kuramoto L, et al. Effectiveness of educational interventions on asthma self-management in Punjabi and Chinese asthma patients: a randomized controlled trial. *Journal of Asthma* 2012;49(5):542-51.
209. Barbanel D, Eldridge S, Griffiths C. Can a self-management programme delivered by a community pharmacist improve asthma control? A randomised trial. *Thorax* 2003;58(10):851-4.
210. Nokela M, Arnlin M, Ehlers P-O, Krakau I, Forslund L, Jonsson EW. The influence of structured information and monitoring on the outcome of asthma treatment in primary care: a cluster randomized study. *Respiration* 2010;79(5):388-94.
211. Partridge MR, Caress AL, Brown C, Hennings J, Luker K, Woodcock A, et al. Can lay people deliver asthma self-management education as effectively as primary care based practice nurses? *Thorax* 2008;63(9):778-83.
212. Guendelman S, Meade K, Benson M, Chen YQ, Samuels S. Improving asthma outcomes and self-management behaviors of inner-city children: a randomized trial of the Health Buddy interactive device and an asthma diary. *Arch Pediatr Adolesc Med* 2002;156(2):114-20.
213. Delaronde S, Peruccio DL, Bauer BJ. Improving asthma treatment in a managed care population. *Am J Manag Care* 2005;11(6):361-8.
214. Feifer RA, Verbrugge RR, Khalid M, Levin R, O'Keefe GB, Aubert RE. Improvements in asthma pharmacotherapy and self-management: An example of a population-based disease management program. *Dis Manag Health Outcomes* 2004;12(2):93-102.
215. Glasgow NJ, Ponsonby AL, Yates R, Beilby J, Dugdale P. Proactive asthma care in childhood: general practice based randomised controlled trial. *British Medical Journal* 2003;327(7416):659.
216. Homer CJ, Forbes P, Horvitz L, Peterson LE, Wypij D, Heinrich P. Impact of a quality improvement program on care and outcomes for children with asthma. *Arch Pediatr Adolesc Med* 2005;159(5):464-9.
217. Cleland JA, Hall S, Price D, Lee AJ. An exploratory, pragmatic, cluster randomised trial of practice nurse training in the use of asthma action plans. *Prim Care Respir J* 2007;16(5):311-8.
218. Kew Kayleigh M, Quinn M, Quon Bradley S, Ducharme Francine M. Increased versus stable doses of inhaled corticosteroids for exacerbations of chronic asthma in adults and children. *Cochrane Database of Systematic Reviews* 2016: Issue 6.
219. McKeever T, Mortimer K, Wilson A, Walker S, Brightling C, Skeggs A, et al. Quadrupling Inhaled Glucocorticoid Dose to Abort Asthma Exacerbations. *N Engl J Med* 2018;378(10):902-10.
220. Jackson D, LB B, DT M. Quintupling Inhaled Glucocorticoids to Prevent Childhood Asthma Exacerbations. *N Engl J Med* 2018;378(10):891-901.

221. Thoonen BP, Schermer TR, Van Den Boom G, Molema J, Folgering H, Akkermans RP, et al. Self-management of asthma in general practice, asthma control and quality of life: a randomised controlled trial. *Thorax* 2003;58(1):30-6.
222. Osman LM, Abdalla MI, Beattie JA, Ross SJ, Russell IT, Friend JA, et al. Reducing hospital admission through computer supported education for asthma patients. *British Medical Journal* 1994;308(6928):568-71.
223. Osman LM, Calder C, Godden DJ, Friend JA, McKenzie L, Legge JS, et al. A randomised trial of self-management planning for adult patients admitted to hospital with acute asthma. *Thorax* 2002;57(10):869-74.
224. Yoon R, McKenzie DK, Bauman A, Miles DA. Controlled trial evaluation of an asthma education programme for adults. *Thorax* 1993;48(11):1110-6.
225. Madge P, McColl J, Paton J. Impact of a nurse-led home management training programme in children admitted to hospital with acute asthma: a randomised controlled study. *Thorax* 1997;52(3):223-8.
226. Royal Pharmaceutical Society of Great Britain. From compliance to concordance: achieving shared goals in medicine taking. London: Royal Pharmaceutical Society of Great Britain; 1997. [cited
227. Wilson SR, Strub P, Buist AS, Knowles SB, Lavori PW, Lapidus J, et al. Shared treatment decision making improves adherence and outcomes in poorly controlled asthma. *Am J Respir Crit Care Med* 2010;181(6):566-77.
228. Garrett J, Fenwick JM, Taylor G, Mitchell E, Rea H. Peak expiratory flow meters (PEFMs)--who uses them and how and does education affect the pattern of utilisation? *Aust N Z J Med* 1994;24(5):521-9.
229. Redline S, Wright EC, Kattan M, Kerckmar C, Weiss K. Short-term compliance with peak flow monitoring: results from a study of inner city children with asthma. *Pediatr Pulmonol* 1996;21(4):203-10.
230. Effectiveness of routine self monitoring of peak flow in patients with asthma. Grampian Asthma Study of Integrated Care (GRASSIC). *BMJ* 1994;308(6928):564-7.
231. Burkhart PV, Dunbar-Jacob JM, Fireman P, Rohay J. Children's adherence to recommended asthma self-management. *Pediatr Nurs* 2002;28(4):409-14.
232. Yoos HL, Kitzman H, McMullen A, Henderson C, Sidora K. Symptom monitoring in childhood asthma: a randomized clinical trial comparing peak expiratory flow rate with symptom monitoring. *Annals of Allergy, Asthma, & Immunology* 2002;88(3):283-91.
233. National Collaborating Centre for Primary Care. Clinical Guidelines and Evidence Review for Medicines Adherence: involving patients in decisions about prescribed medicines and supporting adherence. London: NICE; 2009. (NICE guideline GC76). [cited 10 Jul 2014] Available from <http://www.nice.org.uk/Guidance/CG76>
234. Krishnan JA, Bender BG, Wamboldt FS, Szefer SJ, Adkinson Jr NF, Zeiger RS, et al. Adherence to inhaled corticosteroids: An ancillary study of the Childhood Asthma Management Program clinical trial. *Journal of Allergy and Clinical Immunology* 2012;129(1):112-8.
235. Reznik M, Ozuah PO. Measurement of inhaled corticosteroid adherence in inner-city, minority children with persistent asthma by parental report and integrated dose counter. *J Allergy (Cairo)* 2012(570850).
236. Schultz A, Sly PD, Zhang G, Venter A, Devadason SG, le Souef PN. Usefulness of parental response to questions about adherence to prescribed inhaled corticosteroids in young children. *Archives of Disease in Childhood* 2012;97(12):1092-6.
237. Blais L, Kettani FZ, Beauchesne MF, Lemiere C, Perreault S, Forget A. New measure of adherence adjusted for prescription patterns: The case of adults with asthma treated with inhaled corticosteroid monotherapy. *Annals of Pharmacotherapy* 2011;45(3):335-41.
238. Taylor A, Chen L-C, Smith MD. Adherence to inhaled corticosteroids by asthmatic patients: measurement and modelling. *Int J Clin Pharm* 2014;36(1):112-9.
239. Patel M, Perrin K, Pritchard A, Williams M, Wijesinghe M, Weatherall M, et al. Accuracy of patient self-report as a measure of inhaled asthma medication use. *Respirology* 2013;18(3):546-52.
240. Horne R, Weinman J, Hankins M. The beliefs about medicines questionnaire: the development and evaluation of a new method for assessing the cognitive representation of medication. *Psychol Health* 1999;14(1):1-24.
241. Van Steenis M, Driesenaar J, Bensing J, Van Hulten R, Souverein P, Van Dijk L, et al. Relationship between medication beliefs, self-reported and refill adherence, and symptoms in patients with asthma using inhaled corticosteroids. *Patient Prefer Adherence* 2014;8:83-91.
242. Horne R, Chapman SC, Parham R, Freemantle N, Forbes A, Cooper V. Understanding patients' adherence-related beliefs about medicines prescribed for long-term conditions: a meta-analytic review of the Necessity-Concerns Framework. *PloS one* 2013;8(12):e80633.
243. Vollmer WM, Xu M, Feldstein A, Smith D, Waterbury A, Rand C. Comparison of pharmacy-based measures of medication adherence. *BMC health services research* 2012; 155.
244. McNicholl DM, Stevenson M, McGarvey LP, Heaney LG. The utility of fractional exhaled nitric oxide suppression in the identification of nonadherence in difficult asthma. *Am J Respir Crit Care Med* 2012;186(11):1102-8.
245. Hagan JB, Netzel BC, Matthews MR, Korpi-Steiner NL, Singh RJ. Urinary fluticasone propionate-17beta-carboxylic acid to assess asthma therapy adherence. *Allergy Asthma Proc* 2012;33(4):e35-9.
246. Geryk LL, Roberts CA, Carpenter DM. A systematic review of school-based interventions that include inhaler technique education. *Respir Med* 2017;132:21-30.

247. Hui CY, Walton R, McKinstry B, Jackson T, Parker R, Pinnock H. The use of mobile applications to support self-management for people with asthma: a systematic review of controlled studies to identify features associated with clinical effectiveness and adherence. *J Am Med Inform Assoc* 2017;24(3): 619-32.
248. Miller L, Schuz B, Walters J, Walters EH. Mobile Technology Interventions for Asthma Self-Management: Systematic Review and Meta-Analysis. *JMIR Mhealth Uhealth* 2017;5(5):e57.
249. Moullec G, Gour-Provençal G, Bacon SL, Campbell TS, Lavoie KL. Efficacy of interventions to improve adherence to inhaled corticosteroids in adult asthmatics: Impact of using components of the chronic care model. *Respiratory Medicine* 2012;106(9):1211-25.
250. Normansell R, Kew KM, Stovold E. Interventions to improve adherence to inhaled steroids for asthma (Cochrane Review). In: *The Cochrane Library*, 2017.
251. Britto MT, Rohan JM, Dodds CM, Byczkowski TL. A Randomized Trial of User-Controlled Text Messaging to Improve Asthma Outcomes: A Pilot Study. *Clin Pediatr (Phila)* 2017;56(14):1336-44.
252. Johnson KB, Patterson BL, Ho YX, Chen Q, Nian H, Davison CL, et al. The feasibility of text reminders to improve medication adherence in adolescents with asthma. *J Am Med Inform Assoc* 2016;23(3):449-55.
253. Wells KE, Peterson EL, Ahmedani BK, Williams LK. Real-world effects of once vs greater daily inhaled corticosteroid dosing on medication adherence. *Annals of Allergy, Asthma, & Immunology* 2013;111(3):216-20.
254. Bender BG, Cvietusa PJ, Goodrich GK, Lowe R, Nuanes HA, Rand C, et al. Pragmatic trial of health care technologies to improve adherence to pediatric asthma treatment: a randomized clinical trial. *Jama, Pediatr* 2015;169(4):317-23.
255. Foster JM, Usherwood T, Smith L, Sawyer SM, Xuan W, Rand CS, et al. Inhaler reminders improve adherence with controller treatment in primary care patients with asthma. *Journal of Allergy & Clinical Immunology* 2014;134(6):1260-8.e3.
256. Kripalani S, Yao X, Haynes RB. Interventions to enhance medication adherence in chronic medical conditions: a systematic review. *Arch Intern Med* 2007;167(6):540-50.
257. Kahana S, Drotar D, Frazier T. Meta-analysis of psychological interventions to promote adherence to treatment in pediatric chronic health conditions. *J Pediatr Psychol* 2008;33(6):590-611.
258. Viswanathan M, Golin CE, Jones CD, Ashok M, Blalock SJ, Wines RCM, et al. Interventions to improve adherence to self-administered medications for chronic diseases in the United States: a systematic review. *Annals of Internal Medicine* 2012;157(11):785-95.
259. Brooks TL, Leventhal H, Wolf MS, O'Connor R, Morillo J, Martynenko M, et al. Strategies used by older adults with asthma for adherence to inhaled corticosteroids. *J Gen Intern Med* 2014;29(11):1506-12.
260. Kenyon CC, Gruschow SM, Quarshie WO, Griffis H, Leach MC, Zorc JJ, et al. Controller adherence following hospital discharge in high risk children: A pilot randomized trial of text message reminders. *J Asthma* 2018;1-9.
261. Wiener-Ogilvie S, Pinnock H, Huby G, Sheikh A, Partridge MR, Gillies J. Do practices comply with key recommendations of the British Asthma Guideline? If not, why not? *Prim Care Respir J* 2007;16(6): 369-77.
262. Asthma UK. Compare your care. Available from <http://www.asthma.org.uk/support-us/campaigns/publications/survey>: [Accessed. 10 Jul 2014].
263. Ring N, Malcolm C, Wyke S, Macgillivray S, Dixon D, Hoskins G, et al. Promoting the use of Personal Asthma Action Plans: a systematic review. *Prim Care Respir J* 2007;16(5):271-83.
264. Bunik M, Federico MJ, Beaty B, Rannie M, Olin JT, Kempe A. Quality improvement for asthma care within a hospital-based teaching clinic. *Acad Pediatr* 2011;11(1):58-65.
265. Bunting BA, Cranor CW. The Asheville Project: long-term clinical, humanistic, and economic outcomes of a community-based medication therapy management program for asthma. *J Am Pharm Assoc* (2003) 2006;46(2):133-47.
266. Gerald LB, Redden D, Wittich AR, Hains C, Turner-Henson A, Hemstreet MP, et al. Outcomes for a comprehensive school-based asthma management program. *J Sch Health* 2006;76(6):291-6.
267. Vollmer WM, Kirshner M, Peters D, Drane A, Stibolt T, Hickey T, et al. Use and impact of an automated telephone outreach system for asthma in a managed care setting. *Am J Manag Care* 2006;12(12):725-33.
268. Forshee JD, Whalen EB, Hackel R, Butt LT, Smeltzer PA, Martin J, et al. The effectiveness of one-on-one nurse education on the outcomes of high-risk adult and pediatric patients with asthma. *Manag Care Interface* 1998;11(12):82-92.
269. Findley SE, Thomas G, Madera-Reese R, McLeod N, Kintala S, Andres Martinez R, et al. A community-based strategy for improving asthma management and outcomes for preschoolers. *J Urban Health* 2011;88 Suppl 1:85-99.
270. Polivka BJ, Chaudry RV, Crawford J, Bouton P, Sweet L. Impact of an urban healthy homes intervention. *J Environ Health* 2011;73(9):16-20.
271. Kempe T, Rogers C. A mailed personalised self-management plan improves attendance and increases patients' understanding of asthma. *Prim Care Respir J* 2003;12(4):110-4.
272. Swanson V, Wright S, Power KG, Duncan B, Morgan J, Turner E, et al. The impact of a structured programme of asthma care in general practice. *Int J Clin Pract* 2000;54(9):573-80.
273. Lindberg M, Ahlner J, Ekstrom T, Jonsson D, Moller M. Asthma nurse practice improves outcomes and reduces costs in primary health care. *Scand J Caring Sci* 2002;16(1):73-8.

274. Haahtela T, Tuomisto LE, Pietinalho A, Klaukka T, Erhola M, Kaila M, et al. A 10 year asthma programme in Finland: major change for the better. *Thorax* 2006;61(8):663-70.
275. Kauppi P, Linna M, Martikainen J, Makela MJ, Haahtela T. Follow-up of the Finnish Asthma Programme 2000-2010: reduction of hospital burden needs risk group rethinking. *Thorax* 2013;68(3):292-3.
276. Chini L, Iannini R, Chianca M, Corrente S, Graziani S, La Rocca M, et al. Happy air®, a successful school-based asthma educational and interventional program for primary school children. *J Asthma* 2011;48(4):419-26.
277. Andrade WC, Camargos P, Lasmar L, Bousquet J. A pediatric asthma management program in a low-income setting resulting in reduced use of health service for acute asthma. *Allergy* 2010;65(11):1472-7.
278. Souza-Machado C, Souza-Machado A, Franco R, Ponte EV, Barreto ML, Rodrigues LC, et al. Rapid reduction in hospitalisations after an intervention to manage severe asthma. *Eur Respir J* 2010;35(3):515-21.
279. Maas T, Kaper J, Sheikh A, Knottnerus JA, Wesseling G, Dompeling E, et al. Mono and multifaceted inhalant and/or food allergen reduction interventions for preventing asthma in children at high risk of developing asthma (Cochrane Review). In: *The Cochrane Library*, 2009.
280. Maas T, Dompeling E, Muris J, Wesseling G, Knottnerus J, van Schayck OC. Prevention of asthma in genetically susceptible children: a multifaceted intervention trial focussed on feasibility in general practice. *Pediatr Allergy Immunol* 2011;22(8):794-802.
281. Wahn U, Lau S, Bergmann R, Kulig M, Forster J, Bergmann K, et al. Indoor allergen exposure is a risk factor for sensitization during the first three years of life. *J Allergy Clin Immunol* 1997;99(6 Pt 1):763-9.
282. Corver K, Kerkhof M, Brussee JE, Brunekreef B, van Strien RT, Vos AP, et al. House dust mite allergen reduction and allergy at 4 yr: follow up of the PIAMA-study. *Pediatr Allergy Immunol* 2006;17(5):329-36.
283. Lau S, Illi S, Sommerfeld C, Niggemann B, Bergmann R, von Mutius E, et al. Early exposure to house-dust mite and cat allergens and development of childhood asthma: a cohort study. Multicentre Allergy Study Group. *Lancet* 2000;356(9239):1392-7.
284. Arshad SH, Bateman B, Matthews SM. Primary prevention of asthma and atopy during childhood by allergen avoidance in infancy: a randomised controlled study. *Thorax* 2003;58(6):489-93.
285. Sporik R, Holgate ST, Platts-Mills TA, Cogswell JJ. Exposure to house-dust mite allergen (Der p I) and the development of asthma in childhood. A prospective study. *N Engl J Med* 1990;323(8):502-7.
286. Cullinan P, MacNeill SJ, Harris JM, Moffat S, White C, Mills P, et al. Early allergen exposure, skin prick responses, and atopic wheeze at age 5 in English children: a cohort study. *Thorax* 2004;59(10):855-61.
287. Chan-Yeung M, Ferguson A, Watson W, Dimich-Ward H, Rousseau R, Lilley M, et al. The Canadian Childhood Asthma Primary Prevention Study: outcomes at 7 years of age. *J Allergy Clin Immunol* 2005;116(1):49-55.
288. Horak F Jr, Matthews S, Ihorst G, Arshad SH, Frischer T, Kuehr J, et al. Effect of mite-impermeable mattress encasings and an educational package on the development of allergies in a multinational randomized, controlled birth-cohort study -- 24 months results of the Study of Prevention of Allergy in Children in Europe. *Clin Exp Allergy* 2004;34(8):1220-5.
289. Custovic A, Simpson BM, Simpson A, Kissen P, Woodcock A. Effect of environmental manipulation in pregnancy and early life on respiratory symptoms and atopy during the first year of life: a randomised trial. *Lancet* 2001;358(9277):188-93.
290. Woodcock A, Lowe LA, Murray CS, Simpson BM, Pipis SD, Kissen P, et al. Early life environmental control: effect on symptoms, sensitization, and lung function at age 3 years. *Am J Respir Crit Care Med* 2004;170(4):433-9.
291. Takkouche B, Gonzalez-Barcala FJ, Etminan M, Fitzgerald M. Exposure to furry pets and the risk of asthma and allergic rhinitis: a meta-analysis. *Allergy* 2008;63(7):857-64.
292. Lodge CJ, Allen KJ, Lowe AJ, Hill DJ, Hosking CS, Abramson MJ, et al. Perinatal cat and dog exposure and the risk of asthma and allergy in the urban environment: a systematic review of longitudinal studies. *Clin Dev Immunol* 2012;2012:176484.
293. Chen C, Tischer C, Schnappinger M, Heinrich J. The role of cats and dogs in asthma and allergy--a systematic review. *Int J Hyg Environ Health* 2010;213(1):1-31.
294. Lødrup Carlsen KC, Roll S, Carlsen K, Mowinckel P, Wijga A, Brunekreef B, et al. Does pet ownership in infancy lead to asthma or allergy at school age? Pooled analysis of individual participant data from 11 European birth cohorts. *PLoS ONE* 2012;7(8):e43214.
295. Muraro A, Dreborg S, Halken S, Høst A, Niggemann B, Aalberse R, et al. Dietary prevention of allergic diseases in infants and small children. Part I: immunologic background and criteria for hypoallergenicity. *Pediatr Allergy Immunol* 2004;15(2):103-11.
296. Muraro A, Dreborg S, Halken S, Høst A, Niggemann B, Aalberse R, et al. Dietary prevention of allergic diseases in infants and small children. Part III: Critical review of published peer-reviewed observational and interventional studies and final recommendations. *Pediatr Allergy Immunol* 2004;15(4):291-307.
297. Kramer MS, Kakuma R. Maternal dietary antigen avoidance during pregnancy or lactation, or both, for preventing or treating atopic disease in the child (Cochrane Review). In: *The Cochrane Library*, 2006.

298. Vance GH, Grimshaw KE, Briggs R, Lewis SA, Mullee MA, Thornton CA, et al. Serum ovalbumin-specific immunoglobulin G responses during pregnancy reflect maternal intake of dietary egg and relate to the development of allergy in early infancy. *Clin Exp Allergy* 2004;34(12):1855-61.
299. van Odiijk J, Kull I, Borres MP, Brandtzaeg P, Edberg U, Hanson LA, et al. Breast feeding and allergic disease: a multi-disciplinary review of the literature (1996-2001) on the mode of early feeding in infancy and its impact on later atopic manifestations. *Allergy* 2003;58(9):833-43.
300. Sears MR, Greene JM, Willan AR, Taylor DR, Flannery EM, Cowan JO, et al. Long-term relation between breastfeeding and development of atopy and asthma in children and young adults: a longitudinal study. *Lancet* 2002;360(9337):901-7.
301. Osborn DA, Sinn J. Formulas containing hydrolysed protein for prevention of allergy and food intolerance in infants (Cochrane Review). In: The Cochrane Library, 2006.
302. Osborn DA, Sinn J. Soy formula for prevention of allergy and food intolerance in infants (Cochrane Review). In: The Cochrane Library, 2006.
303. Tricon S, Willers S, Smit HA, Burney PG, Devereux G, Frew AJ, et al. Nutrition and allergic disease. *Clin Exp Allergy Rev* 2006;6(5):117-88.
304. Zutavern A, von Mutius E, Harris J, Mills P, Moffatt S, White C, et al. The introduction of solids in relation to asthma and eczema. *Arch Dis Child* 2004;89(4):303-8.
305. Dunstan JA, Mori TA, Barden A, Beilin LJ, Taylor AL, Holt PG, et al. Fish oil supplementation in pregnancy modifies neonatal allergen-specific immune responses and clinical outcomes in infants at high risk of atopy: a randomized, controlled trial. *J Allergy Clin Immunol* 2003;112(6):1178-84.
306. Mahrshahi S, Peat JK, Webb K, Oddy W, Marks GB, Mellis CM. Effect of omega-3 fatty acid concentrations in plasma on symptoms of asthma at 18 months of age. *Pediatr Allergy Immunol* 2004;15(6):517-22.
307. Shaheen SO, Newson RB, Henderson AJ, Emmett PM, Sherriff A, Cooke M. Umbilical cord trace elements and minerals and risk of early childhood wheezing and eczema. *Eur Respir J* 2004;24(2):292-7.
308. Devereux G, Turner SW, Craig LC, McNeill G, Martindale S, Harbour PJ, et al. Low maternal vitamin E intake during pregnancy is associated with asthma in 5-year-old children. *Am J Respir Crit Care Med* 2006;174(5):499-507.
309. Flaherman V, Rutherford GW. A meta-analysis of the effect of high weight on asthma. *Arch Dis Child* 2006;91(4):334-9.
310. Beuther DA, Sutherland ER. Overweight, obesity, and incident asthma: a meta-analysis of prospective epidemiologic studies. *Am J Respir Crit Care Med* 2007;175(7):661-6.
311. Chen YC, Dong GH, Lin KC, Lee YL. Gender difference of childhood overweight and obesity in predicting the risk of incident asthma: a systematic review and meta-analysis. *Obes Rev* 2013;14(3):222-31.
312. Egan KB, Ettinger AS, Bracken MB, Egan KB, Ettinger AS, Bracken MB. Childhood body mass index and subsequent physician-diagnosed asthma: a systematic review and meta-analysis of prospective cohort studies. *BMC Pediatr* 2013;13:121.
313. Forno E, Young OM, Kumar R, Simhan H, Celdon JC. Maternal obesity in pregnancy, gestational weight gain, and risk of childhood asthma. *Pediatrics* 2014;134(2):e535-46.
314. Holt PG, Sly PD, Bjorksten B. Atopic versus infectious diseases in childhood: a question of balance? *Pediatr Allergy Immunol* 1997;8(2):53-8.
315. Strachan DP. Family size, infection and atopy: the first decade of the "hygiene hypothesis". *Thorax* 2000;55(Suppl 1):S2-10.
316. Kalliomaki M, Salminen S, Arvilommi H, Kero P, Koskinen P, Isolauri E. Probiotics in primary prevention of atopic disease: a randomised placebo-controlled trial. *Lancet* 2001;357(9262):1076-9.
317. Moro G, Arslanoglu S, Stahl B, Jelinek J, Wahn U, Boehm G. A mixture of prebiotic oligosaccharides reduces the incidence of atopic dermatitis during the first six months of age. *Arch Dis Child* 2006;91(10):814-9.
318. Cook DG, Strachan DP. Health effects of passive smoking-10: Summary of effects of parental smoking on the respiratory health of children and implications for research. *Thorax* 1999;54(4):357-66.
319. Dezateau C, Stocks J, Dundas I, Fletcher ME. Impaired airway function and wheezing in infancy: the influence of maternal smoking and a genetic predisposition to asthma. *Am J Respir Crit Care Med* 1999;159(2):403-10.
320. Gilliland FD, Berhane K, McConnell R, Gauderman WJ, Vora H, Rappaport EB, et al. Maternal smoking during pregnancy, environmental tobacco smoke exposure and childhood lung function. *Thorax* 2000;55(4):271-6.
321. Lodrup Carlsen KC, Carlsen KH, Nafstad P, Bakkevig L. Perinatal risk factors for recurrent wheeze in early life. *Pediatr Allergy Immunol* 1999;10(2):89-95.
322. Arshad SH, Kurukulaaratchy RJ, Fenn M, Matthews S. Early life risk factors for current wheeze, asthma, and bronchial hyperresponsiveness at 10 years of age. *Chest* 2005;127(2):502-8.
323. Jaakkola JJ, Gissler M. Maternal smoking in pregnancy, fetal development, and childhood asthma. *Am J Public Health* 2004;94(1):136-40.
324. Kabisch M, Hoefler C, Carr D, Leupold W, Weiland SK, von Mutius E. Glutathione S transferase deficiency and passive smoking increase childhood asthma. *Thorax* 2004;59(7):569-73.
325. Belanger K, Beckett W, Triche E, Bracken MB, Holford T, Ren P, et al. Symptoms of wheeze and persistent cough in the first year of life: associations with indoor allergens, air contaminants, and maternal history of asthma. *Am J Epidemiol* 2003;158(3):195-202.

326. Lee YL, Lin YC, Lee YC, Wang JY, Hsiue TR, Guo YL. Glutathione S-transferase P1 gene polymorphism and air pollution as interactive risk factors for childhood asthma. *Clin Exp Allergy* 2004;34(11):1707-13.
327. Miller RL, Garfinkel R, Horton M, Camann D, Perera FP, Whyatt RM, et al. Polycyclic aromatic hydrocarbons, environmental tobacco smoke, and respiratory symptoms in an inner-city birth cohort. *Chest* 2004;126(4):1071-8.
328. Romieu I, Sienra-Monge JJ, Ramirez-Aguilar M, Moreno-Macias H, Reyes-Ruiz NI, Estela del Rio-Navarro B, et al. Genetic polymorphism of GSTM1 and antioxidant supplementation influence lung function in relation to ozone exposure in asthmatic children in Mexico City. *Thorax* 2004;59(1):8-10.
329. Kemp A, Bjorksten B. Immune deviation and the hygiene hypothesis: a review of the epidemiological evidence. *Pediatr Allergy Immunol* 2003;14(2):74-80.
330. Martignon G, Oryszczyn MP, Annesi-Maesano I. Does childhood immunization against infectious diseases protect from the development of atopic disease? *Pediatr Allergy Immunol* 2005;16(3):193-200.
331. Leas BF, D'Anci KE, Apter AJ, Bryant-Stephens T, Lynch MP, Kaczmarek JL, et al. Effectiveness of Indoor Allergen Reduction in the Management of Asthma: A Systematic Review. *J Allergy Clin Immunol* 2018;13:13.
332. Wood RA, Chapman MD, Adkinson NF Jr, Eggleston PA. The effect of cat removal on allergen content in household-dust samples. *J Allergy Clin Immunol* 1989;83(4):730-4.
333. Wood RA, Johnson EF, Van Natta ML, Chen PH, Eggleston PA. A placebo-controlled trial of a HEPA air cleaner in the treatment of cat allergy. *Am J Respir Crit Care Med* 1998;158(1):115-20.
334. Platts-Mills T, Vaughan J, Squillace S, Woodfolk J, Sporik R. Sensitisation, asthma, and a modified Th2 response in children exposed to cat allergen: a population-based cross-sectional study. *Lancet* 2001;357(9258):752-6.
335. Singh M, Jaiswal N. Dehumidifiers for chronic asthma. *Cochrane Database of Systematic Reviews* 2013: Issue 6.
336. Chalmers GW, MacLeod KJ, Little SA, Thomson LJ, McSharry CP, Thomson NC. Influence of cigarette smoking on inhaled corticosteroid treatment in mild asthma. *Thorax* 2002;57(3):226-30.
337. Ehrlich R, Jordaan E, Du Toit D, Potter P, Volmink J, Zwarenstein M, et al. Household smoking and bronchial hyperresponsiveness in children with asthma. *J Asthma* 2001;38(3):239-51.
338. Gallefoss F, Bakke PS. Does smoking affect the outcome of patient education and self-management in asthmatics? *Patient Educ Couns* 2003;49(1):91-7.
339. Mannino DM, Homa DM, Redd SC. Involuntary smoking and asthma severity in children: Data from the Third National Health and Nutrition Examination Survey. *Chest* 2002;122(2):409-15.
340. Dick S, Doust E, Cowie H, Ayres JG, Turner S. Associations between environmental exposures and asthma control and exacerbations in young children: a systematic review. *BMJ Open* 2014;4(2):e003827.
341. Behbod B, Sharma M, Baxi R, Roseby R, Webster P. Family and carer smoking control programmes for reducing children's exposure to environmental tobacco smoke. *Cochrane Database of Systematic Reviews* 2018: Issue 1.
342. Tonnesen P, Pisinger C, Hvidberg S, Wennike P, Bremann L, Westin A, et al. Effects of smoking cessation and reduction in asthmatics. *Nicotine Tob Res* 2005;7(1):139-48.
343. Rasmussen F, Siersted HC, Lambrechtsen J, Hansen HS, Hansen NC. Impact of airway lability, atopy, and tobacco smoking on the development of asthma-like symptoms in asymptomatic teenagers. *Chest* 2000;117(5):1330-5.
344. Devalia JL, Rusznak C, Herdman MJ, Trigg CJ, Tarraf H, Davies RJ. Effect of nitrogen dioxide and sulphur dioxide on airway response of mild asthmatic patients to allergen inhalation. *Lancet* 1994;344(8938):1668-71.
345. Molfino NA, Wright SC, Katz I, Tarlo S, Silverman F, McClean PA, et al. Effect of low concentrations of ozone on inhaled allergen responses in asthmatic subjects. *Lancet* 1991;338(8761):199-203.
346. Department of Health, Committee on the Medical Effects of Air Pollutants. *Asthma and outdoor air pollution*. London: HMSO; 1995. [cited
347. Lin M, Chen Y, Burnett RT, Villeneuve PJ, Krewski D. Effect of short-term exposure to gaseous pollution on asthma hospitalisation in children: a bi-directional case-crossover analysis. *J Epidemiol Community Health* 2003;57(1):50-5.
348. Norbäck D, Björnsson E, Janson C, Widstrom J, Boman G. Asthmatic symptoms and volatile organic compounds, formaldehyde, and carbon dioxide in dwellings. *Occup Environ Med* 1995;52(6):388-95.
349. Tunnicliffe WS, Burge PS, Ayres JG. Effect of domestic concentrations of nitrogen dioxide on airway responses to inhaled allergen in asthmatic patients. *Lancet* 1994;344(8939-40):1733-6.
350. Burney P. A diet rich in sodium may potentiate asthma. *Epidemiologic evidence for a new hypothesis*. *Chest* 1987;91(6 Suppl):143S-8S.
351. Burney PG. The causes of asthma--does salt potentiate bronchial activity? Discussion paper. *J R Soc Med* 1987;80(6):364-7.
352. Mickleborough TD, Lindley MR, Ray S. Dietary salt, airway inflammation, and diffusion capacity in exercise-induced asthma. *Med Sci Sports Exerc* 2005;37(6):904-14.
353. Arden KD, Ram FS. Dietary salt reduction or exclusion for allergic asthma (Cochrane Review). In: *The Cochrane Library*, 2001.
354. Britton J, Pavord I, Richards K, Wisniewski A, Knox A, Lewis S, et al. Dietary magnesium, lung function, wheezing, and airway hyperreactivity in a random adult population sample. *Lancet* 1994;344(8919):357-62.

355. Blitz M, Blitz S, Beasley R, Diner BM, Hughes R, Knopp JA, et al. Inhaled magnesium sulfate in the treatment of acute asthma (Cochrane Review). In: The Cochrane Library, 2005.
356. Bede O, Suranyi A, Pinter K, Szlavik M, Gyurkovits K. Urinary magnesium excretion in asthmatic children receiving magnesium supplementation: a randomized, placebo-controlled, double-blind study. *Magnes Res* 2003;16(4):262-70.
357. Fogarty A, Lewis SA, Scrivener SL, Antoniak M, Pacey S, Pringle M, et al. Oral magnesium and vitamin C supplements in asthma: a parallel group randomized placebo-controlled trial. *Clin Exp Allergy* 2003;33(10):1355-9.
358. Hill J. Magnesium and airway reactivity. *Clin Sci (Lon)* 1998;95(2):111-2.
359. Prescott SL, Calder PC. N-3 polyunsaturated fatty acids and allergic disease. *Curr Opin Clin Nutr Metab Care* 2004;7(2):123-9.
360. Stephensen CB. Fish oil and inflammatory disease: is asthma the next target for n-3 fatty acid supplements? *Nutr Rev* 2004;62(12):486-9.
361. Thien FC, De Luca S, Woods RK, Abramson MJ. Dietary marine fatty acids (fish oil) for asthma (Cochrane Review). In: The Cochrane Library, 2002.
362. Allam MF, Lucane RA. Selenium supplementation for asthma (Cochrane Review). In: The Cochrane Library, 2004.
363. Pearson PJ, Lewis SA, Britton J, Fogarty A. Vitamin E supplements in asthma: a parallel group randomised placebo controlled trial. *Thorax* 2004;59(8):652-6.
364. Ram FS, Rowe BH, Kaur B. Vitamin C supplementation for asthma (Cochrane Review). In: The Cochrane Library, 2004.
365. Butland BK, Strachan DP, Anderson HR. Fresh fruit intake and asthma symptoms in young British adults: confounding or effect modification by smoking? *Eur Respir J* 1999;13(4):744-50.
366. Carey IM, Strachan DP, Cook DG. Effects of changes in fresh fruit consumption on ventilatory function in healthy British adults. *Am J Respir Crit Care Med* 1998;158(3):728-33.
367. Cook DG, Carey IM, Whincup PH, Papacosta O, Chirico S, Bruckdorfer KR, et al. Effect of fresh fruit consumption on lung function and wheeze in children. *Thorax* 1997;52(7):628-33.
368. Ellwood P, Asher MI, Bjorksten B, Burr M, Pearce N, Robertson CF. Diet and asthma, allergic rhinoconjunctivitis and atopic eczema symptom prevalence: an ecological analysis of the International Study of Asthma and Allergies in Childhood (ISAAC) data. ISAAC Phase One Study Group. *Eur Respir J* 2001;17(3):436-43.
369. Gilliland FD, Berhane KT, Li YF, Gauderman WJ, McConnell R, Peters J. Children's lung function and antioxidant vitamin, fruit, juice, and vegetable intake. *Am J Epidemiol* 2003;158(6):576-84.
370. Heinrich J, Holscher B, Bolte G, Winkler G. Allergic sensitization and diet: ecological analysis in selected European cities. *Eur Respir J* 2001;17(3):395-402.
371. Strachan DP, Cox BD, Erzincinoglu SW, Walters DE, Whichelow MJ. Ventilatory function and winter fresh fruit consumption in a random sample of British adults. *Thorax* 1991;46(9):624-9.
372. Martineau AR, Cates CJ, Urashima M, Jensen M, Griffiths AP, Nurmatov U, et al. Vitamin D for the management of asthma. *Cochrane Database of Systematic Reviews* 2016: Issue 9.
373. Jolliffe DA, Greenberg L, Hooper RL, Griffiths CJ, Camargo CA, Jr., Kerley CP, et al. Vitamin D supplementation to prevent asthma exacerbations: a systematic review and meta-analysis of individual participant data. *Lancet Respir Med* 2017;5(11):881-90.
374. Adeniyi FB, Young T. Weight loss interventions for chronic asthma (Cochrane Review). In: The Cochrane Library, 2012.
375. Jensen ME, Gibson PG, Collins CE, Hilton JM, Wood LG. Diet-induced weight loss in obese children with asthma: A randomized controlled trial. *Clinical and Experimental Allergy* 2013;43(7):775-84.
376. Ma J, Strub P, Xiao L, Lavori PW, Camargo CA, Jr., Wilson SR, et al. Behavioral weight loss and physical activity intervention in obese adults with asthma. A randomized trial. *Ann Am Thorac Soc* 2015;12(1):1-11.
377. Scott HA, Gibson PG, Garg ML, Pretto JJ, Morgan PJ, Callister R, et al. Dietary restriction and exercise improve airway inflammation and clinical outcomes in overweight and obese asthma: A randomized trial. *Clinical and Experimental Allergy* 2013;43(1):36-49.
378. Bjorksten B, Sepp E, Julge K, Voor T, Mikelsaar M. Allergy development and the intestinal microflora during the first year of life. *J Allergy Clin Immunol* 2001;108(4):516-20.
379. Helin T, Haahtela S, Haahtela T. No effect of oral treatment with an intestinal bacterial strain, *Lactobacillus rhamnosus* (ATCC 53103), on birch-pollen allergy: a placebo-controlled double-blind study. *Allergy* 2002;57(3):243-6.
380. Isolauri E, Arvola T, Sutas Y, Moilanen E, Salminen S. Probiotics in the management of atopic eczema. *Clin Exp Allergy* 2000;30(11):1604-10.
381. Wheeler JG, Shema SJ, Bogle ML, Shirrell MA, Burks AW, Pittler A, et al. Immune and clinical impact of *Lactobacillus acidophilus* on asthma. *Ann Allergy Asthma Immunol* 1997;79(3):229-33.
382. Gruber C, Illi S, Lau S, Nickel R, Forster J, Kamin W, et al. Transient suppression of atopy in early childhood is associated with high vaccination coverage. *Pediatrics* 2003;111(3):e282-8.
383. Gruber C, Meinlschmidt G, Bergmann R, Wahn U, Stark K. Is early BCG vaccination associated with less atopic disease? An epidemiological study in German preschool children with different ethnic backgrounds. *Pediatr Allergy Immunol* 2002;13(3):177-81.
384. Henderson J, North K, Griffiths M, Harvey I, Golding J. Pertussis vaccination and wheezing illnesses in young children: prospective cohort study. The Longitudinal Study of Pregnancy and Childhood Team. *British Medical Journal* 1999;318(7192):1173-6.

385. Nilsson L, Kjellman NI, Björkstén B. A randomized controlled trial of the effect of pertussis vaccines on atopic disease. *Arch Pediatr Adolesc Med* 1998;152(8):734-8.
386. Choi IS, Koh YI. Therapeutic effects of BCG vaccination in adult asthmatic patients: a randomized, controlled trial. *Ann Allergy Asthma Immunol* 2002;88(6): 584-91.
387. Arikani C, Bahceciler NN, Deniz G, Akdis M, Akkoc T, Akdis CA, et al. Bacillus Calmette-Guérin-induced interleukin-12 did not additionally improve clinical and immunologic parameters in asthmatic children treated with sublingual immunotherapy. *Clinical & Experimental Allergy* 2004;34(3):398-405.
388. Tsai JJ, Peng HJ, Shen HD. Therapeutic effect of Bacillus Calmette-Guérin with allergen on human allergic asthmatic patients. *J Microbiol Immunol Infect* 2002;35(2):99-102.
389. Nicholson KG, Nguyen-Van-Tam JS, Ahmed AH, Wiselka MJ, Leese J, Ayres J, et al. Randomised placebo-controlled crossover trial on effect of inactivated influenza vaccine on pulmonary function in asthma. *Lancet* 1998;351(9099):326-31.
390. Bueving HJ, Bernsen RM, de Jongste JC, van Suijlekom-Smit LW, Rimmelzwaan GF, Osterhaus AD, et al. Influenza vaccination in children with asthma: randomized double-blind placebo-controlled trial. *American Journal of Respiratory & Critical Care Medicine* 2004;169(4):488-93.
391. Bueving HJ, van der Wouden JC, Raat H, Bernsen RM, de Jongste JC, van Suijlekom-Smit LW, et al. Influenza vaccination in asthmatic children: Effects on quality of life and symptoms. *Eur Respir J* 2004;24(6):925-31.
392. Hanania NA, Sockrider M, Castro M, Holbrook JT, Tonascia J, Wise R, et al. Immune response to influenza vaccination in children and adults with asthma: effect of corticosteroid therapy. *Journal of Allergy & Clinical Immunology* 2004;113(4):717-24.
393. Sheikh A, Alves B, Dhami S. Pneumococcal vaccine for asthma (Cochrane Review). In: The Cochrane Library, 2002.
394. Linde K, Jobst K, Panton J. Acupuncture for chronic asthma (Cochrane Review). In: The Cochrane Library, 2000.
395. Martin J, Donaldson AN, Villarroel R, Parmar MK, Ernst E, Higginson IJ. Efficacy of acupuncture in asthma: systematic review and meta-analysis of published data from 11 randomised controlled trials. *Eur Respir J* 2002;20(4):846-52.
396. Gruber W, Eber E, Malle-Scheid D, Pflieger A, Weinhandl E, Dorfer L, et al. Laser acupuncture in children and adolescents with exercise induced asthma. *Thorax* 2002;57(3):222-5.
397. Malmstrom M, Ahlner J, Carlsson C, Schmekel B. No effect of Chinese acupuncture on isocapnic hyperventilation with cold air in asthmatics, measured with impulse oscillometry. *Acupunct Med* 2002;20(2-3):66-73.
398. Blackhall K, Appleton S, Cates Christopher J. Inhalers for chronic asthma. *Cochrane Database of Systematic Reviews* 2012: Issue 9.
399. Bruton A, Lee A, Yardley L, Raftery J, Arden-Close E, Kirby S, et al. Physiotherapy breathing retraining for asthma: a randomised controlled trial. *Lancet Respir Med* 2018;6(1):19-28.
400. O'Connor E, Patnode CD, Burda BU, Buckley DI, Whitlock EP. Breathing Exercises and/or Retraining Techniques in the Treatment of Asthma: Comparative Effectiveness. Rockville (MD): Agency for Healthcare Research and Quality (US); 2012. (Report number 12-EHC092-EF). [cited 10 Jul 2014] Available from <http://effectivehealthcare.ahrq.gov/search-for-guides-reviews-and-reports/?pageaction=n=displayproduct&productID=1251>
401. Freitas Diana A, Holloway Elizabeth A, Bruno Selma S, Chaves Gabriela SS, Fregonezi Guilherme AF, Mendonça Karla MPP. Breathing exercises for adults with asthma. *Cochrane Database of Systematic Reviews* 2013: Issue 10.
402. Prem V, Sahoo RC, Adhikari P. Comparison of the effects of Buteyko and pranayama breathing techniques on quality of life in patients with asthma - a randomized controlled trial. *Clin Rehabil* 2013;27(2):133-41.
403. Silva IS, Fregonezi GA, Dias FA, Ribeiro CT, Guerra RO, Ferreira GM. Inspiratory muscle training for asthma (Cochrane Review). In: The Cochrane Library, (9) 2013.
404. Yang Z-Y, Zhong H-B, Mao C, Yuan J-Q, Huang Y-F, Wu X-Y, et al. Yoga for asthma. *Cochrane Database of Systematic Reviews* 2016: Issue 4.
405. Cramer H, Posadzki P, Dobos G, Langhorst J. Yoga for asthma: A systematic review and meta-analysis. *Ann Allergy Asthma Immunol* 2014;112(6): 503-10.e5.
406. Macêdo Thalita MF, Freitas Diana A, Chaves Gabriela SS, Holloway Elizabeth A, Mendonça Karla MPP. Breathing exercises for children with asthma. *Cochrane Database of Systematic Reviews* 2016: Issue 4.
407. Huntley A, Ernst E. Herbal medicines for asthma: a systematic review. *Thorax* 2000;55(11):925-9.
408. Chan CK, Kuo ML, Shen JJ, See LC, Chang HH, Huang JL. Ding Chuan Tang, a Chinese herb decoction, could improve airway hyper-responsiveness in stabilized asthmatic children: a randomized, double-blind clinical trial. *Pediatr Allergy Immunol* 2006;17(5):316-22.
409. Hsu CH, Lu CM, Chang TT. Efficacy and safety of modified Mai-Men-Dong-Tang for treatment of allergic asthma. *Pediatr Allergy Immunol* 2005;16(1):76-81.
410. Linde K, Jobst KA. Homeopathy for chronic asthma (Cochrane Review). In: The Cochrane Library, 2000.
411. White A, Slade P, Hunt C, Hart A, Ernst E. Individualised homeopathy as an adjunct in the treatment of childhood asthma: a randomised placebo controlled trial. *Thorax* 2003;58(4):317-21.
412. Huntley A, White AR, Ernst E. Relaxation therapies for asthma: a systematic review. *Thorax* 2002;57(2):127-31.

-
413. Hondras MA, Linde K, Jones AP. Manual therapy for asthma (Cochrane Review). In: The Cochrane Library, 2001.
414. Holloway E, Ram FS. Breathing exercises for asthma (Cochrane Review). In: The Cochrane Library, (CD001277) 2004.
415. Panton J, Barley EA. Family therapy for asthma in children (Cochrane Review). In: The Cochrane Library, 2000.
416. Eccles M, Rousseau N, Higgins B, Thomas L. Evidence-based guideline on the primary care management of asthma. *Fam Pract* 2001;18(2):223-9.
417. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.2: ipratropium bromide. Scottish Intercollegiate Guidelines Network; 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.2.html>: [Accessed. 10 Jul 2014].
418. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.4a: inhaled corticosteroid vs theophylline. Scottish Intercollegiate Guidelines Network; 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.4a.html>: [Accessed. 10 July 2014].
419. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.4c: inhaled corticosteroid vs leukotriene receptor antagonists. Scottish Intercollegiate Guidelines Network; 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.4c.html>: [Accessed. 10 Jul 2014].
420. Adams N, Bestall J, Jones P.W. Inhaled fluticasone propionate for chronic asthma (Cochrane Review). In: The Cochrane Library, 2000.
421. Calpin C, Macarthur C, Stephens D, Feldman W, Parkin PC. Effectiveness of prophylactic inhaled steroids in childhood asthma: a systemic review of the literature. *J Allergy Clin Immunol* 1997;100(4):452-7.
422. Carlsen KC, Stick S, Kamin W, Cirule I, Hughes S, Wixon C. The efficacy and safety of fluticasone propionate in very young children with persistent asthma symptoms. *Respir Med* 2005;99(11):1393-402.
423. Teper AM, Colom AJ, Kofman CD, Maffey AF, Vidaurreta SM, Bergada I. Effects of inhaled fluticasone propionate in children less than 2 years old with recurrent wheezing. *Pediatr Pulmonol* 2004;37(2):111-5.
424. Teper AM, Kofman CD, Szulman GA, Vidaurreta SM, Maffey AF. Fluticasone improves pulmonary function in children under 2 years old with risk factors for asthma. *Am J Respir Crit Care Med* 2005;171(6):587-90.
425. Bisgaard H, Allen D, Milanowski J, Kalev I, Willits L, Davies P. Twelve-month safety and efficacy of inhaled fluticasone propionate in children aged 1 to 3 years with recurrent wheezing. *Pediatrics* 2004;113(2):e87-94.
426. Busse WW, Pedersen S, Pauwels RA, Tan WC, Chen YZ, Lamm CJ, et al. The Inhaled Steroid Treatment As Regular Therapy in Early Asthma (START) study 5-year follow-up: effectiveness of early intervention with budesonide in mild persistent asthma. *J Allergy Clin Immunol* 2008;121(5):1167-74.
427. Castro-Rodriguez JA, Rodrigo GJ. Efficacy of inhaled corticosteroids in infants and preschoolers with recurrent wheezing and asthma: a systematic review with meta-analysis. *Pediatrics* 2009;123(3):e519-25.
428. Kerwin EM, Pearlman DS, de Guia T, Carlsson LG, Gillen M, Uryniak T, et al. Evaluation of efficacy and safety of budesonide delivered via two dry powder inhalers. *Curr Med Res Opin* 2008;24(5):1497-510.
429. Knuffman JE, Sorkness CA, Lemanske RF Jr, Mauger DT, Boehmer SJ, Martinez FD, et al. Phenotypic predictors of long-term response to inhaled corticosteroid and leukotriene modifier therapies in pediatric asthma. *Journal of Allergy and Clinical Immunology* 2009;123(2):411-6.
430. Papi A, Nicolini G, Baraldi E, Boner AL, Cutrera R, Rossi GA, et al. Regular vs prn nebulized treatment in wheeze preschool children. *Allergy* 2009;64(10):1463-71.
431. Rachelefsky G. Inhaled corticosteroids and asthma control in children: assessing impairment and risk. *Pediatrics* 2009;123(1):353-66.
432. O'Byrne PM, Barnes PJ, Rodriguez-Roisin R, Runnerstrom E, Sandstrom T, Svensson K, et al. Low dose inhaled budesonide and formoterol in mild persistent asthma: the OPTIMA randomized trial. *Am J Respir Crit Care Med* 2001;164(8 Pt 1):1392-7.
433. Pauwels RA, Pedersen S, Busse WW, Tan WC, Chen YZ, Ohlsson SV, et al. Early intervention with budesonide in mild persistent asthma: a randomised, double-blind trial. *Lancet* 2003;361(9363):1071-6.
434. Berger WE. Budesonide inhalation suspension for the treatment of asthma in infants and children. *Drugs* 2005;65(14):1973-89.
435. Sorkness CA, Lemanske RF Jr, Mauger DT, Boehmer SJ, Chinchilli VM, Martinez FD, et al. Long-term comparison of 3 controller regimens for mild-moderate persistent childhood asthma: the Pediatric Asthma Controller Trial. *J Allergy Clin Immunol* 2007;119(1):64-72.
436. Szefer SJ, Baker JW, Uryniak T, Goldman M, Silkoff PE. Comparative study of budesonide inhalation suspension and montelukast in young children with mild persistent asthma. *J Allergy Clin Immunol* 2007;120(5):1043-50.
437. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.7: high dose step down. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.7.html>: [Accessed. 11 Jul 2014].
438. Hodges IG, Netherway TA. Once-Daily Fluticasone Propionate is as Effective as Twice-Daily Treatment in Stable, Mild-to-Moderate Childhood Asthma. *Clin Drug Investig* 2005;25(1):13-22.
-

439. Chen YZ, Busse WW, Pedersen S, Tan W, Lamm CJ, O'Byrne PM. Early intervention of recent onset mild persistent asthma in children aged under 11 yrs: the Steroid Treatment As Regular Therapy in early asthma (START) trial. *Pediatr Allergy Immunol* 2006;17 Suppl 17:7-13.
440. Martinez FD, Chinchilli VM, Morgan WJ, Boehmer SJ, Lemanske RF, Jr., Mauger DT, et al. Use of beclomethasone dipropionate as rescue treatment for children with mild persistent asthma (TREXA): a randomised, double-blind, placebo-controlled trial. *Lancet* 2011;377(9766):650-7.
441. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.25: budesonide vs beclometasone. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.25.html>: [Accessed. 10 Jul 2014].
442. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.15: mometasone furoate dry powder inhalation. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.15.html>: [Accessed. 10 July 2014].
443. Woodcock A, Bateman ED, Busse WW, Lotvall J, Snowise NG, Forth R, et al. Efficacy in asthma of once-daily treatment with fluticasone furoate: a randomized, placebo-controlled trial. *Respiratory Research* 2011;12:132.
444. Woodcock A, Bleecker ER, Busse WW, Lotvall J, Snowise NG, Frith L, et al. Fluticasone furoate: once-daily evening treatment versus twice-daily treatment in moderate asthma. *Respiratory Research* 2011;12:160.
445. London Respiratory Team. Inhaled corticosteroid safety information for adults. NHS London; c2012. Available from <http://www.networks.nhs.uk/nhs-networks/london-respiratory-network/key-documents/responsible-respiratory-prescribing/LRT%20Inhaled%20steroid%20safety%20card.pdf>: [Accessed. 11 Jul 2014].
446. Fay JK, Jones A, Ram FS. Primary care based clinics for asthma (Cochrane Review). In: *The Cochrane Library*, 2002.
447. Sharek PJ, Bergman DA. The effect of inhaled steroids on the linear growth of children with asthma: a meta-analysis. *Pediatrics* 2000;106(1):E8.
448. Dunlop KA, Carson DJ, Steen HJ, McGovern V, McNaboe J, Shields MD. Monitoring growth in asthmatic children treated with high dose inhaled glucocorticoids does not predict adrenal suppression. *Arch Dis Child* 2004;89(8):713-6.
449. Covar RA, Spahn JD, Martin RJ, Silkoff PE, Sundstrom DA, Murphy J, et al. Safety and application of induced sputum analysis in childhood asthma. *J Allergy Clin Immunol* 2004;114(3):575-82.
450. Bernstein DI, Allen DB. Evaluation of tests of hypothalamic-pituitary-adrenal axis function used to measure effects of inhaled corticosteroids. *Ann Allergy Asthma Immunol* 2007;98(2):118-27.
451. Kelly A, Tang R, Becker S, Stanley CA. Poor specificity of low growth hormone and cortisol levels during fasting hypoglycemia for the diagnoses of growth hormone deficiency and adrenal insufficiency. *Pediatrics* 2008;122(3):e522-8.
452. Tomlinson JE, McMahon AD, Chaudhuri R, Thompson JM, Wood SF, Thomson NC. Efficacy of low and high dose inhaled corticosteroid in smokers versus non-smokers with mild asthma. *Thorax* 2005;60(4):282-7.
453. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.4d: leukotriene receptor antagonists with short-acting beta-agonists. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.4d.html>: [Accessed. 11 Jul 2014].
454. Ducharme FM. Inhaled glucocorticoids versus leukotriene receptor antagonists as single agent asthma treatment: systematic review of current evidence. *BMJ* 2003;326(7390):621.
455. Knorr B, Franchi LM, Bisgaard H, Vermeulen JH, LeSouef P, Santanello N, et al. Montelukast, a leukotriene receptor antagonist, for the treatment of persistent asthma in children aged 2 to 5 years. *Pediatrics* 2001;108(3):E48.
456. Valovirta E, Boza ML, Robertson CF, Verbruggen N, Smugar SS, Nelsen LM, et al. Intermittent or daily montelukast versus placebo for episodic asthma in children. *Annals of Allergy, Asthma, & Immunology* 2011;106(6):518-26.
457. Edwards AM, Stevens MT. The clinical efficacy of inhaled nedocromil sodium (Tilade) in the treatment of asthma. *Eur Respir J* 1993;6(1):35-41.
458. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.24a: Other preventor therapies - Chromones in children aged 5-12. Edinburgh: SIGN; 2005. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/index.html>: [Accessed. 10 Jul 2014].
459. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.4j: do chromones work as first line preventor in children >5 years? 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.4j.html>: [Accessed. 11 Jul 2014].
460. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.24b: other preventor therapies - chromones in children aged <5. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.24b.html>: [Accessed. 10 Jul 2014].
461. Van Ganse E, Kaufman L, Derde MP, Yernault JC, Delaunois L, Vincken W. Effects of antihistamines in adult asthma: a meta-analysis of clinical trials. *Eur Respir J* 1997;10(10):2216-24.
462. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.11b: add-on drugs for inhaled steroids: long acting or oral B2 agonists. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.11b.html>: [Accessed. 11 Jul 2014].

463. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.11d: add-on drugs for inhaled steroids: theophylline, beclometasone dipropionate, budesonide. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.11d.html>: [Accessed. 11 Jul 2014].
464. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.11c: add-on drugs for inhaled steroids - anticholinergics. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.11c.html>: [Accessed. 11 Jul 2014].
465. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.11a: add on drugs for inhaled steroids - chromones. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.11a.html>: [Accessed. 11 Jul 2014].
466. Becker AB, Simons FE. Formoterol, a new long-acting selective beta 2-adrenergic receptor agonist: double-blind comparison with salbutamol and placebo in children with asthma. *J Allergy Clin Immunol* 1989;84(6 Pt 1):891-5.
467. Kips JC, Pauwels RA. Long-acting inhaled beta(2)-agonist therapy in asthma. *Am J Respir Crit Care Med* 2001;164(6):923-32.
468. de Blic J, Ogorodova L, Klink R, Sidorenko I, Valiulis A, Hofman J, et al. Salmeterol/fluticasone propionate vs. double dose fluticasone propionate on lung function and asthma control in children. *Pediatr Allergy Immunol* 2009;20(8):763-71.
469. Gappa M, Zachgo W, von Berg A, Kamin W, Stern-Strater C, Steinkamp G, et al. Add-on salmeterol compared to double dose fluticasone in pediatric asthma: A double-blind, randomized trial (VIAPAE). *Pediatr Pulmonol* 2009;44(11):1132-42.
470. Morice AH, Peterson S, Beckman O, Kukova Z. Efficacy and safety of a new pressurised metered-dose inhaler formulation of budesonide/formoterol in children with asthma: a superiority and therapeutic equivalence study. *Pulm Pharmacol Ther* 2008;21(1):152-9.
471. Pearlman D, Qaqundah P, Matz J, Yancey SW, Stempel DA, Ortega HG. Fluticasone propionate/salmeterol and exercise-induced asthma in children with persistent asthma. *Pediatr Pulmonol* 2009;44(5):429-35.
472. Chauhan Bhupendrasinh F, Ducharme Francine M. Addition to inhaled corticosteroids of long-acting beta2-agonists versus anti-leukotrienes for chronic asthma. *Cochrane Database of Systematic Reviews* 2014(1).
473. Medicines and Healthcare products Regulatory Agency. Salmeterol (Severant) and formoterol (Oxis) in asthma management. *Current Problems in Pharmacovigilance* 2003; 29. [cited 11 Jul 2014] Available from <http://www.mhra.gov.uk/Publications/Safetyguidance/CurrentProblemsinPharmacovigilance/CON007449>
474. Medicines and Healthcare products Regulatory Agency. Long-acting β_2 -agonists: reminder for use in children and adults. *Drug Safety Update* 2010;4(2):H2.
475. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.22: combined therapy of inhaled steroids and long acting B2 agonists. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.22.html>: [Accessed. 10 Jul 2014].
476. Cates Christopher J, Karner C. Combination formoterol and budesonide as maintenance and reliever therapy versus current best practice (including inhaled steroid maintenance), for chronic asthma in adults and children. *Cochrane Database of Systematic Reviews* 2013(4).
477. Sobieraj DM. Association of Inhaled Corticosteroids and Long-Acting [beta]-Agonists as Controller and Quick Relief Therapy With Exacerbations and Symptom Control in Persistent Asthma: A Systematic Review and Meta-analysis. *JAMA* 2018.
478. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.8c: children with poor asthma control on ICS - is addition of leukotriene receptor antagonists helpful? 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.8c.html>: [Accessed. 11 Jul 2014].
479. Joos S, Miksch A, Szecsenyi J, Wieseler B, Grouven U, Kaiser T, et al. Montelukast as add-on therapy to inhaled corticosteroids in the treatment of mild to moderate asthma: a systematic review. *Thorax* 2008;63(5):453-62.
480. Cao Y, Wang J, Bunjhoo H, Xie M, Xu Y, Fang H. Comparison of leukotriene receptor antagonists in addition to inhaled corticosteroid and inhaled corticosteroid alone in the treatment of adolescents and adults with bronchial asthma: a meta-analysis (Provisional abstract). *Asian Pacific Journal of Allergy and Immunology* 2012(2):130-8.
481. Kew KM, Dahri K. Long-acting muscarinic antagonists (LAMA) added to combination long-acting beta2-agonists and inhaled corticosteroids (LABA/ICS) versus LABA/ICS for adults with asthma. *Cochrane Database Syst Rev* 2016(1):CD011721.
482. Kew KM, Evans DJ, Allison DE, Boyter AC. Long-acting muscarinic antagonists (LAMA) added to inhaled corticosteroids (ICS) versus addition of long-acting beta2-agonists (LABA) for adults with asthma. *Cochrane Database Syst Rev* 2015(6):CD011438.
483. Anderson DE, Kew KM, Boyter AC. Long-acting muscarinic antagonists (LAMA) added to inhaled corticosteroids (ICS) versus the same dose of ICS alone for adults with asthma. *Cochrane Database Syst Rev* 2015(8):CD011397.
484. Bateman ED, Kornmann O, Schmidt P, Pivovarova A, Engel M, Fabbri LM. Tiotropium is noninferior to salmeterol in maintaining improved lung function in B16-Arg/Arg patients with asthma. *J Allergy Clin Immunol* 2011;128(2):315-22.

485. Evans DJ, Kew KM, Anderson DE, Boyter AC. Long-acting muscarinic antagonists (LAMA) added to inhaled corticosteroids (ICS) versus higher dose ICS for adults with asthma. *Cochrane Database Syst Rev* 2015(7):CD011437.
486. Westby M, Benson M, Gibson P. Anticholinergic agents for chronic asthma in adults (Cochrane Review). In: *The Cochrane Library*, 2004.
487. Scottish Intercollegiate Guidelines Network. Management of osteoporosis and the prevention of fragility fractures. 2015. Available from www.sign.ac.uk: [Accessed 17 July 2019].
488. Bachrach LK, Sills IN. Clinical report-bone densitometry in children and adolescents. *Pediatrics* 2011;127(1):189-94.
489. Norman G, Faria R, Paton F, Llewellyn A, Fox D, Palmer S, et al. Omalizumab for the treatment of severe persistent allergic asthma: a systematic review and economic evaluation. *Health Technol Assess* 2013;17(52):1-342.
490. Normansell R, Walker S, Milan Stephen J, Walters EH, Nair P. Omalizumab for asthma in adults and children. *Cochrane Database of Systematic Reviews* 2014: Issue 1.
491. Rodrigo GJ, Neffen H, Castro-Rodriguez JA, Neffen H, Castro-Rodriguez JA. Efficacy and safety of subcutaneous omalizumab vs placebo as add-on therapy to corticosteroids for children and adults with asthma: a systematic review. *Chest* 2011;139(1):28-35.
492. Farne Hugo A, Wilson A, Powell C, Bax L, Milan Stephen J. Anti-IL5 therapies for asthma. *Cochrane Database of Systematic Reviews* 2017: Issue 9.
493. Bel EH, Wenzel SE, Thompson PJ, Prazma CM, Keene ON, Yancey SW, et al. Oral glucocorticoid-sparing effect of mepolizumab in eosinophilic asthma. *New England Journal of Medicine* 2014;371(13):1189-97.
494. Bermejo I, Stevenson M, Cooper K, Harnan S, Hamilton J, Clowes M, et al. Mepolizumab for Treating Severe Eosinophilic Asthma: An Evidence Review Group Perspective of a NICE Single Technology Appraisal. *Pharmacoeconomics* 2018;36(2):131-44.
495. Nachev Z, Krishnan A, Mashtare T, Zhuang T, Mador MJ. Omalizumab versus Mepolizumab as add-on therapy in asthma patients not well controlled on at least an inhaled corticosteroid: A network meta-analysis. *J Asthma* 2018;55(1):89-100.
496. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.13a: immunosuppressive agents. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.13a.html>: [Accessed. 14 Jul 2014].
497. O'Driscoll BR, Ruffles SP, Ayres JG, Cochrane GM. Long term treatment of severe asthma with subcutaneous terbutaline. *Br J Dis Chest* 1988;82(4):360-7.
498. Payne DN, Balfour-Lynn IM, Biggart EA, Bush A, Rosenthal M. Subcutaneous terbutaline in children with chronic severe asthma. *Pediatr Pulmonol* 2002;33(5):356-61.
499. Bremont F, Moisan V, Dutau G. Continuous subcutaneous infusion of beta 2-agonists in infantile asthma. *Pediatr Pulmonol* 1992;12(2):81-3.
500. Berry MA, Hargadon B, Shelley M, Parker D, Shaw DE, Green RH, et al. Evidence of a role of tumor necrosis factor alpha in refractory asthma. *N Engl J Med* 2006;354(7):697-708.
501. Hawkins G, McMahon AD, Twaddle S, Wood SF, Ford I, Thomson NC. Stepping down inhaled corticosteroids in asthma: randomised controlled trial. *BMJ* 2003;326(7399):1115.
502. Abramson MJ, Puy RM, Weiner JM. Injection allergen immunotherapy for asthma (Cochrane Review). In: *The Cochrane Library*, 2010.
503. Shaikh WA. Immunotherapy vs inhaled budesonide in bronchial asthma: an open, parallel, comparative trial. *Clin Exp Allergy* 1997;27(11):1279-84.
504. Durham SR, Walker SM, Varga EM, Jacobson MR, O'Brien F, Noble W, et al. Long-term clinical efficacy of grass-pollen immunotherapy. *N Engl J Med* 1999;341(7):468-75.
505. Normansell R, Kew Kayleigh M, Bridgman A-L. Sublingual immunotherapy for asthma. *Cochrane Database of Systematic Reviews* 2015: Issue 8.
506. Torrego A, Solà I, Munoz Ana M, Roqué i Figuls M, Yepes-Nuñez Juan J, Alonso-Coello P, et al. Bronchial thermoplasty for moderate or severe persistent asthma in adults. *Cochrane Database of Systematic Reviews* 2014: Issue 3.
507. Zhou JP, Feng Y, Wang Q, Zhou LN, Wan HY, Li QY. Long-term efficacy and safety of bronchial thermoplasty in patients with moderate-to-severe persistent asthma: a systemic review and meta-analysis. *J Asthma* 2016;53(1):94-100.
508. Robertson CF, Price D, Henry R, Mellis C, Glasgow N, Fitzgerald D, et al. Short-course montelukast for intermittent asthma in children: a randomized controlled trial. *Am J Manag Care* 2007;175(4):323-9.
509. Adams NP, Bestall JB, Malouf R, Lasserson TJ, Jones PW. Beclomethasone versus placebo for chronic asthma (Cochrane Review). In: *The Cochrane Library*, 2005.
510. Henriksen JM, Agertoft L, Pedersen S. Protective effect and duration of action of inhaled formoterol and salbutamol on exercise-induced asthma in children. *J Allergy Clin Immunol* 1992;89(6):1176-82.
511. Raissy HH, Harkins M, Kelly F, Kelly HW. Pretreatment with albuterol versus montelukast for exercise-induced bronchospasm in children. *Pharmacotherapy* 2008;28(3):287-94.
512. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.3a: long acting B2 agonists in exercise induced asthma. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.3a.html>: [Accessed. 14 Jul 2014].
513. Nassif EG, Weinberger M, Thompson R, Huntley W. The value of maintenance theophylline in steroid-dependent asthma. *New England Journal of Medicine* 1981;304(2):71-5.

514. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.3c: theophyllines in exercise-induced asthma. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.3c.html>: [Accessed. 14 Jul 2014].
515. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.3d: leukotriene receptor antagonists in exercise induced asthma. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.3d.html>: [Accessed. 14 Jul 2014].
516. Kelly K, Spooner CH, Rowe BH. Nedocromil sodium vs. sodium cromoglycate for preventing exercise-induced bronchoconstriction in asthmatics (Cochrane Review). In: The Cochrane Library, 2000.
517. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.3f: anticholinergic therapy for exercise-induced asthma. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.3f.html>: [Accessed. 14 Jul 2014].
518. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.3b: ketotifen for exercise-induced asthma. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.3b.html>: [Accessed. 14 Jul 2014].
519. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.3e: antihistamines for exercise-induced asthma. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.3e.html>: [Accessed. 14 Jul 2014].
520. Stelmach I, Grzelewski T, Majak P, Jerzynska J, Stelmach W, Kuna P. Effect of different antiasthmatic treatments on exercise-induced bronchoconstriction in children with asthma. *J Allergy Clin Immunol* 2008;121(2):383-9.
521. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.10: rhinitis. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.10.html>: [Accessed. 14 Jul 2014].
522. Pedroletti C, Lundahl J, Alving K, Hedlin G. Effect of nasal steroid treatment on airway inflammation determined by exhaled nitric oxide in allergic schoolchildren with perennial rhinitis and asthma. *Pediatr Allergy Immunol* 2008;19(3):219-26.
523. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.19: allergic bronchopulmonary aspergillosis. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.19.html>: [Accessed. 14 Jul 2014].
524. Scottish Intercollegiate Guidelines Network (SIGN). Pharmacological management of asthma. Evidence table 4.21: aspirin intolerant asthma. 2002. Available from <http://www.sign.ac.uk/guidelines/published/support/guideline63/table4.21.html>: [Accessed. 10 Jul 2014].
525. Wark PA, Gibson PG, Wilson AJ. Azoles for allergic bronchopulmonary aspergillosis associated with asthma (Cochrane Review). In: The Cochrane Library, 2004.
526. Coughlan JL, Gibson PG, Henry RL. Medical treatment for reflux oesophagitis does not consistently improve asthma control: a systematic review. *Thorax* 2001;56(3):198-204.
527. Gibson PG, Henry RL, Coughlan JL. Gastro-oesophageal reflux treatment for asthma in adults and children (Cochrane Review). In: The Cochrane Library, 2003.
528. Sopo SM, Radzik D, Calvani M. Does treatment with proton pump inhibitors for gastroesophageal reflux disease (GERD) improve asthma symptoms in children with asthma and GERD? A systematic review. *J Investig Allergol Clin Immunol* 2009;19(1):1-5.
529. Chan WW, Chiou E, Obstein KL, Tignor AS, Whitlock TL. The efficacy of proton pump inhibitors for the treatment of asthma in adults: a meta-analysis. *Arch Intern Med* 2011;171(7):620-9.
530. Brocklebank D, Ram F, Wright J, Barry P, Cates C, Davies L, et al. Comparison of the effectiveness of inhaler devices in asthma and chronic obstructive airways disease: a systematic review of the literature. *Health Technol Assess* 2001;5(26):1-149.
531. Cates Christopher J, Welsh Emma J, Rowe Brian H. Holding chambers (spacers) versus nebulisers for beta-agonist treatment of acute asthma. *Cochrane Database of Systematic Reviews* 2013(9).
532. Leversha AM, Campanella SG, Aickin RP, Asher MI. Costs and effectiveness of spacer versus nebuliser in young children with moderate and severe acute asthma. *J Pediatr* 2000;136(4):497-502.
533. Closa RM, Ceballos JM, Gomez-Papi A, Galiana AS, Gutierrez C, Marti-Henneber C. Efficacy of bronchodilators administered by nebulizers versus spacer devices in infants with acute wheezing. *Pediatr Pulmonol* 1998;26(5):344-8.
534. Delgado A, Chou KJ, Silver EJ, Crain EF. Nebulizers vs metered-dose inhalers with spacers for bronchodilator therapy to treat wheezing in children aged 2 to 24 months in a pediatric emergency department. *Archives of Pediatrics & Adolescent Medicine* 2003;157(1):76-80.
535. Ram FS, Wright J, Brocklebank D, White JE. Systematic review of clinical effectiveness of pressurised metered dose inhalers versus other hand held inhaler devices for delivering beta (2) agonists bronchodilators in asthma. *BMJ* 2001;323(7318):901-5.
536. Broeders ME, Molema J, Hop WC, Vermue NA, Folgering HT. Does the inhalation device affect the bronchodilatory dose response curve of salbutamol in asthma and chronic obstructive pulmonary disease patients? *Eur J Clin Pharmacol* 2003;59(5-6):449-55.
537. Hughes DA, Woodcock A, Walley T. Review of therapeutically equivalent alternatives to short acting beta(2) adrenoceptor agonists delivered via chlorofluorocarbon-containing inhalers. *Thorax* 1999;54(12):1087-92.

538. Farmer IS, Middle M, Savic J, Perri VL, Herdman MJ. Therapeutic equivalence of inhaled beclomethasone dipropionate with CFC and non-CFC (HFA 134a) propellants both delivered via the Easibreathe inhaler for the treatment of paediatric asthma. *Respir Med* 2000;94(1):57-63.
539. Cates CJ, Adams N, Bestall J. Holding chambers versus nebulisers for inhaled steroids in chronic asthma (Cochrane Review). In: *The Cochrane Library*, 2001.
540. Alotaibi S, Hassan WM, Alhashimi H. Concurrent use of metered dose inhalers without spacer and dry powder inhalers by asthmatic children adversely affect proper inhalation technique. *Internet J Pediatr Neonatol* 2011;13(1).
541. van der Palen J, Klein JJ, van Herwaarden CL, Zielhuis GA, Seydel ER. Multiple inhalers confuse asthma patients. *Eur Respir J* 1999;14(5):1034-7.
542. Myrdal PB, Sheth P, Stein SW. Advances in metered dose inhaler technology: formulation development. *AAPS PharmSciTech* 2014;15(2):434-55.
543. Environmental Audit Committee. UK progress on reducing F-Gas emissions inquiry. House of Commons Environmental Audit Committee; 2018. Available from <https://www.parliament.uk/business/committees/committees-a-z/commons-select/environmental-audit-committee/inquiries/parliament-2017/uk-progress-on-reducing-f-gas-emissions-17-19/publications/>. [Accessed. 03 May 2019].
544. Lavorini F, Corrigan CJ, Barnes PJ, Dekhuijzen PR, Levy ML, Pedersen S, et al. Retail sales of inhalation devices in European countries: So much for a global policy. *Respir Med* 2011;105(7):1099-103.
545. Accuracy of death certificates in bronchial asthma. Accuracy of certification procedures during the confidential inquiry by the British Thoracic Association. A subcommittee of the BTA Research Committee. *Thorax* 1984;39(7):505-9.
546. Bucknall CE, Slack R, Godley CC, Mackay TW, Wright SC. Scottish Confidential Inquiry into Asthma Deaths (SCIAD), 1994-6. *Thorax* 1999;54(11):978-84.
547. Burr ML, Davies BH, Hoare A, Jones A, Williamson IJ, Holgate SK, et al. A confidential inquiry into asthma deaths in Wales. *Thorax* 1999;54(11):985-9.
548. Mohan G, Harrison BD, Badminton RM, Mildenhall S, Wareham NJ. A confidential enquiry into deaths caused by asthma in an English health region: implications for general practice. *Br J Gen Pract* 1996;46(410):529-32.
549. Wareham NJ, Harrison BD, Jenkins PF, Nicholls J, Stableforth DE. A district confidential enquiry into deaths due to asthma. *Thorax* 1993;48(11):1117-20.
550. Royal College of Physicians. Why asthma still kills: the national review of asthma deaths (NRAD); confidential enquiry report 2014. London: Royal College of Physicians; 2014. [cited 15 Jul 2014] Available from <http://www.rcplondon.ac.uk/sites/default/files/why-asthma-still-kills-full-report.pdf>
551. Harrison B, Slack R, Berrill WT, Burr ML, Stableforth DE, Wright SC. Results of a national confidential enquiry into asthma deaths. *Asthma J* 2000;5(4):180-6.
552. Spitzer WO, Suissa S, Ernst P, Horwitz RI, Habbick B, Cockcroft D, et al. The use of beta-agonists and the risk of death and near death from asthma. *N Engl J Med* 1992;326(8):501-6.
553. Suissa S, Blais L, Ernst P. Patterns of increasing beta-agonist use and the risk of fatal or near-fatal asthma. *Eur Respir J* 1994;7(9):1602-9.
554. Jalaludin BB, Smith MA, Chey T, Orr NJ, Smith WT, Leeder SR. Risk factors for asthma deaths: a population-based, case-control study. *Aust NZ J Public Health* 1999;23(6):595-600.
555. Rea HH, Scragg R, Jackson R, Beaglehole R, Fenwick J, Sutherland DC. A case-control study of deaths from asthma. *Thorax* 1986;41(11):833-9.
556. Campbell MJ, Cogman GR, Holgate ST, Johnston SL. Age specific trends in asthma mortality in England and Wales, 1983-95: results of an observational study. *BMJ* 1997;314(7092):1439-41.
557. Richards GN, Kolbe J, Fenwick J, Rea HH. Demographic characteristics of patients with severe life threatening asthma: comparison with asthma deaths. *Thorax* 1993;48(11):1105-9.
558. Innes NJ, Reid A, Halstead J, Watkin SW, Harrison BD. Psychosocial risk factors in near-fatal asthma and in asthma deaths. *J R Coll Physicians Lond* 1998;32(5):430-4.
559. Khot A, Evans N, Lenney W. Seasonal trends in childhood asthma in south east England. *Br Med J (Clin Res Ed)* 1983;287(6401):1257-8.
560. Barr RG, Woodruff PG, Clark S, Camargo CA Jr. Sudden-onset asthma exacerbations: clinical features, response to therapy, and 2-week follow-up. Multicenter Airway Research Collaboration (MARC) investigators. *Eur Respir J* 2000;15(2):266-73.
561. Kolbe J, Fergusson W, Garrett J. Rapid onset asthma: a severe but uncommon manifestation. *Thorax* 1998;53(4):241-7.
562. Kolbe J, Fergusson W, Vámos M, Garrett J. Case-control study of severe life threatening asthma (SLTA) in adults: demographics, health care, and management of the acute attack. *Thorax* 2000;55(12):1007-15.
563. Rodrigo GJ, Rodrigo C. Rapid-onset asthma attack: a prospective cohort study about characteristics and response to emergency department treatment. *Chest* 2000;118(6):1547-52.
564. Turner MO, Noertjojo K, Vedral S, Bai T, Crump S, Fitzgerald JM. Risk factors for near-fatal asthma. A case-control study in hospitalized patients with asthma. *Am J Respir Crit Care Med* 1998;157(6 Pt 1):1804-9.
565. Woodruff PG, Emond SD, Singh AK, Camargo CA Jr. Sudden-onset severe acute asthma: clinical features and response to therapy. *Acad Emerg Med* 1998;5(7):695-701.

566. British Thoracic Society, National Asthma Campaign, Royal College of Physicians of London in association with the General Practitioner in Asthma Group, The British Association of Accident and Emergency Medicine, The British Paediatric Respiratory Society, Royal College of Paediatrics and Child Health. The British guidelines on asthma management 1995 review and position statement. *Thorax* 1997;52(Suppl 1):S1-S21.
567. Scottish Intercollegiate Guidelines Network (SIGN). Emergency management of acute asthma. Edinburgh: SIGN; 1999. (SIGN publication no. 38). [cited
568. International consensus report on the diagnosis and treatment of asthma. National Heart, Lung, and Blood Institute, National Institutes of Health. Bethesda, Maryland 20892. Publication no. 92-3091, March 1992. *Eur Respir J* 1992;5(5):601-41.
569. Neville E, Gribbin H, Harrison BD. Acute severe asthma. *Respir Med* 1991;85(6):463-74.
570. Brenner B, Kohn MS. The acute asthmatic patient in the ED: to admit or discharge. *Am J Emerg Med* 1998;16(1):69-75.
571. Boulet LP, Becker A, Berube D, Beveridge R, Ernst P. Canadian asthma consensus report, 1999. Canadian asthma consensus group. *CMAJ* 1999;161(11 Suppl):S1-61.
572. Nunn AJ, Gregg I. New regression equations for predicting peak expiratory flow in adults. *BMJ* 1989;298(6680):1068-70.
573. Abramson MJ, Bailey MJ, Couper FJ, Driver JS, Drummer OH, Forbes AB, et al. Are asthma medications and management related to deaths from asthma? *Am J Respir Crit Care Med* 2001;163(1):12-8.
574. Robinson SM, Harrison BD, Lambert MA. Effect of a preprinted form on the management of acute asthma in an accident and emergency department. *J Accid Emerg Med* 1996;13(2):93-7.
575. Arnold DH, Gebretsadik T, Minton PA, Higgins S, Hartert TV. Clinical measures associated with FEV₁ in persons with asthma requiring hospital admission. *Am J Emerg Med* 2007;25(4):425-9.
576. Shim CS, Williams MH Jr. Evaluation of the severity of asthma: patients versus physicians. *Am J Med* 1980;68(1):11-3.
577. Emerman CL, Cydulka RK. Effect of pulmonary function testing on the management of acute asthma. *Arch Intern Med* 1995;155(20):2225-8.
578. O'Driscoll BR, Howard LS, Davison AG, British Thoracic Society. BTS guideline for emergency oxygen use in adult patients. *Thorax* 2008;63(suppl. 6):vi1-68.
579. Carruthers D, Harrison BD. Arterial blood gas analysis or oxygen saturation in the assessment of acute asthma? *Thorax* 1995;50(2):186-8.
580. Pearson MG, Spence DP, Ryland I, Harrison BD. Value of pulsus paradoxus in assessing acute severe asthma. British Thoracic Society Standards of Care Committee. *BMJ* 1993;307(6905):659.
581. McFadden ER Jr, Lyons HA. Arterial-blood gas tension in asthma. *N Engl J Med* 1968;278(19):1027-32.
582. Rebuck AS, Read J. Assessment and management of severe asthma. *Am J Med* 1971;51(6):788-98.
583. Jenkins PF, Benfield GF, Smith AP. Predicting recovery from acute severe asthma. *Thorax* 1981;36(11):835-41.
584. Molfino NA, Nannini LJ, Martelli AN, Slutsky AS. Respiratory arrest in near-fatal asthma. *N Engl J Med* 1991;324(5):285-8.
585. Perrin K, Wijesinghe M, Healy B, Wadsworth K, Bowditch R, Bibby S, et al. Randomised controlled trial of high concentration versus titrated oxygen therapy in severe exacerbations of asthma. *Thorax* 2011;66(11):937-41.
586. McFadden ER Jr. Critical appraisal of the therapy of asthma--an idea whose time has come. *Am Rev Respir Dis* 1986;133(5):723-4.
587. Rossing TH, Fanta CH, Goldstein DH, Snapper JR, McFadden ER Jr. Emergency therapy of asthma: comparison of the acute effects of parenteral and inhaled sympathomimetics and infused aminophylline. *Am Rev Respir Dis* 1980;122(3):365-71.
588. Siegel D, Sheppard D, Gelb A, Weinberg PF. Aminophylline increases the toxicity but not the efficacy of an inhaled beta-adrenergic agonist in the treatment of acute exacerbations of asthma. *Am Rev Respir Dis* 1985;132(2):283-6.
589. Rodrigo G, Nannini L. Comparison between nebulized adrenaline and beta2 agonists for the treatment of acute asthma. A meta-analysis of randomized trials. *Am J Emerg Med* 2006;24(2):217-22.
590. Travers A, Jones AP, Kelly K, Barker SJ, Camargo CA, Rowe BH. Intravenous beta2-agonists for acute asthma in the emergency department (Cochrane Review). In: The Cochrane Library, 2001.
591. Lewis L, Ferguson I, House SL, Aubuchon K, Schneider J, Johnson K, et al. Albuterol administration is commonly associated with increases in serum lactate in patients with asthma treated for acute exacerbation of asthma. *Chest* 2014(1):53-9.
592. Gleeson JG, Green S, Price JF. Air or oxygen as driving gas for nebulised salbutamol. *Arch Dis Child* 1988;63(8):900-4.
593. Douglas JG, Rafferty P, Fergusson RJ, Prescott RJ, Crompton GK, Grant IW. Nebulised salbutamol without oxygen in severe acute asthma: how effective and how safe? *Thorax* 1985;40(3):180-3.
594. Lin RY, Sauter D, Newman T, Sirleaf J, Walters J, Tavakol M. Continuous versus intermittent albuterol nebulization in the treatment of acute asthma. *Ann Emerg Med* 1993;22(12):1847-53.
595. Rudnitsky GS, Eberlein RS, Schoffstall JM, Mazur JE, Spivey WH. Comparison of intermittent and continuously nebulized albuterol for treatment of asthma in an urban emergency department. *Ann Emerg Med* 1993;22(12):1842-6.
596. Shrestha M, Bidadi K, Gourlay S, Hayes J. Continuous vs intermittent albuterol, at high and low doses, in the treatment of severe acute asthma in adults. *Chest* 1996;110(1):42-7.

597. Camargo CA Jr, Spooner C, Rowe BH. Continuous versus intermittent beta-agonists for acute asthma (Cochrane Review). In: The Cochrane Library, (CD001115) 2009.
598. Rowe BH, Spooner C, Ducharme FM, Bretzlaff JA, Bota GW. Early emergency department treatment of acute asthma with systemic corticosteroids (Cochrane Review). In: The Cochrane Library, 2001.
599. Rowe BH, Spooner CH, Ducharme FM, Bretzlaff JA, Bota GW. Corticosteroids for preventing relapse following acute exacerbations of asthma (Cochrane Review). In: The Cochrane Library, 2001.
600. Manser R, Reid D, Abramson M. Corticosteroids for acute severe asthma in hospitalised patients (Cochrane Review). In: The Cochrane Library, 2001.
601. Lahn M, Bijur P, Gallagher EJ. Randomized clinical trial of intramuscular vs oral methylprednisolone in the treatment of asthma exacerbations following discharge from an emergency department. *Chest* 2004;126(2):362-8.
602. Hatton MQ, Vathenen AS, Allen MJ, Davies S, Cooke NJ. A comparison of 'abruptly stopping' with 'tailing off' oral corticosteroids in acute asthma. *Respir Med* 1995;89(2):101-4.
603. O'Driscoll BR, Kalra S, Wilson M, Pickering CA, Carroll KB, Woodcock AA. Double-blind trial of steroid tapering in acute asthma. *Lancet* 1993;341(8841):324-7.
604. Edmonds ML, Camargo CA, Pollack CV Jr, Rowe BH. Early use of inhaled corticosteroids in the emergency department treatment of acute asthma (Cochrane Review). In: The Cochrane Library, (3) 2003.
605. Rodrigo GJ. Rapid effects of inhaled corticosteroids in acute asthma: an evidence-based evaluation. *Chest* 2006;130(5):1301-11.
606. Lanes SF, Garrett JE, Wentworth CE 3rd, Fitzgerald JM, Karpel JP. The effect of adding ipratropium bromide to salbutamol in the treatment of acute asthma: a pooled analysis of three trials. *Chest* 1998;114(2):365-72.
607. Rodrigo G, Rodrigo C, Burschtin O. A meta-analysis of the effects of ipratropium bromide in adults with acute asthma. *Am J Med* 1999;107(4):363-70.
608. Stoodley RG, Aaron SD, Dales RE. The role of ipratropium bromide in the emergency management of acute asthma exacerbation: a metaanalysis of randomized clinical trials. *Ann Emerg Med* 1999;34(1):8-18.
609. Knightly R, Milan Stephen J, Hughes R, Knopp-Sihota Jennifer A, Rowe Brian H, Normansell R, et al. Inhaled magnesium sulfate in the treatment of acute asthma. *Cochrane Database of Systematic Reviews* 2017: Issue 11.
610. Goodacre S, Cohen J, Bradburn M, Gray A, Bengner J, Coats T, et al. Intravenous or nebulised magnesium sulphate versus standard therapy for severe acute asthma (3Mg trial): a double-blind, randomised controlled trial. *Lancet* 2013;1(4):293-300.
611. Blitz M, Blitz S, Hughes R, Diner B, Beasley R, Knopp J, et al. Aerosolized magnesium sulfate for acute asthma: a systematic review. *Chest* 2005;128(1):337-44.
612. Rowe BH, Bretzlaff JA, Bourdon C, Bota GW, Camargo CA, Jr. Magnesium sulfate for treating exacerbations of acute asthma in the emergency department. *Cochrane Database Syst Rev* 2000(2):CD001490.
613. Kew Kayleigh M, Kirtchuk L, Michell Clare I. Intravenous magnesium sulfate for treating adults with acute asthma in the emergency department. *Cochrane Database of Systematic Reviews* 2014(5).
614. Parameswaran K, Belda J, Rowe BH. Addition of intravenous aminophylline to beta2-agonists in adults with acute asthma (Cochrane Review). In: The Cochrane Library, 2000.
615. Watts K, Chavasse Richard JPG. Leukotriene receptor antagonists in addition to usual care for acute asthma in adults and children. *Cochrane Database of Systematic Reviews* 2012(5).
616. Graham VA, Milton AF, Knowles GK, Davies RJ. Routine antibiotics in hospital management of acute asthma. *Lancet* 1982;1(8269):418-20.
617. Long W, Li LJ, Huang GZ, Zhang XM, Zhang YC, Tang JG, et al. Procalcitonin guidance for reduction of antibiotic use in patients hospitalized with severe acute exacerbations of asthma: A randomized controlled study with 12-month follow-up. *Critical Care* 2014;18(5).
618. Tang J, Long W, Yan L, Zhang Y, Xie J, Lu G, et al. Procalcitonin guided antibiotic therapy of acute exacerbations of asthma: a randomized controlled trial. *BMC Infect Dis* 2013;13:596.
619. Kass JE, Terregino CA. The effect of heliox in acute severe asthma: a randomized controlled trial. *Chest* 1999;116(2):296-300.
620. Henderson SO, Acharya P, Kilagbhan T, Perez J, Korn CS, Chan LS. Use of heliox-driven nebulizer therapy in the treatment of acute asthma. *Ann Emerg Med* 1999;33(2):141-6.
621. Rodrigo GJ, Pollack CV, Rodrigo C, Rowe BH. Heliox for nonintubated acute asthma patients (Cochrane Review). In: The Cochrane Library, (4) 2006.
622. Rodrigo GJ, Rodrigo C, Pollack CV, Rowe B. Use of helium-oxygen mixtures in the treatment of acute asthma: a systematic review. *Chest* 2003;123(3):891-6.
623. Yen ZS, Chen SC. Best evidence topic report. Nebulised furosemide in acute adult asthma. *Emergency Medicine Journal* 2005;22(9):654-5.
624. Goyal S, Agrawal A. Ketamine in status asthmaticus: A review. *Indian J* 2013;17(3):154-61.
625. Yeo HJ, Kim D, Jeon D, Kim YS, Rycus P, Cho WH. Extracorporeal membrane oxygenation for life-threatening asthma refractory to mechanical ventilation: analysis of the Extracorporeal Life Support Organization registry. *Crit Care* 2017;21(1):297.
626. Silverman RA, Foley F, Dalipi R, Kline M, Lesser M. The use of rhDNase in severely ill, non-intubated adult asthmatics refractory to bronchodilators: A pilot study. *Respiratory Medicine* 2012;106(8):1096-102.
627. Meduri GU, Cook TR, Turner RE, Cohen M, Leeper KV. Noninvasive positive pressure ventilation in status asthmaticus. *Chest* 1996;110(3):767-74.

628. Lim Wei J, Mohammed Akram R, Carson Kristin V, Mysore S, Labiszewski Nadina A, Wedzicha Jadwiga A, et al. Non-invasive positive pressure ventilation for treatment of respiratory failure due to severe acute exacerbations of asthma. *Cochrane Database of Systematic Reviews* 2012: Issue 12.
629. Galindo-Filho VC, Brandao DC, Ferreira RCS, Menezes MJC, Almeida-Filho P, Parreira VF, et al. Noninvasive ventilation coupled with nebulization during asthma crises: A randomized controlled trial. *Respiratory Care* 2013;58(2):241-9.
630. Pallin M, Hew M, Naughton MT. Is non-invasive ventilation safe in acute severe asthma? *Respirology* 2015;20(2):251-7.
631. Lim KL, Harrison BD. A criterion based audit of inpatient asthma care. Closing the feedback loop. *J R Coll Physicians Lond* 1992;26(1):71-5.
632. Goldberg R, Chan L, Haley P, Harmata-Booth J, Bass G. Critical pathway for the emergency department management of acute asthma: effect on resource utilization. *Ann Emerg Med* 1998;31(5):562-7.
633. Udawadia ZF, Harrison BD. An attempt to determine the optimal duration of hospital stay following a severe attack of asthma. *J R Coll Physicians Lond* 1990;24(2):112-4.
634. Pearson MG, Ryland I, Harrison BD. National audit of acute severe asthma in adults admitted to hospital. Standards of Care Committee, British Thoracic Society. *Qual Health Care* 1995;4(1):24-30.
635. Emerman CL, Woodruff PG, Cydulka RK, Gibbs MA, Pollack CV Jr, Camargo CA Jr. Prospective multicenter study of relapse following treatment for acute asthma among adults presenting to the emergency department. MARC investigators. Multicenter Asthma Research Collaboration. *Chest* 1999;115(4):919-27.
636. Cowie RL, Revitt SG, Underwood MF, Field SK. The effect of a peak flow-based action plan in the prevention of exacerbations of asthma. *Chest* 1997;112(6):1534-8.
637. Nathan JA, Pearce L, Field C, Dotesio-Eyres N, Sharples LD, Cafferty F, et al. A randomized controlled trial of follow-up of patients discharged from the hospital following acute asthma: best performed by specialist nurse or doctor? *Chest* 2006;130(1):51-7.
638. Baren JM, Boudreaux ED, Brenner BE, Cydulka RK, Rowe BH, Clark S, et al. Randomized controlled trial of emergency department interventions to improve primary care follow-up for patients with acute asthma. *Chest* 2006;129(2):257-65.
639. Davies G, Paton JY, Beaton SJ, Young D, Lenney W. Children admitted with acute wheeze/asthma during November 1998-2005: a national UK audit. *Arch Dis Child* 2008 93(11):952-8.
640. Connett GJ, Lenney W. Use of pulse oximetry in the hospital management of acute asthma in childhood. *Pediatr Pulmonol* 1993;15(6):345-9.
641. Geelhoed GC, Landau LI, Le Souef PN. Evaluation of SaO₂ as a predictor of outcome in 280 children presenting with acute asthma. *Ann Emerg Med* 1994;23(6):1236-41.
642. Schuh S, Johnson D, Stephens D, Callahan S, Canny G. Hospitalization patterns in severe acute asthma in children. *Pediatr Pulmonol* 1997;23(3):184-92.
643. Wright RO, Santucci KA, Jay GD, Steele DW. Evaluation of pre- and posttreatment pulse oximetry in acute childhood asthma. *Acad Emerg Med* 1997;4(2):114-7.
644. Brooks LJ, Cloutier MM, Afshani E. Significance of roentgenographic abnormalities in children hospitalized for asthma. *Chest* 1982;82(3):315-8.
645. Gershel JC, Goldman HS, Stein RE, Shelov SP, Ziprkowski M. The usefulness of chest radiographs in first asthma attacks. *N Engl J Med* 1983;309(6):336-9.
646. Cunningham S, Logan C, Lockerbie L, Dunn MJ, McMurray A, Prescott RJ. Effect of an integrated care pathway on acute asthma/wheeze in children attending hospital: cluster randomized trial. *Journal of Pediatrics* 2008;152(3):315-20.
647. Schuh S, Parkin P, Rajan A, Canny G, Healy R, Rieder M, et al. High-versus low-dose, frequently administered, nebulized albuterol in children with severe, acute asthma. *Pediatrics* 1989;83(4):513-8.
648. Schuh S, Reider MJ, Canny G, Pender E, Forbes T, Tan YK, et al. Nebulized albuterol in acute childhood asthma: comparison of two doses. *Pediatrics* 1990;86(4):509-13.
649. Robertson CF, Smith F, Beck R, Levison H. Response to frequent low doses of nebulized salbutamol in acute asthma. *J Pediatr* 1985;106(4):672-4.
650. Schuh S, Johnson DW, Stephens D, Callahan S, Winders P, Canny GJ. Comparison of albuterol delivered by a metered dose inhaler with spacer versus a nebulizer in children with mild acute asthma. *J Pediatr* 1999;135(1):22-7.
651. Khine H, Fuchs SM, Saville AL. Continuous vs intermittent nebulized albuterol for emergency management of asthma. *Acad Emerg Med* 1996;3(11):1019-24.
652. Papo MC, Frank J, Thompson AE. A prospective, randomized study of continuous versus intermittent nebulized albuterol for severe status asthmaticus in children. *Crit Care Med* 1993;21(10):1479-86.
653. Plotnick LH, Ducharme FM. Combined inhaled anticholinergic agents and beta-2-agonists for initial treatment of acute asthma in children (Cochrane Review). In: *The Cochrane Library*, (2) 2000.
654. Altamimi S, Robertson G, Jastaniah W, Davey A, Dehghani N, Chen R, et al. Single-dose oral dexamethasone in the emergency management of children with exacerbations of mild to moderate asthma. *Pediatric Emergency Care* 2006;22(12):786-93.
655. Gordon S, Tompkins T, Dayan PS. Randomized trial of single-dose intramuscular dexamethasone compared with prednisolone for children with acute asthma. *Pediatric Emergency Care* 2007;23(8):521-7.
656. Greenberg RA, Kerby G, Roosevelt GE. A comparison of oral dexamethasone with oral prednisone in pediatric asthma exacerbations treated in the emergency department. *Clinical Pediatrics* 2008;47(8):817-23.

657. Panickar J, Lakhanpaul M, Lambert PC, Kenia P, Stephenson T, Smyth A, et al. Oral prednisolone for preschool children with acute virus-induced wheezing. *N Engl J Med* 2009;360(4):329-38.
658. Becker JM, Arora A, Scarfone RJ, Spector ND, Fontana-Penn ME, Gracely E, et al. Oral versus intravenous corticosteroids in children hospitalized with asthma. *J Allergy Clin Immunol* 1999;103(4):586-90.
659. Barnett PL, Caputo GL, Baskin M, Kuppermann N. Intravenous versus oral corticosteroids in the management of acute asthma in children. *Ann Emerg Med* 1997;29(2):212-7.
660. Langton Hewer S, Hobbs J, Reid F, Lenney W. Prednisolone in acute childhood asthma: clinical responses to three dosages. *Respir Med* 1998;92(3):541-6.
661. Edmonds ML, Brenner BE, Camargo CA, Rowe BH. Inhaled steroids in acute asthma following emergency department discharge (Cochrane Review). In: *The Cochrane Library*, (3) 2000.
662. McKean MC, Ducharme F. Inhaled steroids for episodic viral wheeze of childhood (Cochrane Review). In: *The Cochrane Library*, 2000.
663. Schuh S, Reisman J, Alshehri M, Dupuis A, Corey M, Arseneault R, et al. A comparison of inhaled fluticasone and oral prednisone for children with severe acute asthma. *N Engl J Med* 2000;343(10):689-94.
664. Ducharme FM, Lemire C, Noya FJ, Davis GM, Alos N, Leblond H, et al. Preemptive use of high-dose fluticasone for virus-induced wheezing in young children. *New England Journal of Medicine* 2009;360(4):339-53.
665. Papi A, Nicolini G, Boner AL, Baraldi E, Cutrera R, Fabbri LM, et al. Short term efficacy of nebulized beclomethasone in mild-to-moderate wheezing episodes in pre-school children. *Italian Journal of Pediatrics* 2011;37:39.
666. Schuh S, Dick PT, Stephens D, Hartley M, Khaikin S, Rodrigues L, et al. High-dose inhaled fluticasone does not replace oral prednisolone in children with mild to moderate acute asthma. *Pediatrics* 2006;118(2):644-50.
667. Upham BD, Mollen CJ, Scarfone RJ, Seiden J, Chew A, Zorc JJ. Nebulized budesonide added to standard pediatric emergency department treatment of acute asthma: a randomized, double-blind trial. *Academic Emergency Medicine* 2011;18(7):665-73.
668. Volovitz B, Bilavsky E, Nussinovitch M. Effectiveness of high repeated doses of inhaled budesonide or fluticasone in controlling acute asthma exacerbations in young children. *Journal of Asthma* 2008;45(7):561-7.
669. Harmanci K, Bakirtas A, Turkas I, Degim T. Oral montelukast treatment of preschool-aged children with acute asthma. *Annals of allergy, asthma & immunology* 2006;96(5):731-5.
670. Powell C, Kolamunnage-Dona R, Lowe J, Boland A, Petrou S, Doull I, et al. Magnesium sulphate in acute severe asthma in children (MAGNETIC): a randomised, placebo-controlled trial.[Erratum appears in *Lancet Respir Med*. 2013 Jun;1(4):285]. *The Lancet Respiratory Medicine* 2013;1(4):301-8.
671. Singhi S, Grover S, Bansal A, Chopra K. Randomised comparison of intravenous magnesium sulphate, terbutaline and aminophylline for children with acute severe asthma. *Acta Paediatr* 2014;103(12):1301-6.
672. Travers AH, Jones AP, Camargo CA, Milan SJ, Rowe BH. Intravenous beta2-agonists versus intravenous aminophylline for acute asthma (Cochrane Review). In: *The Cochrane Library*, 2012.
673. Ciarallo L, Brousseau D, Reinert S. Higher-dose intravenous magnesium therapy for children with moderate to severe acute asthma. *Arch Pediatr Adolesc Med* 2000;154(10):979-83.
674. Goodman DC, Littenberg B, O'Connor GT, Brooks JG. Theophylline in acute childhood asthma: a meta-analysis of its efficacy. *Pediatr Pulmonol* 1996;21(4):211-8.
675. Yung M, South M. Randomised controlled trial of aminophylline for severe acute asthma. *Arch Dis Child* 1998;79(5):405-10.
676. Griffiths B, Kew Kayleigh M. Intravenous magnesium sulfate for treating children with acute asthma in the emergency department. *Cochrane Database of Systematic Reviews* 2016: Issue 4.
677. Jat Kana R, Chawla D. Ketamine for management of acute exacerbations of asthma in children. *Cochrane Database of Systematic Reviews* 2012: Issue 11.
678. Schutte D, Zwitterloot AM, Houmes R, De Hoog M, Draaisma JM, Lemson J. Sevoflurane therapy for life-threatening asthma in children. *British Journal of Anaesthesia* 2013;111(6):967-70.
679. Korang SK, Feinberg J, Wetterslev J, Jakobsen JC. Non-invasive positive pressure ventilation for acute asthma in children. *Cochrane Database of Systematic Reviews* 2016;2016 Issue 9.
680. Mayordomo-Colunga J, Medina A, Rey C, Concha A, Menendez S, Arcos ML, et al. Non-invasive ventilation in pediatric status asthmaticus: a prospective observational study. *Pediatric Pulmonology* 2011;46(10):949-55.
681. Stormon MO, Mellis CM, Van Asperen PP, Kilham HA. Outcome evaluation of early discharge of asthmatic children from hospital: a randomized control trial. *J Qual Clin Pract* 1999;19(3):149-54.
682. Chung KF, Godard P, Adelroth E, Ayres J, Barnes N, Barnes P, et al. Difficult/therapy-resistant asthma: The need for an integrated approach to define clinical phenotypes, evaluate risk factors, understand pathophysiology and find novel therapies. *European Respiratory Journal* 1999;13(5):1198-208.
683. Prys-Picard CO, Campbell SM, Ayres JG, Miles JF, Niven RM, Consensus on Difficult Asthma Consortium UK. Defining and investigating difficult asthma: developing quality indicators. *Respiratory Medicine* 2006;100(7):1254-61.

684. Bratton DL, Price M, Gavin L, Glenn K, Brenner M, Gelfand EW, et al. Impact of a multidisciplinary day program on disease and healthcare costs in children and adolescents with severe asthma: a two-year follow-up study. *Pediatr Pulmonol* 2001;31(3): 177-89.
685. Robinson DS, Campbell DA, Durham SR, Pfeffer J, Barnes PJ, Chung KF, et al. Systematic assessment of difficult-to-treat asthma. *European Respiratory Journal* 2003;22(3):478-83.
686. Weinstein AG, McKee L, Stapleford J, Faust D. An economic evaluation of short-term inpatient rehabilitation for children with severe asthma. *Journal of Allergy & Clinical Immunology* Vol 1996;98(2):264-73.
687. Gamble J, Stevenson M, McClean E, Heaney LG. The prevalence of nonadherence in difficult asthma. *American Journal of Respiratory & Critical Care Medicine* 2009;180(9):817-22.
688. Murphy AC, Proeschal A, Brightling CE, Wardlaw AJ, Pavord I, Bradding P, et al. The relationship between clinical outcomes and medication adherence in difficult-to-control asthma. *Thorax* 2012;67(8): 751-3.
689. Bracken M, Fleming L, Hall P, Van Stiphout N, Bossley C, Biggart E, et al. The importance of nurse-led home visits in the assessment of children with problematic asthma. *Arch Dis Child* 2009;94(10):780-4.
690. Ranganathan SC, Payne DN, Jaffe A, McKenzie SA. Difficult asthma: defining the problems. *Pediatr Pulmonol* 2001;31(2):114-20.
691. Apter AJ, Wang X, Bogen DK, Rand CS, McElligott S, Polsky D, et al. Problem solving to improve adherence and asthma outcomes in urban adults with moderate or severe asthma: A randomized controlled trial. *Journal of Allergy and Clinical Immunology* 2011;128(3):516-23.e5.
692. Vamos M, Kolbe J. Psychological factors in severe chronic asthma. *Australian & New Zealand Journal of Psychiatry* 1999;33(4):538-44.
693. Vila G, Nollet-Clemencon C, de Blic J, Mouren-Simeoni MC, Scheinmann P. Asthma severity and psychopathology in a tertiary care department for children and adolescent. *European Child & Adolescent Psychiatry* 1998;7(3):137-44.
694. Wainwright NW, Surtees PG, Wareham NJ, Harrison BD. Psychosocial factors and incident asthma hospital admissions in the EPIC-Norfolk cohort study. *Allergy* 2007;62(5):554-60.
695. Wamboldt MZ, Weintraub P, Krafchick D, Wamboldt FS. Psychiatric family history in adolescents with severe asthma. *Journal of the American Academy of Child & Adolescent Psychiatry* 1996;35(8):1042-9.
696. Miles JF, Garden GM, Tunnicliffe WS, Cayton RM, Ayres JG. Psychological morbidity and coping skills in patients with brittle and non-brittle asthma: a case-control study. *Clinical & Experimental Allergy* 1997;27(10):1151-9.
697. ten Brinke A, Ouwerkerk ME, Bel EH, Spinhoven P. Similar psychological characteristics in mild and severe asthma. *Journal of Psychosomatic Research* 2001;50(1):7-10.
698. Wamboldt MZ, Fritz G, Mansell A, McQuaid EL, Klein RB. Relationship of asthma severity and psychological problems in children. *J Amer Acad Child Adolescent Psychiatry* 1998;37(9):943-50.
699. McQuaid EL, Kopel SJ, Nassau JH. Behavioral adjustment in children with asthma: A meta-analysis. *Journal of Developmental & Behavioral Pediatrics* 2001;22(6):430-9.
700. Brown ES, Vigil L, Khan DA, Liggin JD, Carmody TJ, Rush AJ. A randomized trial of citalopram versus placebo in outpatients with asthma and major depressive disorder: a proof of concept study. *Biological psychiatry* 2005;58(11):865-70.
701. Godding V, Kruth M, Jamart J. Joint consultation for high-risk asthmatic children and their families, with pediatrician and child psychiatrist as co-therapists: model and evaluation. *Family Process* 1997;36(3):265-80.
702. Smith JR, Mildenhall S, Noble MJ, Shepstone L, Koutantji M, Mugford M, et al. The Coping with Asthma Study: a randomised controlled trial of a home based, nurse led psychoeducational intervention for adults at risk of adverse asthma outcomes. *Thorax* 2005;60(12):1003-11.
703. Smith JR, Mugford M, Holland R, Candy B, Noble MJ, Harrison BD, et al. A systematic review to examine the impact of psycho-educational interventions on health outcomes and costs in adults and children with difficult asthma. *Health Technology Assessment* 2005;9(23):iii-iv,1-167.
704. Position statement. Environmental allergen avoidance in allergic asthma. Ad Hoc Working Group on Environmental Allergens and Asthma. *J Allergy Clin Immunol* 1999;103(2 Pt 1):203-5.
705. O'Driscoll BR, Hopkinson LC, Denning DW. Mold sensitization is common amongst patients with severe asthma requiring multiple hospital admissions. *BMC Pulmonary Medicine* 2005;5:4.
706. Zureik M, Neukirch C, Leynaert B, Liard R, Bousquet J, Neukirch F, et al. Sensitisation to airborne moulds and severity of asthma: cross sectional study from European Community respiratory health survey. *British Medical Journal* 2002;325(7361):411-4.
707. Black PN, Udy AA, Brodie SM. Sensitivity to fungal allergens is a risk factor for life-threatening asthma. *Allergy* 2000;55(5):501-4.
708. O'Hollaren MT, Yunginger JW, Offord KP, Somers MJ, O'Connell EJ, Ballard DJ, et al. Exposure to an aeroallergen as a possible precipitating factor in respiratory arrest in young patients with asthma. *N Engl J Med* 1991;324(6):359-63.
709. Green RH, Brightling CE, McKenna S, Hargadon B, Parker D, Bradding P, et al. Asthma exacerbations and sputum eosinophil counts: a randomised controlled trial. *Lancet* 2002;360(9347):1715-21.
710. Jayaram L, Pizzichini MM, Cook RJ, Boulet LP, Lemiere C, Pizzichini E, et al. Determining asthma treatment by monitoring sputum cell counts: Effect on exacerbations. *Eur Respir J* 2006;27(3):483-94.
711. Chlumsky J, Striz I, Terl M, Vondracek J. Strategy aimed at reduction of sputum eosinophils decreases exacerbation rate in patients with asthma. *J Int Med Res* 2006;34(2):129-39.

712. Fahy JV, Boushey HA, Lazarus SC, Mauger EA, Cherniack RM, Chinchilli VM, et al. Safety and reproducibility of sputum induction in asthmatic subjects in a multicenter study. *Am J Respir Crit Care Med* 2001;163(6):1470-5.
713. Grootendorst DC, van den Bos JW, Romeijn JJ, Veselic-Charvat M, Duiverman EJ, Vrijlandt EJ, et al. Induced sputum in adolescents with severe stable asthma. Safety and the relationship of cell counts and eosinophil cationic protein to clinical severity. *Eur Respir J* 1999;13(3):647-53.
714. Loh LC, Kanabar V, D'Amato M, Barnes NC, O'Connor BJ. Sputum induction in corticosteroid-dependant asthmatics: risks and airway cellular profile. *Asian Pacific Journal of Allergy & Immunology* 2005;23(4):189-96.
715. Tarodo de la Fuente P, Romagnoli M, Carlsson L, Godard P, Bousquet J, Chanez P. Eosinophilic inflammation assessed by induced sputum in corticosteroid-dependent asthma. *Respiratory Medicine* 1999;93(3):183-9.
716. Pijnenburg MW, Hofhuis W, Hop WC, De Jongste JC. Exhaled nitric oxide predicts asthma relapse in children with clinical asthma remission. *Thorax* 2005;60(3):215-8.
717. Smith AD, Cowan JO, Brasset KP, Herbison GP, Taylor DR. Use of exhaled nitric oxide measurements to guide treatment in chronic asthma. *N Engl J Med* 2005;352(21):2163-73.
718. English A, Park MJ, Shafer MA, Kreipe RE, D'Angelo LJ. Health care reform and adolescents-an agenda for the lifespan: a position paper of the Society for Adolescent Medicine. *J Adolesc Health* 2009;45(3):310-5.
719. Royal Australasian College of Physicians. National standards for the care of children and adolescents. Sydney: 2008. [cited 02 Jul 2014] Available from <http://www.racp.edu.au/index.cfm?objectid=393E4ADA-CDAA-D1AF-0D543B5DC13C7B46>
720. Royal College of Paediatrics and Child Health. Bridging the gaps: health care for adolescents. London: Royal College of Paediatrics and Child Health; 2003. [cited 02 Jul 2014] Available from <http://www.rcpsych.ac.uk/files/pdfversion/cr114.pdf>
721. Royal Australasian College of Physicians Joint Adolescent Health Committee. Confidential Health Care for Adolescents and Young People. 2010. [cited 11th April 2011] Available from <http://www.racp.edu.au/index.cfm?objectid=655B70C1-A0F2-D4A4-6DB6505DCA1AB937>
722. Lai CK, Beasley R, Crane J, Foliaki S, Shah J, Weiland S, et al. Global variation in the prevalence and severity of asthma symptoms: phase three of the International Study of Asthma and Allergies in Childhood (ISAAC). *Thorax* 2009;64(6):476-83.
723. Siersted HC, Boldsen J, Hansen HS, Mostgaard G, Hyldebrandt N. Population based study of risk factors for underdiagnosis of asthma in adolescence: Odense schoolchild study. *BMJ* 1998;316(7132):651-5.
724. Yeatts KB, Shy CM. Prevalence and consequences of asthma and wheezing in African-American and white adolescents. *J Adolesc Health* 2001;29(5):314-9.
725. Yeatts K, Davis KJ, Sotir M, Herget C, Shy C. Who gets diagnosed with asthma? Frequent wheeze among adolescents with and without a diagnosis of asthma. *Pediatrics* 2003;111(5 Pt 1):1046-54.
726. Yeatts K, Johnston Davis K, Peden D, Shy C. Health consequences associated with frequent wheezing in adolescents without asthma diagnosis. *Eur Respir J* 2003;22(5):781-6.
727. Yeatts K, Shy C, Sotir M, Music S, Herget C. Health consequences for children with undiagnosed asthma-like symptoms. *Arch Pediatr Adolesc Med* 2003;157(6):540-4.
728. Abramson JM, Wollan P, Kurland M, Yawn BP. Feasibility of school-based spirometry screening for asthma. *J Sch Health* 2003;73(4):150-3.
729. Yawn BP. Asthma screening, case identification and treatment in school-based programs. *Curr Opin Pulm Med* 2006;12(1):23-7.
730. Henriksen AH, Tveit KH, Holmen TL, Sue-Chu M, Bjermer L. A study of the association between exercise-induced wheeze and exercise versus methacholine-induced bronchoconstriction in adolescents. *Pediatr Allergy Immunol* 2002;13(3):203-8.
731. Seear M, Wensley D, West N. How accurate is the diagnosis of exercise induced asthma among Vancouver schoolchildren? *Arch Dis Child* 2005;90(9):898-902.
732. Mallol J, Castro-Rodriguez JA. Differences in prevalence of asthma, rhinitis, and eczema between parental and self-completed questionnaires in adolescents. *Pediatr Pulmonol* 2006;41(5):482-7.
733. Raat H, Mangunkusumo RT, Mohangoo AD, Juniper EF, Van Der Lei J. Internet and written respiratory questionnaires yield equivalent results for adolescents. *Pediatr Pulmonol* 2007;42(4):357-61.
734. Juniper EF, Svensson K, Mork AC, Stahl E. Modification of the asthma quality of life questionnaire (standardised) for patients 12 years and older. *Health Qual Life Outcomes* 2005;3:58.
735. Burkhart PV, Svavarsdottir EK, Rayens MK, Oakley MG, Orlygsdottir B. Adolescents with asthma: predictors of quality of life. *J Adv Nurs* 2009;65(4):860-6.
736. Obase Y, Shimoda T, Kawano T, Saeki S, Tomari S, Izaki K, et al. Bronchial hyperresponsiveness and airway inflammation in adolescents with asymptomatic childhood asthma. *Allergy* 2003;58(3):213-20.
737. Rodriguez MA, Winkleby MA, Ahn D, Sundquist J, Kraemer HC. Identification of population subgroups of children and adolescents with high asthma prevalence: findings from the Third National Health and Nutrition Examination Survey. *Arch Pediatr Adolesc Med* 2002;156(3):269-75.

738. Duse M, Donato F, Porteri V, Pirali F, Spinoni V, Tosoni C, et al. High prevalence of atopy, but not of asthma, among children in an industrialized area in North Italy: the role of familial and environmental factors--a population-based study. *Pediatr Allergy Immunol* 2007;18(3):201-8.
739. Del-Rio-Navarro B, Berber A, Blandon-Vijil V, Ramirez-Aguilar M, Romieu I, Ramirez-Chanona N, et al. Identification of asthma risk factors in Mexico City in an International Study of Asthma and Allergy in Childhood survey. *Allergy Asthma Proc* 2006;27(4):325-33.
740. Hyvarinen MK, Kotaniemi-Syrjanen A, Reijonen TM, Korhonen K, Korppi MO. Teenage asthma after severe early childhood wheezing: an 11-year prospective follow-up. *Pediatr Pulmonol* 2005;40(4):316-23.
741. Anand D, Stevenson CJ, West CR, Pharoah PO. Lung function and respiratory health in adolescents of very low birth weight. *Arch Dis Child* 2003;88(2):135-8.
742. Fagan JK, Scheff PA, Hryhorczuk D, Ramakrishnan V, Ross M, Persky V. Prevalence of asthma and other allergic diseases in an adolescent population: association with gender and race. *Ann Allergy Asthma Immunol* 2001;86(2):177-84.
743. Debley JS, Redding GJ, Critchlow CW. Impact of adolescence and gender on asthma hospitalization: a population-based birth cohort study. *Pediatr Pulmonol* 2004;38(6):443-50.
744. Nicolai T, Pereszlenyiova-Bliznakova L, Illi S, Reinhardt D, von Mutius E. Longitudinal follow-up of the changing gender ratio in asthma from childhood to adulthood: role of delayed manifestation in girls. *Pediatr Allergy Immunol* 2003;14(4):280-3.
745. Bernard A, Nickmilder M, Voisin C. Outdoor swimming pools and the risks of asthma and allergies during adolescence. *Eur Respir J* 2008;32(4):979-88.
746. Bernard A, Carbonnelle S, de Burbure C, Michel O, Nickmilder M. Chlorinated pool attendance, atopy, and the risk of asthma during childhood. *Environ Health Perspect* 2006;114(10):1567-73.
747. Goodwin RD, Fergusson DM, Horwood LJ. Asthma and depressive and anxiety disorders among young persons in the community. *Psychol Med* 2004;34(8):1465-74.
748. Richardson LP, Lozano P, Russo J, McCauley E, Bush T, Katon W. Asthma symptom burden: relationship to asthma severity and anxiety and depression symptoms. *Pediatrics* 2006;118(3):1042-51.
749. Hommel KA, Chaney JM, Wagner JL, McLaughlin MS. Asthma-specific quality of life in older adolescents and young adults with long-standing asthma: the role of anxiety and depression. *J Clin Psychol Med Settings* 2002;9(3):185-92.
750. Powell C, Brazier A. Psychological approaches to the management of respiratory symptoms in children and adolescents. *Paediatr Respir Rev* 2004;5(3):214-24.
751. Katon W, Russo J, Richardson L, McCauley E, Lozano P. Anxiety and depression screening for youth in a primary care population. *Ambul Pediatr* 2008;8(3):182-8.
752. Brenner JS, Kelly CS, Wenger AD, Brich SM, Morrow AL. Asthma and obesity in adolescents: is there an association? *J Asthma* 2001;38(6):509-15.
753. Mai XM, Nilsson L, Axelsson O, Braback L, Sandin A, Kjellman NI, et al. High body mass index, asthma and allergy in Swedish schoolchildren participating in the International Study of Asthma and Allergies in Childhood: Phase II. *Acta Paediatr* 2003;92(10):1144-8.
754. Gilliland FD, Berhane K, Islam T, McConnell R, Gauderman WJ, Gilliland SS, et al. Obesity and the risk of newly diagnosed asthma in school-age children. *Am J Epidemiol* 2003;158(5):406-15.
755. Debley JS, Carter ER, Redding GJ. Prevalence and impact of gastroesophageal reflux in adolescents with asthma: a population-based study. *Pediatr Pulmonol* 2006;41(5):475-81.
756. Thakkar K, Boatright RO, Gilger MA, El-Serag HB. Gastroesophageal reflux and asthma in children: a systematic review. *Pediatrics* 2010;125(4):e925-30.
757. Rasmussen F, Taylor DR, Flannery EM, Cowan JO, Greene JM, Herbison GP, et al. Outcome in adulthood of asymptomatic airway hyperresponsiveness in childhood: a longitudinal population study. *Pediatr Pulmonol* 2002;34(3):164-71.
758. Orbon KH, van der Gulden JW, Schermer TR, van den Nieuwenhof L, Boot CR, van den Hoogen H, et al. Vocational and working career of asthmatic adolescents is only slightly affected. *Respir Med* 2006;100(7):1163-73.
759. Gerald LB, Gerald JK, Gibson L, Patel K, Zhang S, McClure LA. Changes in environmental tobacco smoke exposure and asthma morbidity among urban school children. *Chest* 2009;135(4):911-6.
760. Precht DH, Keiding L, Madsen M. Smoking patterns among adolescents with asthma attending upper secondary schools: a community-based study. *Pediatrics* 2003;111(5 Pt 1):e562-8.
761. Annesi-Maesano I, Oryszczyn MP, Raheison C, Kopferschmitt C, Pauli G, Taytard A, et al. Increased prevalence of asthma and allied diseases among active adolescent tobacco smokers after controlling for passive smoking exposure. A cause for concern? *Clin Exp Allergy* 2004;34(7):1017-23.
762. Genuneit J, Weinmayr G, Radon K, Dressel H, Windstetter D, Rzehak P, et al. Smoking and the incidence of asthma during adolescence: results of a large cohort study in Germany. *Thorax* 2006;61(7):572-8.
763. Larsson L. Incidence of asthma in Swedish teenagers: relation to sex and smoking habits. *Thorax* 1995;50(3):260-4.
764. Hedman L, Bjerg A, Sundberg S, Forsberg B, Ronmark E. Both environmental tobacco smoke and personal smoking is related to asthma and wheeze in teenagers. *Thorax* 2011;66(1):20-5.
765. Lombardi C, Gani F, Landi M, Boner A, Canonica GW, Passalacqua G. Clinical and therapeutic aspects of allergic asthma in adolescents. *Pediatr Allergy Immunol* 2003;14(6):453-7.

766. Reznik M, Ozuah PO, Franco K, Cohen R, Motlow F. Use of complementary therapy by adolescents with asthma. *Arch Pediatr Adolesc Med* 2002;156(10):1042-4.
767. Juntunen-Backman K, Kajosaari M, Laurikainen K, Malinen A, Kaila M, Mustala L, et al. Comparison of Easyhaler(R) metered-dose, dry powder Inhaler and a pressurised metered-dose inhaler plus spacer in the treatment of asthma in children. *Clin Drug Invest* 2002;22(12):827-35.
768. Adler LM, Anand C, Wright FG deL, Barret CF, McKeith D, Clark WI, et al. Efficacy and tolerability of beclomethasone dipropionate delivered by a novel multidose dry powder inhaler (Clickhaler®) versus a metered-dose inhaler in children with asthma. *Current Therapeutic Research* 2001;62(11):758-69.
769. Brennan VK, Osman LM, Graham H, Critchlow A, Everard ML. True device compliance: the need to consider both competence and contrivance. *Respir Med* 2005;99(1):97-102.
770. Edgecombe K, Latter S, Peters S, Roberts G. Health experiences of adolescents with uncontrolled severe asthma. *Arch Dis Child* 2010;95(12):985-91.
771. Salisbury C, Francis C, Rogers C, Parry K, Thomas H, Chadwick S, et al. A randomised controlled trial of clinics in secondary schools for adolescents with asthma. *Br J Gen Pract* 2002;52(485):988-96.
772. Shah S, Peat JK, Mazurski EJ, Wang H, Sindhusake D, Bruce C, et al. Effect of peer led programme for asthma education in adolescents: cluster randomised controlled trial. *BMJ* 2001;322(7286):583-5.
773. Henry RL, Lough S, Mellis C. National policy on asthma management for schools. *J Paediatr Child Health* 2006;42(9):491-5.
774. Royal College of Physicians of Edinburgh Steering Group. Think Transition: developing the essential link between paediatric and adult care. Edinburgh: Royal College of Physicians of Edinburgh; 2008. [cited 09 Jul 2014] Available from <http://www.cen.scot.nhs.uk/files/160-think-transition-edinburgh.pdf>
775. Scal P, Davern M, Ireland M, Park K. Transition to adulthood: delays and unmet needs among adolescents and young adults with asthma. *J Pediatr* 2008;152:471-5.
776. Sawyer S, Drew S, Duncan R. Adolescents with chronic disease--the double whammy. *Aust Fam Physician* 2007;36(8):622-7.
777. Cordina M, Hughes CM, McElroy JC. Health-related issues of importance to school children with asthma - a qualitative study. *J Soc Adm Pharm* 2002;19(5):162-70.
778. Cohen R, Franco K, Motlow F, Reznik M, Ozuah PO. Perceptions and attitudes of adolescents with asthma. *J Asthma* 2003;40(2):207-11.
779. Kyngas H. Patient education: perspective of adolescents with a chronic disease. *J Clin Nurs* 2003;12(5):744-51.
780. Bender BG, Rankin A, Tran ZV, Wamboldt FS. Brief-interval telephone surveys of medication adherence and asthma symptoms in the Childhood Asthma Management Program Continuation Study. *Ann Allergy Asthma Immunol* 2008;101(4):382-6.
781. Buston KM, Wood SF. Non-compliance amongst adolescents with asthma: listening to what they tell us about self-management. *Fam Pract* 2000;17(2):134-8.
782. Kyngas HA. Compliance of adolescents with asthma. *Nurs Health Sci* 1999;1(3):195-202.
783. Bender BG. Risk taking, depression, adherence, and symptom control in adolescents and young adults with asthma. *Am J Respir Crit Care Med* 2006;173(9):953-7.
784. Sawyer S, Bowes G. Caring for adolescents with asthma: do we know how to? *Med J Aust* 1996;165(9):463-4.
785. Goldenring JM, Cohen E. Getting into adolescent heads. *Contemp Pediatr* 1988;5(7):75-90.
786. Kyngas HA, Kroll T, Duffy ME. Compliance in adolescents with chronic diseases: a review. *J Adolesc Health* 2000;26(6):379-88.
787. Gerald LB, McClure LA, Mangan JM, Harrington KF, Gibson L, Erwin S, et al. Increasing adherence to inhaled steroid therapy among schoolchildren: randomized, controlled trial of school-based supervised asthma therapy. *Pediatrics* 2009;123(2):466-74.
788. Schatz M, Harden K, Forsythe A, Chilingar L, Hoffman C, Sperling W, et al. The course of asthma during pregnancy, post partum, and with successive pregnancies: a prospective analysis. *J Allergy Clin Immunol* 1988;81(3):509-17.
789. Dombrowski MP, Schatz M, Wise R, Momirova V, Landon M, Mabie W, et al. Asthma during pregnancy. *Obstetrics and Gynecology* 2004;103(1):5-12.
790. Juniper E F, Newhouse M T. Effect of pregnancy on asthma: a systematic review and meta-analysis. In: Schatz M, Zeiger RS, Claman HN, editors. *Asthma and immunological diseases in pregnancy and early infancy*. New York: Marcel Dekker; 1998. p.401-25.
791. Gluck JC, Gluck PA. The effect of pregnancy on the course of asthma. *Immunology & Allergy Clinics of North America* 2006;26(1):63-80.
792. Kwon HL, Belanger K, Bracken MB. Effect of pregnancy and stage of pregnancy on asthma severity: a systematic review. *American Journal of Obstetrics and Gynecology* 2004;190(5):1201-10.
793. Schatz M, Zeiger RS, Hoffman CP, Harden K, Forsythe A, Chilingar L, et al. Perinatal outcomes in the pregnancies of asthmatic women: a prospective controlled analysis. *Am J Respir Crit Care Med* 1995;151(4):1170-4.
794. Wendel PJ, Ramin SM, Barnett-Hamm C, Rowe TF, Cunningham FG. Asthma treatment in pregnancy: a randomized controlled study. *Am J Obstet Gynecol* 1996;175(1):150-4.
795. Stenius-Aarniala BS, Hedman J, Teramo KA. Acute asthma during pregnancy. *Thorax* 1996;51(4):411-4.
796. Schatz M. Interrelationships between asthma and pregnancy: a literature review. *J Allergy Clin Immunol* 1999;103(2 Pt 2):S330-6.
797. Stenius-Aarniala B, Piirila P, Teramo K. Asthma and pregnancy: a prospective study of 198 pregnancies. *Thorax* 1988;43(1):12-8.

798. Fitzsimons R, Greenberger PA, Patterson R. Outcome of pregnancy in women requiring corticosteroids for severe asthma. *J Allergy Clin Immunol* 1986;78(2):349-53.
799. Perlow JH, Montgomery D, Morgan MA, Towers CV, Porto M. Severity of asthma and perinatal outcome. *Am J Obstet Gynecol* 1992;167(4 Pt 1):963-7.
800. Schatz M, Zeiger RS, Hoffman CP. Intrauterine growth is related to gestational pulmonary function in pregnant asthmatic women. Kaiser-Permanente Asthma and Pregnancy Study Group. *Chest* 1990;98(2):389-92.
801. Demissie K, Breckenridge MB, Rhoads GG. Infant and maternal outcomes in the pregnancies of asthmatic women. *Am J Respir Crit Care Med* 1998;158(4):1091-5.
802. Kallen B, Rydhstroem H, Aberg A. Asthma during pregnancy--a population based study. *Eur J Epidemiol* 2000;16(2):167-71.
803. Bracken MB, Triche EW, Belanger K, Saftlas A, Beckett WS, Leaderer BP. Asthma symptoms, severity, and drug therapy: a prospective study of effects on 2205 pregnancies. *Obstetrics & Gynecology* 2003;102(4):739-52.
804. Murphy VE, Clifton VL, Gibson PG. Asthma exacerbations during pregnancy: incidence and association with adverse pregnancy outcomes. *Thorax* 2006;61(2):169-76.
805. Schatz M, Dombrowski MP, Wise R, Momirova V, Landon M, Mabie W, et al. Spirometry is related to perinatal outcomes in pregnant women with asthma. *American Journal of Obstetrics & Gynecology* 2006;194(1):120-6.
806. Cydulka RK, Emerman CL, Schreiber D, Molander KH, Woodruff PG, Camargo CA Jr. Acute asthma among pregnant women presenting to the emergency department. *Am J Respir Crit Care Med* 1999;160(3):887-92.
807. Department of Health. Why mothers die: report on confidential enquiries into maternal deaths in the United Kingdom 1994-1996. London: Stationery Office; 1998. [cited
808. Lewis G, Drife J. Why mothers die, 1997-1999. The fifth report of the Confidential Enquiries into Maternal Deaths in the United Kingdom London: RCOG Press; 2001. [cited
809. Lewis G, editor. The Confidential Enquiry into Maternal and Child Health (CEMACH). Why mothers die: the sixth report of the confidential enquiries into maternal deaths in the UK. London: RCOG Press; 2004. [cited
810. Lewis G, editor. The Confidential Enquiry into Maternal and Child Health (CEMACH). Saving mothers' lives: reviewing maternal deaths to make motherhood safer 2003-2005. The seventh report on confidential enquiries into maternal deaths in the UK. London: CEMACH; 2007. [cited
811. Cantwell R, Clutton-Brock T, Cooper G, Dawson A, Drife J, Garrod D, et al. Saving Mothers' Lives: Reviewing maternal deaths to make motherhood safer: 2006-2008. The Eighth Report of the Confidential Enquiries into Maternal Deaths in the United Kingdom. *Bjog* 2011;118 Suppl 1:1-203.
812. Intensive Care National Audit and Research Centre. Female admissions (aged 16-50 years) to adult, general critical care units in England, Wales and Northern Ireland reported as 'currently pregnant' or 'recently pregnant'. London: ICNARC; 2013. [cited 09 Jul 2014] Available from http://www.oaa-anaes.ac.uk/assets/_managed/cms/files/Obstetric%20admissions%20to%20critical%20care%202009-2012%20-%20FINAL.pdf
813. Campbell LA, Klocke RA. Implications for the pregnant patient. *Am J Respir Crit Care Med* 2001;163(5):1051-4.
814. Templeton A, Kelman GR. Maternal blood-gases, (PAo2--Pao2), physiological shunt and VD/VT in normal pregnancy. *Br J Anaesth* 1976;48(10):1001-4.
815. Van Hook J, Harvey CJ, Anderson GD. Effect of pregnancy on maternal oxygen saturation values: use of reflectance pulse oximetry during pregnancy. *South Med J* 1996;89(12):1188-92.
816. Altman D, Carroli G, Duley L, Farrell B, Moodley J, Neilson J, et al. Do women with pre-eclampsia, and their babies, benefit from magnesium sulphate? The magpie trial: a randomised placebo-controlled trial. *Lancet* 2002;359(9321):1877-90.
817. Gee J, Packer B, Millen J, Robin E. Pulmonary mechanics during pregnancy. *J Clin Invest* 1967;46(6):945-52.
818. Izci B, Riha R, Martin S, Vennelle M, Liston W, Dundas K, et al. The upper airway in pregnancy and pre-eclampsia. *Am J Respir Crit Care Med* 2003;167(2):137-40.
819. Schatz M, Zeiger RS, Harden K, Hoffman CC, Chilingar L, Petitti D. The safety of asthma and allergy medications during pregnancy. *J Allergy Clin Immunol* 1997;100(3):301-6.
820. Chambers C. Safety of asthma and allergy medications in pregnancy. *Immunology & Allergy Clinics of North America* 2006;26(1):13-28.
821. Tata L, Lewis S, McKeever T, Smith C, Doyle P, Smeeth L, et al. Effect of maternal asthma, exacerbations and asthma medication use on congenital malformations in offspring: a UK population-based study. *Thorax* 2008;63(11).
822. Rayburn WF, Atkinson BD, Gilbert K, Turnbull GL. Short-term effects of inhaled albuterol on maternal and fetal circulations. *Am J Obstet Gynecol* 1994;171(3):770-3.
823. Schatz M, Dombrowski M, Wise R, Momirova V, Landon M, Mabie W, et al. The relationship of asthma medication use to perinatal outcomes. *J Allergy Clin Immunol* 2004 113(6):1040-5.
824. Schatz M, Zeiger RS, Harden KM, Hoffman CP, Forsythe AB, Chilingar LM, et al. The safety of inhaled beta-agonist bronchodilators during pregnancy. *J Allergy Clin Immunol* 1988;82(4):686-95.
825. Mann RD, Kubota K, Pearce G, Wilton L. Salmeterol: a study by prescription-event monitoring in a UK cohort of 15,407 patients. *J Clin Epidemiol* 1996;49(2):247-50.

826. Wilton L, Shakir SA. A post-marketing surveillance study of formoterol (Foradil): its use in general practice in England. *Drug Safety* 2002;25(3):213-23.
827. Gluck JC, Gluck PA. Asthma controller therapy during pregnancy. *American Journal of Obstetrics & Gynecology* 2005;192(2):369-80.
828. Nelson H, Weiss S, Bleecker E, Yancey S, Dorinsky P. The salmeterol multicenter asthma research trial (SMART Study Group): a comparison of usual pharmacotherapy for asthma or usual pharmacotherapy plus salmeterol. *Chest* 2006;129(1):15-26.
829. Perrio MJ, Wilton LV, Shakir SA. A modified prescription-event monitoring study to assess the introduction of Seretide Evohaler in England: an example of studying risk monitoring in pharmacovigilance. *Drug Saf* 2007;30(8):681-95.
830. Greenberger PA, Patterson R. Beclomethasone dipropionate for severe asthma during pregnancy. *Ann Intern Med* 1983;98(4):478-80.
831. Dombrowski M, Thom E, McNellis D. Maternal-Fetal Medicine Units (MFMU) studies of inhaled corticosteroids during pregnancy. *J Allergy Clin Immunol* 1999;103(2 Pt 2):S356-9.
832. Dombrowski MP, Brown CL, Berry SM. Preliminary experience with triamcinolone acetonide during pregnancy. *J Matern Fetal Med* 1996;5(6):310-3.
833. Kallen B, Rydhstroem H, Aberg A. Congenital malformations after the use of inhaled budesonide in early pregnancy. *Obstet Gynecol* 1999;93(3):392-5.
834. Silverman M, Sheffer A, Diaz PV, Lindmark B, Radner F, Broddene M, et al. Outcome of pregnancy in a randomized controlled study of patients with asthma exposed to budesonide. *Annals of Allergy, Asthma, & Immunology* 2005;95(6):566-70.
835. Christensson C, Thoren A, Lindberg B. Safety of inhaled budesonide: clinical manifestations of systemic corticosteroid-related adverse effects. *Drug Safety* 2008;31(11):965-88.
836. Lim A, Stewart K, Konig K, George J. Systematic review of the safety of regular preventive asthma medications during pregnancy. *Annals of Pharmacotherapy* 2011;45(7-8):931-45.
837. Breton MC, Beauchesne MF, Lemire C, Rey E, Forget A, Blais L. Risk of perinatal mortality associated with inhaled corticosteroid use for the treatment of asthma during pregnancy. *Journal of Allergy and Clinical Immunology* 2010;126(4):772-7.e2.
838. Lin S, Munsie JPW, Herdt-Losavio ML, Druschel CM, Campbell K, Browne ML, et al. Maternal asthma medication use and the risk of selected birth defects. *Pediatrics* 2012;129(2):e317-24.
839. Rahimi R, Nikfar S, Abdollahi M. Meta-analysis finds use of inhaled corticosteroids during pregnancy safe: a systematic meta-analysis review. *Human & Experimental Toxicology* 2006;25(8):447-52.
840. Stenius-Aarniala B, Riikonen S, Teramo K. Slow-release theophylline in pregnant asthmatics. *Chest* 1995;107(3):642-7.
841. Schatz M. Asthma during pregnancy: interrelationships and management. *Ann Allergy* 1992;68(2):123-33.
842. Czeizel AE, Rockenbauer M. Population-based case-control study of teratogenic potential of corticosteroids. *Teratology* 1997;56(5):335-40.
843. Källén B. Maternal drug use and infant cleft lip/palate with special reference to corticoids. *Cleft Palate Craniofac J* 2003;40(6):624-8.
844. Rodriguez-Pinilla E, Martinez-Frias ML. Corticosteroids during pregnancy and oral clefts: a case-control study. *Teratology* 1998;58(1):2-5.
845. Park-Wyllie L, Mazzotta P, Pastuszak A, Moretti ME, Beique L, Hunnisett L, et al. Birth defects after maternal exposure to corticosteroids: prospective cohort study and meta-analysis of epidemiological studies. *Teratology* 2000;62(6):385-92.
846. Carmichael SL, Shaw GM, Ma C, Werler MM, Rasmussen SA, Lammer EJ. Maternal corticosteroid use and orofacial clefts. *Am J Obstet Gynecol* 2007;197(6):585 e1-7.
847. Bakhireva LN, Schatz M, Chambers CD. Effect of maternal asthma and gestational asthma therapy on fetal growth. *Journal of Asthma* 2007;44(2):71-6.
848. Bakhireva LN, Jones KL, Schatz M, Klonoff-Cohen HS, Johnson D, Slymen DJ, et al. Safety of leukotriene receptor antagonists in pregnancy. *J Allergy Clin Immunol* 2007;119(3):618-25.
849. Nelsen LM, Shields KE, Cunningham ML, Stoler JM, Bamshad MJ, Eng PM, et al. Congenital malformations among infants born to women receiving montelukast, inhaled corticosteroids, and other asthma medications. *Journal of Allergy & Clinical Immunology* 2012;129(1):251-4.e1-6.
850. Sarkar M, Koren G, Kalra S, Ying A, Smorlesi C, De Santis M, et al. Montelukast use during pregnancy: a multicentre, prospective, comparative study of infant outcomes. *European Journal of Clinical Pharmacology* 2009;65(12):1259-64.
851. Mabie W C, Barton J R, Wasserstrum N, Sibai B M. Clinical observations on asthma in pregnancy. *J Matern Fetal Med* 1992;1(1):45-50.
852. Lao TT, Huengsborg M. Labour and delivery in mothers with asthma. *Eur J Obstet Gynecol Reprod Biol* 1990;35(2-3):183-90.
853. Arad I, Landau H. Adrenocortical reserve of neonates born of long-term, steroid-treated mothers. *Eur J Pediatr* 1984;142(4):279-80.
854. Turner ES, Greenberger PA, Patterson R. Management of the pregnant asthmatic patient. *Ann Intern Med* 1980;93(6):905-18.
855. Ost L, Wettrell G, Bjorkhem I, Rane A. Prednisolone excretion in human milk. *J Pediatr* 1985;106(6):1008-11.
856. McKenzie SA, Selley JA, Agnew JE. Secretion of prednisolone into breast milk. *Arch Dis Child* 1975;50(11):894-6.
857. Greenberger PA, Odeh YK, Frederiksen MC, Atkinson AJ Jr. Pharmacokinetics of prednisolone transfer to breast milk. *Clin Pharmacol Ther* 1993;53(3):324-8.

858. Meredith S, Nordman H. Occupational asthma: measures of frequency from four countries. *Thorax* 1996;51(4):435-40.
859. Blanc PD, Toren K. How much adult asthma can be attributed to occupational factors? *Am J Med* 1999;107(6):580-7.
860. Balmes J, Becklake M, Blanc P, Henneberger P, Kreiss K, Mapp C, et al. American Thoracic Society Statement: Occupational contribution to the burden of airway disease. *Am J Respir Crit Care Med* 2003;167(5):787-97.
861. Ross DJ. Ten years of the SWORD project. Surveillance of Work-related and Occupational Respiratory Disease. *Clin Exp Allergy* 1999;29(6):750-3.
862. Hendrick DJ. Occupational disorders of the lung: recognition, management and prevention. Gulf Professional Publishing; 2002.
863. Banks DE, Wang ML. Occupational asthma: "the big picture". *Occup Med* 2000;15(2):335-58.
864. Ameille J, Pauli G, Calastrenge-Crinquand A, Vervloet D, Iwatsubo Y, Popin E, et al. Reported incidence of occupational asthma in France, 1996-99: the ONAP programme. *Occup Environ Med* 2003;60(2):136-41.
865. Brhel P. Occupational respiratory diseases in the Czech Republic. *Ind Health* 2003;41(2):121-3.
866. Cortona G, Pisati G, Dellabianca A, Moscato G. Respiratory occupational allergies: the experience of the Hospital Operative Unit of Occupational Medicine in Lombardy from 1990 to 1998 [Italian]. *G Ital Med Lav Ergon* 2001;23(1):64-70.
867. Gannon PF, Burge PS. The SHIELD scheme in the West Midlands Region, United Kingdom. Midland Thoracic Society Research Group. *Br J Ind Med* 1993;50(9):791-6.
868. Hnizdo E, Esterhuizen TM, Rees D, Laloo UG. Occupational asthma as identified by the Surveillance of Work-related and Occupational Respiratory Diseases programme in South Africa. *Clin Exp Allergy* 2001;31(1):32-9.
869. McDonald JC, Keynes HL, Meredith SK. Reported incidence of occupational asthma in the United Kingdom, 1989-97. *Occup Environ Med* 2000;57(12):823-9.
870. Meyer JD, Holt DL, Cherry NM, McDonald JC. SWORD '98: surveillance of work-related and occupational respiratory disease in the UK. *Occup Med (Oxf)* 1999;49(8):485-9.
871. Sallie BA, Ross DJ, Meredith SK, McDonald JC. SWORD '93. Surveillance of work-related and occupational respiratory disease in the UK. *Occup Med (Oxf)* 1994;44(4):177-82.
872. Toren K, Jarvholm B, Brisman J, Hagberg S, Hermansson BA, Lillienberg L. Adult-onset asthma and occupational exposures. *Scand J Work Environ Health* 1999;25(5):430-5.
873. Meredith SK, Taylor VM, McDonald JC. Occupational respiratory disease in the United Kingdom 1989: a report to the British Thoracic Society and the Society of Occupational Medicine by the SWORD project group. *Br J Ind Med* 1991;48(5):292-8.
874. Karjalainen A, Kurppa K, Martikainen R, Karjalainen J, Klaukka T. Exploration of asthma risk by occupation--extended analysis of an incidence study of the Finnish population. *Scand J Work Environ Health* 2002;28(1):49-57.
875. Reijula K, Haahtela T, Klaukka T, Rantanen J. Incidence of occupational asthma and persistent asthma in young adults has increased in Finland. *Chest* 1996;110(1):58-61.
876. Jaakkola JJ, Piipari R, Jaakkola MS. Occupation and asthma: a population-based incident case-control study. *Am J Epidemiol* 2003;158(10):981-7.
877. Johnson AR, Dimich-Ward HD, Manfreda J, Becklake MR, Ernst P, Sears MR, et al. Occupational asthma in adults in six Canadian communities. *Am J Respir Crit Care Med* 2000;162(6):2058-62.
878. Kogevinas M, Anto JM, Soriano JB, Tobias A, Burney P. The risk of asthma attributable to occupational exposures. A population-based study in Spain. Spanish Group of the European Asthma Study. *Am J Respir Crit Care Med* 1996;154(1):137-43.
879. Kogevinas M, Anto JM, Sunyer J, Tobias A, Kromhout H, Burney P. Occupational asthma in Europe and other industrialised areas: a population-based study. European Community Respiratory Health Survey Study Group. *Lancet* 1999;353(9166):1750-4.
880. Fishwick D, Barber CM, Bradshaw LM, Harris-Roberts J, Francis M, Naylor S, et al. Standards of care for occupational asthma. *Thorax* 2008;63(3):240-50.
881. Nicholson P J, Cullinan P, Burge P S, Boyle C. Occupational asthma: Prevention, identification & management: Systematic review & recommendations. London: British Occupational Health Research Foundation; 2010. [cited 08 Jul 2014] Available from <http://www.bohrf.org.uk/projects/asthma.html>
882. Chiry S, Boulet L-P, Lepage J, Forget A, Begin D, Chaboillez S, et al. Frequency of work-related respiratory symptoms in workers without asthma. *American Journal of Industrial Medicine* 2009;52(6):447-54.
883. Burge PS, Pantin CF, Newton DT, Gannon PF, Bright P, Belcher J, et al. Development of an expert system for the interpretation of serial peak expiratory flow measurements in the diagnosis of occupational asthma. Midlands Thoracic Society Research Group. *Occup Environ Med* 1999;56(11):758-64.
884. Bright P, Newton DT, Gannon PF, Pantin CF, Burge PS. OASYS-3: improved analysis of serial peak expiratory flow in suspected occupational asthma. *Monaldi Arch Chest Dis* 2001;56(3):281-8.
885. Burge PS. Occupational asthma in electronics workers caused by colophony fumes: follow-up of affected workers. *Thorax* 1982;37(5):348-53.
886. Cote J, Kennedy S, Chan-Yeung M. Sensitivity and specificity of PC20 and peak expiratory flow rate in cedar asthma. *J Allergy Clin Immunol* 1990;85(3):592-8.
887. Leroyer C, Perfetti L, Trudeau C, L'Archeveque J, Chan-Yeung M, Malo JL. Comparison of serial monitoring of peak expiratory flow and FEV₁ in the diagnosis of occupational asthma. *Am J Respir Crit Care Med* 1998;158(3):827-32.

888. Liss GM, Tarlo SM. Peak expiratory flow rates in possible occupational asthma. *Chest* 1991;100(1):63-9.
889. Malo JL, Cote J, Cartier A, Boulet LP, L'Archeveque J, Chan-Yeung M. How many times per day should peak expiratory flow rates be assessed when investigating occupational asthma? *Thorax* 1993;48(12):1211-7.
890. Moore VC, Jaakkola MS, Burge PS. A Systematic Review of Serial Peak Expiratory Flow Measurements in the Diagnosis of Occupational Asthma. *Ann Respir Med* 2009;1(1):31-44.
891. Malo JL, Ghezzi H, L'Archeveque J, Lagier F, Perrin B, Cartier A. Is the clinical history a satisfactory means of diagnosing occupational asthma? *Am Rev Respir Dis* 1991;143(3):528-32.
892. Anees W, Gannon P F, Huggins V, Pantin C F, Burge P S. Effect of peak expiratory flow data quantity on diagnostic sensitivity and specificity in occupational asthma. *Eur Respir J* 2004;23(5):730-4.
893. Moore VC, Jaakkola MS, Burge CB, Pantin CF, Robertson AS, Vellore AD, et al. PEF analysis requiring shorter records for occupational asthma diagnosis. *Occupational Medicine (Oxford)* 2009;59(6):413-7.
894. Burge CBSG, Moore VC, Pantin CFA, Robertson AS, Burge PS. Diagnosis of occupational asthma from time point differences in serial PEF measurements. *Thorax* 2009;64(12):1032-6.
895. Malo JL, Cardinal S, Ghezzi H, L'Archeveque J, Castellanos L, Maghni K. Association of bronchial reactivity to occupational agents with methacholine reactivity, sputum cells and immunoglobulin E-mediated reactivity. *Clinical & Experimental Allergy* 2011;41(4):497-504.
896. Vandenplas O, Suojalehto H, Aasen TB, Baur X, Burge PS, de Blay F, et al. Specific inhalation challenge in the diagnosis of occupational asthma: consensus statement. *Eur Respir J* 2014;43(6):1573-87.
897. D'Alpaos V, Vandenplas O, Evrard G, Jamart J. Inhalation challenges with occupational agents: threshold duration of exposure. *Respir Med* 2013;107(5):739-44.
898. Stenton SC, Avery AJ, Walters EH, Hendrick DJ. Statistical approaches to the identification of late asthmatic reactions. *Eur Respir J* 1994;7(4):806-12.
899. Vandenplas O, D'Alpaos V, Heymans J, Jamart J, Thimpont J, Huaux F, et al. Sputum eosinophilia: an early marker of bronchial response to occupational agents. *Allergy* 2009;64(5):754-61.
900. Munoz X, Velasco MI, Culebras M, Roca O, Morell F, Cruz MJ. Utility of exhaled breath condensate pH for diagnosing occupational asthma. *International archives of allergy and immunology* 2012;159(3):313-20.
901. Sanchez-Vidaurre S, Cruz MJ, Gomez-Olles S, Morell F, Munoz X. Diagnostic utility of exhaled breath condensate analysis in conjunction with specific inhalation challenge in individuals with suspected work-related asthma. *Ann Allergy Asthma Immunol* 2012;108(3):151-6.
902. Chan-Yeung M, Lam S, Koener S. Clinical features and natural history of occupational asthma due to western red cedar (*Thuja plicata*). *Am J Med* 1982;72(3):411-5.
903. Merget R, Schulte A, Gebler A, Breitstadt R, Kulzer R, Berndt ED, et al. Outcome of occupational asthma due to platinum salts after transferral to low-exposure areas. *Int Arch Occup Environ Health* 1999;72(1):33-9.
904. Moscato G, Dellabianca A, Perfetti L, Brame B, Galdi E, Niniano R, et al. Occupational asthma: a longitudinal study on the clinical and socioeconomic outcome after diagnosis. *Chest* 1999;115(1):249-56.
905. Pisati G, Baruffini A, Zedda S. Toluene diisocyanate induced asthma: outcome according to persistence or cessation of exposure. *Br J Ind Med* 1993;50(1):60-4.
906. Rosenberg N, Garnier R, Rousselin X, Mertz R, Gervais P. Clinical and socio-professional fate of isocyanate-induced asthma. *Clin Allergy* 1987;17(1):55-61.
907. Tarlo SM, Banks D, Liss G, Broder I. Outcome determinants for isocyanate induced occupational asthma among compensation claimants. *Occup Environ Med* 1997;54(10):756-61.
908. Valentino M, Pizzichini MA, Monaco F, Governa M. Latex-induced asthma in four healthcare workers in a regional hospital. *Occup Med (Lond)* 1994;44(3):161-4.
909. Valentino M, Rapisarda V. Course of isocyanate-induced asthma in relation to exposure cessation: longitudinal study of 50 subjects [Italian]. *G Ital Med Lav Ergon* 2002;24(1):26-31.
910. Vandenplas O, Delwiche JP, Depelchin S, Sibille Y, Vande Weyer R, Delaunois L. Latex gloves with a lower protein content reduce bronchial reactions in subjects with occupational asthma caused by latex. *Am J Respir Crit Care Med* 1995;151(3 Pt 1):887-91.
911. Chan-Yeung M, MacLean L, Paggiaro PL. Follow-up study of 232 patients with occupational asthma caused by western red cedar (*Thuja plicata*). *J Allergy Clin Immunol* 1987;79(5):792-6.
912. Malo JL, Cartier A, Ghezzi H, Lafrance M, McCants M, Lehrer SB. Patterns of improvement in spirometry, bronchial hyperresponsiveness, and specific IgE antibody levels after cessation of exposure in occupational asthma caused by snow-crab processing. *Am Rev Respir Dis* 1988;138(4):807-12.
913. Gannon PF, Weir DC, Robertson AS, Burge PS. Health, employment, and financial outcomes in workers with occupational asthma. *Brit J Ind Med* 1993;50(6):491-6.
914. Axon EJ, Beach JR, Burge PS. A comparison of some of the characteristics of patients with occupational and non-occupational asthma. *Occup Med (Lond)* 1995;45(2):109-11.
915. Cannon J, Cullinan P, Newman Taylor A. Consequences of occupational asthma. *British Medical Journal* 1995;311(7005):602-3.
916. Larbanois A, Jamart J, Delwiche JP, Vandenplas O. Socioeconomic outcome of subjects experiencing asthma symptoms at work. *Eur Respir J* 2002;19(6):1107-13.

917. Ross DJ, McDonald JC. Health and employment after a diagnosis of occupational asthma: a descriptive study. *Occup Med (Lond)* 1998;48(4):219-25.
918. Ameille J, Pairon JC, Bayeux MC, Brochard P, Choudat D, Conso F, et al. Consequences of occupational asthma on employment and financial status: a follow-up study. *Eur Respir J* 1997;10(1):55-8.
919. Marabini A, Dimich-Ward H, Kwan SY, Kennedy SM, Waxler-Morrison N, Chan-Yeung M. Clinical and socioeconomic features of subjects with red cedar asthma. A follow-up study. *Chest* 1993;104(3):821-4.
920. Vandenplas O, Jamart J, Delwiche JP, Evrard G, Larbanois A. Occupational asthma caused by natural rubber latex: outcome according to cessation or reduction of exposure. *J Allergy Clin Immunol* 2002;109(1):125-30.
921. Venables KM, Davison AG, Newman Taylor AJ. Consequences of occupational asthma. *Respir Med* 1989;83(5):437-40.
922. Rotter T, Kinsman L, James E L, Machotta A, Gothe H, Willis J, et al. Clinical pathways: effects on professional practice, patient outcomes, length of stay and hospital costs (Cochrane Review). In: *The Cochrane Library*, 2010.
923. Smith JR, Noble MJ, Musgrave S, Murdoch J, Price GM, Barton GR, et al. The at-risk registers in severe asthma (ARRISA) study: A cluster-randomised controlled trial examining effectiveness and costs in primary care. *Thorax* 2012;67(12):1052-60.
924. Mitchell EA, Didsbury PB, Kruithof N, Robinson E, Milmine M, Barry M, et al. A randomized controlled trial of an asthma clinical pathway for children in general practice. *Acta Paediatrica* 2005;94(2):226-33.
925. Doherty SR, Jones PD. Use of an 'evidence-based implementation' strategy to implement evidence-based care of asthma into rural district hospital emergency departments. *Rural Remote Health* 2006;6(1):529.
926. Johnson KB, Blaisdell CJ, Walker A, Eggleston P. Effectiveness of a clinical pathway for inpatient asthma management. *Pediatrics* 2000;106(5):1006-12.
927. Zorc JJ, Chew A, Allen JL, Shaw K. Beliefs and barriers to follow-up after an emergency department asthma visit: a randomized trial. *Pediatrics* 2009;124(4):1135-42.
928. O'Brien MA, Rogers S, Jamtvedt G, Oxman AD, Odgaard-Jensen J, Kristoffersen DT, et al. Educational outreach visits: effects on professional practice and health care outcomes (Cochrane Review). In: *The Cochrane Library*, 2007.
929. Forsetlund L, Bjørndal A, Rashidian A, Jamtvedt G, O'Brien MA, Wolf F, et al. Continuing education meetings and workshops: effects on professional practice and health care outcomes (Cochrane Review). In: *The Cochrane Library*, 2009.
930. Cabana MD, Slish KK, Evans D, Mellins RB, Brown RW, Lin X, et al. Impact of physician asthma care education on patient outcomes. *Pediatrics* 2006;117(6):2149-57.
931. Shah S, Sawyer SM, Toelle BG, Mellis CM, Peat JK, Lagleva M, et al. Improving paediatric asthma outcomes in primary health care: A randomised controlled trial. *Medical Journal of Australia* 2011;195(7):405-9.
932. Lozano P, Finkelstein JA, Carey VJ, Wagner EH, Inui TS, Fuhlbrigge AL, et al. A multisite randomized trial of the effects of physician education and organizational change in chronic-asthma care: health outcomes of the Pediatric Asthma Care Patient Outcomes Research Team II Study. *Archives of Pediatrics & Adolescent Medicine* 2004;158(9):875-83.
933. Smelee IJ, Grol RP, van Schayck CP, van den Bosch WJ, van den Hoogen HJ, Muris JW. Can small group education and peer review improve care for patients with asthma/chronic obstructive pulmonary disease? *Qual Health Care* 1999;8(2):92-8.
934. Witt K, Knudsen E, Ditlevsen S, Holtnagel H. Academic detailing has no effect on prescribing of asthma medication in Danish general practice: a 3-year randomized controlled trial with 12-monthly follow-ups. *Family Practice* 2004;21(3):248-53.
935. Liaw ST, Sulaiman ND, Barton CA, Chondros P, Harris CA, Sawyer S, et al. An interactive workshop plus locally adapted guidelines can improve general practitioners asthma management and knowledge: a cluster randomised trial in the Australian setting. *BMC Family Practice* 2008;9:22.
936. Goeman DP, Sanci LA, Scharf SL, Bailey M, O'Hehir RE, Jenkins CR, et al. Improving general practice consultations for older people with asthma: a cluster randomised control trial. *Medical Journal of Australia* 2009;191(2):113-7.
937. Stout JW, Smith K, Zhou C, Solomon C, Dozor AJ, Garrison MM, et al. Learning from a distance: Effectiveness of online spirometry training in improving asthma care. *Academic pediatrics* 2012;12(2):88-95.
938. Charlton I, Charlton G, Broomfield J, Mullee MA. Audit of the effect of a nurse run asthma clinic on workload and patient morbidity in a general practice. *Br J Gen Pract* 1991;41(347):227-31.
939. Hoskins G, Neville RG, Smith B, Clark RA. The link between nurse training and asthma outcomes. *Br J Comm Nursing* 1999;4(5):222-8.
940. Heard AR, Richards IJ, Alpers JH, Pilotto LS, Smith BJ, Black JA. Randomised controlled trial of general practice based asthma clinics. *Med J Aust* 1999;171(2):68-71.
941. Feder G, Griffiths C, Highton C, Eldridge S, Spina M, Southgate L. Do clinical guidelines introduced with practice based education improve care of asthmatic and diabetic patients? A randomised controlled trial in general practitioners in east London. *British Medical Journal* 1995;311(7018):1473-8.
942. Bryce FP, Neville RG, Crombie IK, Clark RA, McKenzie P. Controlled trial of an audit facilitator in diagnosis and treatment of childhood asthma in general practice. *British Medical Journal* 1995;310(6983):838-42.

943. Dickinson J, Hutton S, Atkin A, Jones K. Reducing asthma morbidity in the community: the effect of a targeted nurse-run asthma clinic in an English general practice. *Respir Med* 1997;91(10):634-40.
944. Lindberg M, Ahlner J, Moller M, Ekstrom T. Asthma nurse practice - a resource-effective approach in asthma management. *Respir Med* 1999;93(8):584-8.
945. Baishnab E, Karner C. Primary care based clinics for asthma (Cochrane Review). In: *The Cochrane Library*, 2012.
946. Tran N, Coffman JM, Sumino K, Cabana MD. Patient reminder systems and asthma medication adherence: a systematic review. *Journal of Asthma* 2014;51(5):536-43.
947. McPherson AC, Glazebrook C, Forster D, James C, Smyth A. A randomized, controlled trial of an interactive educational computer package for children with asthma. *Pediatrics* 2006;117(4):1046-54.
948. Hieftje K, Edelman EJ, Camenga DR, Fiellin LE. Electronic media-based health interventions promoting behavior change in youth: a systematic review. *JAMA Pediatr* 2013;167(6):574-80.
949. Joseph CLM, Ownby DR, Havstad SL, Saltzgaber J, Considine S, Johnson D, et al. Evaluation of a Web-Based Asthma Management Intervention Program for Urban Teenagers: Reaching the Hard to Reach. *Journal of Adolescent Health* 2013;52(4):419-26.
950. Pare G, Moqadem K, Pineau G, St-Hilaire C. Clinical effects of home telemonitoring in the context of diabetes, asthma, heart failure and hypertension: a systematic review. *Journal of Medical Internet Research* 2010;12(2):e21.
951. Marcano Belisario JS, Huckvale K, Greenfield G, Car J, Gunn LH. Smartphone and tablet self management apps for asthma. *Cochrane Database of Systematic Reviews* 2013;11:CD010013.
952. Gustafson D, Wise M, Bhattacharya A, Pulvermacher A, Shanovich K, Phillips B, et al. The effects of combining web-based eHealth with telephone nurse case management for pediatric asthma control: a randomized controlled trial. *Journal of Medical Internet Research* 2012;14(4):e101.
953. Ryan D, Price D, Musgrave SD, Malhotra S, Lee AJ, Ayansina D, et al. Clinical and cost effectiveness of mobile phone supported self monitoring of asthma: multicentre randomised controlled trial. *BMJ* 2012;344:e1756.
954. van der Meer V, Bakker MJ, van den Hout WB, Rabe KF, Sterk PJ, Kievit J, et al. Internet-based self-management plus education compared with usual care in asthma: a randomized trial. *Annals of Internal Medicine* 2009;151(2):110-20.
955. van der Meer V, van Stel HF, Bakker MJ, Roldaan AC, Assendelft WJ, Sterk PJ, et al. Weekly self-monitoring and treatment adjustment benefit patients with partly controlled and uncontrolled asthma: an analysis of the SMASHING study. *Respiratory Research* 2010;11:74.
956. DiBello K, Boyar K, Abrenica S, Worral PS. The effectiveness of text messaging programs on adherence to treatment regimens among adults aged 18 to 45 years diagnosed with asthma: a systematic review (Provisional abstract). *Database of Abstracts of Reviews of Effects* 2014(1):485-532.
957. Morrison D, Wyke S, Agur K, Cameron EJ, Docking RI, Mackenzie AM, et al. Digital asthma self-management interventions: a systematic review. *Journal of Medical Internet Research* 2014;16(2):e51.
958. de Jong CC, Ros WJ, Schrijvers G. The effects on health behavior and health outcomes of Internet-based asynchronous communication between health providers and patients with a chronic condition: a systematic review. *Journal of Medical Internet Research* 2014;16(1):e19.
959. Deshpande A, Khoja S, Lorca J, McKibbin A, Rizo C, Huserau D, et al. Asynchronous telehealth: a scoping review of analytic studies. *Open Med* 2009;3(2):e69-91.
960. Garbutt JM, Banister C, Highstein G, Sterkel R, Epstein J, Bruns J, et al. Telephone coaching for parents of children with asthma: impact and lessons learned. *Archives of Pediatrics & Adolescent Medicine* 2010;164(7):625-30.
961. Pinnock H, Bawden R, Proctor S, Wolfe S, Scullion J, Price D, et al. Accessibility, acceptability, and effectiveness in primary care of routine telephone review of asthma: pragmatic, randomised controlled trial. *British Medical Journal* 2003;326(7387):477-9.
962. Matui P, Wyatt JC, Pinnock H, Sheikh A, McLean S. Computer decision support systems for asthma: a systematic review. *NPJ Prim Care Respir Med* 2014;24:14005.
963. Fathima M, Peiris D, Naik-Panvelkar P, Saini B, Armour CL. Effectiveness of computerized clinical decision support systems for asthma and chronic obstructive pulmonary disease in primary care: a systematic review. *BMC Pulmonary Medicine* 2014;14:189.
964. Clark NM, Shah S, Dodge JA, Thomas LJ, Andridge RR, Little RJA. An evaluation of asthma interventions for preteen students. *Journal of School Health* 2010;80(2):80-7.
965. Halterman JS, Szilagyi PG, Fisher SG, Fagnano M, Tremblay P, Conn KM, et al. Randomized controlled trial to improve care for urban children with asthma: results of the school-based asthma therapy trial. *Archives of Pediatrics & Adolescent Medicine* 2011;165(3):262-8.
966. Bruzzese JM, Sheares BJ, Vincent EJ, Du Y, Sadeghi H, Levison MJ, et al. Effects of a school-based intervention for urban adolescents with asthma: a controlled trial. *American Journal of Respiratory & Critical Care Medicine* 2011;183(8):998-1006.
967. Foster G, Taylor SJC, Eldridge S, Ramsay J, Griffiths CJ. Self-management education programmes by lay leaders for people with chronic conditions (Cochrane Review). In: *The Cochrane Library*, (4) 2007.

968. Pande S, Hiller JE, Nkansah N, Bero L. The effect of pharmacist-provided non-dispensing services on patient outcomes, health service utilisation and costs in low- and middle-income countries (Cochrane Review). In: The Cochrane Library, 2013.
969. Benavides S, Rodriguez JC, Maniscalco-Feichtl M. Pharmacist involvement in improving asthma outcomes in various healthcare settings: 1997 to present. *Annals of Pharmacotherapy* 2009;43(1): 85-97.
970. Basheti IA, Armour CL, Bosnic-Anticevich SZ, Reddel HK. Evaluation of a novel educational strategy, including inhaler-based reminder labels, to improve asthma inhaler technique. *Patient Education & Counseling* 2008;72(1):26-33.
971. Hammerlein A, Muller U, Schulz M. Pharmacist-led intervention study to improve inhalation technique in asthma and COPD patients. *Journal of Evaluation in Clinical Practice* 2011;17(1):61-70.
972. Mehuys E, Van Bortel L, De Bolle L, Van Tongelen I, Annemans L, Remon JP, et al. Effectiveness of pharmacist intervention for asthma control improvement. *European Respiratory Journal* 2008;31(4):790-9.
973. Elliott RA, Barber N, Clifford S, Horne R, Hartley E. The cost effectiveness of a telephone-based pharmacy advisory service to improve adherence to newly prescribed medicines. *Pharmacy World & Science* 2008;30(1):17-23.
974. Bereznicki BJ, Peterson G, Jackson S, Walters EH, George J, Stewart K, et al. Uptake and effectiveness of a community pharmacy intervention programme to improve asthma management. *Journal of Clinical Pharmacy and Therapeutics* 2013;38(3):212-8.



Healthcare Improvement Scotland

Edinburgh Office

Gyle Square
1 South Gyle Crescent
Edinburgh
EH12 9EB
0131 623 4300

Glasgow Office

Delta House
50 West Nile Street
Glasgow
G1 2NP
0141 225 6999