

Can open prescribing data detect medicine effects on tuberculosis? An ecological study of primary care and hospital prescribing and TB notifications in England

Draft 4 (revised after peer review, round 2), 14 September 2026

Abstract

Background. - Corticosteroids, biologics and other immunosuppressants increase individual risk of tuberculosis (TB). - Openly published English prescribing and TB surveillance data could, in principle, be linked to study these relationships at population level. - We assessed whether such ecological analyses can detect associations between medicines and TB, and why they might not.

Methods. - Exposures. - Primary care prescribing, 2011–2024: practice-level data apportioned to local authorities by where registered patients live. Drug groups were defined at presentation level (systemic oral glucocorticoids, conventional DMARDs, transplant immunosuppressants and others), including a negative-control exposure. - Hospital medicines, 2019–2024: NHS Secondary Care Medicines Data in WHO defined daily doses, apportioned by trust catchment. Controls were a disease-specific positive control (active-TB treatment) and negative-control exposures. - **Outcomes.** Annual UKHSA TB notifications for 294 lower-tier and 151 upper-tier authorities, and by region, place of birth and age. - **Analysis.** - Poisson panel models with area and year fixed effects, exposure in the previous year, and falsification (lead) tests. - Regional models with randomisation inference. - Minimum detectable effects (MDE), compared with expected population effects derived from published relative risks and UKHSA-recorded drug-associated TB. - A permutation-based power simulation on the real notification counts.

Results. - Linkage. Hospital active-TB treatment tracked TB notifications between areas (Spearman $\rho = 0.81$) and within areas in the same year (incidence rate ratio [IRR] 1.035, 95% CI 1.009–1.061, per 10% increase). The within-area elasticity was only 0.36–0.40, indicating substantial attenuation even for a disease-specific drug. - **Primary care.** No drug group was associated with notifications in the following year. For systemic oral glucocorticoids the IRR was 0.969 (0.929–1.012) per 10% within-area increase. - **Hospital medicines.** TNF inhibitors (0.989, 0.972–1.007), JAK inhibitors and other immunosuppressants were null. Hospital systemic glucocorticoids had a small nominal association (1.024, 1.001–1.046) that was not robust to clustering by trust or to excluding COVID-19 years. - **Falsification and negative control.** The negative-control exposure showed inverse estimates similar in size to metformin, and following-year prescribing was inversely associated with notifications for several groups. With collinear adjacent-year exposures these tests are weak, but the pattern is compatible with residual confounding by area-specific trends. - **Power.** - For oral glucocorticoids, the MDE (6.3% per 10% increase) was about 18 times the effect expected from published relative risks (0.34%), and about 70–140 times that implied by UKHSA-recorded steroid-associated TB (0.04–0.09%). - In simulations on the real counts, the null hypothesis was rejected for the published effect in 8.3% of replicates, no more often than with no effect (7.8%); simulated power exceeded analytic power at the real standard error. - **UK-born TB.** A regional association between prednisolone dose and UK-born TB was implausibly large, not significant by randomisation inference, dependent on London, and accompanied by failed negative-control and falsification tests.

Conclusions. - Hospital medicines data could be linked to TB notifications only with heavy attenuation, and no working positive control was available for primary care. Ecological analyses of these data cannot detect plausible population-level effects of medicines on TB. - Apparent associations are compatible with residual confounding, chance or measurement error, not with drug effects of plausible size. - Causal questions require individual-level linked data analysed with target trial emulation. Surveillance of drug-associated TB is better served by more complete recording of immunosuppression in national TB surveillance.

Introduction

Tuberculosis (TB) remains a public health problem in England.

Recent trends. - **Decline:** notifications fell by 44% between 2011 and 2018 (8,282 to 4,609). For 2011–2015, most of the fall reflected declining TB rates in almost all populations, with smaller contributions from fewer recent non-EU migrants and from pre-entry screening of long-stay visa applicants from high-incidence countries [Thomas 2018; Aldridge 2016]. - **2020:** notifications dropped to 4,125 during the COVID-19 pandemic, a fall thought to reflect service disruption rather than less disease [UKHSA 2021; Morrison 2023]. - **2021–2024 rise:** notifications then increased, reaching 5,490 in 2024 (9.4 per 100,000). - 81.9% were in people born outside the UK [UKHSA 2025, Supplementary Table 12]. - 41% of those were notified within five years of arrival. - The rise has been attributed mainly to migration from higher-incidence countries [UKHSA 2025]. - **2025:** national data show 5,424 notifications, with 2024 revised to 5,487 [UKHSA 2026].

TB in UK-born people. UK-born notifications fell by 48% between 2014 and 2022 (1,756 to 916), then rose to 995 in 2024, still 43% below 2014 [UKHSA 2025]. UK-born TB is heterogeneous: it includes reactivation in older adults and transmission linked to social risk factors [Davidson 2018].

Medicines that change TB risk. Host factors also modify TB risk, and several are shaped by commonly prescribed medicines. - **Oral glucocorticoids:** current use is associated with an odds ratio of about 5 for active TB, higher at doses ≥ 15 mg/day [Jick 2006]. - **Inhaled corticosteroids and conventional DMARDs:** smaller increases [Brassard 2011; Castellana 2019; Brassard 2006]. - **Anti-TNF biologics:** TB typically presents within months of starting treatment [Keane 2001]. Latent infection screening before treatment greatly reduces this risk [BTS 2005; Carmona 2005]. - **Proton pump inhibitors:** associations with increased risk [Song 2019]. - **Statins and metformin:** associations with lower risk [Li 2020; Zhang 2020].

Many of these estimates come from high-incidence settings or designs vulnerable to confounding by indication and protopathic bias.

Open data and their problems. England publishes monthly prescribing for every general practice, hospital medicines use by NHS trust, and TB notifications by local authority. Linking these offers a low-cost way to generate or test hypotheses about medicines and TB. Such ecological analyses face well-known problems: - confounding by area characteristics; - cross-level bias, which covariate adjustment cannot remove when effects differ between groups [Greenland 1989; Morgenstern 1995]; - limited power, which depends on the number of areas [Sheppard 1996].

The data carry specific problems too: - practice lists are inflated in ways that track population mobility [Hudson 2025; Venkatesan 2025]; - practice locations do not match where patients live; - prescribing is patterned by deprivation and was disrupted by COVID-19 [Richards 2025]; - hospital-prescribed medicines are recorded separately from primary care [Bacon 2020].

The gap. We found no previous study linking area-level prescribing to TB. In analogous settings, drugs with large individual effects produced weak population signals [Hermans 2015], and between-area and within-area associations had opposite signs [Költringer 2023].

This study. We asked whether open English prescribing data can detect associations between medicines and TB notifications at population level, and why they might not. We combined: - cross-sectional and fixed-effects panel designs at two geographic scales; - primary care prescribing apportioned to where registered patients live; - hospital medicines apportioned by trust catchment, with a disease-specific positive control; - negative-control exposures and falsification tests [Lipsitch 2010; Prasad 2013]; - regional analyses of TB in UK-born and older people.

We compared the smallest effects these designs could detect, both analytically and by simulation, with the population effects implied by published individual-level relative risks and by UKHSA surveillance records of drug-associated TB.

Methods

Design and reporting

This study uses aggregate, openly published data in three linked ecological designs:

- a cross-sectional analysis;
- a longitudinal panel analysis of local authorities with area and year fixed effects;
- regional analyses by place of birth and age.

It was reported with reference to the RECORD and STROBE guidance. All data were public, aggregate and anonymised, so ethical approval was not required. Code, processed data and outputs are available at <https://github.com/drcjar/tb-prescribing-england>.

Pre-specification. The analysis developed in stages, and we report the chronology.

1. **First pass.** Before any results were seen, we specified the drug groups, including a negative-control exposure (levothyroxine) and a positive control (primary care antituberculosis prescribing), and the cross-sectional design. The first panel design was specified after the cross-sectional results but before any panel results; it used three-year rolling notifications and exposure in years $t-5$ to $t-3$. The pre-specified positive control failed: primary care antituberculosis prescribing did not track notifications, because TB treatment is delivered by specialist services. It was reclassified as descriptive.
2. **Later additions, made after those analyses gave null results:**
 - annual outcomes;
 - lower-tier authorities;
 - hospital medicines and their controls;
 - regional analyses by birthplace and age;
 - power simulation;
 - spatial diagnostics.
3. **Revisions after peer review:**
 - drug groups redefined;
 - prescribing apportioned by patient residence;
 - exposure in year $t-1$ as primary;
 - lag and lead estimated jointly;
 - randomisation inference;
 - additional covariates;
 - after the second round: correction of the GP practice restriction, redefinition of the hospital positive control and glucocorticoid exposure, joint models of previous, same and following years, and randomisation inference on the t statistic.

All analyses after the first pass are exploratory. Estimates from the original panel design are given in Table S8.

Assumed causal structure

Figure 1 sets out the assumed causal structure. It shows which factors are absorbed by area and year fixed effects, which are measured and adjusted for, and which remain unmeasured. The primary exposure is prescribing in the year before the outcome year, because same-year prescribing may be prescribing for undiagnosed TB (protopathic bias).




Figure 1: **Figure 1.** Assumed causal structure for area-level prescribing and TB notifications. Area fixed effects absorb stable area characteristics; year fixed effects absorb national shocks; time-varying measured factors are adjusted for; dashed nodes are unmeasured. Same-year prescribing for undiagnosed TB (protopathic bias) motivates the use of prescribing in the previous year.

TB notifications

TB notifications are cases reported to UKHSA surveillance: the Enhanced TB Surveillance system (ETS) until 2021, and the National TB Surveillance System (NTBS) since, into which data from 2018 onwards were migrated. Cases are culture-confirmed or clinically diagnosed, and are assigned to areas by residential postcode at notification.

We used four sources:

1. Annual counts by local authority, 2001-2024.

- *Source:* UKHSA TB regional reports 2024, supplementary tables.
- *Coverage:* upper-tier authorities (UTLAs) and lower-tier authorities (LTLAs) on April 2023 boundaries. UKHSA combines the City of London with Hackney and the Isles of

Scilly with Cornwall. We combined population, covariates and prescribing in the same way at both levels, giving 151 UTLAs and 294 LTLAs.

- *Consistency*: summed over three years, these counts matched the Fingertips three-year counts ($r = 0.996$). Local authority sums slightly exceed national totals (5,539 vs 5,490 in 2024), possibly reflecting different extract dates or residence-assignment rules.
 - *Details not given in the tables*: counts are by year of notification. How people without a residential postcode (for example, no fixed abode) are assigned is not described. Area-years with no notifications were retained.
2. **Region × place of birth × year, 2000–2024**, with Labour Force Survey denominators: *TB in England 2025*, Supplementary Table 12.
 3. **Region × place of birth × age group × year**: regional reports, Table 9.
 4. **National notifications with recorded immunosuppression by cause, 2024**: *TB in England 2025*, chapter 1.

The 2020 fall in notifications is thought to reflect service disruption rather than a true reduction in incidence.

Primary care prescribing

Sources. - **2010–2013**: HSCIC practice-level prescribing (PDPI), August 2010–December 2013.
- **2014–2024**: NHSBSA English Prescribing Dataset (EPD).

Both record items dispensed in the community from primary care prescriptions. We aggregated each month to practice level using identical drug-group rules. For the EPD this was done with server-side SQL.

Practices included. We restricted to standard GP practices: practices with NHS ODS prescribing setting RO76, plus practice codes absent from the current ODS file (which omits practices closed before 2017) that have the standard GP practice code format (a letter and five digits; this format matches 94% of RO76 codes and few codes in other settings). Without this addition, prescribing by practices that later closed would have been dropped unevenly over time: 3.8% of items in 2011, 1.9% in 2014 and none from 2016.

Drug groups. Groups were defined at BNF presentation level:

Group	Definition
Systemic oral glucocorticoids	Prednisolone, prednisone, methylprednisolone, deflazacort and dexamethasone. Injectables, hydrocortisone (mainly replacement therapy) and betamethasone soluble tablets excluded. Oral hydrocortisone and oral dexamethasone are also reported separately.
Inhaled corticosteroids	—
Conventional DMARDs	Rheumatology methotrexate (including subcutaneous), leflunomide, azathioprine, mercaptopurine.
Transplant immunosuppressants	Tacrolimus, ciclosporin, mycophenolate, sirolimus, everolimus.
Other groups	Proton pump inhibitors; statins; metformin; insulins; fluoroquinolones; all antibacterials; vitamin D.

Group	Definition
Descriptive only	Antituberculosis drugs: TB treatment in England is delivered by specialist services, so primary care prescribing of these drugs is not a valid positive control.
Negative control exposure	Levothyroxine.

Eye, ear, nose and skin preparations were excluded.

Exposure measures. - Items per 1,000 residents per year (primary). - Prednisolone-equivalent milligrams for systemic oral glucocorticoids: strength × quantity × potency, with dexamethasone 6.67, methylprednisolone 1.25 and deflazacort 0.83. - Average daily quantities (ADQ) for the groups where the EPD populates them.

Apportionment to areas. Each practice's annual prescribing was shared across local authority districts in proportion to where its registered patients lived, using NHS Digital counts of patients registered at each practice by LSOA: - April releases from 2014 to 2024 were used, each practice-year taking that practice's nearest release in time (so 2014 shares were applied to 2011–2013); - practices in no release (closed before April 2014) were assigned to the district of their postcode; - LSOAs were mapped to April 2023 districts.

Assigning each practice to the district containing its postcode was a sensitivity analysis. In 2024, 7.3% of registered patients lived outside the district of their practice.

Denominators. ONS mid-year resident population estimates.

Hospital medicines

Source and period. NHS Secondary Care Medicines Data (SCMD; NHSBSA), which gives trust × month × product quantities. We used January 2019 to December 2024.

Classification. Products were grouped by mechanism, with quantities converted to DDD-years using WHO defined daily doses (DDD), or documented maintenance doses where WHO gives none. The groups were:

- TNF inhibitors;
- IL-6 inhibitors and abatacept;
- JAK inhibitors used in rheumatology (ruxolitinib reported separately);
- rituximab;
- calcineurin and mTOR inhibitors;
- antiproliferatives;
- systemic glucocorticoids (prednisolone and methylprednisolone), measured in prednisolone-equivalent mg and excluding intra-articular, depot and topical forms. Dexamethasone and hydrocortisone, used mainly in oncology, as antiemetics, for COVID-19 [RECOVERY 2021] and as replacement therapy, were reported separately as a descriptive group.

Controls. - **Positive control:** active-TB treatment, measured as pyrazinamide defined daily doses (1.5 g) from all pyrazinamide-containing products, including the fixed-dose combinations Rifater and Voractiv, so that a day of intensive-phase treatment counts once whatever the formulation. - **Reported separately:** rifamycin and isoniazid products, which are also used for latent TB and, in the case of rifampicin, other infections. - **Negative-control exposures:** IL-17, IL-12/23 and $\alpha 4\beta 7$ inhibitors and anakinra (low TB risk), and levetiracetam.

Trust mergers. Trusts that merged during 2019–2024 were reassigned to their successors using NHS ODS records. With this, 98–100% of DDD in every group and year came from trusts with catchment data.

Apportionment to areas. Trust quantities were shared across local authorities using OHID acute trust catchment populations (2024, all admissions). Elective-admission catchments were a sensitivity analysis.

Covariates

Time-varying (panel analyses): - ONS mid-year population estimates by age (% aged ≥ 65 ; % aged 15–44); - international in-migration per 1,000 (ONS components of population change); - diagnosed HIV prevalence; - QOF diabetes prevalence; - the latent TB infection (LTBI) testing and treatment programme for new migrants, introduced from 2015/16 through primary care in high-incidence areas for recent entrants aged 16–35 from high-incidence countries [Loutet 2018; Berrocal-Almanza 2022]. We used an indicator of programme activity, mapped from CCGs to local authorities (active in 83 LTLAs). Area-level reporting ended in 2019/20, so the indicator was carried forward unchanged and disruption in 2020 was not modelled. Because the programme targets young recent entrants, an all-age area indicator is a weak control for it; - people receiving asylum support per 1,000 (Home Office; available from 2014, so used for outcome years from 2015).

Cross-sectional analyses: - Census 2021 percentage born outside the UK.

Statistical analysis

Panel analyses. - Model: Poisson pseudo-maximum-likelihood regression [Santos Silva 2006] of annual notifications, with area and year fixed effects, a log population offset and SEs clustered by area. - **Primary exposure:** log prescribing rate in year $t-1$. Drug-associated TB typically presents within months of exposure [Keane 2001], and same-year prescribing is affected by prescribing for undiagnosed TB (protopathic bias). - **Outcome years:** 2014–2024. - **Covariates:** the time-varying set above. Diabetes prevalence was not adjusted for in the metformin and insulin models. - **Effect measure:** incidence rate ratio (IRR) per 10% within-area increase in prescribing, with Benjamini–Hochberg false discovery rate (FDR) correction across the 13 non-control drug groups in the primary model, including the descriptive groups (oral hydrocortisone, oral dexamethasone, all antibacterials, vitamin D). Levothyroxine and antituberculosis drugs were excluded. - **Falsification:** prescribing in years $t-1$ and $t+1$ estimated jointly on a common sample, and again with year t added, with a Wald test of the difference between the $t-1$ and $t+1$ coefficients. A true effect should load on $t-1$. Prescribing in adjacent years is highly correlated within areas, so the two estimates are strongly negatively correlated and opposite signs can arise by chance; we treat these tests as weak diagnostics. - **Sensitivity analyses:** - no covariates; - a distributed lag (t , $t-1$, $t-2$); - the original three-year window ($t-3$ to $t-1$); - area-specific linear trends; - log population as a covariate instead of an offset (shared-denominator bias); - adjustment for the LTBI programme; - adjustment for asylum support; - excluding outcome years 2020–21 and exposure years 2020–21; - outcome years 2018–2024 only (NTBS-era data); - outcome years 2015–2024 only, so that all exposure comes from the EPD (splice sensitivity); - ADQ instead of items, for groups where the EPD populates ADQ; - practice-postcode apportionment; - UTLA geography. - **Spatial dependence:** Moran’s I of year-specific Pearson residuals (5-nearest-neighbour weights, 999 permutations). - **Missing data:** complete-case analysis. Area-years with missing covariates, or with zero exposure (log undefined), were excluded model by model; the numbers excluded are recorded in the result files. - **Bounds:** for each estimate, the largest population attributable fraction compatible with the upper 90% confidence limit and, for drugs expected to protect, the largest prevented fraction compatible with the lower limit, both under the model used for expected effects. As a sensitivity analysis, the upper limit was divided by the negative control’s estimate from the same model, as if its bias applied equally to every drug.

Hospital analyses. The same model was fitted for 2019–2024, with exposure in: - the same year (for the positive control, where treatment follows diagnosis); - the previous year (for candidate drugs); - the following year.

SEs were also clustered by each area's principal trust, because apportioned exposures share trust-level variation. The within-area elasticity of the positive control (log IRR per log unit of exposure) was used as an attenuation factor.

Regional analyses. - **Units:** 9 regions $\times$ year. - **Outcomes:** all TB notifications; UK-born and non-UK-born notifications; UK-born notifications at age ≥ 65 (denominator: all residents aged ≥ 65). - **Model:** Poisson regression with region and year fixed effects, exposure lagged or led 1–3 years, adjusted for in-migration and age structure. - **Inference:** with only nine clusters, cluster-robust $t(8)$ intervals and randomisation p-values. The randomisation test compared the cluster-robust t statistic with 499 permutations of whole regional exposure histories across regions; it is valid under exchangeability of those histories, and $p = (1 + \text{count}) / (1 + 499)$. - **Falsification:** lag 2 and lead 2 estimated jointly, with London excluded as a sensitivity analysis for UK-born notifications. - **Long differences:** across UTLAs with at least 10 notifications in each period, change in log prescribing (2014–16 to 2019–21) against change in log notification rate (2014–16 to 2022–24), weighted by notifications and adjusted for change in in-migration.

Cross-sectional analyses. Negative binomial regression of TB notifications on standardised log prescribing rates for the same period, adjusted for percentage born outside the UK, age structure, in-migration, HIV and diabetes prevalence.

Expected effects and minimum detectable effects (MDEs).

Expected effects. The population effect of a 10% increase in use was calculated as $[1 + 1.1p(RR - 1)] / [1 + p(RR - 1)]$, where p is prevalence of use and RR the individual-level relative risk (odds ratios where only these were available). The calculation assumes that:

1. prevalence of use scales with the prescribing measure;
2. extra prescribing reaches new users rather than lengthening existing courses;
3. baseline risk is homogeneous;
4. timing is aligned.

Assumptions 3 and 4 favour detection. The direction of any bias from assumptions 1 and 2 is unknown: prednisolone-equivalent mg per item fell over time, so prescribing volume and the number of people treated need not move together. Alternative expected effects came from three sources:

- recorded drug-associated notifications, converted to attributable fractions (exposed share $\times (RR - 1)/RR$), assuming complete recording or 50% recording;
- stratification by age (the same age-specific prevalence of use for UK-born and non-UK-born people);
- SCMD-derived prevalence of hospital drug use, including screening-attenuated relative risks for TNF inhibitors and attenuation by the positive-control elasticity.

MDEs. The MDE was calculated as $\exp(2.80 \times SE)$. For each MDE we report:

- the minimum detectable population attributable fraction (PAF);
- the largest PAF compatible with the upper 90% confidence limit.

Power simulation. A permutation-based simulation kept the real TB counts, so real overdispersion, serial correlation and trends were preserved. Whole exposure trajectories were permuted across areas, and known effects were injected by adding cases. This gave 500 null and 300 effect replicates per scenario, including scenarios where the effect acts through same-year exposure. We also report analytic power for each scenario at the real-data standard error.

Analyses used Python 3.14 (pandas 3.0, statsmodels).

Results

Prescribing, TB notifications and within-area variation

TB notifications in England - Fell from 6,474 in 2014 to a low of 4,123 in 2020 (Fingertips national series). - Rose to 5,490 in 2024, when 81.9% of people notified were born outside the UK.

Within-area variation in prescribing - Once prescribing was apportioned by where patients live, it varied little within areas over time. - For systemic oral glucocorticoids, the SD of the log rate after removing area and year means was 0.042, against 0.29 between areas; after removing area and year means, 95% of area-years were within $\pm 9\%$. - PPIs (0.036), metformin (0.041) and levothyroxine (0.042) were similar; conventional DMARDs varied more (0.088). Prednisolone-equivalent mg (0.041) and ADQ measures (0.026–0.097) varied as little as items. - A 10% within-area change in prescribing is therefore at the edge of the observed data (Figure 2).




Figure 2: **Figure 2.** Within-area versus between-area variation in log prescribing rates, 294 lower-tier authorities, 2014–2024. Orange line: a 10% increase.

Primary care prescribing and TB notifications (panel analyses)

Primary analysis (294 lower-tier authorities, 3,233 area-years, outcome years 2014–2024) - No drug group was associated with notifications in the following year after correction for multiple testing (Table 1; all FDR $q \geq 0.51$). - Systemic oral glucocorticoids: IRR 0.969 (95% CI 0.929–1.012) per 10% within-area increase.

Other groups (same scale, 95% CI):

Drug group	IRR
Oral hydrocortisone	0.998 (0.985–1.011)
Inhaled corticosteroids	0.966 (0.920–1.014)
Conventional DMARDs	1.002 (0.983–1.021)
Transplant immunosuppressants	0.999 (0.993–1.005)
Proton pump inhibitors	0.996 (0.955–1.039)
Statins	0.992 (0.952–1.034)
Metformin	0.966 (0.934–0.998)
Levothyroxine (negative control)	0.962 (0.933–0.992)

- Metformin (nominal $p = 0.04$) was inversely associated to the same extent as the negative control, levothyroxine ($p = 0.01$), which has no plausible effect on TB.

Table 1. Primary care prescribing and TB notifications: incidence rate ratio per 10% within-area increase (95% CI)

Drug group	Previous year: residence (294 areas, 2014–2024)	Previous year: UTLA, residence	Previous year: LTLA, practice postcode	Joint model: previous year (t–1)	Joint model: following year (t+1)	p, t–1 vs t+1	Largest compatible PAF / prevented fraction (PAF after negative-control shift)
Antituberculosis (descriptive)	1.000 (0.997–1.003)	1.000 (0.996–1.004)	1.001 (0.998–1.003)	1.000 (0.996–1.003)	1.002 (0.998–1.005)	0.423	3% / 2% (42%)
Systemic oral glucocorticoids	0.969 (0.929–1.012)	0.974 (0.928–1.023)	0.974 (0.939–1.010)	0.990 (0.941–1.041)	0.952 (0.912–0.994)	0.334	5% / 39% (44%)
Oral hydrocortisone	0.998 (0.985–1.011)	0.999 (0.985–1.014)	0.999 (0.987–1.011)	0.995 (0.978–1.012)	1.007 (0.991–1.023)	0.416	9% / 11% (49%)
Oral dexamethasone	1.003 (0.992–1.014)	1.004 (0.992–1.016)	1.001 (0.991–1.011)	1.002 (0.991–1.013)	1.003 (0.993–1.013)	0.907	12% / 6% (52%)
Inhaled corticosteroids	0.966 (0.920–1.014)	0.965 (0.916–1.017)	0.973 (0.934–1.014)	1.058 (0.995–1.126)	0.879 (0.840–0.921)	0.000	7% / 42% (46%)
Conventional DMARDs	1.002 (0.983–1.021)	1.001 (0.978–1.023)	1.004 (0.987–1.021)	1.011 (0.982–1.042)	0.988 (0.954–1.024)	0.446	18% / 13% (58%)
Transplant immunosuppressants	0.999 (0.993–1.005)	0.998 (0.992–1.005)	1.000 (0.995–1.006)	0.999 (0.990–1.009)	0.999 (0.992–1.006)	0.954	4% / 6% (44%)

Drug group	Previous year: LTLA, residence (294 areas, 2014-2024)	Previous year: UTLA, residence	Previous year: LTLA, practice postcode	Joint model: previous year (t-1)	Joint model: following year (t+1)	p, t-1 vs t+1	Largest compatible PAF / prevented fraction (PAF after negative-control shift)
Proton pump inhibitors	0.996 (0.955-1.039)	0.997 (0.953-1.043)	0.992 (0.955-1.029)	1.033 (0.979-1.090)	0.966 (0.900-1.037)	0.253	32% / 28% (72%)
Statins	0.992 (0.952-1.034)	0.985 (0.943-1.030)	0.989 (0.954-1.025)	0.996 (0.940-1.055)	0.992 (0.922-1.068)	0.954	27% / 29% (68%)
Metformin	0.966 (0.934-0.998)	0.958 (0.924-0.992)	0.964 (0.935-0.994)	0.967 (0.913-1.024)	0.999 (0.939-1.062)	0.576	0% / 38% (32%)
Insulins	0.966 (0.929-1.005)	0.971 (0.929-1.014)	0.972 (0.940-1.005)	1.028 (0.974-1.085)	0.921 (0.866-0.980)	0.038	0% / 39% (38%)
Fluoroquinolones	1.005 (0.992-1.017)	1.004 (0.991-1.018)	1.003 (0.991-1.016)	1.010 (0.995-1.026)	0.993 (0.979-1.009)	0.185	15% / 6% (55%)
All antibacterials	0.996 (0.955-1.039)	1.003 (0.957-1.052)	0.995 (0.958-1.033)	1.000 (0.957-1.045)	0.998 (0.967-1.031)	0.945	32% / 28% (73%)
Vitamin D	1.007 (0.987-1.027)	1.009 (0.987-1.031)	1.007 (0.988-1.025)	1.001 (0.975-1.028)	1.007 (0.980-1.033)	0.827	24% / 9% (64%)
Levothyroxine (negative control)	0.962 (0.933-0.992)	0.956 (0.926-0.988)	0.965 (0.939-0.993)	0.983 (0.928-1.042)	0.972 (0.908-1.040)	0.846	0% / 38% (26%)

Poisson PML with area and year fixed effects, adjusted for age structure, international immigration, HIV and diabetes prevalence (diabetes not adjusted for metformin and insulins); SEs clustered by area. Joint model: prescribing in t-1 and t+1 in one model on a common sample; because the two are highly correlated within areas, their estimates are negatively correlated and opposite signs can arise by chance. PAF, largest population attributable fraction compatible with the upper 90% confidence limit; prevented fraction, the same from the lower 90% limit; in brackets, the PAF after dividing the upper limit by the negative control (levothyroxine) estimate, as if its bias applied to every drug.

Largest attributable fraction compatible with the data (upper 90% confidence limit) - Systemic oral glucocorticoids 5%; conventional DMARDs 18%; transplant immunosuppressants 4%. - These bounds depend on point estimates that happened to fall below 1, and they are not robust to the bias revealed by the negative control. Dividing each upper limit by the levothyroxine estimate raised them to 44%, 58% and 44%, close to the minimum detectable attributable fractions (Table 2). - For drugs expected to protect, the data were compatible with prevented fractions of up to 38% (metformin) and 29% (statins).

Robustness - Upper-tier geography (151 areas) gave similar results: - systemic oral glucocorticoids 0.974 (0.928-1.023); all FDR $q \geq 0.22$ - the negative control, levothyroxine, was non-

TB notifications and primary care prescribing: lower-tier local authorities, England

Annual notifications 2014-2024, 294 areas, up to 3,234 area-years; prescribing apportioned by residence. Poisson PML with area and year fixed effects; adjusted for age structure, international in-migration, HIV and diabetes prevalence (not diabetes for metformin and insulins); SEs clustered by area.




Figure 3: **Figure 4.** Primary care prescribing and TB notifications, lower-tier authorities, prescribing apportioned by patient residence. Blue: prescribing in the previous year (primary). Orange: prescribing in the following year, estimated jointly (falsification).

inally inverse at 0.956 (0.926–0.988), the same size as metformin (0.958, 0.924–0.992). This suggests residual confounding by trends. - Apportioning prescribing by practice postcode gave similar results: lower-tier oral glucocorticoids 0.974 (0.939–1.010). - For oral glucocorticoids, estimates were also materially unchanged by (Table S1): - a three-year prior exposure window, area-specific linear trends, or log population as a covariate - adjustment for the LTBI programme or asylum support - excluding COVID-affected years - restricting to outcome years 2018–2024 (0.983, 0.941–1.028) or to EPD-era exposure (0.979, 0.939–1.021). - Inhaled corticosteroids were nominally inverse in several sensitivity analyses (for example 0.938 for outcome years 2018–2024), the direction opposite to any plausible harmful effect.

Falsification tests (prescribing in years $t-1$ and $t+1$ estimated jointly; Table 1 and Table S2b) - Following-year prescribing was inversely associated with notifications for some groups: - inhaled corticosteroids: $t-1$ 1.058 (0.995–1.126), $t+1$ 0.879 (0.840–0.921); difference $p < 0.001$ - insulins: $t-1$ 1.028 (0.974–1.085), $t+1$ 0.921 (0.866–0.980); difference $p = 0.04$ - systemic oral glucocorticoids: $t-1$ 0.990 (0.941–1.041), $t+1$ 0.952 (0.912–0.994); difference $p = 0.33$. - Within areas, prescribing in nearby years is highly correlated, so in the two-year model the $t-1$ and $t+1$ estimates were negatively correlated (-0.16 to -0.83 across groups) and opposite signs can arise by chance. - With same-year prescribing added, previous-year terms were null for every group. Same-year terms were positive for several groups (for example proton pump inhibitors 1.128, 1.035–1.229), the direction expected from prescribing for undiagnosed TB, and following-year terms remained inverse. - Area-specific linear trends gave inhaled corticosteroids 1.033 (0.981–1.087). - **Multiplicity.** Of all 277 estimates in the lower-tier residence models, 30 were nominally significant, more than the 5% expected by chance. They were concentrated in the negative control (levothyroxine, 6, all inverse), metformin (5, inverse) and inhaled corticosteroids (8, mostly inverse or following-year terms), a pattern that points to shared trends rather than drug effects.

Spatial dependence - Raw TB notification rates were strongly spatially clustered in every year (Moran's I 0.25–0.39 lower-tier; 0.40–0.49 upper-tier). - Residual dependence after the primary model was weak and concentrated in 2014: - lower-tier: median I 0.026 (range -0.004 to 0.163), nominally significant in 2 of 11 years (2014, $I = 0.163$; 2023, $I = 0.079$) - upper-tier: median I 0.009, nominally significant only in 2014 ($I = 0.209$). - 2014 is the only outcome year whose previous-year exposure comes from the pre-2014 HSCIC series.

Oral glucocorticoid prescribing trends and changes in TB notifications

National trends, 2011–2024 (standard GP practices) - Systemic oral glucocorticoid items rose from 119 to 132 per 1,000 residents in 2011–2016, then fell to 111 in 2024. Most of the fall came in 2020–21 (Figure 3). - Prednisolone-equivalent mg per item fell from 225 to 192, so courses became smaller. - Oral hydrocortisone items, mainly replacement therapy, rose from 5.5 to 9.1 per 1,000.

Correlation with national TB trends - National notification rates and prescribing were not correlated in levels (14 years; $r = 0.33$, $p = 0.25$). - Year-on-year changes were not correlated at lags of 0–2 years. - At a 3-year lag, one of eight correlations was nominally significant and inverse ($r = -0.75$).

Regional and local changes - In the regional panel (9 regions, 2012–2024), adjusted estimates were null: - items, lag 1: 1.034 (0.774–1.382); randomisation $p = 0.80$ - prednisolone mg, lag 1: 1.094 (0.845–1.416); randomisation $p = 0.57$. - Across 140 upper-tier authorities, changes in prescribing (2014–16 to 2019–21) and in notifications (2014–16 to 2022–24) were weakly inversely related before adjustment (Spearman $\rho = -0.19$). After adjustment for change in in-migration they were unrelated (ratio per 10% 0.983, 0.917–1.054).

Oral glucocorticoid prescribing in English primary care, 2011–2024

Standard GP practices; HSCIC practice-level prescribing (2011–13) and NHSBSA English Prescribing Dataset (2014–24); per resident using ONS mid-year estimates. Systemic = prednisolone, prednisone, methylprednisolone, deflazacort, dexamethasone.




Figure 4: **Figure 3.** Oral glucocorticoid prescribing in English primary care, 2011–2024: systemic oral glucocorticoid items, oral hydrocortisone items and prednisolone-equivalent mg per resident.

Hospital medicines

Coverage - After reassigning merged trusts to their successors, 98–100% of DDD in every drug group and year came from trusts with catchment data, and 95% for levetiracetam (Table S7).

Positive control: active-TB treatment (pyrazinamide DDD, upper-tier authorities) - Between areas: tracked notifications (Spearman $\rho = 0.81$; adjusted IRR per SD 1.23, 1.10–1.36). - Within areas, same year: 1.035 (1.009–1.061) per 10% increase. - Previous year: 1.001 (0.993–1.010). Following year, estimated jointly: 0.997 (0.986–1.007). - The within-area elasticity was 0.36 (95% CI 0.10–0.62; lower-tier 0.40, 0.14–0.66). It reflects apportionment error, timing (treatment of people notified late in a year continues into the next) and variation in drug volume per person treated, not linkage error alone. - Rifamycin/isoniazid products, which also cover latent TB and other infections, had the same elasticity (0.37, 0.22–0.51). The more specific measure gained nothing, so the attenuation is dominated by apportionment rather than drug specificity.




Figure 5: **Figure 5.** Positive control: hospital active-TB treatment (pyrazinamide DDD-years per 1,000 residents, including fixed-dose combinations) and TB notifications between areas (left) and within areas over time (right).

Candidate drugs (prescribing in the previous year, IRR per 10% increase; Table 3 and Table S3b)

Drug group	Upper-tier	Lower-tier
TNF inhibitors	0.989 (0.972–1.007)	0.992 (0.976–1.008)
IL-6 inhibitors/abatacept	1.001 (0.992–1.009)	1.001 (0.993–1.010)
JAK inhibitors	0.994 (0.987–1.002)	0.995 (0.989–1.002)
Rituximab	0.998 (0.984–1.012)	1.001 (0.986–1.016)
Calcineurin/mTOR inhibitors	1.008 (0.996–1.020)	1.008 (0.997–1.020)
Antiproliferatives	0.996 (0.978–1.014)	0.997 (0.982–1.013)
Systemic glucocorticoids (prednisolone-equivalent mg)	1.024 (1.001–1.046)	1.021 (1.000–1.043)
Dexamethasone and hydrocortisone (descriptive)	1.008 (0.980–1.036)	1.010 (0.984–1.036)

- **Systemic glucocorticoids** were nominally associated with notifications in the following

year at upper-tier level. The association:

- was not robust to clustering by principal trust (1.024, 0.999–1.049) or to excluding outcome years 2020–21 (1.018, 0.969–1.069);
- would imply, under the model used for expected effects, that hospital glucocorticoids account for about a quarter of all notifications (an IRR of 1.024 per 10% corresponds to an attributable fraction of 24%), whereas steroid-associated immunosuppression was recorded for 0.5% of notifications [UKHSA 2025]. We interpret it as chance or confounding by hospital activity, not a drug effect.
- The negative controls were null: low-TB-risk biologics 1.002 (0.986–1.018); levetiracetam 1.014 (0.986–1.042).
- Clustering by principal trust or using elective catchments barely changed the other estimates (Table S3b).
- JAK inhibitors showed small inverse estimates (same year 0.993, 0.987–0.999; previous year excluding outcome years 2020–21 0.978, 0.961–0.996), consistent with trends in uptake rather than protection.

Table 3. Hospital medicines (SCMD) and TB notifications, 2019–2024

Level	Drug group	Cross-sectional, adjusted (per SD)	Within-area, same year (per 10%)	Elasticity (same year)	Within-area, previous year (per 10%)	Previous year, SE clustered by principal trust	Following year (joint with previous)
UTLA	Active-TB treatment (pyrazinamide DDD, including fixed-dose combinations; positive control)	1.225 (1.101–1.362)	1.035 (1.009–1.061)	0.36	1.001 (0.993–1.010)	1.001 (0.993–1.010)	0.997 (0.986–1.007)
UTLA	Rifamycin/isoniazid (active and latent TB, other infections)	1.158 (1.038–1.292)	1.036 (1.021–1.050)	0.37	1.005 (0.993–1.018)	1.005 (0.993–1.018)	1.018 (0.996–1.041)
UTLA	TNF inhibitors	0.957 (0.907–1.010)	0.996 (0.979–1.013)	-0.05	0.989 (0.972–1.007)	0.989 (0.970–1.009)	0.978 (0.949–1.008)
UTLA	IL-6 inhibitors and abatacept	0.971 (0.923–1.022)	1.000 (0.990–1.010)	-0.00	1.001 (0.992–1.009)	1.001 (0.993–1.009)	1.001 (0.984–1.018)
UTLA	JAK inhibitors (rheumatology)	0.972 (0.916–1.031)	0.993 (0.987–0.999)	-0.07	0.994 (0.987–1.002)	0.994 (0.986–1.003)	0.998 (0.984–1.011)

Level	Drug group	Cross-sectional, adjusted (per SD)	Within-area, same year (per 10%)	Elasticity (same year)	Within-area, previous year (per 10%)	Previous year, SE clustered by principal trust	Following year (joint with previous)
UTLA	Rituximab	0.942 (0.886–1.002)	0.992 (0.978–1.005)	-0.09	0.998 (0.984–1.012)	0.998 (0.987–1.009)	1.006 (0.989–1.024)
UTLA	Calcineurin/mTOR inhibitors	1.049 (0.995–1.106)	0.994 (0.985–1.004)	-0.06	1.008 (0.996–1.020)	1.008 (0.996–1.020)	0.985 (0.966–1.005)
UTLA	Antiproliferatives	1.025 (0.970–1.082)	0.993 (0.978–1.008)	-0.08	0.996 (0.978–1.014)	0.996 (0.979–1.013)	0.993 (0.974–1.012)
UTLA	Systemic glucocorticoids without dexamethasone/hydrocortisone (prednisolone-equivalent mg)	0.970 (0.929–1.012)	1.009 (0.986–1.032)	0.09	1.024 (1.001–1.046)	1.024 (0.999–1.049)	0.991 (0.953–1.030)
UTLA	Dexamethasone and hydrocortisone (descriptive)	0.934 (0.893–0.976)	1.005 (0.977–1.034)	0.05	1.008 (0.980–1.036)	1.008 (0.979–1.037)	0.994 (0.943–1.047)
UTLA	Low-TB-risk biologics (negative control)	1.014 (0.961–1.070)	1.004 (0.990–1.019)	0.05	1.002 (0.986–1.018)	1.002 (0.983–1.021)	1.010 (0.982–1.040)
UTLA	Levetiracetam (negative control)	1.017 (0.970–1.066)	0.998 (0.975–1.021)	-0.03	1.014 (0.986–1.042)	1.014 (0.986–1.042)	0.987 (0.948–1.028)
LTLA	Active-TB treatment (pyrazinamide DDD, including fixed-dose combinations; positive control)	1.154 (1.080–1.232)	1.039 (1.013–1.065)	0.40	1.000 (0.991–1.008)	1.000 (0.992–1.008)	0.995 (0.985–1.005)

Level	Drug group	Cross-sectional, adjusted (per SD)	Within-area, same year (per 10%)	Elasticity (same year)	Within-area, previous year (per 10%)	Previous year, SE clustered by principal trust	Following year (joint with previous)
LTLA	Rifamycin/isoniazid (active and latent TB, other infections)	1.065 (1.000–1.135)	1.038 (1.025–1.052)	0.39	1.004 (0.992–1.016)	1.004 (0.992–1.016)	1.018 (0.997–1.040)
LTLA	TNF inhibitors	0.968 (0.927–1.012)	0.994 (0.979–1.010)	-0.06	0.992 (0.976–1.008)	0.992 (0.976–1.008)	0.975 (0.946–1.006)
LTLA	IL-6 inhibitors and abatacept	0.944 (0.906–0.984)	1.001 (0.991–1.011)	0.01	1.001 (0.993–1.010)	1.001 (0.993–1.010)	0.997 (0.981–1.013)
LTLA	JAK inhibitors (rheumatology)	0.948 (0.905–0.992)	0.994 (0.988–0.999)	-0.06	0.995 (0.989–1.002)	0.995 (0.988–1.003)	0.998 (0.984–1.012)
LTLA	Rituximab	0.974 (0.929–1.021)	0.990 (0.976–1.004)	-0.10	1.001 (0.986–1.016)	1.001 (0.988–1.014)	1.005 (0.988–1.023)
LTLA	Calcineurin/mTOR inhibitors	0.992 (0.959–1.046)	0.994 (0.985–1.003)	-0.07	1.008 (0.997–1.020)	1.008 (0.998–1.019)	0.983 (0.961–1.005)
LTLA	Antiproliferatives	0.983 (0.936–1.031)	0.997 (0.983–1.012)	-0.03	0.997 (0.982–1.013)	0.997 (0.980–1.015)	0.993 (0.974–1.012)
LTLA	Systemic glucocorticoids without dexamethasone/hydrocortisone (prednisolone-equivalent mg)	0.976 (0.945–1.008)	1.008 (0.990–1.026)	0.08	1.021 (1.000–1.043)	1.021 (0.999–1.044)	0.986 (0.952–1.020)
LTLA	Dexamethasone and hydrocortisone (descriptive)	0.978 (0.938–1.019)	1.008 (0.980–1.037)	0.08	1.010 (0.984–1.036)	1.010 (0.982–1.037)	0.989 (0.940–1.041)
LTLA	Low-TB-risk biologics (negative control)	0.991 (0.948–1.036)	1.005 (0.991–1.019)	0.05	1.003 (0.988–1.019)	1.003 (0.988–1.019)	1.012 (0.985–1.040)

Level	Drug group	Cross-sectional, adjusted (per SD)	Within-area, same year (per 10%)	Elasticity (same year)	Within-area, previous year (per 10%)	Previous year, SE clustered by principal trust	Following year (joint with previous)
LTLA	Levetiracetam (negative control)	1.016 (0.979–1.054)	1.003 (0.981–1.025)	0.03	1.013 (0.988–1.039)	1.013 (0.989–1.037)	0.992 (0.955–1.031)

Exposure: DDD-years per 1,000 residents (positive control: pyrazinamide DDD including fixed-dose combinations; systemic glucocorticoids: prednisolone-equivalent mg, without dexamethasone and hydrocortisone), apportioned by 2024 acute trust catchments. Cross-sectional: negative binomial regression of 2019–24 notifications on the standardised log mean rate (per SD), adjusted for % born outside the UK, age structure, in-migration, HIV and diabetes prevalence. Within-area: Poisson PML with area and year fixed effects and time-varying covariates, per 10% increase; SEs clustered by area unless stated. Elasticity: log IRR per log unit of exposure in the same-year model, shown for all groups but interpretable as linkage attenuation only for the positive control. Following year: for the positive control an association is possible because treatment continues into the next year; for candidate drugs it is a falsification test.

Minimum detectable effects and expected population effects

Primary care, oral glucocorticoids (lower-tier panel) - Minimum detectable effect (MDE): 6.3% per 10% increase in prescribing. - Expected effect under published relative risks and prevalence: 0.34% (Table 2). - Stratified by age: 0.29%. - Based on UKHSA-recorded steroid-associated notifications, converted to an attributable fraction: 0.043% with complete recording, 0.087% if only half are recorded. - The design could therefore detect only effects about 18 to 140 times larger than expected. - The MDE corresponds to a drug responsible for 63% of all notifications. - For the other candidate groups, MDEs exceeded expected effects by factors of 11 (statins) to 272 (conventional DMARDs).

Hospital medicines - MDEs were smaller (1.0–2.9% per 10% increase), but so were expected effects. - TNF inhibitors (SCMD DDD-years per resident 0.34%): - An illustrative RR of 4 before latent TB screening implies an expected change of 0.10% (MDE 26 times larger at upper-tier level). - An RR of 1.5 with screening, which probably overstates post-screening risk, implies 0.017% (MDE 154 times larger). - Dividing by the positive-control elasticity widens these gaps to 72 and 426 times (Table S3). - Across hospital scenarios, MDEs exceeded expected effects by 26 to 520 times before attenuation. - UKHSA-recorded biological-therapy notifications imply 0.08–0.16%.

Simulation (permutation-based, real notification counts; Table S4) - **Calibration.** With no effect, the two-sided test rejected in 7.8% of replicates at lower-tier level (7.6% upper-tier), against a nominal 5%. Within the simulation, replicate SEs (median 0.0161) were close to the spread of null estimates (0.0166), so the clustered test was mildly anti-conservative, not conservative. - **Real-data precision.** The real-data SE (0.0218) was larger than in the simulation, because permuting exposure trajectories across areas breaks their alignment with each area's own notification trends. Simulated power is therefore optimistic, so we also give analytic power at the real-data SE. - **Published glucocorticoid effect (RR 4.9).** Rejection 8.3% (upper-tier 8.0%), no more often than with no effect; power in the correct direction 6.0%, against 3.8% with no effect. - **Larger individual relative risks.** Rejection 17.7% for RR 25 and 71.0% for RR 100 (analytic power at the real SE for RR 100: 52.1%). - **Direct population effects.** Rejection 21.7%, 83.0% and 100.0% for IRR 1.02, 1.05 and 1.10 per 10%; analytic power at the real SE 14.3%, 64.6% and 99.4% (upper-tier, IRR 1.05: 51.0%). - **Effect acting through same-year**

prescribing (analysed with previous-year exposure). Rejection 61.7% for 1.05 and 99.3% for 1.10.

Table 2. Minimum detectable effects (LTLA panel, residence apportionment, exposure t–1) versus expected population effects of a 10% increase in use

Drug group	Individual RR/OR	Prevalence of use	PAF (negative: prevented fraction)	SE of log IRR per 10%	Expected change	MDE	MDE ÷ expected	Power	Minimum detectable PAF	RR for 80% power
Systemic oral glucocorticoids	4.9	0.9%	3.4%	0.0218	0.34%	6.3%	18	3.6%	63%	190
Inhaled corticosteroids	1.27	5.2%	1.4%	0.0249	0.14%	7.2%	50	2.8%	72%	51
Conventional DMARDs	1.2	0.5%	0.1%	0.0097	0.01%	2.8%	272	2.6%	28%	77
Transplant immunosuppressants	2	0.1%	0.0%	0.0031	0.00%	0.9%	172	2.6%	9%	190
Proton pump inhibitors	1.28	14.2%	3.8%	0.0213	0.38%	6.2%	16	3.7%	62%	12
Statins	0.6	12.8%	-5.4%	0.0210	-0.54%	6.1%	11	4.4%	61%	not attainable
Metformin	0.51	4.4%	-2.2%	0.0169	-0.22%	4.8%	21	3.4%	48%	not attainable

Expected changes use published individual-level relative risks (ORs for oral glucocorticoids and PPIs) and UK prevalence of use, assuming that prevalence of use scales with prescribing, that extra prescribing reaches new users, a homogeneous baseline risk and aligned timing. Negative PAF values are prevented fractions. $MDE = \exp(2.80 \times SE) - 1$. The ratio of MDE to expected change is essentially independent of the 10% contrast, because for small changes both scale in proportion to it. Power is one-sided power to detect the expected effect in the correct direction. Benchmarks: recorded cases (steroids), UKHSA 2024, 100% recorded, RR 4.9: 0.043%; recorded cases (steroids), UKHSA 2024, 50% recorded, RR 4.9: 0.087%; recorded cases (biological therapy), UKHSA 2024, 100% recorded, RR 4.9: 0.078%; recorded cases (biological therapy), UKHSA 2024, 50% recorded, RR 4.9: 0.157%; age-stratified (oral corticosteroids, RR 4.9): 0.293%; homogeneous (oral corticosteroids, RR 4.9, prevalence 0.9%): 0.339%.

Regional analyses by place of birth and age

Precision - Regional models were very imprecise. MDEs for glucocorticoid prescribing were 45–82% per 10% increase for UK-born notifications and 46–99% for UK-born notifications at age ≥ 65 . - Under the model used for expected effects, a 10% increase in use can raise notifications by at most 10%. The expected effect for UK-born people aged ≥ 65 (prevalence of use 2.5%, RR 4.9) is 0.9%. - Any nominally significant regional estimate must therefore reflect chance or bias.

Non-UK-born notifications - Oral glucocorticoid prescribing in earlier years was not associated with notifications (lag 1–3: 0.95–1.03). - The falsification test failed: prescribing three years later predicted earlier notifications (items 1.49, 1.17–1.89, randomisation $p = 0.01$; prednisolone mg 1.29, 1.03–1.60, $p = 0.04$).

UK-born notifications - Prednisolone-equivalent mg: lagged estimates were imprecise positive values (lag 1: 1.25, 0.95–1.65, randomisation $p = 0.12$; lag 2: 1.33, 0.96–1.84, $p = 0.16$). - With lag 2 and lead 2 estimated jointly: lag 1.45 and lead 0.88 (difference $p = 0.04$; randomisation p for lag = 0.23). Excluding London: 1.12 and 0.90 (difference $p = 0.43$; randomisation $p = 0.73$) (Table S6). - Oral glucocorticoid items: null (joint lag 2: 1.27; randomisation $p = 0.46$).

UK-born notifications at age ≥ 65 - Glucocorticoid prescribing in earlier years was not associated with notifications. - The falsification test failed for glucocorticoid items (lead 2: 1.68, 1.04–2.70; randomisation $p = 0.04$). - The negative control, levothyroxine, was associated with notifications in the following year (lead 1: 1.38, 1.05–1.80; randomisation $p = 0.04$), and imprecisely with the previous year (lag 1: 1.23, 1.00–1.51; $p = 0.07$).

Summary - Of 81 adjusted regional estimates, 4 had randomisation $p \leq 0.05$, all of them falsification (lead) terms. - The UK-born association with prednisolone mg was implausibly large, not significant by randomisation inference, and dependent on London.

Discussion

Principal findings

We linked open English data on primary care and hospital prescribing with TB notifications, using several ecological designs at two geographic scales. The linkage worked only partly.

- **The positive control.** Hospital active-TB treatment tracked notifications between areas and, weakly, within areas. Its elasticity of 0.36–0.40, with wide confidence intervals, reflects apportionment error, timing and variation in drug volume per person treated.
- **Candidate drugs.** No medicine plausibly affecting TB risk was robustly associated with subsequent notifications. This held for systemic oral glucocorticoids, inhaled corticosteroids, DMARDs, transplant immunosuppressants, TNF, IL-6 and JAK inhibitors, and rituximab. Hospital systemic glucocorticoids had a small nominal association that was not robust and implied an implausibly large attributable fraction.
- **Signs of confounding.**
 - The negative-control exposure showed inverse estimates of similar size to some candidate drugs.
 - Following-year prescribing was inversely associated with notifications for several groups, although adjacent-year exposures are highly collinear.
 - The regional association with UK-born TB was implausibly large, not significant by randomisation inference, and dependent on London.

These patterns are compatible with residual confounding by area-specific trends, chance or measurement error, and not with drug effects of plausible size.

- **Power.** The analytic calculations and the simulation both showed that plausible effects are undetectable: the designs could detect only effects roughly 10 to several hundred times larger than those implied by published relative risks or by UKHSA records of drug-associated TB. Simulated power exceeded analytic power at the real standard error, because permuted exposures lost their alignment with local notification trends.

Oral glucocorticoid prescribing and UK-born TB notifications by region, indexed to 2014 = 100

Prescribing: standard GP practices, per 1,000 residents. UK-born TB: UKHSA TB in England 2025, Supplementary Table 12 (LFS denominators; small annual counts in some regions).




Figure 6: **Figure 6.** Oral glucocorticoid prescribing and UK-born TB notification rates by region, indexed to 2014 = 100.

Why population effects of these medicines are undetectable

Expected effects are small. The population effect of a medicine is roughly its attributable fraction scaled by the relative change in use.

- **Oral glucocorticoids.** Current use carries an odds ratio of about 5 [Jick 2006], but only about 1% of people use them at any time [van Staa 2000; Fardet 2011]. A 10% change in use should therefore change TB notifications by about 0.3%.
 - Stratifying by age lowers this to 0.29%, because use is concentrated in older people while most notifications are in younger adults.
 - UKHSA recorded steroid-associated immunosuppression in 30 of 5,490 people notified in 2024 [UKHSA 2025]. Converted to an attributable fraction, that implies 0.04% per 10% increase with complete recording, or 0.09% if only half of drug-associated immunosuppression is recorded.
- **Biologics.** Latent TB screening before anti-TNF therapy has been standard in the UK since 2005 [BTS 2005] and reduced TB rates in treated patients by about 78% in registry data [Carmona 2005]; risk was about seven times higher when the recommendations were not followed [Gómez-Reino 2007]. Growth in biologic use during 2019-2024 therefore took place under screening, so the relative risk relevant to a marginal increase in use is small. Screening is less consistently done before oral glucocorticoids.

The data carry little information. - **Little within-area variation.** Within-area variation in prescribing was very small (SD of log rate 0.04 for glucocorticoids, against 0.29 between areas), so a 10% change sits at the edge of the observed data. Fixed effects remove the between-area variation where most information lies [Gunasekara 2014]. - **Few areas.** Power in aggregate studies depends mainly on the number of areas [Sheppard 1996], which cannot be increased. - **Linkage attenuation.** For hospital medicines, the positive control shows that apportionment would shrink any true effect further. No working positive control was available for primary care, so the validity of that linkage is untested. - **Simulation.** Simulations on the real notification counts rejected the null for the published glucocorticoid effect no more often than when there was no effect.

Confounding, falsification and negative controls

Crude versus within-area associations. Crude cross-sectional associations were strongly inverse and were explained by country of birth and age. Within areas, residual confounding remained visible: - **Negative control.** Levothyroxine, which has no plausible effect on TB, showed inverse estimates (0.96) similar to metformin. - **Lag and lead patterns.** Following-year prescribing was inversely associated with notifications for inhaled corticosteroids and insulins, and in models with previous, same and following years, same-year terms were positive for several groups. Adjacent-year exposures are highly collinear, so their estimates are strongly negatively correlated and these seesaw patterns are weak evidence on their own. - **Likely cause.** Both patterns are compatible with prescribing and notifications sharing area-specific trends, for example migration-driven changes in the age and origin of the population (with accompanying changes in practice registration and in vitamin D prescribing in South Asian-born communities), changes in diagnostic activity, and prescribing policy. Year fixed effects cannot remove these trends.

Metformin. Its estimates should be read against the high prevalence of both diabetes and latent TB infection among South Asian-born people; diabetes itself increases TB risk [Pealing 2015].

Ecological bias. Area-level adjustment cannot remove bias arising from effect modification or from differences in baseline risk between people within areas [Greenland 1989; Morgenstern 1995]. For example, steroid users are mostly older and UK-born, whereas most notifications are in younger people born abroad.

The UK-born association. Regionally, prednisolone dose in an earlier year was imprecisely associated with UK-born TB (joint lag estimate 1.45 per 10% increase). We do not interpret this

as a drug effect: - The regional MDEs (about 50% per 10%) are far above the 10% maximum possible under the model used for expected effects, so any significant regional estimate must reflect chance or bias. - It was not significant by randomisation inference ($p = 0.23$), and it largely disappeared when London was excluded (1.12). - The same nine-region design produced failed falsification tests for non-UK-born TB and a failed negative control (levothyroxine) for older UK-born people. - UK-born TB fell by 48% between 2014 and 2022 [UKHSA 2025], while oral glucocorticoid dosing also declined. Any two declining regional series will tend to correlate, so coincident trends are the most plausible explanation.

Protopathic bias. Glucocorticoids and antibacterials prescribed for undiagnosed TB, and glucocorticoids used in treating TB meningitis and pericarditis [Thwaites 2004], create reverse associations in the same year. This is why the primary exposure was prescribing in the previous year. The same bias probably inflates the individual-level estimates used to derive expected effects [Jick 2006]; if so, the true gap between expected and detectable effects is even wider.

Strengths and limitations

Strengths. - **Open and reproducible.** All data are public, and code and processed data are openly available. - **Exposure measurement.** - Primary care exposure was defined at presentation level, with dose measures, and apportioned by where registered patients live. - Hospital medicines were classified by mechanism using defined daily doses, with trust mergers resolved. - **Checks on validity.** - A disease-specific positive control, negative-control exposures and falsification tests estimated jointly. - Randomisation inference for the nine-region analyses. - Year-specific spatial diagnostics. - Power assessed both analytically and by simulation on the real data. - **Independent peer review.** Three independent reviews prompted substantial revision.

Limitations: exposure data. - **Prescribing measures.** Exposure was measured as prescribing volume per resident, not as people treated. - **Apportionment.** Apportioning by patient residence used annual April snapshots, with 2014 shares applied to 2011–2013. - **Pre-2014 data.** The 2011–2013 series come from a different release, and overlap-period concordance could not be assessed. National year-on-year changes were continuous across the 2013/2014 splice for most groups, but not for oral glucocorticoids (+3.3% at the splice against +0.6% and +1.0% either side) or all antibacterials (−0.1% against −4.5% and −5.6%). Restricting to EPD-era exposure did not change the conclusions (Table S1). - **Hospital catchments.** Hospital quantities were apportioned using admission-based catchments, which may misallocate specialist services, and the 2024 catchments were applied to every year from 2019. - **Hospital data capture.** Capture of homecare-delivered biologics in SCMD could not be verified. - **Hospital data period.** The hospital series spans only 2019–2024, including the pandemic. Hospital dexamethasone use rose with its adoption for severe COVID-19 [RECOVERY 2021], a marker of COVID-19 severity patterned by deprivation and ethnicity, so dexamethasone and hydrocortisone were separated from the candidate glucocorticoid exposure. - **Hospital prescriptions dispensed in the community.** Prescriptions written in hospitals and dispensed by community pharmacies (FP10(HP)) are attributed to hospital prescribers, not GP practices, and are not in SCMD. Some TB treatment and specialist medicines reach patients this way, so neither data source captures them. - **Drug group definitions.** Conventional DMARDs exclude hydroxychloroquine and sulfasalazine, which carry little TB risk. Mercaptopurine includes leukaemia maintenance therapy, and the primary care “transplant immunosuppressant” group includes shared-care use of mycophenolate and ciclosporin for autoimmune disease. Dexamethasone, despite its high potency, contributed only 5.6–6.7% of prednisolone-equivalent mg in primary care (2015, 2019 and 2024 samples), so it does not drive the dose measure.

Limitations: outcome and confounder data. - **Notifications, not incidence.** The outcome is notified TB, subject to diagnostic delay, residence assignment, the 2021 transition from ETS to NTBS, and 2020 disruption. - **Birthplace data.** TB by place of birth is published only by region, and UK-born population denominators by age are not published regionally. The model for UK-born notifications at age ≥ 65 therefore uses all residents aged ≥ 65 as the denominator, which can create spurious trends as the older non-UK-born population grows. - **Screening of**

migrants. Pre-entry screening of long-stay visa applicants from high-incidence countries (UK-wide from 2012) and the LTBI programme for new migrants reduce notifications among recent entrants in the areas where they settle. Neither is fully captured by our covariates. - **Outcome definitions.** We did not analyse culture-confirmed pulmonary TB or drug-resistant TB, which were not available by local authority and year in the published tables we used. - **Unmeasured confounders.** - Social risk factors and diagnostic intensity. - The area-level share of recent arrivals. - Latent TB programme activity after 2019/20.

Limitations: design and analysis. - **Pre-specification.** Many analyses were added after initial null results and are exploratory. - **Expected-effect inputs.** Relative risks for PPIs, statins and metformin come mainly from high-incidence settings, and prevalence inputs for hospital drugs are derived or illustrative. Neither would change the conclusions unless the true effects were an order of magnitude larger. - **Positive control.** The hospital positive control is a disease-specific drug with a large same-year signal. We did not test whether a small lagged effect injected through the apportioned hospital exposure would be recovered, and primary care had no working positive control.

Implications

For research. - Open prescribing data are valuable for describing prescribing [Bacon 2020; OpenPrescribing 2026] but cannot estimate the effects of medicines on rare outcomes such as TB. - Ecological studies of rare outcomes should report the following alongside any associations: - minimum detectable effects; - a positive control with its elasticity; - negative controls; - falsification tests. - Causal questions need individual-level linked data analysed with target trial emulation (Box). Even national cohorts give wide intervals for modest relative risks [Peeling 2015]. The only drug-TB target trial emulation we identified concerned DPP-4 inhibitors [Chen 2025].

For TB programmes. - **The null results do not mean these drugs are safe.** Glucocorticoids, biologics and JAK inhibitors remain important for individual patients. Drug-associated TB is serious, often extrapulmonary [Keane 2001], and largely preventable. - **Surveillance.** Monitoring is better served by more complete and detailed immunosuppression fields in NTBS than by ecological analysis of prescribing. Useful fields would record drug class, time since starting, and whether latent TB screening was done and treated. - **Audit.** The practical lever is auditing latent TB screening before biologics, JAK inhibitors and high-dose, long-term glucocorticoids. - **Aggregate data from UKHSA.** Aggregate UKHSA tables would allow population monitoring in the groups where drug-associated reactivation matters, namely: - notifications with recorded biologic or steroid immunosuppression; - notifications by area, year, place of birth, age and time since entry.

Conclusion

Open English prescribing and TB notification data can be linked, although hospital medicines were linked only with heavy attenuation and the linkage of primary care prescribing could not be validated. Ecological analyses of these data cannot detect the population-level effects of medicines on TB: plausible effects lie roughly 10 to several hundred times below what the designs can resolve, and the associations that do arise are compatible with residual confounding, chance or measurement error rather than drug effects.

Box. Individual-level target trial emulation (example: oral glucocorticoids and TB)

Data - CPRD Aurum (primary care prescribing with dose and quantity). - Hospital Episode Statistics (admitted patient care and outpatients) and ONS deaths. - Small-area deprivation. - A bespoke linkage to the UKHSA National TB Surveillance System, which provides notification-validated TB, site of disease, culture confirmation, and country of birth and year of entry. - Hospital-only drugs (biologics, JAK inhibitors) are not captured in CPRD. For these, use NHS

England high-cost drugs data (e.g. via OpenSAFELY) or disease registries (BSRBR-RA, BADBIR, UK IBD Registry) with the same TB linkage.

Eligibility - Adults aged ≥ 18 years with ≥ 12 months' registration. - No previous TB, TB or latent TB treatment, or oral glucocorticoid use in the previous 12 months. - No HIV or solid organ transplant. - Defined indication cohorts (e.g. polymyalgia rheumatica/giant cell arteritis, rheumatoid arthritis, COPD, asthma, inflammatory bowel disease).

Treatment strategies - Initiate and sustain oral glucocorticoids for ≥ 3 months, by prednisolone-equivalent dose (< 7.5 , 7.5 – < 15 , ≥ 15 mg/day), versus no initiation. - An active comparator where available.

Assignment and time zero - Sequential monthly nested trials. - Clone-censor-weight methods for sustained-use and dose strategies.

Outcome - Incident active TB: the first of an NTBS notification, a HES diagnosis (ICD-10 A15–A19), or a CPRD diagnosis with treatment. - Secondary: pulmonary vs extrapulmonary, culture-confirmed TB, TB death.

Follow-up - Up to 5 years, or until death, deregistration or end of linkage.

Estimands and analysis - Intention-to-treat and per-protocol cumulative incidence differences and ratios at 1, 2 and 5 years. - Pooled logistic regression with inverse probability of treatment and censoring weights.

Confounders - Age, sex, ethnicity, country of birth and time since entry, deprivation. - Diabetes, chronic kidney disease, smoking, alcohol, BMI. - TB contact or latent TB screening codes. - Indication severity, healthcare use, co-prescribed immunosuppressants.

Effect modification and bias checks - Effect modification by country of birth, specified in advance. - Negative-control exposure (levothyroxine initiation) and negative-control outcome. - E-values.

Feasibility - The TB notification rate in UK-born people was 2.1 per 100,000 in 2024, all ages [UKHSA 2025, Supplementary Table 12]; rates in older UK-born adults are of a similar order. Tens of thousands of long-term initiators followed for several years would yield only tens of TB events, so the individual-level study is itself constrained by sample size.

Data and code availability

All inputs are publicly available (NHSBSA Open Data Portal: English Prescribing Dataset and Secondary Care Medicines Data; NHS Digital practice-level prescribing and registered patients by LSOA; UKHSA TB reports and Fingertips; OHID acute trust catchment populations; ONS/Nomis; Home Office asylum statistics; NHS England ODS; MHCLG English Indices of Deprivation). Code, processed datasets and outputs: <https://github.com/drcjar/tb-prescribing-england>. Data were obtained in September 2026: the EPD and SCMD through the NHSBSA API (SCMD months use final data where available, otherwise provisional), and the NHS England ODS epraccr file, practice registration releases (April 2014–2024), OHID acute trust catchments (2024 catchment year), and the UKHSA TB in England 2025 and regional 2024 supplementary tables as published at that time. These portals revise data between releases.

References

Each journal entry was checked against PubMed metadata (via DOI→PMID conversion or citation lookup) and Crossref DOI metadata. Each DOI below resolves to the stated article. The grey literature was checked on GOV.UK, in the report PDF, or through search results where a site blocks automated access. Corrections relative to references_v3.md are listed at the end.

- Aldridge RW, Zenner D, White PJ, et al. Tuberculosis in migrants moving from high-incidence to low-incidence countries: a population-based cohort study of 519 955 migrants screened

- before entry to England, Wales, and Northern Ireland. *Lancet* 2016;388(10059):2510–8. doi:10.1016/S0140-6736(16)31008-X. PMID 27742165
- Bacon S, Goldacre B. Barriers to working with National Health Service England’s open data. *J Med Internet Res* 2020;22(1):e15603. doi:10.2196/15603. PMID 31929101
 - Berrocal-Almanza LC, Harris RJ, Collin SM, et al. Effectiveness of nationwide programmatic testing and treatment for latent tuberculosis infection in migrants in England: a retrospective, population-based cohort study. *Lancet Public Health* 2022;7(4):e305–15. doi:10.1016/S2468-2667(22)00031-7. PMID 35338849
 - Brassard P, Kezouh A, Suissa S. Antirheumatic drugs and the risk of tuberculosis. *Clin Infect Dis* 2006;43(6):717–22. doi:10.1086/506935. PMID 16912945
 - Brassard P, Suissa S, Kezouh A, Ernst P. Inhaled corticosteroids and risk of tuberculosis in patients with respiratory diseases. *Am J Respir Crit Care Med* 2011;183(5):675–8. doi:10.1164/rccm.201007-1099OC. PMID 20889902
 - British Thoracic Society Standards of Care Committee. BTS recommendations for assessing risk and for managing *Mycobacterium tuberculosis* infection and disease in patients due to start anti-TNF- α treatment. *Thorax* 2005;60(10):800–5. doi:10.1136/thx.2005.046797. PMID 16055611
 - Carmona L, Gómez-Reino JJ, Rodríguez-Valverde V, et al. Effectiveness of recommendations to prevent reactivation of latent tuberculosis infection in patients treated with tumor necrosis factor antagonists. *Arthritis Rheum* 2005;52(6):1766–72. doi:10.1002/art.21043. PMID 15934089
 - Castellana G, Castellana M, Castellana C, et al. Inhaled corticosteroids and risk of tuberculosis in patients with obstructive lung diseases: a systematic review and meta-analysis of non-randomized studies. *Int J Chron Obstruct Pulmon Dis* 2019;14:2219–27. doi:10.2147/COPD.S209273. PMID 31576118
 - Chen YG, et al. Target trial emulation of DPP-4 inhibitors in patients with T2DM for pulmonary tuberculosis: a nationwide observational data. *BMC Med* 2025;23:587. doi:10.1186/s12916-025-04423-1. PMID 41137015
 - Davidson JA, Thomas HL, Maguire H, et al. Understanding tuberculosis transmission in the United Kingdom: findings from 6 years of mycobacterial interspersed repetitive unit-variable number tandem repeats strain typing, 2010–2015. *Am J Epidemiol* 2018;187(10):2233–42. doi:10.1093/aje/kwy119. PMID 29878041
 - Fardet L, Petersen I, Nazareth I. Prevalence of long-term oral glucocorticoid prescriptions in the UK over the past 20 years. *Rheumatology (Oxford)* 2011;50(11):1982–90. doi:10.1093/rheumatology/ker017. PMID 21393338
 - Gómez-Reino JJ, Carmona L, Angel Descalzo M, et al. Risk of tuberculosis in patients treated with tumor necrosis factor antagonists due to incomplete prevention of reactivation of latent infection. *Arthritis Rheum* 2007;57(5):756–61. doi:10.1002/art.22768. PMID 17530674
 - Greenland S, Morgenstern H. Ecological bias, confounding, and effect modification. *Int J Epidemiol* 1989;18(1):269–74. doi:10.1093/ije/18.1.269. PMID 2656561
 - Gunasekara FI, Richardson K, Carter K, Blakely T. Fixed effects analysis of repeated measures data. *Int J Epidemiol* 2014;43(1):264–9. doi:10.1093/ije/dyt221. PMID 24366487
 - Hermans S, Boule A, Caldwell J, et al. Temporal trends in TB notification rates during ART scale-up in Cape Town: an ecological analysis. *J Int AIDS Soc* 2015;18(1):20240. doi:10.7448/IAS.18.1.20240. PMID 26411694
 - Hudson SM, Hudson C. Is GP practice bowel, breast and cervical cancer screening coverage correlated with GP practice list inflation? *J Med Screen* 2026;33(1):1–8 (epub 17 June 2025). doi:10.1177/09691413251347408. PMID 40525516
 - Jick SS, Lieberman ES, Rahman MU, Choi HK. Glucocorticoid use, other associated factors, and the risk of tuberculosis. *Arthritis Rheum* 2006;55(1):19–26. doi:10.1002/art.21705. PMID 16463407
 - Keane J, Gershon S, Wise RP, et al. Tuberculosis associated with infliximab, a tumor necrosis factor α -neutralizing agent. *N Engl J Med* 2001;345(15):1098–104. doi:10.1056/NEJMoa011110. PMID 11596589

- Költringer FA, et al. The social determinants of national tuberculosis incidence rates in 116 countries: a longitudinal ecological study between 2005–2015. *BMC Public Health* 2023;23:337. doi:10.1186/s12889-023-15213-w. PMID 36793018
- Li X, Sheng L, Lou L. Statin use may be associated with reduced active tuberculosis infection: a meta-analysis of observational studies. *Front Med (Lausanne)* 2020;7:121. doi:10.3389/fmed.2020.00121. PMID 32391364
- Lipsitch M, Tchetgen Tchetgen E, Cohen T. Negative controls: a tool for detecting confounding and bias in observational studies. *Epidemiology* 2010;21(3):383–8. doi:10.1097/EDE.0b013e3181d61eeb. PMID 20335814
- Loutet MG, Burman M, Jayasekera N, et al. National roll-out of latent tuberculosis testing and treatment for new migrants in England: a retrospective evaluation in a high-incidence area. *Eur Respir J* 2018;51(1):1701226. doi:10.1183/13993003.01226-2017. PMID 29326327
- Morgenstern H. Ecologic studies in epidemiology: concepts, principles, and methods. *Annu Rev Public Health* 1995;16:61–81. doi:10.1146/annurev.pu.16.050195.000425. PMID 7639884
- Morrison H, et al. Impact of COVID-19 on NHS tuberculosis services: results of a UK-wide survey. *J Infect* 2023;87(1):59–61. doi:10.1016/j.jinf.2023.04.004. PMID 37044162
- National Institute for Health and Care Excellence. Tuberculosis. NICE guideline [NG33]. London: NICE; published 13 January 2016, last updated 16 February 2024. <https://www.nice.org.uk/guidance/ng33> [dates confirmed from search results only; nice.org.uk returned HTTP 403 to automated access. The wording of the LTBI testing recommendation still needs checking by hand against the current text.]
- OpenPrescribing.net, Bennett Institute for Applied Data Science, University of Oxford. Frequently asked questions. 2026. <https://openprescribing.net/faq/> [accessed date to be added; the FAQ asks academic users to cite “OpenPrescribing.net, Bennett Institute for Applied Data Science, University of Oxford, 2026”, confirmed via search results because a direct fetch failed]
- Pealing L, Wing K, Mathur R, et al. Risk of tuberculosis in patients with diabetes: population based cohort study using the UK Clinical Practice Research Datalink. *BMC Med* 2015;13:135. doi:10.1186/s12916-015-0381-9. PMID 26048371
- Prasad V, Jena AB. Prespecified falsification end points: can they validate true observational associations? *JAMA* 2013;309(3):241–2. doi:10.1001/jama.2012.96867. PMID 23321761
- RECOVERY Collaborative Group; Horby P, Lim WS, Emberson JR, et al. Dexamethasone in hospitalized patients with Covid-19. *N Engl J Med* 2021;384(8):693–704. doi:10.1056/NEJMoa2021436. PMID 32678530
- Richards TC, et al. Using OpenPrescribing.net to evaluate neighbourhood-level prescribing of inhalers for asthma and COPD. *Sci Rep* 2025;15:18089. doi:10.1038/s41598-025-02969-x. PMID 40413267
- Santos Silva JMC, Tenreiro S. The log of gravity. *Rev Econ Stat* 2006;88(4):641–58. doi:10.1162/rest.88.4.641 (not indexed in PubMed)
- Sheppard L, Prentice RL, Rossing MA. Design considerations for estimation of exposure effects on disease risk, using aggregate data studies. *Stat Med* 1996;15(17-18):1849–58. doi:10.1002/(SICI)1097-0258(19960915)15:17<1849::AID-SIM396>3.0.CO;2-4. PMID 8888477
- Song HJ, Park H, Park S, Kwon JW. The association between proton pump inhibitor use and the risk of tuberculosis: a case-control study. *Pharmacoepidemiol Drug Saf* 2019;28(6):830–9. doi:10.1002/pds.4773. PMID 30920070
- Thomas HL, Harris RJ, Muzyamba MC, et al. Reduction in tuberculosis incidence in the UK from 2011 to 2015: a population-based study. *Thorax* 2018;73(8):769–75. doi:10.1136/thoraxjnl-2017-211074. PMID 29674389
- Thwaites GE, Nguyen DB, Nguyen HD, et al. Dexamethasone for the treatment of tuberculous meningitis in adolescents and adults. *N Engl J Med* 2004;351(17):1741–51. doi:10.1056/NEJMoa040573. PMID 15496623
- UK Health Security Agency. Tuberculosis in England: 2021 report (presenting data to

end of 2020). London: UKHSA; 2021 (added to GOV.UK 28 October 2021; corrected PDF 30 March 2022). https://assets.publishing.service.gov.uk/media/62441310e90e075f124018e8/TB_annual-report-2021.pdf [The PDF reports “In 2020, 4,125 people were notified with TB in England, with a rate of 7.3 per 100,000 population”. Its suggested citation is “UK Health Security Agency. (2021) Tuberculosis in England: 2020. UK Health Security Agency, London.”]

- UK Health Security Agency. Tuberculosis in England: 2025 report (data up to end of 2024), chapter 1 and supplementary tables 5, 12 and 22; and TB regional reports 2024, supplementary data. London: UKHSA; first published 8 October 2025 (GOV.UK change history; the content API timestamp reads 9 October 2025 BST), last updated 15 July 2026. <https://www.gov.uk/government/publications/tuberculosis-in-england-2025-report> [the GOV.UK page title reads “Tuberculosis in England, 2025 report”. The specific supplementary table numbers and the regional reports publication were not individually checked.]
- UK Health Security Agency. Tuberculosis notifications in England stabilise in 2025. News story, 29 January 2026. <https://www.gov.uk/government/news/tuberculosis-notifications-in-england-stabilise-in-2025> [confirmed: “5,424 people notified compared to 5,487 in 2024 – a decrease of 1.1%”; rate 9.4 per 100,000]
- van Staa TP, Leufkens HG, Abenham L, et al. Use of oral corticosteroids in the United Kingdom. *QJM* 2000;93(2):105–11. doi:10.1093/qjmed/93.2.105. PMID 10700481
- Venkatesan S, et al. Correcting for the inflated adult population denominator in an English nationwide health care cohort: database analysis study. *JMIR Public Health Surveill* 2025;11:e64788. doi:10.2196/64788. PMID 41144579
- Zhang M, He JQ. Impacts of metformin on tuberculosis incidence and clinical outcomes in patients with diabetes: a systematic review and meta-analysis. *Eur J Clin Pharmacol* 2020;76(2):149–59. doi:10.1007/s00228-019-02786-y. PMID 31786617

Supplementary tables

Table S1. Sensitivity analyses, LTLA panel (residence apportionment): IRR per 10% increase (95% CI)

Drug group	primary (t-1)	FE only (t-1)	prior 3 years (t-3..t-1)	+ area trends (t-1)	population + co-vari-ate (t-1)	+ LTBI pro-gramme (t-1)	+ asy-lum sup-port (t-1)	exclude COVID years (t-1)	outcome years 2018-2024 (t-1)	EPD expo-sure only (t-1)	ADQ mea-sure (t-1)
Antituber- (de- scrip- tive)	1.000 (0.997- 1.003)	1.001 (0.998- 1.005)	1.001 (0.997- 1.006)	0.999 (0.996- 1.003)	1.000 (0.997- 1.004)	1.000 (0.997- 1.003)	1.001 (0.997- 1.004)	1.000 (0.996- 1.004)	1.001 (0.998- 1.005)	1.001 (0.997- 1.004)	—
Systemic oral glu- co- corti- coids	0.969 (0.929- 1.012)	0.964 (0.917- 1.014)	0.976 (0.930- 1.025)	0.985 (0.932- 1.042)	0.972 (0.931- 1.014)	0.971 (0.929- 1.015)	0.982 (0.938- 1.028)	0.974 (0.932- 1.018)	0.983 (0.941- 1.028)	0.979 (0.939- 1.021)	—
Oral hy- dro- corti- sone	0.998 (0.985- 1.011)	0.995 (0.980- 1.010)	0.995 (0.979- 1.012)	0.999 (0.982- 1.017)	1.000 (0.987- 1.012)	0.998 (0.986- 1.011)	0.992 (0.979- 1.006)	0.997 (0.982- 1.012)	0.994 (0.978- 1.011)	0.994 (0.982- 1.006)	—

Drug group	primary (t-1)	FE only (t-1)	prior 3 years (t-1)	+ area trends (t-1)	population co-vari-ate (t-1)	+ LTBI pro-gramme (t-1)	+ asy-lum sup-port (t-1)	exclude COVID years (t-1)	outcome years 2018-2024 (t-1)	EPD expo-sure only (t-1)	ADQ mea-sure (t-1)
Oral dex-am-etha-sone	1.003 (0.992-1.014)	1.005 (0.993-1.018)	1.007 (0.992-1.022)	0.995 (0.985-1.005)	1.003 (0.992-1.014)	1.003 (0.993-1.014)	1.002 (0.991-1.014)	1.007 (0.996-1.018)	0.998 (0.986-1.010)	1.003 (0.992-1.013)	—
Inhaled corti-cos-teroids	0.966 (0.920-1.014)	0.940 (0.892-0.990)	0.983 (0.929-1.039)	1.033 (0.981-1.087)	0.969 (0.922-1.018)	0.967 (0.921-1.016)	0.948 (0.905-0.993)	0.961 (0.916-1.009)	0.938 (0.892-0.985)	0.954 (0.912-0.997)	0.972 (0.951-0.993)
Conventional DMARDs	1.002 (0.983-1.021)	1.004 (0.983-1.024)	1.003 (0.982-1.024)	1.015 (0.979-1.053)	1.001 (0.982-1.020)	1.001 (0.983-1.021)	0.996 (0.979-1.014)	1.004 (0.985-1.023)	1.003 (0.979-1.026)	1.000 (0.983-1.016)	—
Transplan-tim-muno-sup-pres-sants	0.999 (0.993-1.005)	0.997 (0.991-1.003)	1.000 (0.993-1.007)	1.000 (0.993-1.008)	0.999 (0.993-1.005)	0.998 (0.993-1.004)	0.999 (0.993-1.005)	1.001 (0.994-1.007)	1.001 (0.994-1.008)	1.000 (0.994-1.006)	—
Proton pump in-hibitors	0.996 (0.955-1.039)	0.968 (0.921-1.018)	0.997 (0.954-1.042)	0.962 (0.903-1.024)	0.998 (0.955-1.042)	1.002 (0.960-1.046)	0.982 (0.939-1.027)	0.992 (0.952-1.035)	0.936 (0.885-0.991)	0.984 (0.943-1.028)	0.980 (0.917-1.047)
Statins	0.992 (0.952-1.034)	0.968 (0.924-1.013)	0.993 (0.950-1.037)	0.973 (0.919-1.029)	0.991 (0.951-1.033)	0.998 (0.957-1.040)	1.000 (0.956-1.045)	0.983 (0.944-1.024)	0.986 (0.929-1.047)	0.999 (0.958-1.041)	0.983 (0.946-1.023)
Metformin	0.966 (0.934-0.998)	0.947 (0.914-0.981)	0.970 (0.934-1.007)	0.947 (0.906-0.990)	0.967 (0.935-0.999)	0.971 (0.938-1.005)	0.971 (0.935-1.009)	0.963 (0.932-0.994)	0.975 (0.926-1.027)	0.970 (0.937-1.005)	0.982 (0.931-1.037)
Insulins	0.966 (0.929-1.005)	0.964 (0.923-1.007)	0.974 (0.931-1.018)	1.016 (0.945-1.093)	0.971 (0.931-1.013)	0.969 (0.930-1.008)	0.968 (0.921-1.017)	0.962 (0.923-1.002)	0.979 (0.912-1.051)	0.973 (0.930-1.017)	—
Fluoroquinolones	1.005 (0.992-1.017)	1.008 (0.994-1.023)	1.007 (0.993-1.020)	1.006 (0.991-1.020)	1.006 (0.993-1.019)	1.004 (0.992-1.017)	1.003 (0.990-1.017)	1.000 (0.987-1.013)	0.993 (0.973-1.012)	1.002 (0.989-1.015)	1.001 (0.988-1.013)
All anti-bac-teri-als	0.996 (0.955-1.039)	1.001 (0.952-1.052)	1.009 (0.963-1.058)	0.979 (0.940-1.020)	1.000 (0.959-1.041)	0.993 (0.950-1.037)	0.990 (0.950-1.033)	1.011 (0.965-1.059)	0.951 (0.909-0.995)	0.992 (0.953-1.033)	0.990 (0.952-1.029)
Vitamin D	1.007 (0.987-1.027)	1.010 (0.988-1.033)	1.008 (0.987-1.029)	0.983 (0.959-1.007)	1.009 (0.990-1.029)	1.004 (0.984-1.025)	1.006 (0.983-1.030)	1.016 (0.995-1.037)	0.990 (0.959-1.021)	1.005 (0.985-1.026)	—
Levothyroxine (negative control)	0.962 (0.933-0.992)	0.943 (0.914-0.973)	0.963 (0.933-0.995)	0.954 (0.902-1.009)	0.960 (0.930-0.990)	0.967 (0.936-1.000)	0.968 (0.935-1.002)	0.961 (0.928-0.994)	0.968 (0.924-1.014)	0.968 (0.936-1.001)	0.950 (0.909-0.994)

Table S2. Distributed lag model (t, t-1, t-2 in one model), LTLA residence

Drug group	t (same year)	t-1	t-2
Antituberculosis (descriptive)	0.999 (0.996-1.003)	1.001 (0.997-1.004)	1.000 (0.997-1.003)
Systemic oral glucocorticoids	0.974 (0.933-1.016)	0.984 (0.933-1.038)	1.006 (0.961-1.053)
Oral hydrocortisone	1.009 (0.993-1.026)	0.990 (0.970-1.010)	1.001 (0.982-1.019)
Oral dexamethasone	1.000 (0.991-1.009)	1.003 (0.993-1.013)	1.000 (0.990-1.010)
Inhaled corticosteroids	0.922 (0.874-0.973)	1.001 (0.927-1.082)	1.037 (0.962-1.118)
Conventional DMARDs	1.015 (0.978-1.053)	0.981 (0.924-1.041)	1.009 (0.966-1.054)
Transplant immunosuppressants	1.001 (0.990-1.011)	0.995 (0.983-1.008)	1.004 (0.995-1.013)
Proton pump inhibitors	1.002 (0.926-1.084)	1.012 (0.928-1.103)	0.981 (0.905-1.063)
Statins	1.028 (0.946-1.118)	0.978 (0.892-1.073)	0.989 (0.920-1.064)
Metformin	1.021 (0.952-1.096)	0.955 (0.885-1.031)	0.993 (0.940-1.049)
Insulins	0.966 (0.901-1.035)	0.988 (0.907-1.076)	1.007 (0.943-1.076)
Fluoroquinolones	1.002 (0.986-1.019)	0.998 (0.981-1.016)	1.005 (0.990-1.020)
All antibacterials	1.032 (0.998-1.066)	0.950 (0.908-0.994)	1.030 (0.982-1.080)
Vitamin D	1.017 (0.981-1.054)	0.982 (0.937-1.030)	1.011 (0.979-1.044)
Levothyroxine (negative control)	1.006 (0.934-1.084)	0.979 (0.894-1.072)	0.976 (0.905-1.053)

Table S3. Hospital medicines: minimum detectable effects (previous-year exposure) versus illustrative expected effects

Level	Drug group	Scenario (illustrative RR)	Prevalence (SCMD DDD-years per resident)	Expected change per 10%	MDE	MDE ÷ expected	MDE ÷ expected, attenuated by positive-control elasticity
UTLA	TNF inhibitors	RR 4 (illustrative, before LTBI screening)	0.34%	0.100%	2.6%	26	72
UTLA	TNF inhibitors	RR 1.5 (illustrative, with LTBI screening; probably nearer 1, so favours detection)	0.34%	0.017%	2.6%	154	426

Level	Drug group	Scenario (illustrative RR)	Prevalence (SCMD DDD-years per resident)	Expected change per 10%	MDE	MDE ÷ expected	MDE ÷ expected, attenuated by positive-control elasticity
UTLA	IL-6 inhibitors and abatacept	RR 2	0.02%	0.002%	1.2%	520	1443
UTLA	JAK inhibitors (rheumatology)	RR 2	0.04%	0.004%	1.0%	235	652
UTLA	Rituximab	RR 1.5	0.09%	0.005%	2.0%	420	1165
UTLA	Calcineurin/mTOR inhibitors	RR 10 (solid organ transplant)	0.07%	0.066%	1.7%	26	72
UTLA	Antiproliferatives	RR 2	0.16%	0.016%	2.6%	158	439
LTLA	TNF inhibitors	RR 4 (illustrative, before LTBI screening)	0.34%	0.100%	2.3%	23	58
LTLA	TNF inhibitors	RR 1.5 (illustrative, with LTBI screening; probably nearer 1, so favours detection)	0.34%	0.017%	2.3%	137	343
LTLA	IL-6 inhibitors and abatacept	RR 2	0.02%	0.002%	1.2%	522	1305
LTLA	JAK inhibitors (rheumatology)	RR 2	0.04%	0.004%	1.0%	218	544
LTLA	Rituximab	RR 1.5	0.09%	0.005%	2.2%	470	1174
LTLA	Calcineurin/mTOR inhibitors	RR 10 (solid organ transplant)	0.07%	0.066%	1.6%	25	61
LTLA	Antiproliferatives	RR 2	0.16%	0.016%	2.2%	138	345

Relative risks are illustrative values, not estimates. DDD-years per resident approximate the proportion treated only for continuous maintenance therapy. The attenuated column divides the expected effect by the positive control's same-year elasticity (UTLA 0.36, 95% CI 0.10–0.62; LTLA 0.40, 95% CI 0.14–0.66). This assumes the same linkage attenuation for previous-year exposure and for drugs with other supply routes (homecare, specialist centres), and the elasticity also reflects variation in drug volume per person treated, so the attenuated ratios are illustrative.

Table S4. Permutation-based power simulation (systemic oral glucocorticoids, exposure t–1)

Level	Scenario	Replicates	Two-sided rejection	Power, correct direction (95% MC interval)	Analytic power at real-data SE	Median estimated IRR per 10%	Median replicate SE	Null empirical SD	Real-data SE
LTLA	No effect	500	7.8%	3.8% (±1.7)	5.1%	1.002	0.0161	0.0166	0.0218
LTLA	Published RR 4.9	300	8.3%	6.0% (±2.7)	5.2%	1.003	0.0161	0.0166	0.0218
LTLA	RR 25	300	17.7%	17.7% (±4.3)	12.2%	1.017	0.0161	0.0166	0.0218
LTLA	RR 100	300	71.0%	71.0% (±5.1)	52.1%	1.045	0.0161	0.0166	0.0218
LTLA	IRR 1.02 per 10%	300	21.7%	21.7% (±4.7)	14.3%	1.019	0.0161	0.0166	0.0218
LTLA	IRR 1.05 per 10%	300	83.0%	83.0% (±4.3)	64.6%	1.052	0.0161	0.0166	0.0218
LTLA	IRR 1.10 per 10%	300	100.0%	100.0% (±0.0)	99.4%	1.102	0.0161	0.0166	0.0218
LTLA	IRR 1.05 per 10%, acting via concurrent year	300	61.7%	61.7% (±5.5)	41.0%	1.039	0.0161	0.0166	0.0218
LTLA	IRR 1.10 per 10%, acting via concurrent year	300	99.3%	99.3% (±0.9)	94.2%	1.080	0.0161	0.0166	0.0218

Level	Scenario	Replicates	Two-sided rejection	Power, correct direction (95% MC interval)	Analytic power at real-data SE	Median estimated IRR per 10%	Median replicate SE	Null empirical SD	Real-data SE
UTLA	No effect	500	7.6%	4.0% (± 1.7)	5.0%	1.000	0.0167	0.0171	0.0249
UTLA	Published RR 4.9	300	8.0%	6.7% (± 2.8)	5.1%	1.003	0.0167	0.0171	0.0249
UTLA	RR 25	300	20.0%	19.3% (± 4.5)	11.5%	1.019	0.0167	0.0171	0.0249
UTLA	RR 100	300	66.7%	66.7% (± 5.3)	46.2%	1.047	0.0167	0.0171	0.0249
UTLA	IRR 1.02 per 10%	300	24.7%	24.7% (± 4.9)	13.8%	1.022	0.0167	0.0171	0.0249
UTLA	IRR 1.05 per 10%	300	78.0%	78.0% (± 4.7)	51.0%	1.051	0.0167	0.0171	0.0249
UTLA	IRR 1.10 per 10%	300	100.0%	100.0% (± 0.0)	96.9%	1.100	0.0167	0.0171	0.0249
UTLA	IRR 1.05 per 10%, acting via concurrent year	300	65.7%	65.7% (± 5.4)	38.9%	1.043	0.0167	0.0171	0.0249
UTLA	IRR 1.10 per 10%, acting via concurrent year	300	99.3%	99.3% (± 0.9)	90.7%	1.085	0.0167	0.0171	0.0249

Within the simulation, replicate SEs matched the spread of null estimates, so the test was approximately calibrated (slightly anti-conservative). Real-data SEs were larger than in the simulation because permuting exposure trajectories across areas breaks their alignment with each area's own notification trends; simulated power is therefore optimistic. Analytic power uses the median simulated estimate as the effect and the real-data SE.

Table S5. Regional models (9 regions): TB notifications by place of birth and age vs primary care prescribing

Outcome	Exposure	Analysis	Lag (years; negative = lead)	IRR per 10% (t(8) 95% CI)	p (cluster t8)	p (randomi- sation)
TB, UK born	oral_glucocorticoids	lagged exposure	1	1.19 (0.87–1.62)	0.233	0.254
TB, UK born	oral_glucocorticoids	lagged exposure	2	1.20 (0.80–1.80)	0.323	0.386
TB, UK born	oral_glucocorticoids	lagged exposure	3	1.10 (0.67–1.79)	0.678	0.714
TB, UK born	oral_glucocorticoids	lagged, excluding 2020-21	1	1.19 (0.87–1.63)	0.244	0.280
TB, UK born	oral_glucocorticoids	lagged, excluding 2020-21	2	1.20 (0.80–1.80)	0.323	0.372
TB, UK born	oral_glucocorticoids	lagged, excluding 2020-21	3	1.09 (0.67–1.78)	0.686	0.714
TB, UK born	oral_glucocorticoids	lead (falsifi- cation)	-1	1.20 (0.78–1.84)	0.356	0.374
TB, UK born	oral_glucocorticoids	lead (falsifi- cation)	-2	1.09 (0.67–1.78)	0.695	0.686
TB, UK born	oral_glucocorticoids	lead (falsifi- cation)	-3	1.09 (0.83–1.44)	0.484	0.536
TB, UK born	mg_pred_equivalents	lagged exposure	1	1.25 (0.95–1.65)	0.097	0.122
TB, UK born	mg_pred_equivalents	lagged exposure	2	1.33 (0.96–1.84)	0.077	0.162
TB, UK born	mg_pred_equivalents	lagged exposure	3	1.33 (0.94–1.89)	0.098	0.238
TB, UK born	mg_pred_equivalents	lagged, excluding 2020-21	1	1.27 (0.94–1.72)	0.105	0.114
TB, UK born	mg_pred_equivalents	lagged, excluding 2020-21	2	1.32 (0.97–1.80)	0.072	0.140
TB, UK born	mg_pred_equivalents	lagged, excluding 2020-21	3	1.30 (0.91–1.85)	0.134	0.286
TB, UK born	mg_pred_equivalents	lead (falsifi- cation)	-1	1.17 (0.85–1.60)	0.292	0.274
TB, UK born	mg_pred_equivalents	lead (falsifi- cation)	-2	0.91 (0.70–1.19)	0.446	0.436
TB, UK born	mg_pred_equivalents	lead (falsifi- cation)	-3	0.92 (0.82–1.03)	0.120	0.084
TB, UK born	levothyroxine	lagged exposure	1	0.96 (0.85–1.08)	0.407	0.518
TB, UK born	levothyroxine	lagged exposure	2	0.91 (0.80–1.03)	0.127	0.246
TB, UK born	levothyroxine	lagged exposure	3	0.90 (0.77–1.04)	0.139	0.258
TB, UK born	levothyroxine	lagged, excluding 2020-21	1	0.95 (0.84–1.08)	0.366	0.478

Outcome	Exposure	Analysis	Lag (years; negative = lead)	IRR per 10% (t(8) 95% CI)	p (cluster t8)	p (randomi- sation)
TB, UK born	levothyroxine	lagged, excluding 2020-21	2	0.91 (0.80-1.04)	0.152	0.288
TB, UK born	levothyroxine	lagged, excluding 2020-21	3	0.90 (0.76-1.05)	0.154	0.282
TB, UK born	levothyroxine	lead (falsifi- cation)	-1	0.93 (0.81-1.07)	0.293	0.428
TB, UK born	levothyroxine	lead (falsifi- cation)	-2	0.97 (0.82-1.14)	0.636	0.710
TB, UK born	levothyroxine	lead (falsifi- cation)	-3	0.96 (0.80-1.15)	0.624	0.720
TB, Non-UK born	oral_glucocorticoids	lagged exposure	1	0.95 (0.72-1.25)	0.678	0.716
TB, Non-UK born	oral_glucocorticoids	lagged exposure	2	1.00 (0.81-1.23)	0.982	0.974
TB, Non-UK born	oral_glucocorticoids	lagged exposure	3	1.03 (0.83-1.28)	0.725	0.766
TB, Non-UK born	oral_glucocorticoids	lagged, excluding 2020-21	1	0.98 (0.72-1.32)	0.853	0.874
TB, Non-UK born	oral_glucocorticoids	lagged, excluding 2020-21	2	1.03 (0.84-1.26)	0.785	0.812
TB, Non-UK born	oral_glucocorticoids	lagged, excluding 2020-21	3	1.06 (0.86-1.32)	0.523	0.600
TB, Non-UK born	oral_glucocorticoids	lead (falsifi- cation)	-1	1.13 (0.88-1.44)	0.306	0.358
TB, Non-UK born	oral_glucocorticoids	lead (falsifi- cation)	-2	1.33 (1.00-1.76)	0.049	0.070
TB, Non-UK born	oral_glucocorticoids	lead (falsifi- cation)	-3	1.49 (1.17-1.89)	0.005	0.012
TB, Non-UK born	mg_pred_equivalents	lagged exposure	1	0.99 (0.77-1.26)	0.906	0.934
TB, Non-UK born	mg_pred_equivalents	lagged exposure	2	1.08 (0.91-1.27)	0.343	0.544
TB, Non-UK born	mg_pred_equivalents	lagged exposure	3	1.11 (0.94-1.30)	0.179	0.384
TB, Non-UK born	mg_pred_equivalents	lagged, excluding 2020-21	1	1.02 (0.78-1.33)	0.884	0.918

Outcome	Exposure	Analysis	Lag (years; negative = lead)	IRR per 10% (t(8) 95% CI)	p (cluster t8)	p (randomi- sation)
TB, Non-UK born	mg_pred_equivalent	lagged, excluding 2020-21	2	1.07 (0.92-1.26)	0.327	0.542
TB, Non-UK born	mg_pred_equivalent	lagged, excluding 2020-21	3	1.13 (0.94-1.35)	0.158	0.378
TB, Non-UK born	mg_pred_equivalent	lead (falsifi- cation)	-1	1.15 (0.91-1.46)	0.208	0.332
TB, Non-UK born	mg_pred_equivalent	lead (falsifi- cation)	-2	1.31 (0.94-1.83)	0.101	0.168
TB, Non-UK born	mg_pred_equivalent	lead (falsifi- cation)	-3	1.29 (1.03-1.60)	0.030	0.040
TB, Non-UK born	levothyroxine	lagged exposure	1	1.02 (0.90-1.16)	0.695	0.778
TB, Non-UK born	levothyroxine	lagged exposure	2	1.02 (0.88-1.17)	0.767	0.854
TB, Non-UK born	levothyroxine	lagged exposure	3	1.01 (0.88-1.17)	0.835	0.866
TB, Non-UK born	levothyroxine	lagged, excluding 2020-21	1	1.02 (0.90-1.16)	0.682	0.798
TB, Non-UK born	levothyroxine	lagged, excluding 2020-21	2	1.02 (0.88-1.17)	0.771	0.844
TB, Non-UK born	levothyroxine	lagged, excluding 2020-21	3	1.01 (0.88-1.17)	0.835	0.860
TB, Non-UK born	levothyroxine	lead (falsifi- cation)	-1	1.00 (0.86-1.15)	0.947	0.964
TB, Non-UK born	levothyroxine	lead (falsifi- cation)	-2	1.01 (0.88-1.16)	0.839	0.878
TB, Non-UK born	levothyroxine	lead (falsifi- cation)	-3	1.10 (0.91-1.31)	0.279	0.476
TB, UK born aged 65+	oral_glucocorticoids	lagged exposure	1	1.05 (0.75-1.46)	0.750	0.790
TB, UK born aged 65+	oral_glucocorticoids	lagged exposure	2	1.06 (0.75-1.50)	0.714	0.712
TB, UK born aged 65+	oral_glucocorticoids	lagged exposure	3	0.85 (0.48-1.51)	0.539	0.578

Outcome	Exposure	Analysis	Lag (years; negative = lead)	IRR per 10% (t(8) 95% CI)	p (cluster t8)	p (randomi- sation)
TB, UK born aged 65+	oral_glucocorticoids	lagged, excluding 2020-21	1	1.09 (0.80-1.49)	0.524	0.592
TB, UK born aged 65+	oral_glucocorticoids	lagged, excluding 2020-21	2	1.10 (0.79-1.53)	0.512	0.472
TB, UK born aged 65+	oral_glucocorticoids	lagged, excluding 2020-21	3	0.88 (0.53-1.45)	0.572	0.572
TB, UK born aged 65+	oral_glucocorticoids	lead (falsifi- cation)	-1	1.27 (0.88-1.85)	0.173	0.198
TB, UK born aged 65+	oral_glucocorticoids	lead (falsifi- cation)	-2	1.68 (1.04-2.70)	0.036	0.044
TB, UK born aged 65+	oral_glucocorticoids	lead (falsifi- cation)	-3	1.53 (0.93-2.54)	0.086	0.108
TB, UK born aged 65+	mg_pred_equivalents	lagged exposure	1	1.06 (0.76-1.48)	0.679	0.712
TB, UK born aged 65+	mg_pred_equivalents	lagged exposure	2	1.13 (0.83-1.54)	0.388	0.478
TB, UK born aged 65+	mg_pred_equivalents	lagged exposure	3	0.94 (0.56-1.58)	0.803	0.844
TB, UK born aged 65+	mg_pred_equivalents	lagged, excluding 2020-21	1	1.12 (0.77-1.61)	0.513	0.566
TB, UK born aged 65+	mg_pred_equivalents	lagged, excluding 2020-21	2	1.12 (0.80-1.56)	0.452	0.490
TB, UK born aged 65+	mg_pred_equivalents	lagged, excluding 2020-21	3	0.88 (0.49-1.60)	0.640	0.618
TB, UK born aged 65+	mg_pred_equivalents	lead (falsifi- cation)	-1	1.28 (0.89-1.85)	0.151	0.204
TB, UK born aged 65+	mg_pred_equivalents	lead (falsifi- cation)	-2	1.58 (0.92-2.73)	0.089	0.086
TB, UK born aged 65+	mg_pred_equivalents	lead (falsifi- cation)	-3	1.30 (0.78-2.19)	0.271	0.320
TB, UK born aged 65+	levothyroxine	lagged exposure	1	1.23 (1.00-1.51)	0.051	0.074
TB, UK born aged 65+	levothyroxine	lagged exposure	2	1.19 (0.95-1.48)	0.111	0.168

Outcome	Exposure	Analysis	Lag (years; negative = lead)	IRR per 10% (t(8) 95% CI)	p (cluster t8)	p (randomi- sation)
TB, UK born aged 65+	levothyroxine	lagged exposure	3	1.06 (0.89–1.27)	0.463	0.534
TB, UK born aged 65+	levothyroxine	lagged, excluding 2020-21	1	1.22 (0.99–1.50)	0.055	0.068
TB, UK born aged 65+	levothyroxine	lagged, excluding 2020-21	2	1.20 (0.96–1.50)	0.098	0.152
TB, UK born aged 65+	levothyroxine	lagged, excluding 2020-21	3	1.09 (0.91–1.30)	0.304	0.406
TB, UK born aged 65+	levothyroxine lead (falsifi- cation)		-1	1.38 (1.05–1.80)	0.025	0.038
TB, UK born aged 65+	levothyroxine lead (falsifi- cation)		-2	1.20 (0.85–1.71)	0.262	0.382
TB, UK born aged 65+	levothyroxine lead (falsifi- cation)		-3	1.29 (0.85–1.97)	0.195	0.308

Randomisation p-values compare the cluster-robust t statistic with 499 permutations of whole regional exposure histories across regions, valid under exchangeability of those histories; $p = (1 + \text{count}) / (1 + 499)$, so the minimum is 0.002.

Table S6. Regional models: lag 2 and lead 2 estimated jointly

Outcome	Exposure	Regions ex- cluded	Years	IRR lag (per 10%)	IRR lead (per 10%)	Lag ÷ lead	p differ- ence	p ran- domisa- tion (lag)
TB, UK born	oral_glucocorticoids	none	2013- 2022	1.26	0.95	1.33	0.151	0.460
TB, UK born	oral_glucocorticoids	London	2013- 2022	0.99	0.97	1.02	0.919	0.964
TB, UK born	mg_pred_equivalent	none	2013- 2022	1.45	0.88	1.65	0.035	0.232
TB, UK born	mg_pred_equivalent	London	2013- 2022	1.12	0.90	1.25	0.430	0.730
TB, UK born	levothyroxine	none	2013- 2022	0.94	0.91	1.03	0.953	0.830
TB, UK born	levothyroxine	London	2013- 2022	1.20	1.23	0.97	0.945	0.634
TB, Non-UK born	oral_glucocorticoids	none	2013- 2022	0.98	1.38	0.71	0.181	0.916
TB, Non-UK born	mg_pred_equivalent	none	2013- 2022	1.14	1.27	0.89	0.653	0.502

Outcome	Exposure	Regions ex- cluded	Years	IRR lag (per 10%)	IRR lead (per 10%)	Lag ÷ lead	p differ- ence	p ran- domisa- tion (lag)
TB, Non-UK born	levothyroxine	none	2013- 2022	0.95	1.00	0.94	0.903	0.830
TB, UK born aged 65+	oral_glucocorticoids	none	2013- 2022	0.87	1.79	0.49	0.065	0.506
TB, UK born aged 65+	mg_pred_equivalent	none	2013- 2022	1.08	1.57	0.69	0.306	0.786
TB, UK born aged 65+	levothyroxine	none	2013- 2022	1.00	1.09	0.91	0.833	1.000

Table S7. Hospital medicines: share of DDD from trusts matched to catchments directly, via successor trusts, or unmatched

Drug group	Year	Direct	Successor	Unmatched
Antiproliferatives	2019	90.9%	8.4%	0.7%
Antiproliferatives	2020	93.8%	4.8%	1.4%
Antiproliferatives	2021	96.4%	2.4%	1.2%
Antiproliferatives	2022	97.7%	0.7%	1.6%
Antiproliferatives	2023	97.6%	0.5%	1.9%
Antiproliferatives	2024	97.7%	0.4%	1.9%
Active-TB treatment (pyrazinamide DDD, including fixed-dose combinations; positive control)	2019	90.9%	8.9%	0.3%
Active-TB treatment (pyrazinamide DDD, including fixed-dose combinations; positive control)	2020	94.2%	5.7%	0.1%
Active-TB treatment (pyrazinamide DDD, including fixed-dose combinations; positive control)	2021	97.4%	2.5%	0.1%

Drug group	Year	Direct	Successor	Unmatched
Active-TB treatment (pyrazinamide DDD, including fixed-dose combinations; positive control)	2022	98.8%	1.1%	0.0%
Active-TB treatment (pyrazinamide DDD, including fixed-dose combinations; positive control)	2023	98.9%	1.0%	0.0%
Active-TB treatment (pyrazinamide DDD, including fixed-dose combinations; positive control)	2024	97.4%	1.1%	1.5%
Rifamycin/isoniazid (active and latent TB, other infections)	2019	90.6%	9.0%	0.3%
Rifamycin/isoniazid (active and latent TB, other infections)	2020	95.1%	4.6%	0.3%
Rifamycin/isoniazid (active and latent TB, other infections)	2021	97.7%	2.1%	0.2%
Rifamycin/isoniazid (active and latent TB, other infections)	2022	99.0%	0.8%	0.2%
Rifamycin/isoniazid (active and latent TB, other infections)	2023	99.0%	0.8%	0.2%
Rifamycin/isoniazid (active and latent TB, other infections)	2024	99.2%	0.6%	0.1%
Dexamethasone and hydrocortisone (descriptive)	2019	91.5%	8.1%	0.4%
Dexamethasone and hydrocortisone (descriptive)	2020	95.2%	4.4%	0.4%

Drug group	Year	Direct	Successor	Unmatched
Dexamethasone and hydrocortisone (descriptive)	2021	97.5%	2.1%	0.4%
Dexamethasone and hydrocortisone (descriptive)	2022	98.6%	1.0%	0.3%
Dexamethasone and hydrocortisone (descriptive)	2023	99.0%	0.7%	0.3%
Dexamethasone and hydrocortisone (descriptive)	2024	99.2%	0.5%	0.3%
IL-6 inhibitors and abatacept	2019	90.1%	8.7%	1.2%
IL-6 inhibitors and abatacept	2020	93.9%	4.9%	1.2%
IL-6 inhibitors and abatacept	2021	96.3%	2.8%	0.9%
IL-6 inhibitors and abatacept	2022	97.8%	1.0%	1.2%
IL-6 inhibitors and abatacept	2023	97.9%	0.5%	1.5%
IL-6 inhibitors and abatacept	2024	97.8%	0.4%	1.8%
JAK inhibitors (rheumatology)	2019	90.1%	8.9%	0.9%
JAK inhibitors (rheumatology)	2020	94.8%	4.4%	0.9%
JAK inhibitors (rheumatology)	2021	97.1%	2.5%	0.4%
JAK inhibitors (rheumatology)	2022	98.5%	1.1%	0.4%
JAK inhibitors (rheumatology)	2023	98.9%	0.4%	0.7%
JAK inhibitors (rheumatology)	2024	98.4%	0.3%	1.3%
Low-TB-risk biologics (negative control)	2019	91.9%	7.9%	0.2%
Low-TB-risk biologics (negative control)	2020	96.0%	3.9%	0.1%
Low-TB-risk biologics (negative control)	2021	97.9%	2.0%	0.1%

Drug group	Year	Direct	Successor	Unmatched
Low-TB-risk biologics (negative control)	2022	99.1%	0.8%	0.1%
Low-TB-risk biologics (negative control)	2023	99.5%	0.4%	0.1%
Low-TB-risk biologics (negative control)	2024	99.6%	0.2%	0.2%
Levetiracetam (negative control)	2019	87.6%	7.5%	4.9%
Levetiracetam (negative control)	2020	91.3%	4.2%	4.5%
Levetiracetam (negative control)	2021	93.5%	2.3%	4.2%
Levetiracetam (negative control)	2022	94.7%	1.2%	4.0%
Levetiracetam (negative control)	2023	94.8%	0.9%	4.3%
Levetiracetam (negative control)	2024	95.0%	0.7%	4.3%
Rituximab	2019	91.2%	8.4%	0.4%
Rituximab	2020	95.1%	4.4%	0.5%
Rituximab	2021	97.5%	2.1%	0.5%
Rituximab	2022	98.7%	0.8%	0.5%
Rituximab	2023	99.0%	0.4%	0.5%
Rituximab	2024	99.2%	0.2%	0.6%
Systemic glucocorticoids without dexamethasone/hydrocortisone (prednisolone-equivalent mg)	2019	89.8%	9.5%	0.7%
Systemic glucocorticoids without dexamethasone/hydrocortisone (prednisolone-equivalent mg)	2020	94.1%	5.1%	0.8%

Drug group	Year	Direct	Successor	Unmatched
Systemic glucocorticoids without dexamethasone/hydrocortisone (prednisolone-equivalent mg)	2021	97.1%	2.3%	0.6%
Systemic glucocorticoids without dexamethasone/hydrocortisone (prednisolone-equivalent mg)	2022	98.2%	1.1%	0.6%
Systemic glucocorticoids without dexamethasone/hydrocortisone (prednisolone-equivalent mg)	2023	98.6%	0.8%	0.6%
Systemic glucocorticoids without dexamethasone/hydrocortisone (prednisolone-equivalent mg)	2024	98.8%	0.5%	0.6%
TNF inhibitors	2019	91.2%	8.2%	0.6%
TNF inhibitors	2020	95.3%	4.1%	0.6%
TNF inhibitors	2021	97.6%	2.0%	0.4%
TNF inhibitors	2022	98.5%	0.9%	0.6%
TNF inhibitors	2023	98.6%	0.4%	0.9%
TNF inhibitors	2024	98.9%	0.3%	0.9%
Calcineurin/mTOR inhibitors	2019	92.3%	7.6%	0.1%
Calcineurin/mTOR inhibitors	2020	96.2%	3.6%	0.1%
Calcineurin/mTOR inhibitors	2021	99.2%	0.6%	0.1%
Calcineurin/mTOR inhibitors	2022	99.8%	0.1%	0.1%
Calcineurin/mTOR inhibitors	2023	99.9%	0.0%	0.1%
Calcineurin/mTOR inhibitors	2024	99.9%	0.0%	0.1%

Table S2b. Previous, same and following year in one model, LTLA residence: IRR per 10% (95% CI)

Drug group	t-1	t (same year)	t+1	p, t-1 vs t+1	Correlation of t-1 and t+1 estimates
Antituberculosis (descriptive)	1.000 (0.997-1.004)	0.999 (0.995-1.003)	1.002 (0.999-1.006)	0.434	0.11
Systemic oral glucocorti- coids	0.980 (0.926-1.036)	1.021 (0.971-1.073)	0.943 (0.898-0.990)	0.337	-0.11
Oral hydro- cortisone	0.989 (0.969-1.010)	1.012 (0.990-1.034)	1.001 (0.982-1.020)	0.421	-0.05
Oral dexam- ethasone	1.002 (0.991-1.012)	1.001 (0.991-1.011)	1.003 (0.993-1.012)	0.905	0.01
Inhaled corti- costeroids	1.036 (0.965-1.111)	1.043 (0.965-1.127)	0.861 (0.809-0.917)	0.000	0.04
Conventional DMARDs	0.985 (0.949-1.022)	1.051 (1.005-1.100)	0.964 (0.920-1.010)	0.487	-0.06
Transplant immunosup- pressants	0.999 (0.988-1.011)	1.000 (0.990-1.010)	0.999 (0.990-1.008)	0.954	-0.07
Proton pump inhibitors	0.974 (0.911-1.041)	1.128 (1.035-1.229)	0.910 (0.835-0.991)	0.246	-0.12
Statins	0.956 (0.891-1.026)	1.085 (0.999-1.179)	0.953 (0.874-1.039)	0.962	-0.16
Metformin	0.945 (0.883-1.012)	1.047 (0.971-1.128)	0.976 (0.904-1.053)	0.585	-0.23
Insulins	0.991 (0.924-1.062)	1.079 (1.000-1.164)	0.888 (0.829-0.950)	0.039	-0.14
Fluoroquinolones	1.001 (0.984-1.019)	1.019 (0.999-1.039)	0.985 (0.968-1.001)	0.180	-0.02
All antibacterials	0.969 (0.924-1.015)	1.065 (1.012-1.122)	0.967 (0.926-1.010)	0.962	0.17
Vitamin D	0.995 (0.961-1.030)	1.012 (0.966-1.061)	1.000 (0.963-1.038)	0.842	0.12
Levothyroxine (negative control)	0.954 (0.897-1.015)	1.065 (0.973-1.166)	0.941 (0.856-1.034)	0.818	-0.19

Table S3b. Hospital medicines: sensitivity analyses, IRR per 10% increase (95% CI)

Level	Drug group	lag (t-1)	lag (t-1), excluding outcome years 2020-21	lag (t-1), elective catch- ments	concurrent (t)	concurrent (t), clustered by principal trust	concurrent (t), elective catch- ments
UTLA	Active-TB treat- ment (pyrazi- namide DDD, including fixed- dose combina- tions; positive control)	1.001 (0.993- 1.010)	1.000 (0.986- 1.013)	1.001 (0.993- 1.010)	1.035 (1.009- 1.061)	1.035 (1.009- 1.062)	1.035 (1.009- 1.062)
UTLA	Rifamycin/isoniazid (active and latent TB, other in- fections)	1.005 (0.993- 1.018)	1.000 (0.979- 1.022)	1.006 (0.993- 1.019)	1.036 (1.021- 1.050)	1.036 (1.020- 1.051)	1.036 (1.022- 1.050)
UTLA	TNF inhibitors	0.989 (0.972- 1.007)	0.970 (0.928- 1.013)	0.990 (0.972- 1.008)	0.996 (0.979- 1.013)	0.996 (0.980- 1.012)	0.996 (0.979- 1.013)
UTLA	IL-6 inhibitors and abat- accept	1.001 (0.992- 1.009)	0.993 (0.976- 1.011)	1.001 (0.993- 1.009)	1.000 (0.990- 1.010)	1.000 (0.991- 1.009)	1.000 (0.990- 1.010)
UTLA	JAK inhibitors (rheuma- tology)	0.994 (0.987- 1.002)	0.978 (0.961- 0.996)	0.995 (0.987- 1.002)	0.993 (0.987- 0.999)	0.993 (0.987- 1.000)	0.993 (0.987- 0.999)
UTLA	Rituximab	0.998 (0.984- 1.012)	0.986 (0.959- 1.015)	0.999 (0.985- 1.013)	0.992 (0.978- 1.005)	0.992 (0.979- 1.004)	0.992 (0.979- 1.006)
UTLA	Calcineurin/mTOR inhibitors	0.998 (0.996- 1.020)	1.012 (0.989- 1.035)	1.008 (0.996- 1.020)	0.994 (0.985- 1.004)	0.994 (0.985- 1.004)	0.994 (0.985- 1.004)
UTLA	Antiproliferative	0.996 (0.978- 1.014)	1.004 (0.973- 1.036)	0.996 (0.978- 1.015)	0.993 (0.978- 1.008)	0.993 (0.978- 1.008)	0.993 (0.978- 1.009)
UTLA	Systemic glucocor- ticoids without dexam- etha- sone/hydrocortisone (prednisolone- equivalent mg)	1.024 (1.001- 1.046)	1.018 (0.969- 1.069)	1.025 (1.003- 1.048)	1.009 (0.986- 1.032)	1.009 (0.985- 1.033)	1.010 (0.987- 1.034)

Level	Drug group	lag (t-1)	lag (t-1), excluding outcome years 2020-21	lag (t-1), elective catch- ments	concurrent (t)	concurrent (t), clustered by principal trust	concurrent (t), elective catch- ments
UTLA	Dexamethasone and hydrocortisone (descriptive)	1.008 (0.980- 1.036)	1.021 (0.963- 1.082)	1.009 (0.981- 1.039)	1.005 (0.977- 1.034)	1.005 (0.975- 1.036)	1.008 (0.979- 1.038)
UTLA	Low-TB- risk biologics (negative control)	1.002 (0.986- 1.018)	0.999 (0.958- 1.042)	1.002 (0.985- 1.018)	1.004 (0.990- 1.019)	1.004 (0.990- 1.019)	1.004 (0.989- 1.019)
UTLA	Levetiracetam (negative control)	1.014 (0.986- 1.042)	1.035 (0.978- 1.096)	1.014 (0.986- 1.044)	0.998 (0.975- 1.021)	0.998 (0.974- 1.022)	0.999 (0.976- 1.022)
LTLA	Active-TB treat- ment (pyrazi- namide DDD, including fixed- dose combina- tions; positive control)	1.000 (0.991- 1.008)	0.998 (0.984- 1.012)	1.000 (0.991- 1.009)	1.039 (1.013- 1.065)	1.039 (1.012- 1.066)	1.039 (1.013- 1.066)
LTLA	Rifamycin/isoniazid (active and latent TB, other in- fections)	1.004 (0.992- 1.016)	0.996 (0.976- 1.016)	1.005 (0.992- 1.017)	1.038 (1.025- 1.052)	1.038 (1.023- 1.054)	1.039 (1.026- 1.052)
LTLA	TNF inhibitors	0.992 (0.976- 1.008)	0.976 (0.935- 1.019)	0.993 (0.977- 1.009)	0.994 (0.979- 1.010)	0.994 (0.979- 1.009)	0.995 (0.979- 1.011)
LTLA	IL-6 inhibitors and abat- cept	1.001 (0.993- 1.010)	0.996 (0.980- 1.012)	1.002 (0.994- 1.010)	1.001 (0.991- 1.011)	1.001 (0.991- 1.011)	1.001 (0.991- 1.011)
LTLA	JAK inhibitors (rheuma- tology)	0.995 (0.989- 1.002)	0.980 (0.963- 0.998)	0.996 (0.989- 1.003)	0.994 (0.988- 0.999)	0.994 (0.988- 1.000)	0.994 (0.989- 1.000)
LTLA	Rituximab	1.001 (0.986- 1.016)	0.989 (0.963- 1.016)	1.002 (0.986- 1.017)	0.990 (0.976- 1.004)	0.990 (0.977- 1.003)	0.991 (0.977- 1.005)

Level	Drug group	lag (t-1)	lag (t-1), excluding outcome years 2020-21	lag (t-1), elective catch- ments	concurrent (t)	concurrent (t), clustered by principal trust	concurrent (t), elective catch- ments
LTLA	Calcineurin/mTOR inhibitors	1.008 (0.997- 1.020)	1.015 (0.994- 1.036)	1.008 (0.997- 1.020)	0.994 (0.985- 1.003)	0.994 (0.985- 1.003)	0.994 (0.985- 1.003)
LTLA	Antiproliferative	0.997 (0.982- 1.013)	1.008 (0.982- 1.035)	0.997 (0.981- 1.013)	0.997 (0.983- 1.012)	0.997 (0.986- 1.009)	0.998 (0.983- 1.013)
LTLA	Systemic glucocor- ticoids without dexam- etha- sone/hydrocortisone (prednisolone- equivalent mg)	1.021 (1.000- 1.043)	1.022 (0.979- 1.068)	1.022 (1.001- 1.044)	1.008 (0.990- 1.026)	1.008 (0.988- 1.027)	1.009 (0.991- 1.028)
LTLA	Dexamethasone and hydrocor- tisone (descrip- tive)	1.010 (0.984- 1.036)	1.019 (0.970- 1.070)	1.011 (0.984- 1.039)	1.008 (0.980- 1.037)	1.008 (0.979- 1.038)	1.011 (0.982- 1.040)
LTLA	Low-TB- risk biologics (negative control)	1.003 (0.988- 1.019)	0.996 (0.959- 1.036)	1.003 (0.988- 1.019)	1.005 (0.991- 1.019)	1.005 (0.989- 1.020)	1.004 (0.991- 1.018)
LTLA	Levetiracetam (negative control)	1.013 (0.988- 1.039)	1.021 (0.970- 1.074)	1.013 (0.988- 1.040)	1.003 (0.981- 1.025)	1.003 (0.980- 1.027)	1.004 (0.982- 1.027)

Table S8. Original panel design (UTLA; Fingertips three-year notification windows; practice-postcode assignment; original drug groups): IRR per 10% increase (95% CI)

Drug group	primary: t-5..t-3	lag: t-3 only	concurrent: t-2..t	lead (falsification): t+1..t+3
Antituberculosis (descriptive)	1.006 (0.999-1.012)	1.002 (0.998-1.005)	1.000 (0.996-1.004)	0.999 (0.994-1.005)
oral_corticosteroids	1.001 (0.951-1.055)	0.999 (0.958-1.040)	0.978 (0.931-1.028)	0.992 (0.946-1.040)
Inhaled corticosteroids	1.013 (0.944-1.086)	1.011 (0.962-1.064)	0.963 (0.918-1.011)	0.971 (0.916-1.029)
immunosuppressants	1.006 (0.978-1.034)	1.007 (0.987-1.027)	1.009 (0.988-1.031)	1.003 (0.976-1.030)
Proton pump inhibitors	1.007 (0.941-1.077)	1.002 (0.957-1.048)	1.000 (0.953-1.049)	1.040 (0.985-1.098)

Drug group	Rule
All antibacterials	code: ^0501
Vitamin D	name: ^(:colecalfiferol\ ergocalciferol)
Levothyroxine (negative control)	name: ^levothyroxine

BNF chapters 11-13 (eye, ear and nose, skin) are excluded unless a code rule applies; injectable presentations are excluded from oral groups. Prednisolone-equivalent factors: prednisolone and prednisone 1, methylprednisolone 1.25, deflazacort 0.83, dexamethasone 6.67. The SCMD product classification with DDDs is in outputs/hospital/vmp_classification.csv.