

A rule-based algorithm for tillage detection from Sentinel-1 time series

Dries De Bièvre, Louis Descarpentries, Sophie Bontemps, Pierre Defourny, Bas van Wesemael
UCLouvain – Earth and Life Institute, Louvain-la-Neuve, Belgium
dries.debievre@uclouvain.be

1 Context

Tillage strongly affects **soil health**, influencing **C-dynamics** and **aggregate stability** [1-3]

Earth observation could support monitoring tillage practices, yet **optical indices lack reliability** due to cloud cover and are only sensitive to differences in crop residue cover

Can we use **Sentinel-1 SAR** data to estimate tillage dates and tillage type? Previous studies have shown promising results [4-6]

➔ Leveraging **coherence (**sensitive to **surface change)** and **backscatter** (sensitive to **soil roughness)**


Terminology

Tillage: agricultural preparation of soil by mechanical agitation

Inversion tillage: The use of a mouldboard or disk plough to invert the topsoil, typically to a depth of 20-30 cm

2 Methods

Algorithm



Ground-truth data collection by weekly revisits

Site	Parcels	Observations	Tillage events	Inversion tillage events
Spring 2025	150	2127	167	29
Autumn 2025	106	1702	136	20




Figure 1. Location of the study sites in Belgium with respect to the official agricultural regions

3 Results




Figure 2. Example of the coherence change detections for a parcel with three recorded tillage events

STEP 1
Detect tillage events

0.25 Fixed threshold on coherence change

Average F1 goes from 0.50 to 0.58 by filtering detections

Other causes of coherence drop (snow, vegetation) must be filtered out.

STEP 2
Separate tillage types

Spring 2025 Autumn 2025

VV
VH

Average F1 scores for single orbit, for VV: 0.55, for VH: 0.55 (spring)
VV: 0.32
VH: 0.36 (autumn)

STEP 3
Consolidate & validate

F1-score against in-situ observations

	Spring 2025	Autumn 2025
Tillage detection	0.74	0.66
Inversion tillage	0.66	0.52




Figure 3. Distribution of backscatter values for three types of tillage on both study sites.

4 Conclusions

Coherence change can accurately **detect tillage events** with a threshold of 0.25 and applying filters for false detections

Types of tillage can be **differentiated by backscatter**, performance differs across seasons

Combining detections from **overlapping orbits increases accuracy**

→ **PDF**

Can you help evaluate our algorithm in other regions? Get in touch.

1. Clement, T., Bielders, C. L. & Degré, A. How much do conservation cropping practices mitigate runoff and soil erosion under Western European conditions: A focus on conservation tillage, tied ridging and winter cover crops. *Soil Use and Management* **40**, e13047 (2024).
2. Meurer, K. H. E., Haddaway, N. R., Bolinder, M. A. & Kätterer, T. Tillage intensity affects total SOC stocks in boreo-temperate regions only in the topsoil—A systematic review using an ESM approach. *Earth-Science Reviews* **177**, 613–622 (2018).
3. Six, J., Elliott, E. T. & Paustian, K. Soil macroaggregate turnover and microaggregate formation: a mechanism for C sequestration under no-tillage agriculture. *Soil Biology and Biochemistry* **32**, 2099–2103 (2000).
4. Dingle Robertson, L. *et al.* Monitoring autumn agriculture activities using Synthetic Aperture Radar (SAR) and coherence change detection. *Heliyon* **9**, e17322 (2023).
5. Satalino, G. *et al.* Copernicus Sentinels For Tillage Change Detection. in *IGARSS 2024 - 2024 IEEE International Geoscience and Remote Sensing Symposium* 1249–1252 (2024). doi:10.1109/IGARSS53475.2024.10641126.
6. McNairn, H. *et al.* Coherent Change Detection to Monitor Tillage. in *IGARSS 2022 - 2022 IEEE International Geoscience and Remote Sensing Symposium* 5858–5861 (2022). doi:10.1109/IGARSS46834.2022.9883744.

2nd EO 4 Soil Symposium
ESA SYMPOSIUM ON EARTH OBSERVATION FOR SOIL PROTECTION AND RESTORATION
04–06 November 2026 | ESA–ESRIN | Frascati (Rome), Italy