

Édouard Lucas:

The theory of recurrent sequences is an inexhaustible mine which contains all the properties of numbers; by calculating the successive terms of such sequences, decomposing them into their prime factors and seeking out by experimentation the laws of appearance and reproduction of the prime numbers, one can advance in a systematic manner the study of the properties of numbers and their application to all branches of mathematics.



Applications of Linear Algebra to Graph Theory and Network Science

Daryl DeFord

Washington State University
Department of Mathematics and Statistics

ILAS Vodcast
April 2023



Outline

- **Goals:**

- Introduce graph theory and social network modeling
- Investigate connections with linear algebra
- Inspire interest in exploring further

- **Format:**

- Overview of ideas and connections
- A couple of proofs and examples
- Computational notebooks and resources

- **Topics:**

- Introduction to graphs networks
- Graph Matrices
- Centrality
- Clustering
- Dynamics
- Random Models

- **Resources:**


- https://github.com/ILAS_Networks





Graph Theory

Definition (Graph)

A Graph $G = (V, E)$ is a set of nodes V and a set of edges $E \subseteq V \times V$.



Graph Examples




Example Graph Questions

- Are there Hamiltonian/Eulerian Paths? If so, how many?
- Are there perfect matchings? If so, how many?
- Is it possible color the nodes with k colors so that no neighboring nodes are the same color?
- What is the largest set of nodes that share no edges?
- How many edges/vertices must be removed to disconnect the graph?
- Is it possible to embed the graph in the plane without any edges crossing?
- How many spanning trees does the graph have?
- What is the automorphism group of the graph?
- ...




Adjacency Matrix



$$A = \begin{bmatrix} 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 1 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 0 & 1 & 0 & 1 & 0 \end{bmatrix}$$



Degree Matrix



$$D = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 5 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 6 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 5 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 4 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 4 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 3 \end{pmatrix}$$



Laplacian Matrix (D-A)

$$L = \begin{bmatrix} 1 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\ 0 & 5 & 0 & -1 & -1 & -1 & -1 & -1 \\ 0 & 0 & 2 & -1 & -1 & 0 & 0 & 0 \\ -1 & -1 & -1 & 6 & -1 & -1 & -1 & 0 \\ 0 & -1 & -1 & -1 & 5 & -1 & 0 & -1 \\ 0 & -1 & 0 & -1 & -1 & 4 & -1 & 0 \\ 0 & -1 & 0 & -1 & 0 & -1 & 4 & -1 \\ 0 & -1 & 0 & 0 & -1 & 0 & -1 & 3 \end{bmatrix}$$



Notation and Structural Results

- Notation:
 - $G = (V, E)$
 - Number of nodes: $|V| = n$
 - Number of edges: $|E| = m$
 - There is an edge between u and v : $u \sim v$
 - The degree (number of neighbors of a node): $\deg(u)$
 - The geodesic distance between two nodes: $d(u, v)$
 - The diameter of a graph is the length of a longest path:
 $\text{diam}(G) = \max_{u,v} d(u, v)$.
 - Eigenvalues of A : $|\lambda_1| \geq |\lambda_2| \cdots \geq |\lambda_n|$
- $(A^k)_{i,j}$ counts the number of walks of length k starting at node i and ending at node j .
- When A is a connected graph it is irreducible and hence satisfies the non-negative Perron-Frobenius theorem




Spectral Graph Theory

F. Chung: *Spectral Graph Theory*, AMS, (1997).

“Roughly speaking, half of the main problems of spectral theory lie in deriving bounds on the distributions of eigenvalues. The other half concern the impact and consequences of the eigenvalue bounds as well as their applications.”

R. Brualdi: *The Mutually Beneficial Relationship of Graphs and Matrices*, CBMS, (2011)

Graphs and matrices enjoy a fascinating and mutually beneficial relationship. This interplay has benefited both graph theory and linear algebra. In one direction, knowledge about one of the graphs that can be associated with a matrix can be used to illuminate matrix properties and to get better information about the matrix. [...] Going the other way, linear algebraic properties of one of the matrices associated with a graph can be used to obtain useful combinatorial information about the graph.




Diameter and Multiplicity

Theorem

The diameter of G is bounded above by the number of distinct eigenvalues of A .

Proof.

Let k be the number of distinct eigenvalues of A . Since A is diagonalizable, k is also the degree of the minimal polynomial of A . Thus, the set $\{I, A, A^2, \dots, A^k\}$ is linearly dependent and in particular $A^k = \sum_{i=0}^{k-1} \alpha_i A^i$ for some constants α_i . In order to obtain a contradiction, assume that there is a pair of vertices $u, v \in V$ with $d(u, v) = k$. Then $(A^k)_{u,v} > 0$ but $(A^i)_{u,v} = 0$ for all $0 \leq i \leq k-1$, which is a contradiction.



Bounding λ_1

Theorem

Let δ and Δ be the minimum and maximum degrees of nodes in G and λ_1 be the leading eigenvalue. Then $\delta \leq \lambda_1 \leq \Delta$.

Proof.

Let v be an eigenvector for λ_1 and k be the index of a maximal entry of v , so $v_k = \max_i v_i$. Consider the k th row of $Av = \lambda_1 v$:

$$\lambda_1 v_k = (Av)_k = \sum_{i \sim k} v_i \leq \sum_{i \sim k} v_k = \deg(v_k) v_k \leq \Delta v_k.$$

To see the other inequality consider the quadratic form:

$$\lambda_1 = \sup_{||v||=1} v^T A v \geq \left(\frac{1}{n}\right) \mathbf{1}^T A \mathbf{1} = \left(\frac{1}{n}\right) \sum_{i=1}^n \sum_{j=1}^n A_{i,j} = \frac{2|E|}{n} \geq \delta.$$



Laplacian Properties

- L is positive semi-definite since it can be factored at VV^T where V is an incidence matrix for the graph
- The all ones vector is an eigenvector with eigenvalue 0 for L . The dimension of the 0 eigenspace is the number of connected components of G .
- The smallest non-zero eigenvalue is called the algebraic connectivity or Fiedler value and we'll see this has
- Kirchoff's theorem: The number of spanning trees in a graph is given by the value of any cofactor of the Laplacian.



What is a social network?

Definition (Social Network)

A social network is represented by a collection of “nodes” representing individual actors and a set of “edges” representing a binary relationship between the actors.



What is a social network?

Definition (Social Network)

A social network is represented by a collection of “nodes” representing individual actors and a set of “edges” representing a binary relationship between the actors.

Example

What kinds of systems can social networks describe?

- What could be represented by nodes?



What is a social network?

Definition (Social Network)

A social network is represented by a collection of “nodes” representing individual actors and a set of “edges” representing a binary relationship between the actors.

Example

What kinds of systems can social networks describe?

- What could be represented by nodes?
 - **University Students**



What is a social network?

Definition (Social Network)

A social network is represented by a collection of “nodes” representing individual actors and a set of “edges” representing a binary relationship between the actors.

Example

What kinds of systems can social networks describe?

- What could be represented by nodes?
 - **University Students**
- What type of edges could connect them?



What is a social network?

Definition (Social Network)

A social network is represented by a collection of “nodes” representing individual actors and a set of “edges” representing a binary relationship between the actors.

Example

What kinds of systems can social networks describe?

- What could be represented by nodes?
 - **University Students**
- What type of edges could connect them?
 - **In a class together**
 - **Facebook friends**
 - **Speak at least twice a week**



What is a social network?

Definition (Social Network)

Mathematically, a social network is represented by a collection of “nodes” representing individual actors and a set of “edges” representing a binary relationship between the actors.

Example

What kinds of systems can social networks describe?

- What could be represented by nodes?
 - **Academic Departments**
- What type of edges could connect them?
 - **Located in same building**
 - **Students who major in both**
 - **Crosslisted courses**



Initial Example

Definition (Ego Network)

An ego network is a social network centered at a particular individual containing their connections and the connections between their “friends.”



Initial Example

Definition (Ego Network)

An ego network is a social network centered at a particular individual containing their connections and the connections between their “friends.”

Problem (Draw your ego network)




Initial Example

Definition (Ego Network)

An ego network is a social network centered at a particular individual containing their connections and the connections between their “friends.”

Problem (Draw your ego network)




World Trade Web




D., Multiplex Dynamics on the World Trade Web, Proc. 6th International Conference on Complex Networks and Applications, Studies in Computational Intelligence, Springer, 1111-1123, (2018).



Karnataka Village Data



Village 5



Village 61

Figure: Two of the Karnataka Village networks

A. Banerjee, A.G. Chandrasekhar, E. Duflo, and M.O. Jackson, The Diffusion of Microfinance. Science, (2013).




Common Properties of Social Networks

Example (What features distinguish social networks?)

- Transitivity
- Community structure
- Small average path length
- Long-tailed degree distribution
- Hubs
- ...




Karate Club




Zachary, Wayne W. "An Information Flow Model for Conflict and Fission in Small Groups." *Journal of Anthropological Research*, 33, 452–473, (1977).



Dolphins



Centrality



Clustering

Figure: Dolphin social network¹

¹D. Lusseau, K. Schneider, O. Boisseau, Patti Haase, E. Slooten, and S. Dawson, The bottlenose dolphin community of

Doubtful Sound features a large proportion of long-lasting associations, Behavioral Ecology and Sociobiology 54 (2003), 4, 396–405.




Network Examples


- Internet
 - Nodes: Hardware
 - Edges: Physical Connections
- WWW
 - Nodes: Webpages
 - Edges: Links
- Neuroscience
 - Nodes: Brain areas
 - Edges: Functional Connections
- Food webs
 - Nodes: Species
 - Edges: Predation
- World Trade
 - Nodes: Countries
 - Edges: Trade Agreements
- Banking
 - Nodes: Banks
 - Edges: Loans
- Genetics
 - Nodes: Genes/Proteins
 - Edges: Functional Interactions
- Recommendations
 - Nodes: People/Products
 - Edges: Ratings/Consumption



Network Questions



Centrality



Clustering



Centrality





Figure: Relevant comic by Randall Munroe¹ (emphasis mine).

¹ <https://xkcd.com/451/>



Harder Question Domain



Eigenvector Centrality

- Naively: You are popular if you have lots of friends
- Intuitively: You are popular if your friends are popular
- Formally: Your popularity should be proportional to the sum of your friends' popularities.
- Mathematically: Given a vector v whose entries represent the initial popularity of each node in the network we seek a solution to:


$$v_i = \lambda \sum_{i \sim j} v_j = \sum_{j=1}^n A_{i,j} v_j$$

or equivalently:


$$v = \lambda A v$$



Adjacency Action



Before



After



Eigenvector Centrality Extensions


- Many real-world networks have a directed structure, in which case we might have 'sinks' in the adjacency dynamics that will eventually absorb all of the popularity
- To address this, we might continually pump popularity into the system at each step
- Katz Centrality: For some parameter $\alpha < \lambda_1$ compute

$$K_i = \sum_{k=0}^{\infty} \sum_{j=1}^n \alpha^k (A^k)_{i,j}$$

- PageRank: We can also compute a similar centrality score proportional to the stationary distribution of a random walk on the (di)graph. The pagerank is equivalent to normalizing our adjacency matrix by degree and then allowing some small probability of randomly hopping to any node in the network.



Toy Eigenvector Centrality



$$v = \begin{bmatrix} 0.1052 \\ 0.4470 \\ 0.2022 \\ 0.4508 \\ 0.4155 \\ 0.3926 \\ 0.3683 \\ 0.2873 \end{bmatrix}$$



Clustering Definitions

- Spectral clustering
 - Minimize Inter-Community Edges
 - Minimize $s^T L s$ with $s \in \{\pm 1\}^n$



Clustering Definitions

- Spectral clustering
 - Minimize Inter-Community Edges
 - Minimize $s^T L s$ with $s \in \{\pm 1\}^n$
- Modularity
 - Maximize Intra-Community Edges (compared to expectation)
 - Maximize $s^T B s$ with $s \in \{\pm 1\}^n$ where $B_{i,j} = A_{i,j} - \frac{\deg(i) \deg(j)}{2m}$.



Clustering Definitions

- Spectral clustering
 - Minimize Inter-Community Edges
 - Minimize $s^T L s$ with $s \in \{\pm 1\}^n$
- Modularity
 - Maximize Intra-Community Edges (compared to expectation)
 - Maximize $s^T B s$ with $s \in \{\pm 1\}^n$ where $B_{i,j} = A_{i,j} - \frac{\deg(i) \deg(j)}{2m}$.
- Markov Stability
 - Given a partition $(V_1, V_2, \dots, V_\ell)$ maximize (for fixed t)
$$r(t, V) = \sum_{i=1}^{\ell} \sum_{v_y, v_z \in V_i} C(t)_{y,z}$$
 - Discrete: $C(t) = \Pi S^t - \pi^T \pi$ where π is the steady state vector and Π is the diagonal matrix of π
 - Continuous: $C(t) = \Pi e^{-t(I-S)} - \pi^T \pi$ where π is the steady state vector and Π is the diagonal matrix of π



Spectral Clustering Setup

We can think of this as minimizing the damage done to the network by removing the edges between clusters. Investigating the 2-partition case, we can begin by constructing a vector v to represent the desired partition, with entries of 1 assigned to nodes in one component and entries of -1 assigned to the other component. This gives that $v_i v_j = 1$ if and only if i and j are in the same component. We can now formulate the damage condition algebraically as

$$\text{damage}(v) = \frac{1}{2} \sum_{i,j} \frac{1}{2} (1 - v_i v_j) A_{i,j}.$$

Proceeding by algebraic manipulation, we can reduce this expression to

$$\begin{aligned} \frac{1}{2} \sum_{i,j} \frac{1}{2} (1 - v_i v_j) A_{i,j} &= \frac{1}{4} (\sum_{i,j} A_{i,j} - v_i v_j A_{i,j}) \\ &= \frac{1}{4} \sum_{i,j} v_i \deg(i) v_j \delta_{i,j} - v_i A_{i,j} v_j \\ &= \frac{1}{4} v^T D v - v^T A v \\ &= \frac{1}{4} v^T L v \end{aligned}$$



Spectral Clustering II

- Solving this problem exactly across all binary vectors is:
 - NP-hard
 - usually irrelevant (separating a leaf)
- Instead, we can solve a relaxed problem with $v \in \mathbb{R}^n$ with $\|v\| = 1$ and $v \perp 1$
- In this case our solution is just given by a Rayleigh quotient
- The continuous solution is thus given by the eigenvector corresponding to the Fiedler value and we can recover a binary solution by setting the positive values to 1 and the negative values to -1
- Ratio and Normalized cuts also address the irrelevancy issue
- We'll explore the case of more than two parts on the computational side



Dynamics on Networks

- Adjacency
 - Matrix: A
 - Symmetric, binary
 - Eigenvector centrality – leading eigenvector



Dynamics on Networks

- Adjacency
 - Matrix: A
 - Symmetric, binary
 - Eigenvector centrality – leading eigenvector
- Laplacian
 - Matrix: $L = D - A$
 - Positive semi-definite
 - Discretized version of Laplacian heat diffusion
 - **Fiedler value**: smallest non-zero eigenvalue



Dynamics on Networks

- Adjacency
 - Matrix: A
 - Symmetric, binary
 - Eigenvector centrality – leading eigenvector
- Laplacian
 - Matrix: $L = D - A$
 - Positive semi-definite
 - Discretized version of Laplacian heat diffusion
 - **Fiedler value**: smallest non-zero eigenvalue
- Random Walk
 - Matrix: AD^{-1}
 - Stochastic, regular if G is connected
 - Transition matrix of associated Markov process
 - Convergence governed by second largest eigenvalue



Diffusion Dynamics

Given a vector of initial heat values v the change in temperature at node i with respect to time is given by

$$\frac{dv_i}{dt} = -K \sum_{i \sim j} v_i - v_j$$

or

$$\frac{dv}{dt} = -KLv$$



where L is the graph Laplacian. This is a symmetric, positive semi-definite matrix so the value at time t is

$$v(t) = \sum_{i=1}^n c_i v^i e^{-\lambda_i t}$$



where the (v^i, λ_i) are eigenpairs for L .





Diffusion Animation



Random Walk Dynamics



Random Models



Null Models

Definition (Null Model)

A random network, parameterized to match some features of a given network, used to compare “expected” network measures.



Null Models

Definition (Null Model)

A random network, parameterized to match some features of a given network, used to compare “expected” network measures.
Equivalently, a distribution over binary, symmetric random matrices...



Null Models

Definition (Null Model)

A random network, parameterized to match some features of a given network, used to compare “expected” network measures.
Equivalently, a distribution over binary, symmetric random matrices...

Example (How to grow a network?)

- Ego networks
 - Start with a single node
 - Attach that node to k other nodes
 - Add edges between the each pair of friends with probability p
- ?



Dot Product Graphs

Definition (Dot Product Graph)

G is a dot product graph of dimension d if there exists a map $f : V(G) \rightarrow \mathbb{R}^d$ such that $(i, j) \in E(G)$ if and only if $\langle f(i), f(j) \rangle > 1$.

¹C. Fiduccia, E. Scheinerman, A. Trenk, and J. Zito: Dot Product Representations of Graphs, *Discrete Mathematics*, 181, 1998, 113–138.

²R. Kang, L. Lovasz, T. Muller, and E. Scheinerman: Dot Product Representations of Planar Graphs, *Electronic Journal of Combinatorics*, 18, (2011), 1–14.

³Sphere and Dot Product Representations of Graphs: *Discrete Computational Geometry*, 47, (2012), 548–568.

⁴B. Li and G. Chang: Dot Product Dimension of Graphs, *Discrete Applied Mathematics*, 166, (2014), 159–163



Dot Product Graphs

Definition (Dot Product Graph)

G is a dot product graph of dimension d if there exists a map $f : V(G) \rightarrow \mathbb{R}^d$ such that $(i, j) \in E(G)$ if and only if $\langle f(i), f(j) \rangle > 1$.

- Initial work: Fiduccia et al. (1998)¹
- Planar graphs: Kang et al. (2011)²
- NP-Hard: Kang and Muller (2012)³
- $\frac{n}{2}$ critical graphs: Li and Chang (2014)⁴

¹C. Fiduccia, E. Scheinerman, A. Trenk, and J. Zito: Dot Product Representations of Graphs, *Discrete Mathematics*, 181, 1998, 113–138.

²R. Kang, L. Lovasz, T. Muller, and E. Scheinerman: Dot Product Representations of Planar Graphs, *Electronic Journal of Combinatorics*, 18, (2011), 1–14.

³Sphere and Dot Product Representations of Graphs: *Discrete Computational Geometry*, 47, (2012), 548–568.

⁴B. Li and G. Chang: Dot Product Dimension of Graphs, *Discrete Applied Mathematics*, 166, (2014), 159–163



RDPM

(RDPM 1): Select the number of nodes n .



RDPM

(RDPM 1): Select the number of nodes n .

(RDPM 2): Select the latent dimension d .



RDPM

(RDPM 1): Select the number of nodes n .

(RDPM 2): Select the latent dimension d .

(RDPM 3): Select a distribution W over \mathbb{R}^d with $\mathbb{P}(\langle X, Y \rangle \in (0, 1)) = 1$ where X and Y are drawn independently from W .



RDPM

(RDPM 1): Select the number of nodes n .

(RDPM 2): Select the latent dimension d .

(RDPM 3): Select a distribution W over \mathbb{R}^d with $\mathbb{P}(\langle X, Y \rangle \in (0, 1)) = 1$ where X and Y are drawn independently from W .

(RDPM 4): For each node, $1 \leq j \leq n$, draw a vector, $X_j \in \mathbb{R}^d$ from W .



RDPM

(RDPM 1): Select the number of nodes n .

(RDPM 2): Select the latent dimension d .

(RDPM 3): Select a distribution W over \mathbb{R}^d with $\mathbb{P}(\langle X, Y \rangle \in (0, 1)) = 1$ where X and Y are drawn independently from W .

(RDPM 4): For each node, $1 \leq j \leq n$, draw a vector, $X_j \in \mathbb{R}^d$ from W .

(RDPM 5): Form an adjacency matrix, A , form a network with $A_{j,\ell}$ drawn from $\text{Bernoulli}(\langle X_j, X_\ell \rangle)$ for $j \neq \ell$ and $A_{j,j} = 0$ for all $1 \leq j \leq n$.



Interpretations

- Since each node is associated to a vector, it is natural to try and interpret the properties of the node from the vector



Interpretations

- Since each node is associated to a vector, it is natural to try and interpret the properties of the node from the vector
- $\langle x, y \rangle = ||x|| \cdot ||y|| \cos(x, y)$



Interpretations

- Since each node is associated to a vector, it is natural to try and interpret the properties of the node from the vector
- $\langle x, y \rangle = ||x|| \cdot ||y|| \cos(x, y)$
- Angle – Community assignment




Interpretations


- Since each node is associated to a vector, it is natural to try and interpret the properties of the node from the vector
- $\langle x, y \rangle = ||x|| \cdot ||y|| \cos(x, y)$
- Angle – Community assignment
- Magnitude – Centrality



Angle – Community Assignment




Vectors




Graph



Magnitude – Centrality







Vectors



Graph



Multi-Resolution Communities



Network Properties

- Initial work: Kraetzel et al. (2005)⁵
- General distributions: Young and Scejnerman (2007)⁶

⁵M. KRAETZEL, C. NICKEL, AND E. SCHEINERMAN: *Random Dot Product Networks: A model for social networks*, Preliminary Manuscript, (2005).

⁶S. YOUNG AND E. SCHEINERMAN: *Random Dot Product Models for Social Networks*, Algorithms and Models for the Web-Graph, Lecture Notes in Computer Science, 4863, (2007), 138–149.



Network Properties

- Initial work: Kraetzel et al. (2005)⁵
- General distributions: Young and Scejnerman (2007)⁶
- Small world networks
 - Clustering
 - Small diameter
 - Degree distribution

⁵M. KRAETZEL, C. NICKEL, AND E. SCHEINERMAN: *Random Dot Product Networks: A model for social networks*, Preliminary Manuscript, (2005).

⁶S. YOUNG AND E. SCHEINERMAN: *Random Dot Product Models for Social Networks*, Algorithms and Models for the Web-Graph, Lecture Notes in Computer Science, 4863, (2007), 138–149.



Statistical Applications

- Inverse problem: Scheinerman and Tucker (2010)⁷
 - Iterative SVD for approximating $A_{i,j} = \langle X_i, X_j \rangle$
 - Angular k-means
- MAP/Conic Programming: Wu, Palmer, and DeFord (2022)

⁷E. SCHEINERMAN AND K. TUCKER: *Modeling graphs using dot product representations*, Computational Statistics, 25, (2010), 1–16.

⁸D. SUSSMAN, M. TANG, D. FISHKIND, AND C. PRIEBE: *A consistent adjacency spectral embedding for stochastic blockmodel graphs*, Journal of the American Statistical Association, 107, (2012), 1119–1128.

⁹M. TANG, A. ATHREYA, D. SUSSMAN, V. LYZINSKI, AND C. PRIEBE: *A nonparametric two-sample hypothesis testing problem for random graphs*, Arxiv: 1409.2344v2, (2014), 1–24.

¹⁰M. TANG AND C. PRIEBE: *Limit theorems for eigenvectors of the normalized Laplacian for random graphs*, ArXiv:1607.08601, (2016), 1–52.



Statistical Applications

- Inverse problem: Scheinerman and Tucker (2010)⁷
 - Iterative SVD for approximating $A_{i,j} = \langle X_i, X_j \rangle$
 - Angular k-means
- MAP/Conic Programming: Wu, Palmer, and DeFord (2022)
- Spectral Embedding and Statistics: Priebe Lab (2012–present)
 - Adjacency embedding⁸
 - Hypothesis testing⁹
 - Limit theorems¹⁰

⁷E. SCHEINERMAN AND K. TUCKER: *Modeling graphs using dot product representations*, Computational Statistics, 25, (2010), 1–16.

⁸D. SUSSMAN, M. TANG, D. FISHKIND, AND C. PRIEBE: *A consistent adjacency spectral embedding for stochastic blockmodel graphs*, Journal of the American Statistical Association, 107, (2012), 1119–1128.

⁹M. TANG, A. ATHREYA, D. SUSSMAN, V. LYZINSKI, AND C. PRIEBE: *A nonparametric two-sample hypothesis testing problem for random graphs*, Arxiv: 1409.2344v2, (2014), 1–24.

¹⁰M. TANG AND C. PRIEBE: *Limit theorems for eigenvectors of the normalized Laplacian for random graphs*, ArXiv:1607.08601, (2016), 1–52.



Edge Parameterized Models

Theorem

Let n be a fixed positive integer. For each pair (i, j) with $1 \leq i < j \leq n$ let $a_{i,j} = a_{j,i} \in \mathbb{R}$. Then there exist n real numbers $a_{\ell,\ell}$ for $1 \leq \ell \leq n$ such that the matrix $A_{i,j} = a_{i,j}$ is positive definite.

Proof.

Let the $a_{i,j}$ be selected arbitrarily. For $1 \leq \ell \leq n$ choose $a_{\ell,\ell} \in \mathbb{R}$ so that $a_{\ell,\ell} > \sum_{j \neq \ell} |a_{j,\ell}|$. Form a matrix A with $A_{i,j} = a_{i,j}$. This is a real symmetric matrix and so by the spectral theorem A has real eigenvalues. Applying Gershgorin's Circle Theorem to A gives that the eigenvalues of A lie in the closed disks centered at $a_{\ell,\ell}$ with radius $\sum_{j \neq \ell} |a_{j,\ell}|$. Intersecting these disks with the real line gives that the eigenvalues of A must lie in $\bigcup_{\ell=1}^n \left[a_{\ell,\ell} - \sum_{j \neq \ell} |a_{j,\ell}|, a_{\ell,\ell} + \sum_{j \neq \ell} |a_{j,\ell}| \right] \subseteq \mathbb{R}^+$. Thus, all eigenvalues of A are positive and A is positive definite.



Edge Parameterized Models

Corollary

Any generative network model, on a fixed number of nodes n , where the edge weight between each pair of nodes is drawn independently from a fixed probability distribution, possibly with different parameters for each pair, can be realized under the WRDPN.


Proof.

Let P be the k -parameter distribution from which the edge weights are drawn and for $1 \leq i \leq k$ let $a_{j,\ell}^i = a_{\ell,j}^i$ be the value of the i th parameter between nodes j and ℓ . Applying the previous theorem to the collection $a_{j,\ell}^i = a_{\ell,j}^i$ gives a positive definite matrix A^i . Thus, there exists an $n \times n$ matrix X^i such that $(X^i)^T X^i = A^i$.


To form the WRDPM that matches the given generative model we take $d_i = n$ for all $1 \leq i \leq k$ and to each node $1 \leq j \leq n$ assign the collection of vectors given by the j th columns of the X^i for $1 \leq i \leq k$.




Weighted Collaboration Network



Collaboration Network




Weighted 2-Embedding




Weighted 3-Embedding




Voting Data




Dimension Example




Weighted Network



2-Embedding



3-Embedding



Stress Function



Resources

- Spectral Graph Theory (Chung)
- The Mutually Beneficial Relationship of Graphs and Matrices (Brualdi)
- Networks (Newman)
- Network Analysis and Modeling (Clauset)
- https://github.com/drdeford/ILAS_Networks
- <https://github.com/drdeford/CISER2023NetworkX/>
- <https://github.com/drdeford/JMM2021NetworkX>



Conclusion

Thanks!



Temporary page!

\LaTeX was unable to guess the total number of pages correctly. As there was some unprocessed data that should have been added to the final page, this extra page has been added to receive it.

If you rerun the document (without altering it) this surplus page will go away, because \LaTeX now knows how many pages to expect for this document.