

Collision

YU-CHIA LIN*

Department of Physics & Astronomy, University of New Mexico, Albuquerque, New Mexico 87131, USA

Department of Astronomy, University of Arizona, Tucson, Arizona 85721, USA and

Department of Physics, University of Arizona, Tucson, Arizona 85721, USA

(Dated: July 20, 2022)

We investigate the collision term effects in the neutrino flavor oscillation.

I. INTRODUCTION

A.

1.

II. MODEL

In this work, we consider the two-flavor neutrino flavor oscillation between ν_e and ν_τ , where ν_τ represents a linear combination of the physical μ and τ flavor neutrino. On the other hand, we only consider monochromatic neutrino to simplify the model.

Denoting the (anti-)neutrino flavor density matrix and the corresponding polarization vector as

$$\rho = \begin{bmatrix} \rho_{11} & \rho_{12} \\ \rho_{21} & \rho_{22} \end{bmatrix} \quad (1)$$

$(\bar{\rho})$ and $\mathbf{P}(\bar{\mathbf{P}})$ respectively, or

$$\rho_{ex} \equiv \rho_{12} = \frac{1}{2}(P_x + iP_y) \quad (2)$$

$$\bar{\rho}_{ex} \equiv \bar{\rho}_{12} = \frac{1}{2}(\bar{P}_x + i\bar{P}_y) \quad (3)$$

, where $P_i(\bar{P}_i)$ is the i component of $P(\bar{P})$.

Considering the collision terms, we have equations of motion (EOM) for polarization vectors as follows:

$$\dot{\mathbf{P}} = \omega \mathbf{B} \times \mathbf{P} + \mu(\mathbf{P} - \bar{\mathbf{P}}) \times \mathbf{P} - \Gamma_+^{CC} \mathbf{P}_T + \Gamma_+^{AE}(\mathbf{P}^{AE} - \mathbf{P}) + \Gamma_-^{AE}(P_0^{AE} - P_0)\mathbf{z} \quad (4)$$

$$\dot{\bar{\mathbf{P}}} = -\omega \mathbf{B} \times \bar{\mathbf{P}} + \mu(\mathbf{P} - \bar{\mathbf{P}}) \times \bar{\mathbf{P}} - \bar{\Gamma}_+^{CC} \bar{\mathbf{P}}_T + \bar{\Gamma}_+^{AE}(\bar{\mathbf{P}}^{AE} - \bar{\mathbf{P}}) + \bar{\Gamma}_-^{AE}(\bar{P}_0^{AE} - \bar{P}_0)\mathbf{z} \quad (5)$$

$$\dot{P}_0 = \Gamma_+^{AE}(P_0^{AE} - P_0) + \Gamma_-^{AE}(P_z^{AE} - P_z) \quad (6)$$

$$\dot{\bar{P}}_0 = \bar{\Gamma}_+^{AE}(\bar{P}_0^{AE} - \bar{P}_0) + \bar{\Gamma}_-^{AE}(\bar{P}_z^{AE} - \bar{P}_z) \quad (7)$$

We then choose the parameters

$$\begin{bmatrix} \omega \\ \mathbf{B} \\ \mu \\ \theta \end{bmatrix} = \begin{bmatrix} 0.3 \text{ km}^{-1} \\ -\sin 2\theta \mathbf{e}_x^f + \cos 2\theta \mathbf{e}_z^f \\ 3 \times 10^5 \text{ km}^{-1} \\ 10^{-6} \end{bmatrix} \quad (8)$$

$$\begin{bmatrix} \mathbf{P}^{AE} \\ P_0^{AE} \\ \bar{\mathbf{P}}^{AE} \\ \bar{P}_0^{AE} \end{bmatrix} = \begin{bmatrix} 2 \mathbf{e}_z^f \\ 4 \\ 1.5 \mathbf{e}_z^f \\ 3.5 \end{bmatrix} \quad (11)$$

and the initial condition

$$\begin{bmatrix} \Gamma_+^{CC} \\ \Gamma_+^{AE} \\ \Gamma_-^{AE} \end{bmatrix} = \begin{bmatrix} 0.5/11.4 \text{ km}^{-1} \\ 0.5/0.417 \text{ km}^{-1} \\ 0.5/0.417 \text{ km}^{-1} \end{bmatrix} \quad (9)$$

$$\begin{bmatrix} \mathbf{P} \\ P_0 \\ \bar{\mathbf{P}} \\ \bar{P}_0 \end{bmatrix} = \begin{bmatrix} 2 \mathbf{e}_z^f \\ 4 \\ 1.5 \mathbf{e}_z^f \\ 3.5 \end{bmatrix} \quad (12)$$

III. RESULTS

A. Linearized EOM

$$\begin{bmatrix} \bar{\Gamma}_+^{CC} \\ \bar{\Gamma}_+^{AE} \\ \bar{\Gamma}_-^{AE} \end{bmatrix} = \begin{bmatrix} 0.5/37.2 \text{ km}^{-1} \\ 0.5/4.36 \text{ km}^{-1} \\ 0.5/4.36 \text{ km}^{-1} \end{bmatrix} \quad (10)$$

When $P_x, P_y \ll P_z \sim 1$, nonlinear terms could be ignored, we have

$$i\partial_t \rho_{ex} = \omega \sin 2\theta P_z + \left[-\omega \cos 2\theta - \sqrt{2}G_F(n_{\bar{\nu}_e} - n_{\bar{\nu}_x}) - i\Gamma \right] \rho_{ex} + \sqrt{2}G_F(n_{\nu_e} - n_{\nu_x}) \bar{\rho}_{ex} \quad (13)$$

$$i\partial_t \bar{\rho}_{ex} = -\omega \sin 2\theta \bar{P}_z + \left[+\omega \cos 2\theta + \sqrt{2}G_F(n_{\nu_e} - n_{\nu_x}) - i\bar{\Gamma} \right] \bar{\rho}_{ex} - \sqrt{2}G_F(n_{\bar{\nu}_e} - n_{\bar{\nu}_x}) \rho_{ex} \quad (14)$$

, where

$$\begin{cases} B \equiv (\sin 2\theta, 0, \cos 2\theta) \\ \rho_{ex} \equiv \rho_{21} = P_x + iP_y \\ \bar{\rho}_{ex} \equiv \bar{\rho}_{21} = \bar{P}_x + i\bar{P}_y \\ \Gamma \equiv \Gamma^{AE} + \Gamma^{CC} \\ \bar{\Gamma} \equiv \bar{\Gamma}^{AE} + \bar{\Gamma}^{CC} \end{cases} \quad (15)$$

Assume

$$\begin{cases} \rho_{ex}(t) = \rho_{ex}^0 + Qe^{-i\Omega t} \\ \bar{\rho}_{ex}(t) = \bar{\rho}_{ex}^0 + \bar{Q}e^{-i\Omega t} \end{cases} \quad (19)$$

Let

, where ρ_{ex}^0 and $\bar{\rho}_{ex}^0$ satisfy

$$\begin{cases} \mu_+ \equiv \left(\frac{S+D}{2}\right)\mu \\ \mu_- \equiv \left(\frac{S-D}{2}\right)\mu \end{cases} \quad (16) \quad 0 = \begin{bmatrix} \omega \sin 2\theta \\ -\omega \sin 2\theta \end{bmatrix} + A \begin{bmatrix} \rho_{ex}^0 \\ \bar{\rho}_{ex}^0 \end{bmatrix} \quad (20)$$

and

Then, from equation 18, we have

$$A = \begin{bmatrix} -\omega \cos 2\theta - \mu_- - i\Gamma & \mu_+ \\ -\mu_- & \omega \cos 2\theta + \mu_+ - i\bar{\Gamma} \end{bmatrix} \quad (17) \quad \Omega \begin{bmatrix} Q \\ \bar{Q} \end{bmatrix} = A \begin{bmatrix} Q \\ \bar{Q} \end{bmatrix} \quad (21)$$

, we have

$$i\partial_t \begin{bmatrix} \rho_{ex} \\ \bar{\rho}_{ex} \end{bmatrix} = \begin{bmatrix} \omega \sin 2\theta \\ -\omega \sin 2\theta \end{bmatrix} + A \begin{bmatrix} \rho_{ex} \\ \bar{\rho}_{ex} \end{bmatrix} \quad (18)$$

Notice that, P_z and \bar{P}_z are both close to 1 under this limitation condition.

IV. DISCUSSION

V. CONCLUSIONS

* yuchialin@unm.edu