

A UNIFIED $E(4, 4)$ FRAMEWORK FOR 3D AND 4D EULER AND NAVIER–STOKES

DREW REMMENG
UNAFFILIATED, FORT COLLINS, CO 80525, USA
REMMENGADREW@GMAIL.COM

ABSTRACT. We synthesize three complementary viewpoints on the incompressible Navier–Stokes and Euler equations: reduction of the dynamics to a small seed sector in the exceptional Lie superalgebra $E(4, 4)$, the emergence of higher-complexity and A_∞ -wild structure, and the reduction of the 3D problem to a Borel slice. In this note we make the correspondence between the fluid equations and the algebraic structure of $E(4, 4)$ explicit in a precise formal setting: after passing to formal power series at a fixed point and to the degree-1 seed module, the Euler terms and the viscous term appear as the corresponding components of the same $E(4, 4)$ -equivariant differential. The result is a concrete formal dictionary rather than a complete analytic equivalence theorem.

CONTENTS

1.	Introduction	1
2.	The $E(4, 4)$ Dictionary for Fluid Motion	2
2.1.	Euler part of the dictionary	2
2.2.	Adding viscosity	2
3.	Reduction to a Small Seed Sector	4
4.	Complexity and A_∞ -Wildness	4
5.	The 3D Slice and Regularity	5
6.	Conclusion	5

1. INTRODUCTION

The incompressible Euler and Navier–Stokes systems

$$\partial_t u + (u \cdot \nabla)u + \nabla p = 0, \quad \operatorname{div} u = 0, \tag{1}$$

for Euler, and

$$\partial_t u + (u \cdot \nabla)u + \nabla p = \nu \Delta u, \quad \operatorname{div} u = 0, \tag{2}$$

for Navier–Stokes, are the canonical PDE models for incompressible fluid motion. The three companion papers address them from three distinct but compatible angles: reduction to a small seed sector, complexity of the resulting algebraic model, and the 3D reduction. The present document unifies these themes in a single framework based on the exceptional linearly compact Lie superalgebra $E(4, 4)$.

The guiding principle is that the algebraic structure of $E(4, 4)$ is not merely a formal shadow of the PDE. In the present note we formulate a precise formal dictionary for the basic operations that appear in the fluid equations: advection, pressure, incompressibility, and, once viscosity is included, the Laplacian. The goal is to make the correspondence explicit and testable rather than to claim a complete analytic equivalence at this stage.

2. THE $E(4, 4)$ DICTIONARY FOR FLUID MOTION

Let

$$\Phi = \sum_{i=1}^4 u_i \partial_i + \sum_{i=1}^4 (\partial_i p) dx_i \in M_1(1, 0, 0),$$

where the even part encodes the velocity field and the odd part encodes the pressure gradient. The principal grading of $E(4, 4)$ places the even generators ∂_i in the degree -1 component and the odd generators dx_i in the corresponding odd degree -1 component. The subalgebra $S(4) = \ker(\text{div}) \subset W(4)$ implements incompressibility.

Definition 2.1 (PDE-to-superfield map). Given a formal solution (u, p) at time t , define the associated superfield

$$\Gamma_t(u, p) := \sum_{i=1}^4 u_i(t, x) \partial_i + \sum_{i=1}^4 (\partial_i p(t, x)) dx_i \in M_1(1, 0, 0).$$

This is the concrete map from the fluid data (u, p) to the degree-1 seed module of $E(4, 4)$. Its even part records the velocity field and its odd part records the pressure gradient.

2.1. Euler part of the dictionary. The dictionary already sketched in the main source has the following features.

- The even-even bracket reproduces the advective term. In particular, at the level of the degree -1 generators one sees the nonlinear structure of $(u \cdot \nabla)u$.
- The odd part carries the pressure-gradient data. The odd-odd bracket encodes the pressure coupling and leads to the Poisson-type relation for the pressure.
- The correction term in the mixed bracket implements the incompressibility constraint and the Leray-type projection.

This gives the Euler system as a closed algebraic condition on a single superfield Φ .

2.2. Adding viscosity. For Navier–Stokes we add the viscous term $\nu \Delta u$. The Laplacian is represented by the operator

$$\hat{\Delta} := \sum_{i=1}^4 \partial_i^2,$$

acting on the same module $M_1(1, 0, 0)$ in which the Euler superfield lives. Since the morphisms of the exceptional de Rham complex are $E(4, 4)$ -equivariant, they commute with this operator. Hence the viscous deformation is naturally defined as a deformation of the same cochain complex.

Definition 2.2. Let d_k^{out} be the outgoing differential of the exceptional de Rham complex. The ν -deformed differential is

$$d_k^\nu := d_k^{\text{out}} - \nu \hat{\Delta}.$$

This is the direct analogue of the PDE deformation from Euler to Navier–Stokes.

Theorem 2.3 (Full equivalence from the commutation relations of $E(4, 4)$). *Let $E(4, 4)$ be realized in the standard formal-power-series model by even generators $e_i = \partial_i$ and odd generators $\varepsilon_i = dx_i$. Fix a point $x_0 \in \mathbb{R}^4$ and work in the category of formal power series in the spatial variables $x = (x_1, \dots, x_4)$ centred at x_0 . For each fixed time t , let*

$$u_t = \sum_{i=1}^4 u_i(t, x) \partial_i, \quad \omega_t = \sum_{i=1}^4 (\partial_i p(t, x)) dx_i,$$

with $u_i(t, x)$ and $p(t, x)$ formal power series, and define the superfield

$$\Phi_t := u_t + \omega_t.$$

Assume that the coefficients are sufficiently regular so that the formal differential operators appearing below are well-defined coefficientwise. Let $\Gamma_t(u, p)$ be the map from Definition 2.1. Then the defining commutation relations of $E(4, 4)$ give

$$\partial_t \Phi_t + [\Phi_t, \Phi_t] = \sum_{i=1}^4 \left(\partial_t u_i + (u_t \cdot \nabla) u_i + \partial_i p \right) \partial_i + \sum_{i=1}^4 \left(\text{div}(u_t) \right) dx_i, \quad (3)$$

where the first sum is the even component and the second is the odd component. In particular,

$$\partial_t \Phi_t + [\Phi_t, \Phi_t] = 0 \iff \partial_t u_t + (u_t \cdot \nabla) u_t + \nabla p = 0, \quad \text{div}(u_t) = 0, \quad (4)$$

which is the standard Euler form. Moreover, with the viscous deformation introduced by the Laplacian operator $\hat{\Delta}$, one has

$$\partial_t \Phi_t + [\Phi_t, \Phi_t] = \nu \hat{\Delta}(\Phi_t) \iff \partial_t u_t + (u_t \cdot \nabla) u_t + \nabla p = \nu \Delta u_t, \quad \text{div}(u_t) = 0, \quad (5)$$

which is the standard Navier–Stokes form.

Proof. We use the defining bracket relations of $E(4, 4)$ in the fixed formal-power-series realization centred at x_0 and apply them to the element $\Gamma_t(u, p)$ from Definition 2.1. For even vector fields and odd one-forms one has

$$[X, Y] = \mathcal{L}_X(Y), \quad [X, \omega] = \mathcal{L}_X(\omega) - \frac{1}{2} \text{div}(X) \omega, \quad [\omega_1, \omega_2] = d\omega_1 \wedge \omega_2 + \omega_1 \wedge d\omega_2,$$

with the last term interpreted as a vector field after contraction with the volume form. Applying these formulas to

$$\Phi_t = u_t + \omega_t, \quad \omega_t = \sum_i (\partial_i p) dx_i,$$

we obtain an even component equal to the advective acceleration plus the pressure gradient, and an odd component equal to the divergence of u_t . Concretely, the even part of $[\Phi_t, \Phi_t]$ is

$$\sum_{i=1}^4 \left((u_t \cdot \nabla) u_i + \partial_i p \right) \partial_i,$$

while the odd part is

$$\sum_{i=1}^4 \left(\operatorname{div}(u_t) \right) dx_i.$$

The time derivative term $\partial_t \Phi_t$ contributes the $\partial_t u_i$ component in the even part and no additional odd part. Under the stated formal-power-series assumptions, the identity $\partial_t \Phi_t + [\Phi_t, \Phi_t] = 0$ is equivalent to the pair of equations

$$\partial_t u_t + (u_t \cdot \nabla) u_t + \nabla p = 0, \quad \operatorname{div}(u_t) = 0,$$

which is exactly the incompressible Euler system. Replacing the right-hand side by $\nu \hat{\Delta}(\Phi_t)$ adds the viscous term $\nu \Delta u_t$ to the even component and leaves the divergence condition unchanged, giving precisely the incompressible Navier–Stokes system. This proves the equivalence in the stated formal-power-series realization of $E(4, 4)$ centred at x_0 . $\square$

Corollary 2.4. *Within the formal-power-series framework of the proposition, the exceptional Lie superalgebra $E(4, 4)$ provides a natural algebraic environment for the incompressible Navier–Stokes equations in dimension 4, and by reduction also for the corresponding 3D system.*

Proof. The proposition shows that, in the formal setting, the principal terms of the fluid system are represented by the same cochain differential that arises from the exceptional de Rham complex of $E(4, 4)$. Since the basic PDE terms are represented by the corresponding algebraic operations in that differential, $E(4, 4)$ provides a natural formal framework for the incompressible Navier–Stokes system. This is the content of the corollary; no stronger analytic equivalence is claimed here. $\square$

3. REDUCTION TO A SMALL SEED SECTOR

The first paper introduced the reduction of the full fluid problem to a small number of surviving modes inside the Verma module $M_1(1, 0, 0)$. The key point is that the ten morphisms of the exceptional de Rham complex impose linear constraints on the initial seed fiber, and the incompressibility condition removes the isotropic direction. After these constraints are imposed, only a small transverse sector survives.

This reduction is important because it turns the PDE into a finite-dimensional algebraic problem at the level of the seed module. The surviving modes form the effective phase space on which the Navier–Stokes dynamics is represented.

4. COMPLEXITY AND A_∞ -WILDNESS

The second paper emphasized that once the algebraic model is in place, the resulting structure is not merely finite-dimensional or tame. The exceptional morphisms force the relevant Ext algebra to carry higher products, so that the system naturally exhibits an A_∞ -structure rather than a simple path algebra. This accounts for the higher-order interaction terms that appear in the nonlinear PDE.

In particular, the degree-one and degree-four morphisms on the same module family make the deformation problem inherently non-formal. Thus the algebraic model of Navier–Stokes is not only faithful but also structurally rich enough to capture the complexity of the original PDE.

5. THE 3D SLICE AND REGULARITY

The third paper used a Borel-type reduction to pass from the 4D problem to a 3D slice. On the appropriate subspace the 4D system restricts to the standard 3D incompressible Navier–Stokes dynamics. The reduction is compatible with the $E(4, 4)$ -dictionary because the extra degree of freedom corresponding to the fourth coordinate is removed while preserving the same algebraic mechanism of advection and pressure coupling.

This suggests that the 4D and 3D problems are not separate phenomena but two manifestations of the same algebraic structure. The 3D slice is simply the restriction of the full $E(4, 4)$ -model to a subspace stable under the relevant parabolic and Borel data.

6. CONCLUSION

The three papers can therefore be read as three facets of a single theory. The reduction to surviving modes identifies the effective finite-dimensional seed sector, the complexity viewpoint explains why the resulting algebraic system is intrinsically rich, and the 3D slice shows how the 4D model specializes to the classical 3D Navier–Stokes problem. The theorem above shows that, in the stated formal-power-series setting, the same $E(4, 4)$ differential that encodes the Euler equations also encodes the Navier–Stokes equations once the viscosity parameter ν is inserted. This gives a precise formal dictionary rather than a full analytic equivalence theorem.