

# MATH 498 HW2

Drew Remmenga

1.a.

```
library(splines)
HW2Test<- function(x){
  pgamma( x, shape =10, scale =.02) - .5*x
}
xGrid <- seq( 0,1, length.out=200)
y<-HW2Test(xGrid)
fit1<-lm(y ~ poly(xGrid, 17, raw=TRUE))
lm(y ~ poly(xGrid, 17, raw=TRUE))
```

```
##
## Call:
## lm(formula = y ~ poly(xGrid, 17, raw = TRUE))
##
## Coefficients:
##              (Intercept)      poly(xGrid, 17, raw = TRUE)1
##              7.864e-03              -3.314e+00
## poly(xGrid, 17, raw = TRUE)2      poly(xGrid, 17, raw = TRUE)3
##              1.851e+02              -4.771e+03
## poly(xGrid, 17, raw = TRUE)4      poly(xGrid, 17, raw = TRUE)5
##              6.071e+04              -4.262e+05
## poly(xGrid, 17, raw = TRUE)6      poly(xGrid, 17, raw = TRUE)7
##              1.855e+06              -5.356e+06
## poly(xGrid, 17, raw = TRUE)8      poly(xGrid, 17, raw = TRUE)9
##              1.060e+07              -1.447e+07
## poly(xGrid, 17, raw = TRUE)10     poly(xGrid, 17, raw = TRUE)11
##              1.337e+07              -7.820e+06
## poly(xGrid, 17, raw = TRUE)12     poly(xGrid, 17, raw = TRUE)13
##              2.355e+06              NA
## poly(xGrid, 17, raw = TRUE)14     poly(xGrid, 17, raw = TRUE)15
##              -1.734e+05              NA
## poly(xGrid, 17, raw = TRUE)16     poly(xGrid, 17, raw = TRUE)17
##              1.297e+04              NA
```

```
fit2<-lm(y ~ poly(xGrid, 15, raw=TRUE))
lm(y ~ poly(xGrid, 15, raw=TRUE))
```

```
##
## Call:
## lm(formula = y ~ poly(xGrid, 15, raw = TRUE))
##
## Coefficients:
```

```
##               (Intercept)  poly(xGrid, 15, raw = TRUE)1
##               4.063e-03      -2.274e+00
## poly(xGrid, 15, raw = TRUE)2  poly(xGrid, 15, raw = TRUE)3
##               1.279e+02      -3.457e+03
## poly(xGrid, 15, raw = TRUE)4  poly(xGrid, 15, raw = TRUE)5
##               4.458e+04      -3.062e+05
## poly(xGrid, 15, raw = TRUE)6  poly(xGrid, 15, raw = TRUE)7
##               1.274e+06      -3.451e+06
## poly(xGrid, 15, raw = TRUE)8  poly(xGrid, 15, raw = TRUE)9
##               6.292e+06      -7.762e+06
## poly(xGrid, 15, raw = TRUE)10 poly(xGrid, 15, raw = TRUE)11
##               6.324e+06      -3.162e+06
## poly(xGrid, 15, raw = TRUE)12 poly(xGrid, 15, raw = TRUE)13
##               7.798e+05      NA
## poly(xGrid, 15, raw = TRUE)14 poly(xGrid, 15, raw = TRUE)15
##               -2.989e+04      NA
```

```
fit3<-lm(y ~ poly(xGrid, 13, raw=TRUE))
lm(y ~ poly(xGrid, 13, raw=TRUE))
```

```
##
## Call:
## lm(formula = y ~ poly(xGrid, 13, raw = TRUE))
##
## Coefficients:
##               (Intercept)  poly(xGrid, 13, raw = TRUE)1
##               -8.951e-03      7.375e-01
## poly(xGrid, 13, raw = TRUE)2  poly(xGrid, 13, raw = TRUE)3
##               -1.645e+01      -5.663e+02
## poly(xGrid, 13, raw = TRUE)4  poly(xGrid, 13, raw = TRUE)5
##               1.361e+04      -1.058e+05
## poly(xGrid, 13, raw = TRUE)6  poly(xGrid, 13, raw = TRUE)7
##               4.349e+05      -1.091e+06
## poly(xGrid, 13, raw = TRUE)8  poly(xGrid, 13, raw = TRUE)9
##               1.759e+06      -1.840e+06
## poly(xGrid, 13, raw = TRUE)10 poly(xGrid, 13, raw = TRUE)11
##               1.208e+06      -4.536e+05
## poly(xGrid, 13, raw = TRUE)12 poly(xGrid, 13, raw = TRUE)13
##               7.435e+04      NA
```


```
fit4<-lm(y ~ poly(xGrid, 11, raw=TRUE))
lm(y ~ poly(xGrid, 11, raw=TRUE))
```

```
##
## Call:
## lm(formula = y ~ poly(xGrid, 11, raw = TRUE))
##
## Coefficients:
##               (Intercept)  poly(xGrid, 11, raw = TRUE)1
##               -2.856e-02      4.507e+00
## poly(xGrid, 11, raw = TRUE)2  poly(xGrid, 11, raw = TRUE)3
##               -1.706e+02      2.076e+03
## poly(xGrid, 11, raw = TRUE)4  poly(xGrid, 11, raw = TRUE)5
```

```
##                -1.054e+04                2.688e+04
## poly(xGrid, 11, raw = TRUE)6    poly(xGrid, 11, raw = TRUE)7
##                -3.220e+04                3.614e+02
## poly(xGrid, 11, raw = TRUE)8    poly(xGrid, 11, raw = TRUE)9
##                4.987e+04                -6.389e+04
## poly(xGrid, 11, raw = TRUE)10   poly(xGrid, 11, raw = TRUE)11
##                3.509e+04                -7.484e+03
```

1.b.

```
plot(xGrid, HW2Test(xGrid), col="Red")
pred1<- predict(fit1)
pred2<- predict(fit2)
pred3<- predict(fit3)
pred4<- predict(fit4)
lines(xGrid, pred1, col='Blue', lwd=2)
lines(xGrid, pred2, col='Green', lwd=2)
lines(xGrid, pred3, col='Black', lwd=2)
lines(xGrid, pred4, col='Yellow', lwd=2)
legend("topleft", legend=c("Actual", "Degree 17", "Degree 15", "Degree 13", "Degree 11"),
      col=c("Red", "Blue", "Green", "Black", "Yellow"), lty=1:2, cex=0.8)
title("Different degree Polynomials fit to the Gamma function")
```



1.c. Degrees 17,15,13, and 11.

```
sqrt( mean( ( y - pred1)^2 ))
```

```
## [1] 0.001742046
```

```
sqrt( mean( ( y - pred2)^2 ))
```

```
## [1] 0.002157356
```

```
sqrt( mean( ( y - pred3)^2 ))
```

```
## [1] 0.004626954
```

```
sqrt( mean( ( y - pred4)^2 ))
```


```
## [1] 0.007413881
```

2.a.

```
xGrid<- seq( 0,1,length.out=150)
KN<- seq( 0,1,length.out=10)
naturalSplineBasis <- function(sGrid,
                                sKnots,
                                degree = 3,
                                derivative = 0) {
  boundaryKnots<- c( min(sKnots),max(sKnots))
  sKnots0<- c( rep( boundaryKnots[1],degree),sort(sKnots),
              rep( boundaryKnots[2],degree) )
  testRight<- sGrid < min(sKnots)
  testLeft <- sGrid > max(sKnots)
  if( any(testRight |testLeft) )
  {stop("some points for evaluation outside knot range.")}

  basis <- splineDesign(sKnots0, sGrid,
                        ord= degree+1, outer.ok=TRUE,
                        derivs=derivative)
  # set up constraints to enforce natural BCs.
  const <- splineDesign(sKnots0, boundaryKnots, ord = degree+1,
                        derivs = c(2,2))
  qr.const <- qr(t(const))
  QBasis<- t(qr.qty( qr.const, t(basis) ))
  basis <- QBasis[,-(1:2)]
  basis

  return( basis )
}
B<- naturalSplineBasis( xGrid, sKnots=KN)
matplot( xGrid, B, type="l", lty=1)
```



Just one. 2.b.

```
NBSFit<- function( x, y, xGrid){
N<- length( x)
AData<- naturalSplineBasis( x, sKnots=x)
coef<- solve( AData, y)
AGrid<- naturalSplineBasis( xGrid, sKnots=x )
yFit <-AGrid%*%coef
return( yFit)
}
y <- HW2Test(xGrid)
NBSFit(xGrid,y,xGrid)
```

```
##           [,1]
## [1,] -2.220446e-16
## [2,] -3.355705e-03
## [3,] -6.711407e-03
## [4,] -1.006700e-02
## [5,] -1.342126e-02
## [6,] -1.676781e-02
## [7,] -2.008511e-02
## [8,] -2.331980e-02
## [9,] -2.636568e-02
## [10,] -2.904338e-02
## [11,] -3.108679e-02
## [12,] -3.214119e-02
```

```
## [13,] -3.177501e-02
## [14,] -2.950489e-02
## [15,] -2.483043e-02
## [16,] -1.727415e-02
## [17,] -6.421174e-03
## [18,] 8.046000e-03
## [19,] 2.632126e-02
## [20,] 4.845925e-02
## [21,] 7.437099e-02
## [22,] 1.038300e-01
## [23,] 1.364874e-01
## [24,] 1.718933e-01
## [25,] 2.095226e-01
## [26,] 2.488027e-01
## [27,] 2.891401e-01
## [28,] 3.299456e-01
## [29,] 3.706559e-01
## [30,] 4.107509e-01
## [31,] 4.497671e-01
## [32,] 4.873061e-01
## [33,] 5.230393e-01
## [34,] 5.567090e-01
## [35,] 5.881264e-01
## [36,] 6.171677e-01
## [37,] 6.437677e-01
## [38,] 6.679132e-01
## [39,] 6.896354e-01
## [40,] 7.090017e-01
## [41,] 7.261088e-01
## [42,] 7.410753e-01
## [43,] 7.540354e-01
## [44,] 7.651329e-01
## [45,] 7.745168e-01
## [46,] 7.823371e-01
## [47,] 7.887410e-01
## [48,] 7.938712e-01
## [49,] 7.978632e-01
## [50,] 8.008444e-01
## [51,] 8.029328e-01
## [52,] 8.042372e-01
## [53,] 8.048563e-01
## [54,] 8.048795e-01
## [55,] 8.043867e-01
## [56,] 8.034492e-01
## [57,] 8.021297e-01
## [58,] 8.004837e-01
## [59,] 7.985594e-01
## [60,] 7.963989e-01
## [61,] 7.940386e-01
## [62,] 7.915097e-01
## [63,] 7.888391e-01
## [64,] 7.860498e-01
## [65,] 7.831612e-01
## [66,] 7.801898e-01
```

```
## [67,] 7.771498e-01
## [68,] 7.740527e-01
## [69,] 7.709085e-01
## [70,] 7.677254e-01
## [71,] 7.645103e-01
## [72,] 7.612689e-01
## [73,] 7.580060e-01
## [74,] 7.547254e-01
## [75,] 7.514305e-01
## [76,] 7.481239e-01
## [77,] 7.448078e-01
## [78,] 7.414839e-01
## [79,] 7.381538e-01
## [80,] 7.348186e-01
## [81,] 7.314794e-01
## [82,] 7.281368e-01
## [83,] 7.247916e-01
## [84,] 7.214443e-01
## [85,] 7.180953e-01
## [86,] 7.147449e-01
## [87,] 7.113934e-01
## [88,] 7.080411e-01
## [89,] 7.046880e-01
## [90,] 7.013344e-01
## [91,] 6.979804e-01
## [92,] 6.946260e-01
## [93,] 6.912714e-01
## [94,] 6.879165e-01
## [95,] 6.845614e-01
## [96,] 6.812062e-01
## [97,] 6.778509e-01
## [98,] 6.744955e-01
## [99,] 6.711401e-01
## [100,] 6.677845e-01
## [101,] 6.644290e-01
## [102,] 6.610734e-01
## [103,] 6.577178e-01
## [104,] 6.543622e-01
## [105,] 6.510065e-01
## [106,] 6.476509e-01
## [107,] 6.442952e-01
## [108,] 6.409395e-01
## [109,] 6.375838e-01
## [110,] 6.342281e-01
## [111,] 6.308724e-01
## [112,] 6.275167e-01
## [113,] 6.241610e-01
## [114,] 6.208053e-01
## [115,] 6.174496e-01
## [116,] 6.140939e-01
## [117,] 6.107382e-01
## [118,] 6.073825e-01
## [119,] 6.040268e-01
## [120,] 6.006711e-01
```

```

## [121,] 5.973154e-01
## [122,] 5.939597e-01
## [123,] 5.906040e-01
## [124,] 5.872483e-01
## [125,] 5.838926e-01
## [126,] 5.805369e-01
## [127,] 5.771812e-01
## [128,] 5.738255e-01
## [129,] 5.704698e-01
## [130,] 5.671141e-01
## [131,] 5.637584e-01
## [132,] 5.604027e-01
## [133,] 5.570470e-01
## [134,] 5.536913e-01
## [135,] 5.503356e-01
## [136,] 5.469799e-01
## [137,] 5.436242e-01
## [138,] 5.402685e-01
## [139,] 5.369128e-01
## [140,] 5.335570e-01
## [141,] 5.302013e-01
## [142,] 5.268456e-01
## [143,] 5.234899e-01
## [144,] 5.201342e-01
## [145,] 5.167785e-01
## [146,] 5.134228e-01
## [147,] 5.100671e-01
## [148,] 5.067114e-01
## [149,] 5.033557e-01
## [150,] 5.000000e-01

```

Sines and cosines are mutually orthogonal.  $\int_{-\infty}^{\infty} \sqrt{2}\cos(2\pi x) * \sqrt{2}\cos(2\pi x) = 1$   $\int_{-\infty}^{\infty} \sqrt{2}\sin(2\pi x) * \sqrt{2}\sin(2\pi x) = 1$   $\int_{-\infty}^{\infty} \sqrt{2}\cos(4\pi x) * \sqrt{2}\cos(4\pi x) = 1$   $\int_{-\infty}^{\infty} \sqrt{2}\sin(4\pi x) * \sqrt{2}\sin(4\pi x) = 1$