

Integral Equations Notes: [457]

[Drew Remmenga]

June 18, 2024

Definition 1. *An integral equation is any equation in which the unknown function is inside the integral sign.*

Example 1.

$$\int_a^b k(x, y)u(y)dy = f(x), x \in (a, b)$$

Here k and f are given and $u(x)$ is the unknown function.

Example 2.

$$\int_0^\infty e^{-xt}u(t)dt = f(x), x \in (0, \infty)$$

$f(x)$ is the laplace transform of u .

Remark 1. *There is an inversion formula giving u from f but it involves an integrall of $f(x)$ over complex values of x . In practice we may only know $f(x)$ for real values of x .*

Example 3.

$$\int_{-1}^1 u(t)dt = 1$$

One solution

$$u(t) = \frac{1}{2}, t \in (-1, 1)$$

Clearly there are lots of solutions, e.g.

$$u(t) = \frac{1}{2} +$$

any odd function of t .

or

$$u(t) = \frac{1}{2} + \sum_{n=1}^{\infty} \{a_n \cos(nt\pi) + b_n \sin(nt\pi)\}$$

For any reasonable a_n and b_n .

Example 4.

$$\int_{-1}^1 u(t)dt = f(x), x \in (-1, 1)$$

Now there are no solutions unless $f(x)$ is constant.

Example 5.

$$\alpha u(x) + \beta \int_{-1}^1 u(t)dt = f(x), x \in (-1, 1) \quad (1)$$

Where α and β are given constants with $\alpha \neq 0$ or example 4 put $\int_{-1}^1 u(t)dt = c$ a known constant. Then (1) gives

$$u(x) = \frac{1}{\alpha} \{f(x) - \beta C\} \quad (2)$$

So we will have solved (1) and know C . Integrate (2) gives

$$C = \int_{-1}^1 u(x)dx = \frac{1}{\alpha} \int_{-1}^1 f(x)dx - \frac{\beta C}{\alpha} \int_{-1}^1 dx$$

$$(1 + \frac{2\beta}{\alpha})C = \frac{1}{\alpha} \int_{-1}^1 f(x)dx \quad (3)$$

$$C = \frac{1}{\alpha + 2\beta} \int_{-1}^1 f(x)dx$$

then (2) gives unique solution of (1) real provided $\alpha + 2\beta \neq 0$. What happens if $\alpha + 2\beta = 0$? As before we get to (3) which reduces to

$$0 \int_{-1}^1 f(x)dx \quad (4)$$

If $f(x)$ satisfies (4) eg f could be odd then (2) solves (1) for any choice of the constant C : we have existence but not uniqueness. If $f(x)$ does not satisfy (4) eg $f(x)$ could be x^2 then (1) does not have any solutions $\alpha + 2\beta = 0$. All this is reminiscent of linear algebra (solving $Ax = b$).

Exercise 1. Put $\frac{\beta}{\alpha} = -\lambda$ and $f = 0$ giving

$$u(x) = \int_{-1}^1 u(t)dt = 0$$

A homogenous integral equation. Are there values of λ for which this integral equation has non trivial solution $u \neq 0$?

Example 6.

$$\int_0^t u(\tau)d\tau = f(t), t > 0 \quad (5)$$

Differentiate with respect to $x(t)$:

$$u(x) = f'(x)$$

Two anxieties. 1. (5) suggests that $f(0)=0$ in $\lim_{x \rightarrow 0}$ (5) what happens if $f(0) \neq 0$ eg $f(x) = 1, \forall x$ 2. What happens if f is not differentiable? Try using laplace transform. Let $U(s) = \mathcal{L}\{u\}$

$$\int_0^\infty u(t)e^{-st}dt$$

We have:

$$\begin{aligned}
\mathcal{L}\left\{\int_0^t u(\tau) d\tau\right\} &= \int_{t=0}^{\infty} e^{-st} \int_0^{\tau=t} u(\tau) d\tau \\
&= \int_0^{\infty} u(\tau) \int_{\tau}^{\infty} e^{-st} dt d\tau \\
&= \frac{e^{st}}{s} \Big|_{\tau}^{\infty} = \frac{e^{-s\tau}}{s} \\
&= \int_0^{\infty} u(\tau) \frac{e^{-s\tau}}{s} d\tau \\
&= \frac{1}{s} \int_0^{\infty} u(\tau) e^{-s\tau} d\tau \\
&= \frac{1}{s} \mathcal{L}\{u\} \\
\mathcal{L}\{(5)\} &= \frac{1}{s} U(s) = F(s) \\
&\implies \\
u(s) &= sF(s) \\
\mathcal{L}\{f'\} &= sF(s) - f(0) \\
\mathcal{L}\{u\} &= U = sF(s) + sf(x) - f(0) \\
&= \mathcal{L}\{f\} + f(0)
\end{aligned}$$

Question is is there a $g(t)$ such that $\mathcal{L}\{g\} = 1$ Answer: It depends on what you mean by a function.

$$LHS = \mathcal{L}\{g\} = \int_0^{\infty} g(t) e^{st} dt = G(s)$$

Fact: If g is an ordinary integrable function then we must not have $G(s) \rightarrow 0$ as $s \rightarrow \infty$ As we want $G(s) = 1$. g can't be ordinary. Fact: $\mathcal{L}\{\delta\} = 1$ Where $\delta =$ dirac delta function. SO solution of (5).

$$u(x) = f'(x) + f(0)dx$$

In particular if $f(0) \neq 0$ then (5) has no continuous solutions for $x \geq 0$. Second Anxiety: Consider (5) where f is not a continuous function.

Example 7.

$$f(x) = \begin{cases} 0, & 0 \leq x < c \\ x - c, & x \geq c \end{cases} = (x - c)H(x - c)$$

$H(x)$ is the divided unit function. Then

$$\begin{aligned}
\mathcal{L}\{(5)\} &= \frac{1}{s} u(s) = \mathcal{L}\{f\} = \mathcal{L}\{x - c\}H(x - c) \\
&= \int_0^{\infty} (x - c)H(x - c)e^{-sx} dx \\
&= \int_0^{\infty} (x - c)e^{sx} dx, (x - c) = y \\
&= \int_0^{\infty} ye^{-s(y+c)} dy \\
&= e^{-sc} \int_0^{\infty} yce^{-sy} dy \\
&= e^{-sc} \mathcal{L}\{t\} = \frac{e^{-sc}}{s^2} \\
&\implies U(s) = se^{-sc} \\
u(x) &= H(x - c)
\end{aligned}$$

Example 8.

$$u(x) = \frac{\alpha}{\pi} \int_{-1}^1 \frac{u(t)dt}{(x-2)^2 + \alpha^2} = 1, x \in (-1, 1)$$

The integral contains a parameter α . It turns up in at least two distinct physical applications. One in electro statics. whgere it is called Love's equation and in quantum phases where it is called Lieb's equation. It looks simple byt no explicit solution is known. Soi what is known is that $u(x)$ exists, its continuous, it's unique, and it's even. Numerical solutions are easy to compute unless α is very small.

1 Abel's Equation

A bead starts froim rest at P and slides down a smooth wire riching 0 at time t_0 Find t_p . Motion is under gravity in a vertical plane put point P at (x,y) and Q at (ξ, δ) . Let s denote the length from 0 to Q. Energy at P = Energy at Q. m = mass of the bead.

$$mgy = mg\delta + \frac{1}{2}m\left(\frac{ds}{dt}\right)^2$$
$$\frac{ds}{dt} = -\sqrt{2gz(y-\delta)}$$

integrating

$$t_p = \int_0^{t_p} dt = \int_{\delta=y}^{\delta=0} \frac{ds}{\sqrt{2g(y-\delta)}}$$

Next specify the shape (equation) of the wire. Its convienient to suppost that $s = S(\delta)$ for some function S. Thus $ds = S'(\gamma)d\gamma$

$$t_p = \int_y^0 \frac{s'(\gamma)d\gamma}{\sqrt{2g(y-\gamma)}}$$

This gives t_p if we know the shape of the wire. Abel asked given t_p can we find S?