PH 221 Week 8


Benjamin Bauml

Winter 2025

This material is borrowed/adapted from Chapter 10 of the Student Workbook for Physics for Scientists and Engineers.

R8-1: Interpreting a Potential Energy Graph

A particle with the potential energy shown in the graph is moving to the right at x = 0 m with total energy E.




- (a) At what value or values of x is the particle's speed a maximum?
- (b) At what value or values of x is the particle's speed a minimum?
- (c) At what value or values of x is the potential energy a maximum?
- (d) Does this particle have a turning point in the range of x covered by the graph? If so, where?
- (e) In which intervals of x is the force on the particle to the right?
- (f) In which intervals of x is the force on the particle to the left?
- (g) At what value or values of x is the magnitude of the force a maximum?
- (h) At what value or values of x are positions of stable equilibrium?
- (i) At what value or values of x are positions of unstable equilibrium?
- (j) If the particle is released from rest at x = 0 m, will it reach x = 10m? Explain.

R8-2: Producing a Potential Energy Graph

Below is a set of axes on which you are going to draw a potential energy curve. By doing experiments, you find the following information:

- A particle with energy E_1 oscillates between positions D and E.
- A particle with energy E_2 oscillates between positions C and F.
- A particle with energy E_3 oscillates between positions B and G.
- \bullet A particle with energy E_4 enters from the right, bounces at A, then never returns.

Draw a potential energy curve that is consistent with this information.




R8-3: Potential versus Change in Potential

- (a) If the force on a particle at some point in space is zero, must its potential energy also be zero at that point? Explain.
- (b) If the potential energy of a particle at some point in space is zero, must the force on it also be zero at that point? Explain.

R8-4: Sliding Cube in a Bowl

A small cube of mass m slides back and forth in a frictionless, hemispherical bowl of radius R. Suppose the cube is released at angle θ . What is the cube's speed at the bottom of the bowl?



- (a) Begin by drawing a before-and-after pictorial representation. Let the cube's initial position and speed be y_i and v_i . Use a similar notation for the final position and speed.
- (b) At the initial position, are either K_i or U_{Gi} zero? If so, which?
- (c) At the final position, are either K_f or U_{Gf} zero? If so, which?
- (d) Does thermal energy need to be considered in this situation? Why or why not?
- (e) Write the conservation of energy equation in terms of position and speed variables, omitting any terms that are zero.
- (f) You're given not the initial position but the initial angle. Do the geometry and trigonometry to find y_i in terms of R and θ .
- (g) Use your result of part (f) in the energy conservation equation, and then finish solving the problem.