Math 510 **Fall 2026**
Linear Algebra II **Drew Armstrong**

Contents

1 Inner Product Spaces **2**
1.1 Vector Spaces . . . 2
1.2 Inner Products and Norms . . . 6
1.3 Subspaces and Dimension . . . 9
1.4 Orthogonal Projection . . . 17
1.5 Fourier Series . . . 24
1.6 Inner Product Spaces Over $\mathbb{C}$ . . . 28

2 Linear Functions **33**
2.1 Matrix Multiplication . . . 33
2.2 Linear Functions . . . 37
2.3 Matrix Analysis . . . 43
2.4 Rotations, Reflections, Projections . . . 51
2.5 Unitary and Orthogonal Matrices . . . 56

3 Subspaces Associated to Linear Functions **60**
3.1 The Rank-Nullity Theorem . . . 60
3.2 Subspaces Associated to Matrices . . . 67
3.3 Linear Systems and Affine Subspaces . . . 70
3.4 Matrix Elimination . . . 74
3.5 Least Squares Approximation . . . 83

4 Linear and Bilinear Forms **94**
4.1 Linear Forms . . . 94
4.2 Bilinear Forms . . . 97
4.3 Quadratic Forms . . . 101
4.4 Multivariable Taylor Expansion . . . 105

5 Determinants **109**
5.1 Multilinear Forms . . . 109
5.2 Uniqueness of the Determinant . . . 116
5.3 Algebraic Properties of the Determinant . . . 120
5.4 Formulas for the Determinant . . . 122
5.5 Cramer's Rule (Optional) . . . 125
5.6 Geometric Interpretation . . . 128
5.7 Application to Calculus . . . 135

6 Eigenvalues and Eigenvectors **143**
6.1 A Motivating Example . . . 143
6.2 The Characteristic Polynomial . . . 148
6.3 Diagonalization . . . 153
6.4 Evaluating a Polynomial at a Matrix . . . 156
6.5 The Functional Calculus . . . 164
6.6 Complex Eigenvalues and Eigenvectors of Real Matrices . . . 171
6.7 Normal Operators . . . 174

7 Factorization Theorems **178**
7.1 Gram-Schmidt and QR Factorization . . . 178
7.2 Schur Triangularization . . . 184
7.3 The Spectral Theorem . . . 190
7.4 The Singular Value Decomposition . . . 192
7.5 Jordan Canonical Form . . . 193

8 Applications of Spectral Theory **193**
8.1 The Principal Axes Theorem . . . 193
8.2 Positive Definite Matrices . . . 195
8.3 Differential Equations . . . 195
8.4 Graph Theory . . . 196
8.5 Markov Chains . . . 196
8.6 Singular Value Decomposition . . . 197
8.7 Total Least Squares . . . 197

1 Inner Product Spaces

1.1 Vector Spaces

Let $\mathbb{R}$ be the set of real numbers. A *vector space over* $\mathbb{R}$ consists of a set V (of "vectors"), with two algebraic operations, called *addition* and *scalar multiplication*:

$$\mathbf{u}, \mathbf{v} \in V \rightsquigarrow \mathbf{u} + \mathbf{v} \in V$$
$$a \in \mathbb{R}, \mathbf{v} \in V \rightsquigarrow a\mathbf{v} \in V.$$

[Remark: We could also write scalar multiplication as $\mathbf{v}a$; the order doesn't matter.] These two operations are required to satisfy the following eight axioms:

(1) Axioms of Addition.

(a) $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$

(b) $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$

(c) There exists a vector $\mathbf{0} \in V$ such that $\mathbf{0} + \mathbf{v} = \mathbf{v} + \mathbf{0} = \mathbf{v}$ for all $\mathbf{v} \in V$.

(d) For every vector $\mathbf{v} \in V$ there exists a vector $\mathbf{u} \in V$ such that $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u} = \mathbf{0}$.

Remarks: The vector $\mathbf{0}$ in axiom (1c) is unique. Indeed, if $\mathbf{0}$ and $\mathbf{0}'$ are two vectors satisfying (1c) then we must have

$$\mathbf{0} = \mathbf{0} + \mathbf{0}' = \mathbf{0}'.$$

We call the unique element $\mathbf{0} \in V$ satisfying (1c) the *zero vector*. The vector $\mathbf{u}$ in axiom (1d) is also unique. Indeed, suppose we have two vectors $\mathbf{u}$ and $\mathbf{u}'$ satisfying (1d). Then from axioms (1abc) we must have

$$\mathbf{u} = \mathbf{u} + \mathbf{0} = \mathbf{u} + (\mathbf{v} + \mathbf{u}') = (\mathbf{u} + \mathbf{v}) + \mathbf{u}' = \mathbf{0} + \mathbf{u}' = \mathbf{u}'.$$

The unique element $\mathbf{u}$ satisfying (1d) is called the *additive inverse of* $\mathbf{v}$. We denote it by $-\mathbf{v}$. In other words, we have

$$\mathbf{v} + \mathbf{u} = \mathbf{0} \quad \Longleftrightarrow \quad \mathbf{u} = -\mathbf{v}.$$

Based on this, we define the operation of vector subtraction:

$$\mathbf{u} - \mathbf{v} := \mathbf{u} + (-\mathbf{v}).$$

(2) Axioms of Scalar Multiplication.

(a) For the real number $1 \in \mathbb{R}$ we have $1\mathbf{v} = \mathbf{v}$ for all $\mathbf{v} \in V$.

(b) For all real numbers $a, b \in \mathbb{R}$ and vectors $\mathbf{v} \in V$ we have $a(b\mathbf{v}) = (ab)\mathbf{v}$.[1]

(c) For all $a, b \in \mathbb{R}$ and $\mathbf{v} \in V$ we have $(a + b)\mathbf{v} = a\mathbf{v} + b\mathbf{v}$.[2]

(d) For all $a \in \mathbb{R}$ and $\mathbf{u}, \mathbf{v} \in V$ we have $a(\mathbf{u} + \mathbf{v}) = a\mathbf{u} + a\mathbf{v}$.

Remarks: These eight axioms imply many other basic properties. For example, I claim that the real number $0 \in \mathbb{R}$ satisfies $0\mathbf{v} = \mathbf{0}$ for all vectors $\mathbf{v} \in V$, where $\mathbf{0}$ is the zero vector. Indeed, since $0 + 0 = 0$ as real numbers, we have

$$\begin{aligned}
0 + 0 &= 0 \\
(0 + 0)\mathbf{v} &= 0\mathbf{v} \\
0\mathbf{v} + 0\mathbf{v} &= 0\mathbf{v} \\
(0\mathbf{v} + 0\mathbf{v}) - 0\mathbf{v} &= 0\mathbf{v} - 0\mathbf{v} \\
0\mathbf{v} + (0\mathbf{v} - 0\mathbf{v}) &= \mathbf{0} \\
0\mathbf{v} + \mathbf{0} &= \mathbf{0} \\
0\mathbf{v} &= \mathbf{0}.
\end{aligned}$$

[1] Note that this identity involves two **different** operations: multiplication of real numbers and scalar multiplication in V. This identity is the reason that we use the same notation for both operations.

[2] This identity is the reason that we use the same notation for addition in $\mathbb{R}$ and addition in V.

[We could have taken $0\mathbf{v} = \mathbf{0}$ as another axiom, but we didn't need to.] It follows from this that the additive inverse $-\mathbf{v}$ is the same as $(-1)\mathbf{v}$ for the real number $-1 \in \mathbb{R}$. Indeed, since $1 + (-1) = 0$ as real numbers, we have

$$\begin{aligned}
1 + (-1) &= 0 \\
(1 + (-1))\mathbf{v} &= 0\mathbf{v} \\
1\mathbf{v} + (-1)\mathbf{v} &= 0\mathbf{v} \\
\mathbf{v} + (-1)\mathbf{v} &= \mathbf{0} \\
-\mathbf{v} + (\mathbf{v} + (-1)\mathbf{v}) &= -\mathbf{v} + \mathbf{0} \\
(-\mathbf{v} + \mathbf{v}) + (-1)\mathbf{v} &= -\mathbf{v} \\
\mathbf{0} + (-1)\mathbf{v} &= -\mathbf{v} \\
(-1)\mathbf{v} &= -\mathbf{v}.
\end{aligned}$$

The definition of a vector space has many axioms, but they all "look obvious", so you won't have trouble remembering them. It is important to emphasize that a vector space does **not** include any way to multiply two vectors to get another vector. In the next section we will discuss "inner products", which allow us to multiply two vectors to obtain a scalar.[3]


The Prorotype: Euclidean Space. Let $\mathbb{R}^n$ denote the set of ordered n-tupes of real numbers:

$$\mathbb{R}^n = \{\mathbf{v} = (v_1, v_2, \ldots, v_n) : v_1, v_2, \ldots, v_n \in \mathbb{R}\}.$$

It is easy to check that the following operations make $\mathbb{R}^n$ into a vector space over $\mathbb{R}$:

$$\begin{aligned}
(u_1, \ldots, u_n) + (v_1, \ldots, v_n) &:= (u_1 + v_1, \ldots, u_n + v_n), \\
a(v_1, \ldots, v_n) &:= (av_1, \ldots, av_n).
\end{aligned}$$

We can think of $\mathbf{v} = (v_1, \ldots, v_n)$ as the coordinates of a point in n-dimensional Euclidean space. In this case, the point $\mathbf{0} = (0, \ldots, 0)$ is called the *origin*. The *parallelogram law* says that for any points $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$, the four points $\mathbf{0}$, $\mathbf{u}$, $\mathbf{v}$ and $\mathbf{u} + \mathbf{v}$ are the vertices of a parallelogram. Picture:




[3] In our case the scalars are the real numbers $\mathbb{R}$. The definition of a vector space makes sense over any field of scalars. Later we will consider the field of complex numbers $\mathbb{C}$.

We can also think of an n-tuple $\mathbf{v} = (v_1, \ldots, v_n)$ as a directed line segment (an "arrow") with head at the point $\mathbf{v}$ and tail at the origin $\mathbf{0}$. According to the Pythagorean Theorem, the length $\|\mathbf{v}\|$ of this line segment satisfies

$$\|\mathbf{v}\| = \sqrt{v_1^2 + v_2^2 \cdots + v_n^2}.$$

Geometrically, arrows add "head-to-tail" and subtract "tail-to-tail":



If we let θ denote the angle between arrows $\mathbf{u}$ and $\mathbf{v}$ then the Law of Cosines tells us that

$$\|\mathbf{u} - \mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 - 2\|\mathbf{u}\|\|\mathbf{v}\| \cos\theta.$$

On the other hand, the algebraic formula for the length of an arrow tells us that

$$\begin{aligned}
\|\mathbf{u} - \mathbf{v}\|^2 &= \|(u_1 - v_1, \ldots, u_n - v_n)\|^2 \\
&= (u_1 - v_1)^2 + \cdots + (u_n - v_n)^2 \\
&= (u_1^2 - 2u_1v_1 + v_1^2) + \cdots + (u_n^2 - 2u_nv_n + v_n^2) \\
&= (u_1^2 + \cdots + u_n)^2 + (v_1^2 + \cdots + v_n^2) - 2(u_1v_1 + \cdots + u_nv_n) \\
&= \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2 - 2(u_1v_1 + \cdots + u_nv_n).
\end{aligned}$$

Then comparing the two equations gives the amazing formula

$$u_1v_1 + \cdots + u_nv_n = \|\mathbf{u}\|\|\mathbf{v}\| \cos\theta.$$

This formula allows us to express angles simply in terms of the coordinates. To be precise, we define the *dot product* of two arrows:

$$\mathbf{u} \bullet \mathbf{v} := u_1v_1 + u_2v_2 + \cdots + u_nv_n.$$

Observe that

$$\mathbf{v} \bullet \mathbf{v} = v_1v_1 + \cdots + v_nv_n = v_1^2 + \cdots + v_n^2 = \|\mathbf{v}\|^2.$$

Hence we have

$$\cos\theta = \frac{\mathbf{u} \bullet \mathbf{v}}{\|\mathbf{u}\|\|\mathbf{v}\|} = \frac{\mathbf{u} \bullet \mathbf{v}}{\sqrt{\mathbf{u} \bullet \mathbf{u}}\sqrt{\mathbf{v} \bullet \mathbf{v}}}.$$

The most important case is the following:

$$\mathbf{u} \bullet \mathbf{v} = 0 \quad \Longleftrightarrow \quad \theta \text{ is a right angle.}$$

In this case we say that $\mathbf{u}$ and $\mathbf{v}$ are *orthogonal* (or *perpendicular*).

1.2 Inner Products and Norms

More generally, an *inner product space over* $\mathbb{R}$ consists of a vector space V over $\mathbb{R}$ together with another algebraic operation

$$\mathbf{u}, \mathbf{v} \in V \rightsquigarrow \langle \mathbf{u}, \mathbf{v} \rangle \in \mathbb{R},$$

which must satisfy the following axioms:

(3) Axioms of Inner Products.

(a) $\langle \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{v}, \mathbf{u} \rangle$

(b) $\langle \mathbf{u}, \mathbf{v} + \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{u}, \mathbf{w} \rangle$[4]

(c) For all $a \in \mathbb{R}$ and $\mathbf{u}, \mathbf{v} \in V$ we have $\langle a\mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{u}, a\mathbf{v} \rangle = a\langle \mathbf{u}, \mathbf{v} \rangle$.

(d) For all $\mathbf{v} \in V$ we have $\langle \mathbf{v}, \mathbf{v} \rangle \geq 0$, with $\langle \mathbf{v}, \mathbf{v} \rangle = 0$ if and only if $\mathbf{v} = \mathbf{0}$.

Note that axioms (a) and (b) together imply that

$$\langle \mathbf{u} + \mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{w}, \mathbf{u} + \mathbf{v} \rangle = \langle \mathbf{w}, \mathbf{u} \rangle + \langle \mathbf{w}, \mathbf{v} \rangle = \langle \mathbf{u}, \mathbf{w} \rangle + \langle \mathbf{v}, \mathbf{w} \rangle.$$

More generally, we have

$$\left\langle \sum_{i=1}^{m} a_i \mathbf{u}_i, \sum_{i=1}^{n} b_i \mathbf{v}_i \right\rangle = \sum_{i=1}^{m} \sum_{j=1}^{n} a_i b_i \langle \mathbf{u}_i, \mathbf{v}_j \rangle$$

for any vectors $\mathbf{u}_1, \ldots, \mathbf{u}_m, \mathbf{v}_1, \ldots, \mathbf{v}_n \in V$ and scalars $a_1, \ldots, a_m, b_1, \ldots, b_n \in \mathbb{R}$. Based on axiom (d) it makes sense to define the *norm* (or the *length*) of a vector as

$$\|\mathbf{v}\| := \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}.$$

This is meant to generalize the concept of "length" in Euclidean space. The following important inequality is a direct consequence of the axioms, but its proof is a little bit tricky. I'll give you a hint and have you prove it on the homework.

Cauchy-Schwarz Inequality. For any vectors $\mathbf{u}, \mathbf{v} \in V$ in an inner product space we have

$$|\langle \mathbf{u}, \mathbf{v} \rangle| \leq \|\mathbf{u}\| \|\mathbf{v}\|.$$

[4] By combining (3ab) we also have $\langle \mathbf{u} + \mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{w} \rangle + \langle \mathbf{v}, \mathbf{w} \rangle$.

The Cauchy-Schwarz inequality implies the *triangle inequality*:

$$\|\mathbf{u}+\mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\| \quad \text{for all } \mathbf{u}, \mathbf{v} \in V.$$

Indeed, from the definition of length and the axioms of the inner product we have

$$\begin{aligned}
\|\mathbf{u}+\mathbf{v}\|^2 &= \langle \mathbf{u}+\mathbf{v}, \mathbf{u}+\mathbf{v}\rangle \\
&= \langle \mathbf{u}, \mathbf{u}\rangle + 2\langle \mathbf{u}, \mathbf{v}\rangle + \langle \mathbf{v}, \mathbf{v}\rangle && \text{Axioms (a) and (b)} \\
&\leq \langle \mathbf{u}, \mathbf{u}\rangle + 2\|\mathbf{u}\|\|\mathbf{v}\| + \langle \mathbf{v}, \mathbf{v}\rangle && \text{Cauchy-Schwarz} \\
&= (\|\mathbf{u}\| + \|\mathbf{v}\|)^2.
\end{aligned}$$

Then taking the square root of both sides gives the result.

The Cauchy-Schwarz inequality can be written equivalently as

$$-1 \leq \frac{\langle \mathbf{u}, \mathbf{v}\rangle}{\|\mathbf{u}\|\|\mathbf{v}\|} \leq 1,$$

which allows us to define the *angle* θ between the vectors $\mathbf{u}$ and $\mathbf{v}$ by

$$\theta := \arccos\left(\frac{\langle \mathbf{u}, \mathbf{v}\rangle}{\|\mathbf{u}\|\|\mathbf{v}\|}\right).$$

One should not take this too literally because the elements $\mathbf{u}, \mathbf{v}$ of the abstract vector space V might have nothing to do with "arrows". In general we should think of the angle as a kind of "correlation". A right angle means that $\mathbf{u}$ and $\mathbf{v}$ are "uncorrelated".

Abstraction should never be pursued for its own sake. There are two main motivations for the abstract approach to linear algebra. First of all, there exist important examples of abstract inner product spaces that have nothing to do with arrows or points in Euclidean space.

Example: L^2 Space. Let $L^2[0,1]$ denote the set of real-valued functions $f(x)$ on the interval $[0,1]$ such that the integral of the square of $f(x)$ converges:[5]

$$L^2[0,1] = \{f : [0,1] \to \mathbb{R}, \int_0^1 f(x)^2\, dx < \infty\}.$$

Given functions $f, g \in L^2[0,1]$ and scalar $a \in \mathbb{R}$ we define the new functions $f+g \in L^2[0,1]$ and $af \in L^2[0,1]$ by adding and multiplying their values, as one does in Calculus:

$$\begin{aligned}
(f+g)(x) &:= f(x) + g(x), \\
(af)(x) &:= af(x).
\end{aligned}$$

[5]The letter L stands for Henri Lebesgue. The integral used in the definition is not the typical "Riemann integral" from your Calculus course, but the more general and abstract "Lebesgue integral". The full definition is extremely technical but we will proceed intuitively, just as a physicist would.

One can check that these operations make $L^2[0,1]$ into a vector space over $\mathbb{R}$.[6] Furthermore, one can check that the following operation satisfies the inner product axioms:

$$\langle f(x), g(x) \rangle := \int_0^1 f(x)g(x)\,dx.$$

Such inner product spaces of functions are extremely important in applied mathematics. They are also "infinite dimensional" so they cannot possibly be studied via the language of matrices.

In applications we sometimes need to consider linear combinations in which infinitely many coefficients are nonzero, in which case we have to worry about convergence. We will see an example below when we talk about Fourier series. In order to say anything about convergence of infinite series in a vector space, one needs a way to measure "distance" between vectors.

(5) Axioms of Norms. Let V be a vector space with a function

$$\mathbf{v} \in V \rightsquigarrow \|\mathbf{v}\| \in \mathbb{R}.$$

We call this function a *norm* when it satisfies the following axioms:

(a) $\|\mathbf{v}\| \geq 0$ for all $\mathbf{v} \in V$ with $\|\mathbf{v}\| = 0$ if and only if $\mathbf{v} = \mathbf{0}$.

(b) For all $a \in \mathbb{R}$ and $\mathbf{v} \in V$ we have $\|a\mathbf{v}\| = |a|\|\mathbf{v}\|$.

(c) (Triangle inequality) For all $\mathbf{u}, \mathbf{v} \in V$ we have $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$.

(6) Axioms of Metrics. Let V be a vector space with a function

$$\mathbf{u}, \mathbf{v} \in V \rightsquigarrow \text{dist}(\mathbf{u}, \mathbf{v}) \in \mathbb{R}.$$

We call this function a *metric* when it satisfies the following axioms:

(a) $\text{dist}(\mathbf{u}, \mathbf{v}) = \text{dist}(\mathbf{v}, \mathbf{u})$

(b) $\text{dist}(\mathbf{u}, \mathbf{v}) \geq 0$ for all $\mathbf{u}, \mathbf{v} \in V$ with $\text{dist}(\mathbf{u}, \mathbf{v}) = 0$ if and only if $\mathbf{u} = \mathbf{v}$.

(c) $\text{dist}(\mathbf{u}, \mathbf{v}) \leq \text{dist}(\mathbf{u}, \mathbf{w}) + \text{dist}(\mathbf{w}, \mathbf{v})$ for all $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$.

Every inner product space becomes a normed space by defining $\|\mathbf{v}\| := \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}$, and every normed space becomes a metric space by defining $\text{dist}(\mathbf{u}, \mathbf{v}) := \|\mathbf{u} - \mathbf{v}\|$. The proof is mostly straightforward. The hardest part is to verify the triangle inequality, which we did above.

It is worth mentioning that there exist norms on $\mathbb{R}^n$ that do not come from inner products. The most famous examples are the so-called *p-norms*.

[6] The hardest part of the proof is to show that the sum of square integrable functions is square integrable. This can be shown with the Cauchy-Schwarz inequality.

Example: The p-norm on $\mathbb{R}^n$. Let $p \geq 1$ be a real number. Then for every $\mathbf{x} = (x_1, \ldots, x_n) \in \mathbb{R}^n$ we define

$$\|\mathbf{x}\|_p := \left(\sum_{i=1}^{n} |x_i|^p \right)^{1/p}.$$

It turns out that the function $\| \cdot \|_p$ satisfies the norm axioms, though this is not so easy to prove. Observe that the 2-norm is just the standard Euclidean norm:

$$\|\mathbf{x}\|_2 = \sqrt{|x_1|^2 + \cdots + |x_n|^2} = \sqrt{x_1^2 + \cdots + x_n^2}.$$

The 1-norm and the ∞-norm also have special descriptions in terms of the sum and the maximum of the absolute values of the coordinates:

$$\|\mathbf{x}\|_1 = |x_1| + |x_2| + \cdots + |x_n|,$$
$$\|\mathbf{x}\|_\infty = \max\{|x_1|, |x_2|, \ldots, |x_n|\}.$$

It turns out that the p-norm on $\mathbb{R}^n$ does **not** come from an inner product unless $p = 2$, though this is also a bit tricky to prove.
The p-norm can also be extended to infinite dimensional spaces of functions.

Example: L^p space. Fix some real number p and consider the set

$$L^p[0,1] = \{f : [0,1] \to \mathbb{R}, \int_0^1 |f(x)|^p < \infty\}$$

It turns out that this is a normed vector space with norm function

$$\|f\|_p := \left(\int_0^1 |f(x)|^p \right)^{1/p}.$$

But this norm does not come from an inner product unless $p = 2$.

1.3 Subspaces and Dimension

Another reason for abstraction in linear algebra has to do with "subspaces".

(4) Axioms of Subspaces. Given a vector space V over $\mathbb{R}$ and a subset $U \subseteq V$, we say that U is a *subspace* when it satisfies the following axioms:[7]

(a) If $\mathbf{u}, \mathbf{v} \in U$ then $\mathbf{u} + \mathbf{v} \in U$.

(b) If $a \in \mathbb{R}$ and $\mathbf{v} \in U$ then $a\mathbf{v} \in U$.

[7]We note that (a) is implied by (c) by taking $a = 0$

Note that the second property implies $\mathbf{0} \in U$ by taking $a = 0$, hence every subspace must contain the zero vector. We can state axioms (a) and (b) simultaneously as follows:

$$\text{If } \mathbf{u}, \mathbf{v} \in U \text{ and } a, b \in \mathbb{R} \text{ then } a\mathbf{u} + b\mathbf{v} \in U.$$

In practice we often use a generalized form:

$$\text{If } \mathbf{u}_1, \ldots, \mathbf{u}_n \in U \text{ and } a_1, \ldots, a_n \in \mathbb{R} \text{ then } \sum_{i=1}^{n} a_i \mathbf{u}_i \in U.$$

The sum $a_1\mathbf{u}_1 + \cdots + a_n\mathbf{u}_n$ is called a *linear combination* of the vectors $\mathbf{u}_1, \ldots, \mathbf{u}_n$. Hence we can express the definition of subspaces in words as follows:

A subspace is closed under linear combinations.

It goes without saying that a subspace of a vector space is itself a vector space, with respect to the same operations of addition and scalar multiplication.

We note that Euclidean space $\mathbb{R}^n$ comes with a collection of *standard basis vectors*:

$$\begin{aligned}
\mathbf{e}_1 &= (1, 0, 0, \ldots, 0, 0), \\
\mathbf{e}_2 &= (0, 1, 0, \ldots, 0, 0), \\
&\vdots \\
\mathbf{e}_n &= (0, 0, 0, \ldots, 0, 1).
\end{aligned}$$

By definition, every vector $\mathbf{v} = (v_1, \ldots, v_n) \in \mathbb{R}^n$ has a unique expression as a linear combination of these basis vectors:

$$\begin{aligned}
\mathbf{v} &= (v_1, \ldots, v_n) \\
&= v_1(1, 0, \ldots, 0, 0) + \cdots + v_n(0, 0, \ldots, 0, 1) \\
&= v_1\mathbf{e}_1 + \cdots + v_n\mathbf{e}_n.
\end{aligned}$$

However, **subspaces** of $\mathbb{R}^n$ do not come with standard basis vectors. For example, consider the plane $V \subseteq \mathbb{R}^3$ defined by the equation $x - 2y + z = 0$. In other words,

$$V = \{(x, y, z) \in \mathbb{R}^3 : x - 2y + z = 0\}.$$

This is a subspace of $\mathbb{R}^3$ since for any $(x_1, y_1, z_1), (x_2, y_2, z_3) \in V$ and $a_1, a_2 \in \mathbb{R}$ we have

$$\begin{aligned}
&(a_1x_1 + a_2x_2) - 2(a_1y_1 + a_1y_2) + (a_1z_1 + a_2z_2) \\
&= a_1(x_1 - 2y_1 + z_1) + a_2(x_2 - 2y_2 + z_2) \\
&= a_1 0 + a_2 0 \\
&= 0,
\end{aligned}$$

and hence $a_1(x_1, y_1, z_1)+a_2(x_2, y_2, z_2) = (a1x_1+a_2x_2, a_1y_1+a_2y_2, a_1z_1+a_2z_2) \in V$. The vector space V does not have a "standard basis". However, I claim that every vector $\mathbf{x} = (x, y, z) \in V$ has a unique expression of the form[8]

$$\mathbf{v} = a_1(1, 1, 1) + a_2(1, 2, 3),$$

with $a_1, a_2 \in \mathbb{R}$. In this case we will say that the set $B = \{\mathbf{u}_1, \mathbf{u}_2\}$ with $\mathbf{u}_1 = (1, 1, 1)$ and $\mathbf{u}_2 = (1, 2, 3)$ is a "basis" for the vector space V, and if $\mathbf{x} = a_1\mathbf{u}_1 + a_2\mathbf{u}_2$ we will say that $\mathbf{x} = (a_1, a_2)_B$ are the "coordinates" of $\mathbf{v}$ in the B-basis. Furthermore, since this basis has two elements we will say that the vector space V is "two-dimensional". For example, the vector $\mathbf{v} = (1, -1, -3) \in \mathbb{R}^3$ is in the plane V. It has coordinates $(1, -1, -3)$ as an element of $\mathbb{R}^3$, with respect to the standard basis, but it has coordinates $(3, -2)_B$ as an element of V, with respect to the B-basis, because

$$\begin{aligned}(1, -1 - 3) &= 1\mathbf{e}_1 - 1\mathbf{e}_2 - 3\mathbf{e}_3 \\ &= 3\mathbf{u}_1 - 2\mathbf{u}_2.\end{aligned}$$

But there is nothing special about the basis B. For example, we could equally well choose $\mathbf{u}_1 = (2, 1, 0)$ and $\mathbf{u}_2 = (0, 1, 2)$, in which case we would have $(1, -1, -3) = (1/2, -3/2)_B$. This is why we need the concept of an abstract vector space:

> *Subspaces of $\mathbb{R}^n$ do not come with a standard basis. Therefore we must study them via the axioms of abstract vector spaces.*

We now discuss the technical details behind the concepts of "basis" and "dimension".

Definition (Basis). Let V be a vector space over $\mathbb{R}$ and consider a finite subset $B = \{\mathbf{u}_1, \ldots, \mathbf{u}_n\}$ of vectors in V.

- We say that B is a *spanning set* if for all $\mathbf{v} \in V$ **there exists at least one choice** of scalars $a_1, \ldots, a_n \in \mathbb{R}$ such that

$$\mathbf{v} = a_1\mathbf{u}_1 + \cdots + a_n\mathbf{u}_n.$$

- We say that B is an *independent set* (more commonly called a *linearly independent set* if for all $\mathbf{v} \in V$ **there exists at most one choice** of scalars $a_1, \ldots, a_n \in \mathbb{R}$ such that

$$\mathbf{v} = a_1\mathbf{u}_1 + \cdots + a_n\mathbf{u}_n.$$

- We say that B is a *basis* if it is spanning and independent; that is, if for all $\mathbf{v} \in V$ **there exists a unique choice** of scalars $a_1, \ldots, a_n \in \mathbb{R}$ such that

$$\mathbf{v} = a_1\mathbf{u}_1 + \cdots + a_n\mathbf{u}_n.$$

[8] We won't prove this at the moment.

In this case we say that $a_1, \ldots, a_n \in \mathbb{R}$ are the *B-coordinates* of $\mathbf{v}$, and we write

$$\mathbf{v} = (a_1, \ldots, a_n)_B.$$

After we have chosen a basis, we can work with coordinate vectors and we can pretend that V is $\mathbb{R}^n$, though it is more correct to say that V is *isomorphic* to $\mathbb{R}^n$

Most textbooks use the following equivalent definition of independence, which is more convenient for proofs.

Alternate Definition of Independence. Let V be a real vector space. We say that a set of vectors $\mathbf{u}_1, \ldots, \mathbf{u}_k \in V$ is *linearly independent* if for all scalars $a_1, \ldots, a_k \in \mathbb{R}$ we have

$$a_1\mathbf{u}_1 + \cdots + a_n\mathbf{u}_n = \mathbf{0} \quad \Longrightarrow \quad a_1 = \cdots = a_n = 0.$$

Exercise: Check that the two definitions of independence are equivalent.

The basic theory of dimension is based on two lemmas. The first lemma is a relatively easy consequence of the definitions.

Lemma (Extension and Contraction). Given a set $\{\mathbf{u}_1, \ldots, \mathbf{u}_k\} \subseteq V$ we write

$$\text{Span}\{\mathbf{u}_1, \ldots, \mathbf{u}_k\} := \{a_1\mathbf{u}_1 + \cdots + a_k\mathbf{u}_k : a_1, \ldots, a_k \in \mathbb{R}\}.$$

We have the following two elementary properties:

(a) Let I be an independent set. If $\mathbf{v} \notin \text{Span}(I)$ then $I \cup \{\mathbf{v}\}$ is independent. In this case we say that the vector $\mathbf{v}$ is *independent* from the set I.

(b) Let S be a spanning set. If $\mathbf{v} \in S$ and $\mathbf{v} \in \text{Span}(S \setminus \{\mathbf{v}\})$ then $\text{Span}(S \setminus \{\mathbf{v}\}) = \text{Span}(S)$. In this case we say that $\mathbf{v}$ is a *redundant* element of S.

Proof: Exercise.

The next lemma is tricky to prove. It was first written down by Ernst Steinitz in 1913.

Lemma (Steinitz Exchange). Let V be a vector space over $\mathbb{R}$, let $I \subseteq V$ be a finite independent set and let $S \subseteq V$ be a finite spanning set. Then we must have $|I| \leq |S|$.[9]

Proof. Let $I = \{\mathbf{u}_1, \mathbf{u}_2, \ldots \mathbf{u}_m\}$ be an independent set and let $S = \{\mathbf{v}_1, \ldots, \mathbf{v}_n\}$ be a spanning set. We will show that $m \leq n$. We begin by defining $S_0 := S$. Since S_0 is a spanning set there exist scalars $a_1, \ldots, a_n \in \mathbb{R}$ such that $\mathbf{u}_1 = a_1\mathbf{v}_1 + \cdots + a_n\mathbf{v}_n$. Then since $\mathbf{u}_1 \neq \mathbf{0}$[10]

[9]For this lemma we assume that V has a finite spanning set. In this case, it will follow from the proof that any independent subset of V must also be finite. I didn't make this explicit because I didn't actually define the concepts of infinite spanning sets and infinite independent sets.

[10]The zero vector is not an element of any independent set.

there exists some index i such that $a_i \neq 0$. By relabeling the vectors $\mathbf{v}_i$ if necessary, we may assume that $i = 1$ and hence $a_1 \neq 0$. It follows that

$$\mathbf{v}_1 = \frac{1}{a_1}\mathbf{u}_1 - \sum_{i=2}^{n} \frac{a_i}{a_1}\mathbf{v}_i.$$

Define the set $S_1 := \{\mathbf{u}_1, \mathbf{v}_2, \ldots, \mathbf{v}_n\}$ by "exchanging" $\mathbf{v}_1$ with $\mathbf{u}_1$. I claim that S_1 is a spanning set. Indeed, consider any vector $\mathbf{w}$. Since S_0 is a spanning set we can write $\mathbf{w} = b_1\mathbf{v}_1 + \cdots + b_n\mathbf{v}_n$ for some scalars $b_1, \ldots, b_n \in \mathbb{R}$. But then we have

$$\begin{aligned}\mathbf{w} &= b_1\left(\frac{1}{a_1}\mathbf{u}_1 - \sum_{i=2}^{n}\frac{a_i}{a_k}\mathbf{v}_i\right) + b_2\mathbf{v}_2 + \cdots + b_n\mathbf{v}_n \\ &= \frac{b_1}{a_1}\mathbf{u}_1 + \sum_{i=2}^{n}\left(b_i - \frac{a_i}{a_1}\right)\mathbf{v}_i,\end{aligned}$$

which is in the span of S_1. Now assume for induction that we can relabel the vectors $\mathbf{v}_i$ so that $S_k = \{\mathbf{u}_1, \ldots, \mathbf{u}_k, \mathbf{v}_{k+1}, \ldots, \mathbf{v}_n\}$ is a spanning set for some $k < m$. Then we can write $\mathbf{u}_{k+1} = a_1\mathbf{u}_1 + \cdots + a_k\mathbf{u}_k + b_{k+1}\mathbf{v}_{k+1} + \cdots + b_n\mathbf{v}_n$ for some scalars $a_1, \ldots, a_k, b_{k+1}, \ldots, b_\ell \in \mathbb{R}$. We must have $b_i \neq 0$ for some i, since otherwise the equation $\mathbf{u}_{k+1} = a_1\mathbf{u}_1 + \cdots + a_k\mathbf{u}_k$ contradicts the independence of the $\mathbf{u}_i$. Without loss of generality we may suppose that $b_{k+1} \neq 0$, and then an argument similar to the above shows that $S_{k+1} := \{\mathbf{u}_1, \ldots, \mathbf{u}_{k+1}, \mathbf{v}_{k+2}, \ldots, \mathbf{v}_\ell\}$ is a spanning set. If $m \geq n$ then we conclude by induction that $S_n = \{\mathbf{u}_1, \ldots, \mathbf{u}_n\}$ is a spanning set. But if $m > n$ then this implies that $\mathbf{u}_{n+1} = a_1\mathbf{u}_1 + \ldots + a_n\mathbf{u}_n$ for some scalars $a_1, \ldots, a_n \in \mathbb{R}$, which contradicts the fact that the $\mathbf{u}_i$ are independent. Hence $m \leq n$. □

This result allows us to define the concept of dimension.

Theorem (Existence of Dimension). Let V be a vector space and suppose that V has a finite basis.[11] Any two bases for V have the same number of elements. We call this common number the *dimension* of V:

$$\boxed{\dim V := \text{ the number of vectors in any basis.}}$$

Furthermore, for any independent set I we have $|I| \leq \dim V$[12] and for any spanning set S we have $|S| \geq \dim V$.

Proof. Let $B, C \subseteq V$ be two bases. Since B is independent and C is spanning, Steinitz exchange implies that $|B| \leq |C|$. On the other hand, since C is independent and B is spanning, Steinitz exchange implies $|C| \leq |B|$, hence $|B| = |C|$. Next, let B be any basis, let I be independent and let S be spanning. Since B is spanning we have $|I| \leq |B| = \dim V$ by Steinitz exchange and since B is independent we have $\dim V = |B| \leq |S|$ by Steinitz exchange. □

[11] If V does not have a finite basis we write $\dim V = \infty$.

[12] We don't have to assume that I is finite

Example: Euclidean Space. I claim that

$$\dim \mathbb{R}^n = n.$$

Indeed, the "standard basis vectors" $\mathbf{e}_1, \ldots, \mathbf{e}_n \in \mathbb{R}^n$ are a basis. This implies that **any** basis of $\mathbb{R}^n$ has n vectors. Bases are very common. In fact, any sufficiently random collection of n vectors in $\mathbb{R}^n$ will be a basis. For example, I am almost completely sure that the following set of four vectors is a basis of $\mathbb{R}^4$:

$$\{(1, 4, 3, 2), (3, -7, 4, 1), (100, 89, -72, 36), (23, 24, 25, 26)\}.$$

Not every vector space has a **finite** basis.

Example: Polynomials. Let $\mathbb{R}[x]$ denote the set of polynomials in x with real coefficients. This set carries a natural vector space structure. If we think of polynomials as function, then addition and scalar multiplication are defined "pointwise". That is, for polynomial functions $f(x), g(x) \in \mathbb{R}[x]$, and scalars $\lambda, \mu \in \mathbb{R}$ we define the function $\lambda f + \mu g : \mathbb{R} \to \mathbb{R}$ by setting

$$(\lambda f + \mu g)(x) := \lambda f(x) + \mu g(x) \quad \text{for all } x \in \mathbb{R}.$$

We can also define this operation purely formally in terms of coefficients. We can think of a polynomial as a formal expression

$$f(x) = \sum_{k \geq 0} a_k x^k,$$

where only finitely many of the coefficients $a_0, a_2, \ldots \in \mathbb{R}$ are nonzero. If $f(x) = \sum_{k \geq 0} a_k x^k$ and $g(x) = \sum_{k \geq 0} b_k x^k$ then we define the formal expression

$$\lambda f(x) + \mu g(x) := \sum_{k \geq 0} (\lambda a_k + \mu b_k) x^k,$$

and we note that only finitely many of the real numbers $\lambda a_k + \mu b_k$ are nonzero. The vector space $\mathbb{R}[x]$ does not have a finite basis, so we say that

$$\dim \mathbb{R}[x] = \infty.$$

But it does have a natural **infinite** basis B consisting of the elements[13]

$$B = \{1, x, x^2, x^3, \ldots\}.$$

We list some useful results regarding dimensions of subspaces.

Theorem (Dimension of Subspaces). Let W be a finite dimensional vector space.

[13] By convention we let $x^0 := 1$.

(a) If $V \subseteq W$ is a subspace then $\dim V \leq \dim W$.

(b) If $V \subseteq W$ is a subspace and $\dim V = \dim W$ then $V = W$.

(c) If $U, V \subseteq W$ are subspaces then the following set is also a subspace, called the *sum*:

$$U + V := \{\mathbf{u} + \mathbf{v} : \mathbf{u} \in U \text{ and } \mathbf{v} \in V\}.$$

If $U \cap V = \{\mathbf{0}\}$[14] then we have

$$\dim(U + V) = \dim U + \dim V.$$

In this case we write $U + V = U \oplus V$ we and call this a *direct sum*.

(d) More generally, if $U, V \subseteq W$ are subspaces then the intersection $U \cap V \subseteq W$ is also a subspace, and we have

$$\dim(U + V) = \dim V + \dim U - \dim(U \cap V).$$

Note that agrees with the previous formula if we define $\dim(\{\mathbf{0}\}) = 0$.[15]

Proof. To prove (a), let B be a basis of V. Since B is an independent subset of V it is also an independent subset of W, hence the theorem on the existence of dimension implies that $\dim V = |B| \leq \dim W$.[16] To prove (b), let B be a basis of V, so B is an independent subset of W, and suppose $\dim V = \dim W$. I claim that $\text{Span}(W)$. If not, then choose any $\mathbf{w} \in W \setminus \text{Span}(W)$. By the extension lemma $B \cup \{w\}$ is an independent subset of W with size $|B \cup \{w\}| = |B| + 1 = \dim V + 1 \geq \dim W$. But any independent subset I of W must satisfy $|I| \leq \dim V$ by the theorem on the existence of dimension. We will prove (c) and (d) at the same time, since the dimension formula in (c) is a special case of (d). First we check that $U \cap V$ is a subspace of W. Indeed, for any $\mathbf{u}, \mathbf{v} \in U \cap V$ and $\lambda, \mu \in \mathbb{R}$ we have $\lambda\mathbf{u} + \mu\mathbf{v} \in U$ because U is a subspace and $\lambda\mathbf{u} + \mu\mathbf{v} \in V$ because V is a subspace. Hence $\lambda\mathbf{u} + \mu\mathbf{v} \in U \cap V$. Next we show that $U + V$ is a subspace of W. Indeed, consider any elements $\mathbf{u} + \mathbf{v}$ and $\mathbf{u}' + \mathbf{v}'$ of $U + V$ with $\mathbf{u}, \mathbf{u}' \in U$ and $\mathbf{v}, \mathbf{v}' \in V$ and consider any scalars $\lambda, \mu \in \mathbb{R}$. Then

$$\lambda(\mathbf{u} + \mathbf{v}) + \mu(\mathbf{u}' + \mathbf{v}') = (\lambda\mathbf{u} + \mu\mathbf{u}') + (\lambda\mathbf{v} + \mu\mathbf{v}')$$

is an element of $U + V$ because $\lambda\mathbf{u} + \mu\mathbf{u}'$ is in U and $\lambda\mathbf{v} + \mu\mathbf{v}'$ is in V. To prove the dimension formula, let $B_0 = \{\mathbf{w}_1, \ldots, \mathbf{w}_r\}$ be a basis of $U \cap V$.[17] From the alternate proof of part (a) we may extend this to a basis $B_U = \{\mathbf{w}_1, \ldots, \mathbf{w}_r, \mathbf{u}_1, \ldots, \mathbf{u}_p\}$ of U and a basis $B_V = \{\mathbf{w}_1, \ldots, \mathbf{w}_r, \mathbf{v}_1, \ldots, \mathbf{v}_q\}$ of V, so that $\dim W = r$, $\dim U = r + p$ and $\dim V = r + q$.

[14] Note that every subspace contains the zero vector, so this is the smallest possible intersection.

[15] We could say that the zero subspace $\{\mathbf{0}\}$ has a basis consisting of zero elements, i.e., the empty set.

[16] Alternate proof: Let $\mathbf{v}_1, \ldots, \mathbf{v}_k$ be a basis for V. As long as $\text{Span}\{\mathbf{v}_1, \ldots, \mathbf{v}_j\} \neq W$ we can choose $\mathbf{v}_{j+1} \in W \setminus \text{Span}\{\mathbf{v}_1, \ldots, \mathbf{v}_j\}$. By the extension lemma the set $\{\mathbf{v}_1, \ldots, \mathbf{v}_{j+1}\}$ is still independent. Since the size of any independent set is $\leq \dim W$ by the theorem on the existence of dimension, this process must stop with a basis for W that contains $\mathbf{v}_1, \ldots, \mathbf{v}_k$ as a subset. Hence $\dim V = k \leq \dim W$.

[17] If $U \cap V = \{\mathbf{0}\}$ then we let B_0 be the empty set.

We need to show that $\dim(U+V) = (r+p)+(r+q)-r = r+p+q$. Indeed, I claim that the set $B = \{\mathbf{w}_1, \ldots, \mathbf{w}_r, \mathbf{u}_1, \ldots, \mathbf{u}_p, \mathbf{v}_1, \ldots, \mathbf{v}_q\}$ (which has $r+p+q$ elements) is a basis of $U+V$. It is a spanning set because the $\mathbf{w}_i$ and $\mathbf{u}_i$ span U, while the $\mathbf{w}_i$ and $\mathbf{v}_i$ span V, hence any element $\mathbf{x} = \mathbf{u}+\mathbf{v} \in U+V$ can be written as

$$\begin{aligned}\mathbf{x} = \mathbf{u}+\mathbf{v} &= \Big(\sum a_i\mathbf{w}_i + \sum b_i\mathbf{u}_i\Big) + \Big(\sum a_i'\mathbf{w}_i + \sum c_i\mathbf{v}_i\Big) \\ &= \sum (a_i+a_i')\mathbf{w}_i + \sum b_i\mathbf{u}_i + c_i\mathbf{v}_i.\end{aligned}$$

To prove independence, consider any linear relation

$$\sum a_i\mathbf{w}_i + \sum b_i\mathbf{u}_i + \sum c_i\mathbf{v}_i = \mathbf{0},$$

so that

$$\sum a_i\mathbf{w}_i + \sum b_i\mathbf{u}_i = -\sum c_i\mathbf{v}_i.$$

The vector on the left is in U and the vector on the right is in V, hence both are in $U \cap V$. Since $\mathbf{w}_i$ is a basis for $U \cap V$ this implies that $-\sum c_i\mathbf{v}_i = \sum d_i\mathbf{w}_i$, hence $\sum_i d_i\mathbf{w}_i + \sum c_i\mathbf{v}_i = \mathbf{0}$. Since the $\mathbf{w}_i$ and $\mathbf{v}_i$ together are independent (they are a basis for V) we conclude that $c_i = 0$ and $d_i = 0$ for all i. Then we have $\sum a_i\mathbf{w}_i + \sum b_i\mathbf{u}_i = \mathbf{0}$ which implies that $a_i = 0$ and $b_i = 0$ for all i because the $\mathbf{w}_i$ and $\mathbf{u}_i$ together are independent (they are a basis for U). □

It is worth mentioning that direct sums have a uniqueness property. Suppose, as in part (c) of the theorem, that we have $W = U+V$ and $U \cap V = \{\mathbf{0}\}$, which we express by saying $W = U \oplus V$. This implies that for any vector $\mathbf{w} \in W$ there exists a **unique pair** of vectors $\mathbf{u} \in U$ and $\mathbf{v} \in V$ such that $\mathbf{w} = \mathbf{u}+\mathbf{v}$. Indeed, suppose we had another such expression $\mathbf{w} = \mathbf{u}'+\mathbf{v}'$. Then we would have

$$\begin{aligned}\mathbf{u}+\mathbf{v} &= \mathbf{u}'+\mathbf{v}' \\ \mathbf{u}-\mathbf{u}' &= \mathbf{v}'-\mathbf{v}.\end{aligned}$$

Since $\mathbf{u}-\mathbf{u}' \in U$ and $\mathbf{v}'-\mathbf{v} \in V$, this implies that $\mathbf{u}-\mathbf{u}'$ and $\mathbf{v}'-\mathbf{v}$ are in $U \cap V$. Since $U \cap V = \{\mathbf{0}\}$, this implies that $\mathbf{u}-\mathbf{u}' = \mathbf{0}$ and $\mathbf{v}'-\mathbf{v} = \mathbf{0}$, hence $\mathbf{u} = \mathbf{u}'$ and $\mathbf{v} = \mathbf{v}'$.

Example: Subspaces of Euclidean Space. Every subspace of $\mathbb{R}^n$ looks like a *k-dimensional plane through the origin*. To be precise, for any k independent vectors $\mathbf{u}_1, \ldots, \mathbf{u}_k$ (necessarily with $k \le n$), their span

$$U = \mathrm{Span}\{\mathbf{u}_1, \ldots, \mathbf{u}_k\} = \{a_1\mathbf{u}_1 + \cdots a_k\mathbf{u}_k : a_1\mathbf{u}_k + \cdots + a_k\mathbf{u}_k\} \subseteq \mathbb{R}^n.$$

is called *k-plane*. Note that a k-plane is k-dimensional by definition because it is spanned by a set of k independent vectors. We can also define subspaces of $\mathbb{R}^n$ intrinsically by linear equations. To be precise, for any scalars $a_{ij}, b_i \in \mathbb{R}$ with $1 \le i \le m$ and $1 \le j \le n$, the set of vectors $\mathbf{x} = (x_1, x_2, \ldots, x_n) \in \mathbb{R}^n$ satisfying the following *system of m linear equations in n unknowns* is a subspace of $\mathbb{R}^n$:

$$\left\{\begin{array}{ccccccc} a_{11}x_1 & + & \cdots & + & a_{1n}x_n & = & b_1, \\ \vdots & & & & \vdots & \vdots & \vdots \\ a_{m1}x_1 & + & \cdots & + & a_{mn}x_n & = & b_m. \end{array}\right.$$

We know that the set of solutions must be a k-plane for some k, but it is not easy to read off k just by looking at the equations. We will discuss this in Section ?.

1.4 Orthogonal Projection


The concept of orthogonal projection is central to many areas of applied mathematics, because of its relationship to optimization. Here is a simple example:

Given a function $f : [0, 1] \to \mathbb{R}$, find the polynomial of degree ≤ 2 that is "closest" to f.

There are two issues that we need to address:

- What do we mean by "closest"?
- After choosing a suitable notion of "closest", how do we compute the answer?

Perhaps the most obvious notion of "closest" is to minimize the area between the two curves, as in the following picture, where the solid curve is the function f and the dotted curve is the "best fit parabola":



To be precise, the idea here is to find the polynomial $g(x)$ of degree ≤ 2 that minimizes the integral of the absolute difference:

$$\int_0^1 |f(x) - g(x)|\, dx.$$

Unfortunately, this polynomial is quite difficult to compute, so we choose instead to **minimize the integral of the squared difference**:

$$\int_0^1 (f(x) - g(x))^2\, dx.$$

This is a nice choice because it fits naturally into the theory inner product spaces. Indeed, we recall that the set of "square-integrable functions"

$$L^2[0, 1] = \{f : [0, 1] \to \mathbb{R}, \int_0^1 f(x)^2 < \infty\}$$

is a vector space with inner product

$$\langle f, g\rangle = \int_0^1 f(x)g(x)\,dx.$$

Thus we are looking for the polynomial $g(x)$ of degree ≤ 2 that minimizes the "L^2-distance" between f and g:[18]

$$\|f-g\| = \sqrt{\langle f-g, f-g\rangle} = \sqrt{\int_0^1 (f(x)-g(x))^2\,dx}.$$

We can rephrase the problem as follows. Let $U \subseteq L^2[0,1]$ be the subspace defined by

$$U := \mathrm{Span}\{1, x, x^2\} = \{a_0 + a_1x + a_2x^2 : a_0, a_1, a_2 \in \mathbb{R}\},$$

where we think of each polynomial $a_0 + a_1x + a_2x^2$ as a function $[0,1] \to \mathbb{R}$.[19] Then for any given function $f \in L^2[0,1]$, our goal is to find a function $g \in U$ that minimizes the distance to f. That is, we want

$$\|f-h\| \geq \|f-g\| \quad \text{for all } h \in U.$$

The general concept behind this example is called "orthogonal projection".

Theorem (Orthogonal Projection). Let V be an inner product space (possibly infinite-dimensional) and let $U \subseteq V$ be a finite dimensional subspace. Suppose we can find a basis $\mathbf{u}_1, \ldots, \mathbf{u}_n$ for U satisfying

$$\langle \mathbf{u}_i, \mathbf{u}_j\rangle = \delta_{ij} := \begin{cases} 1 & i = j, \\ 0 & i \neq j. \end{cases}$$

(We call this an *orthonormal basis* for U.) Then for any given vector $\mathbf{v} \in V$, there exists a unique vector $\mathbf{p} \in U$ satisfying

$$\|\mathbf{v}-\mathbf{u}\| \geq \|\mathbf{v}-\mathbf{p}\| \quad \text{for all } \mathbf{u} \in U,$$

and this vector is given by the simple formula

$$\mathbf{p} = \langle \mathbf{v}, \mathbf{u}_1\rangle\mathbf{u}_1 + \langle \mathbf{v}, \mathbf{u}_2\rangle\mathbf{u}_2 + \cdots + \langle \mathbf{v}, \mathbf{u}_n\rangle\mathbf{u}_n.$$

We call $\mathbf{p}$ the *orthogonal projection of* $\mathbf{v}$ *onto* U.

Proof. Let $\mathbf{u}_1, \ldots, \mathbf{u}_n$ be an orthonormal basis for U and let $\mathbf{p}$ be the vector defined by the given formula. Then the vector $\mathbf{v}-\mathbf{p}$ is orthogonal to every vector in U because it is orthogonal to each of the basis vectors $\mathbf{u}_1, \ldots, \mathbf{u}_n$. Indeed, for any j we have

$$\langle \mathbf{v}-\mathbf{p}, \mathbf{u}_j\rangle = \langle \mathbf{v}, \mathbf{u}_j\rangle - \langle \mathbf{p}, \mathbf{u}_j\rangle$$

[18] The integral of the absolute difference is the distance in the vector space $L^1[0,1]$, which is a normed vector space, but it is not an inner product space. In general, there is a normed space $L^p[0,1]$ for each real number $p \geq 1$, and only $L^2[0,1]$ is an inner product space.

[19] Certainly each of these functions is square-integrable.

$$
\begin{aligned}
&= \langle \mathbf{v}, \mathbf{u}_j \rangle - \left\langle \sum_{i=1}^{n} \langle \mathbf{v}, \mathbf{u}_i \rangle \mathbf{u}_i, \mathbf{u}_j \right\rangle \\
&= \langle \mathbf{v}, \mathbf{u}_j \rangle - \sum_{i=1}^{n} \langle \mathbf{v}, \mathbf{u}_i \rangle \langle \mathbf{u}_i, \mathbf{u}_j \rangle \\
&= \langle \mathbf{v}, \mathbf{u}_j \rangle - \langle \mathbf{v}, \mathbf{u}_j \rangle \\
&= 0,
\end{aligned}
$$

and then for any vector $\mathbf{u} = \sum_{j=1}^{n} a_j \mathbf{u}_j$ in U we have

$$
\langle \mathbf{v} - \mathbf{p}, \mathbf{u} \rangle = \left\langle \mathbf{v} - \mathbf{p}, \sum_{j=1}^{n} a_j \mathbf{u}_j \right\rangle = \sum_{j=1}^{n} a_j \langle \mathbf{v} - \mathbf{p}, \mathbf{u}_j \rangle = 0.
$$

But then for any vector $\mathbf{u} \in U$ we have $\mathbf{v} - \mathbf{u} = (\mathbf{v} - \mathbf{p}) + (\mathbf{p} - \mathbf{u})$, and hence

$$
\begin{aligned}
\langle \mathbf{v} - \mathbf{u}, \mathbf{v} - \mathbf{u} \rangle &= \langle \mathbf{v} - \mathbf{p}, \mathbf{v} - \mathbf{p} \rangle + \langle \mathbf{p} - \mathbf{u}, \mathbf{p} - \mathbf{u} \rangle + 2\langle \mathbf{v} - \mathbf{p}, \mathbf{p} - \mathbf{u} \rangle \\
\langle \mathbf{v} - \mathbf{u}, \mathbf{v} - \mathbf{u} \rangle &= \langle \mathbf{v} - \mathbf{p}, \mathbf{v} - \mathbf{p} \rangle + \langle \mathbf{p} - \mathbf{u}, \mathbf{p} - \mathbf{u} \rangle + 0 \\
\|\mathbf{v} - \mathbf{u}\|^2 &= \|\mathbf{v} - \mathbf{p}\|^2 + \|\mathbf{p} - \mathbf{u}\|^2 \\
\|\mathbf{v} - \mathbf{u}\|^2 &\geq \|\mathbf{v} - \mathbf{p}\|^2 \\
\|\mathbf{v} - \mathbf{u}\| &\geq \|\mathbf{v} - \mathbf{p}\|,
\end{aligned}
$$

where we used the fact that $\langle \mathbf{v} - \mathbf{p}, \mathbf{p} - \mathbf{u} \rangle = 0$ because $\mathbf{p} - \mathbf{u}$ is in U. To prove uniqueness, suppose there exists $\mathbf{q} \in U$ with $\|\mathbf{v} - \mathbf{q}\| = \|\mathbf{v} - \mathbf{p}\|$. Then setting $\mathbf{u} = \mathbf{q}$ in the equation $\|\mathbf{v} - \mathbf{u}\|^2 = \|\mathbf{v} - \mathbf{p}\|^2 + \|\mathbf{p} - \mathbf{u}\|^2$ gives $\|\mathbf{p} - \mathbf{q}\|^2 = 0$, which implies $\mathbf{q} = \mathbf{p}$. □

Remark: We don't need to assume that $\langle \mathbf{u}_i, \mathbf{u}_i \rangle = 1$ for all i; only that $\langle \mathbf{u}_i, \mathbf{u}_i \rangle \neq 0$. In the general case we have

$$
\mathbf{p} = \frac{\langle \mathbf{v}, \mathbf{u}_1 \rangle}{\langle \mathbf{u}_1, \mathbf{u}_1 \rangle} \mathbf{u}_1 + \frac{\langle \mathbf{v}, \mathbf{u}_2 \rangle}{\langle \mathbf{u}_2, \mathbf{u}_2 \rangle} \mathbf{u}_2 + \cdots + \frac{\langle \mathbf{v}, \mathbf{u}_n \rangle}{\langle \mathbf{u}_n, \mathbf{u}_n \rangle} \rangle \mathbf{u}_n.
$$

Thus we have reduced the problem to the existence and computation of orthonormal bases. This is accomplished by following important algorithm. Like the Steinitz exchange lemma, the proof that the algorithm works is trickier than you would expect, which is why it has a special name. Before stating the theorem we will prove a small lemma.

Lemma (Orthogonal Sets are Independent). Let V be an inner product space and let $S = \{\mathbf{w}_1, \ldots, \mathbf{w}_n\}$ be a set of nonzero vectors satisfying $\langle \mathbf{w}_i, \mathbf{w}_j \rangle = 0$ for all $i \neq j$. Then S is an independent set.

Proof. If $a_1 \mathbf{w}_1 + \cdots + a_n \mathbf{w}_n = \mathbf{0}$ then taking the inner product of each side with $\mathbf{w}_j$ gives

$$
0 = \langle \mathbf{0}, \mathbf{w}_j \rangle = \left\langle \sum_{i=1}^{n} a_i \mathbf{w}_i, \mathbf{w}_j \right\rangle = \sum_{i=1}^{n} a_i \langle \mathbf{w}_i, \mathbf{w}_j \rangle = a_j \langle \mathbf{w}_j, \mathbf{w}_j \rangle.
$$

Since $\mathbf{w}_j \neq \mathbf{0}$ (by assumption), this implies that $\langle \mathbf{w}_j, \mathbf{w}_j \rangle \neq 0$ and hence $a_j = 0$. $\square$

Theorem (Gram-Schmidt Procedure). Let V be a finite-dimensional inner product space and let $\mathbf{v}_1, \ldots, \mathbf{v}_n$ be any basis for V. Define the vectors $\mathbf{w}_1, \ldots, \mathbf{w}_n \in V$ recursively by the following procedure:

$$\begin{aligned}
\mathbf{w}_1 &:= \mathbf{v}_1, \\
\mathbf{w}_2 &:= \mathbf{v}_2 - \frac{\langle \mathbf{v}_2, \mathbf{w}_1 \rangle}{\langle \mathbf{w}_1, \mathbf{w}_1 \rangle} \mathbf{w}_1, \\
\mathbf{w}_3 &:= \mathbf{v}_3 - \frac{\langle \mathbf{v}_3, \mathbf{w}_1 \rangle}{\langle \mathbf{w}_1, \mathbf{w}_1 \rangle} \mathbf{w}_1 - \frac{\langle \mathbf{v}_3, \mathbf{w}_2 \rangle}{\langle \mathbf{w}_2, \mathbf{w}_2 \rangle} \mathbf{w}_2, \\
&\text{etc.}
\end{aligned}$$

I claim that each $\mathbf{w}_k$ is nonzero (so we are able to divide by $\langle \mathbf{w}_k, \mathbf{w}_k \rangle$), and that

$$\operatorname{Span}\{\mathbf{w}_1, \ldots, \mathbf{w}_k\} = \operatorname{Span}\{\mathbf{v}_1, \ldots, \mathbf{v}_k\} \quad \text{for all } 1 \leq k \leq n.$$

Furthermore, I claim that $\langle \mathbf{w}_i, \mathbf{w}_j \rangle = 0$ for all $i \neq j$. It follows from the previous lemma that $\mathbf{w}_1, \ldots, \mathbf{w}_n$ is an orthogonal basis for V. Then to obtain an orthonormal basis $\mathbf{u}_1, \ldots, \mathbf{u}_n$ we set $\mathbf{u}_k := \mathbf{w}_k / \|\mathbf{w}_k\|$ for all $1 \leq k \leq n$.[20]

Proof. To save space we define the notation

$$\begin{aligned}
W_k &:= \operatorname{Span}\{\mathbf{w}_1, \ldots, \mathbf{w}_k\}, \\
V_k &:= \operatorname{Span}\{\mathbf{v}_1, \ldots, \mathbf{v}_k\}.
\end{aligned}$$

We will use induction to show that the following statement holds for all $1 \leq k \leq n$:

$$W_k = V_k \quad \text{and } \mathbf{w}_i \neq \mathbf{0} \text{ for } 1 \leq i \leq k.$$

The statement holds for $k = 1$ because $\mathbf{w}_1 = \mathbf{v}_1$ and $\mathbf{v}_1 \neq \mathbf{0}$. (No basis contains the zero vector.) Now suppose that the statement holds for some $1 \leq k < n$. In this case we need to show that $W_{k+1} = V_{k+1}$ and $\mathbf{w}_{k+1} \neq \mathbf{0}$. Since $\langle \mathbf{w}_i, \mathbf{w}_i \rangle \neq 0$ for all $1 \leq i \leq k$ we may define

$$\mathbf{w}_{k+1} := \mathbf{v}_{k+1} - \sum_{i=1}^{k} \frac{\langle \mathbf{v}_{k+1}, \mathbf{w}_i \rangle}{\langle \mathbf{w}_i, \mathbf{w}_i \rangle} \mathbf{w}_i.$$

In order to show that $V_{k+1} \subseteq W_{k+1}$ we will show that $\mathbf{v}_1, \ldots, \mathbf{v}_{k+1} \in W_{k+1}$. We have $\mathbf{v}_{k+1} \in W_{k+1}$ because

$$\mathbf{v}_{k+1} = \mathbf{w}_{k+1} + \sum_{i=1}^{k} \frac{\langle \mathbf{v}_{k+1}, \mathbf{w}_i \rangle}{\langle \mathbf{w}_i, \mathbf{w}_i \rangle} \mathbf{w}_i,$$

[20] We could also normalize at each step of the procedure, but this introduces too many square roots into the computation.

and we have $\mathbf{v}_1, \ldots, \mathbf{v}_k \in W_{k+1}$ because $\mathbf{v}_1, \ldots, \mathbf{v}_k \in V_k$ and $V_k = W_k \subseteq W_{k+1}$. In order to show that $W_{k+1} \subseteq V_{k+1}$ we will show that $\mathbf{w}_1, \ldots, \mathbf{w}_{k+1} \in V_{k+1}$. We have $\mathbf{w}_1, \ldots, \mathbf{w}_k \in V_{k+1}$ because $\mathbf{w}_1, \ldots, \mathbf{w}_k \in W_k$ and $W_k = V_k \subseteq V_{k+1}$. And we have $\mathbf{w}_{k+1} \in V_{k+1}$ because

$$\mathbf{z} := \sum_{i=1}^{k} \frac{\langle \mathbf{v}_{k+1}, \mathbf{w}_i \rangle}{\langle \mathbf{w}_i, \mathbf{w}_i \rangle} \mathbf{w}_i \in W_k = V_k \subseteq V_{k+1},$$

hence $\mathbf{w}_{k+1} = \mathbf{v}_{k+1} - \mathbf{z} \in V_{k+1}$. It remains to show that $\mathbf{w}_{k+1} \neq \mathbf{0}$. But if $\mathbf{w}_{k+1} = \mathbf{0}$ then

$$\mathbf{v}_{k+1} = \sum_{i=1}^{k} \frac{\langle \mathbf{v}_{k+1}, \mathbf{w}_i \rangle}{\langle \mathbf{w}_i, \mathbf{w}_i \rangle} \mathbf{w}_i \in W_k = V_k,$$

which contradicts the fact that the set $\mathbf{v}_1, \ldots, \mathbf{v}_k, \mathbf{v}_{k+1}$ is independent. (Any subset of an independent set is independent.)

We have shown that $W_n = V_n = V$, so that $\mathbf{w}_1, \ldots, \mathbf{w}_n$ is a spanning set. To complete the proof we need to show that $\langle \mathbf{w}_i, \mathbf{w}_j \rangle = 0$ for all $i \neq j$. We will do this by showing that the following statement holds for all $1 \leq k \leq n$:

$$\langle \mathbf{w}_i, \mathbf{w}_j \rangle = 0 \quad \text{for all } 1 \leq i < j \leq k.$$

For the case $k = 1$ there is nothing to check. Suppose the statement holds for some $1 \leq k < n$, so that $\langle \mathbf{w}_i, \mathbf{w}_j \rangle = 0$ for all $1 \leq i < j \leq k$. Then for any $1 \leq i \leq k$ we also have

$$\begin{aligned}
\langle \mathbf{w}_i, \mathbf{w}_{k+1} \rangle &= \left\langle \mathbf{w}_i, \mathbf{v}_{k+1} - \sum_{j=1}^{k} \frac{\langle \mathbf{v}_{k+1}, \mathbf{w}_j \rangle}{\langle \mathbf{w}_j, \mathbf{w}_j \rangle} \mathbf{w}_j \right\rangle \\
&= \langle \mathbf{w}_i, \mathbf{v}_{k+1} \rangle - \sum_{j=1}^{k} \frac{\langle \mathbf{v}_{k+1}, \mathbf{w}_j \rangle}{\langle \mathbf{w}_j, \mathbf{w}_j \rangle} \langle \mathbf{w}_i, \mathbf{w}_j \rangle \\
&= \langle \mathbf{w}_i, \mathbf{v}_{k+1} \rangle - \frac{\langle \mathbf{v}_{k+1}, \mathbf{w}_i \rangle}{\langle \mathbf{w}_i, \mathbf{w}_i \rangle} \langle \mathbf{w}_i, \mathbf{w}_i \rangle \\
&= 0.
\end{aligned}$$

□

The Gram-Schmidt procedure looks unmotivated, but it has a natural geometric interpretation in terms of projection.

Theorem (Orthogonal Complement). Let V be an inner product space (finite- or infinite-dimensional) and let $U \subseteq V$ be a finite-dimensional subspace. Define the *orthogonal complement* of U as the set

$$U^{\perp} := \{\mathbf{v} \in V : \langle \mathbf{v}, \mathbf{u} \rangle = 0 \text{ for all } \mathbf{u} \in U\}.$$

I claim that $U^{\perp} \subseteq V$ is a subspace and that $V = U \oplus U^{\perp}$, which means that every vector $\mathbf{v} \in V$ has a unique expression of the form $\mathbf{v} = \mathbf{v}_U + \mathbf{v}_{U^{\perp}}$ with $\mathbf{v}_U \in U$ and $\mathbf{v}_{U^{\perp}} \in U^{\perp}$. I

claim that $\mathbf{v}_U$ is the orthogonal projection of $\mathbf{v}$ onto the subspace U. Furthermore, I claim that $\mathbf{v}_{U^\perp}$ is the unique point in $U^\perp$ that minimizes the distance to $\mathbf{v}$. Thus we will say that

$$\mathbf{v}_{U^\perp} = \text{“the orthogonal projection of } \mathbf{v} \text{ onto the subspace } U^\perp\text{”},$$

even though $U^\perp$ might be infinite-dimensional.[21] Finally, if $\mathbf{u}_1, \ldots, \mathbf{u}_k$ is an orthogonal basis for U then the orthogonal projection of $\mathbf{v}$ onto $U^\perp$ is

$$\mathbf{v}_{U^\perp} = \mathbf{v} - \sum_{i=1}^{k} \frac{\langle \mathbf{v}, \mathbf{u}_i \rangle}{\langle \mathbf{u}_i, \mathbf{u}_i \rangle} \mathbf{u}_i.$$

Proof. To show that $U^\perp \subseteq V$ is a subspace, consider any $\mathbf{x}, \mathbf{y} \in U^\perp$ and $a, b \in \mathbb{R}$. Then for any $\mathbf{u} \in U$ we have

$$\langle a\mathbf{x} + b\mathbf{y}, \mathbf{u} \rangle = a\langle \mathbf{x}, \mathbf{u} \rangle + b\langle \mathbf{y}, \mathbf{u} \rangle = a \cdot 0 + b \cdot 0 = 0,$$

which implies that $a\mathbf{x} + b\mathbf{y} \in U^\perp$, as desired. Now consider any $\mathbf{v} \in V$ and let $\mathbf{v}_U \in U$ be the unique point in U that is closest to $\mathbf{v}$, which exists from the above theorem on orthogonal projection. In the proof of that theorem we saw that $\mathbf{v}_{U^\perp} := \mathbf{v} - \mathbf{v}_U$ is orthogonal to every vector in U. Hence we have $\mathbf{v} = \mathbf{v}_U + \mathbf{v}_{U^\perp}$ with $\mathbf{v}_U \in U$ and $\mathbf{v}_{U^\perp}$, and we have shown that $V = U + U^\perp$. Note that we also have $U \cap U^\perp = \{\mathbf{0}\}$ since any element of the intersection must be orthogonal to itself. Thus we have shown that $V = U \oplus U^\perp$. To show that $\mathbf{v}_{U^\perp}$ minimizes the distance to $U^\perp$, consider any vector $\mathbf{w} \in U^\perp$. Since $\mathbf{v} - \mathbf{w} = \mathbf{v}_U + (\mathbf{v}_{U^\perp} - \mathbf{w})$ with $\langle \mathbf{v}_U, \mathbf{v}_{U^\perp} - \mathbf{w} \rangle = 0$, the Pytharogean theorem gives

$$\|\mathbf{v} - \mathbf{w}\|^2 = \|\mathbf{v}_U\|^2 + \|\mathbf{v}_{U^\perp} - \mathbf{w}\|^2 \geq \|\mathbf{v}_U\|^2 = \|\mathbf{v} - \mathbf{v}_{U^\perp}\|^2,$$

and hence $\|\mathbf{v} - \mathbf{w}\| \geq \|\mathbf{v} - \mathbf{v}_{U^\perp}\|$. Moreover, equality holds when $\|\mathbf{v}_{U^\perp} - \mathbf{w}\| = 0$, i.e., when $\mathbf{w} = \mathbf{v}_{U^\perp}$. Finally, the claimed explicit formula for $\mathbf{v}_{U^\perp}$ in terms of an orthogonal basis for U follows from the known explicit formula for $\mathbf{v}_U$ that we proved earlier. □

Thus we can re-interpret the Gram-Schmidt procedure as follows: Given a basis $\mathbf{v}_1, \ldots, \mathbf{v}_n$ for an inner product space, let $\mathbf{w}_1 := \mathbf{v}_1$, and then recursively define

$$\mathbf{w}_{k+1} := \text{the orthogonal projection of } \mathbf{v}_{k+1} \text{ onto } \mathrm{Span}\{\mathbf{w}_1, \ldots, \mathbf{w}_k\}^\perp.$$

Unfortunately, we couldn't use this idea in the proof of Gram-Schmidt because it would have led to circular reasoning.

We now have all of the necessary ingredients to solve our motivating problem. Again, consider the subspace

$$U = \mathrm{Span}\{1, x, x^2\} \subseteq L^2[0, 1].$$

[21] For a general infinite-dimensional subspace $W \subseteq V$, the “projection onto W” might not exist at all.

The basis $1, x, x^2 \in U$ is not orthogonal, but we may obtain an orthogonal basis $w_0, w_1, w_2 \in U$ by the Gram-Schmidt procedure. First we define $w_0(x) = 1$. Then we define

$$w_1(x) = x - \frac{\langle x, w_0 \rangle}{\langle w_0, w_0 \rangle} \cdot w_0(x) = x - \frac{\int_0^1 x\,dx}{\int_0^1 1\,dx} \cdot 1 = x - \frac{1}{2},$$

and

$$\begin{aligned} w_2(x) &= x^2 - \frac{\langle x^2, w_0 \rangle}{\langle w_0, w_0 \rangle} \cdot w_0(x) - \frac{\langle x^2, w_1 \rangle}{\langle w_1, w_1 \rangle} \cdot w_1(x) \\ &= x^2 - \frac{\int_0^1 x^2\,dx}{\int_0^1 1\,dx} \cdot 1 - \frac{\int_0^1 x^2(x - \frac{1}{2})\,dx}{\int_0^1 (x - \frac{1}{2})^2\,dx} \cdot \left(x - \frac{1}{2}\right) \\ &= x^2 - \frac{1/3}{1} - \frac{1/12}{1/12}\left(x - \frac{1}{2}\right) \\ &= x^2 - \frac{1}{3} - \left(x - \frac{1}{2}\right) \\ &= x^2 - x + \frac{1}{6}. \end{aligned}$$

So the resulting orthogonal basis for the space of quadratic polynomials is[22]

$$w_0(x) = 1, \qquad w_1(x) = x - \frac{1}{2}, \qquad w_2(x) = x^2 - x + \frac{1}{6}.$$

And the resulting orthonormal basis $u_i(x) = w_i(x)/\langle w_i, w_i \rangle$ is

$$u_0(x) = 1, \qquad u_1(x) = \sqrt{3}\,(2x - 1), \qquad u_2(x) = \sqrt{5}\,(6x^2 - 6x + 1).$$

Thus for any square-integrable function $f : [0, 1] \to \mathbb{R}$ the "best fit" quadratic polynomial, in the sense of the L^2 norm, is the orthogonal projection of f onto the subspace U of quadratic polynomials, which is given by the formula

$$g(x) = \langle f, u_0 \rangle\, u_0(x) + \langle f, u_1 \rangle\, u_1(x) + \langle f, u_2 \rangle\, u_2(x).$$

The function $f(x)$ doesn't even have to be continuous.[23] For example, consider the piecewise-defined function

$$f(x) = \begin{cases} 0 & 0 \le x < 3/8, \\ 1 & 3/8 < x < 1/2, \\ 0 & 1/2 < x \le 1. \end{cases}$$


In this case, one may compute $\langle f, u_0 \rangle = 1/8$, $\langle f, u_1 \rangle = -\sqrt{3}/64$ and $\langle f, u_2 \rangle = -15\sqrt{5}/256$, so the best fit parabola (in the L^2 sense) is

$$g(x) = -\frac{225}{128}x^2 + \frac{213}{128}x - \frac{31}{256}.$$

Here is a picture:

[22] These polynomials are closely related to the *Legendre polynomials*.

[23] And it can be undefined at some points, as long as the integrals are not affected.



This was just an introduction to the concept of orthogonal projection, applied to infinite-dimensional spaces of functions. We will have much more to say about projections in later chapters.

1.5 Fourier Series

The most famous and useful application of orthogonality is the theory of Fourier series. The basic ideas were developed in the 1700s by Euler and others, and the method was officially stated by Fourier in 1807, but the full rigorous theory was not complete until 100 years later with the work of several authors on L^2 space.

Let $u(x,t)$ be the deflection of a stretched string with fixed endpoints: $u(0,t) = u(1,t) = 0$ for all t. The wave equation tells us that

$$u_{tt}(x,t) = c^2 u_{xx}(x,t),$$

where c is some constant depending on the string tension. Suppose the string has an initial shape $u(x,0) = f(x)$ for some function $f : [0,1] \to \mathbb{R}$, and that the initial velocity is zero: $u_t(x,0) = 0$ for all $0 \le x \le 1$. In 1753, Daniel Bernoulli claimed that the general solution has the form

$$u(x,t) = \sum_{n \ge 1} a_n \cos(n\pi t) \sin(n\pi x),$$

and the difficulty is to find the coefficients $a_n \in \mathbb{R}$. If Bernoulli's claim is true, then by setting $t = 0$ we should be able to obtain an expansion for any (continuous?) function $f : [0,1] \to \mathbb{R}$ satisfying $f(0) = f(1) = 0$ as an infinite linear combination of sine waves:

$$f(x) = \sum_{n \ge 0} a_n \sin(n\pi x).$$

There was significant controversy about the existence and convergence of such series. In 1807, Fourier gave the following beautiful formula for the coefficients:

$$a_n = 2 \int_0^1 f(x) \sin(n\pi x)\, dx.$$

We now recognize this as an inner product computation in $L^2[0,1]$. In order to manage different boundary conditions apart from just fixed endpoints, it is convenient to also include cosine functions.

Fourier's Observation. For any positive integers $n \geq 1$ we define the functions

$$c_n(x) := \sqrt{2}\,\cos(2\pi n x),$$
$$s_n(x) := \sqrt{2}\,\sin(2\pi n x).$$

Then the infinite set $\{1, c_1(x), s_1(x), c_2(x), s_2(x), \ldots\}$ is **orthonormal** in the inner product space $L^2[0,1]$. (If we want we can view 1 as $c_0(x)/\sqrt{2}$.) For example, we have

$$\langle s_m, s_n\rangle = \int_0^1 \sqrt{2}\,\sin(2\pi m x)\,\sqrt{2}\,\sin(2\pi n x)\,dx = \begin{cases} 1 & m = n, \\ 0 & m \neq n. \end{cases}$$

The proof is a tedious exercise involving trig identities.

This important observation allows us easily to compute "Fourier coefficients". Suppose, for example, that some function $f(x)$ can be expressed in the form

$$f(x) = a_0 + a_1 c_1(x) + b_1 s_1(x) + a_2 c_2(x) + b_2 s_2(x) + \cdots .$$

Then the standard argument with orthonormal sets tells us that

$$a_0 = \langle f, 1\rangle,$$
$$a_n = \langle f, c_n\rangle,$$
$$b_n = \langle f, s_n\rangle.$$

On the other hand, for an arbitrary function $f(x)$ in $L^2[0,1]$ we may consider the infinite series

$$\langle f, 1\rangle + \langle f, c_1\rangle c_1(x) + \langle f, s_1\rangle s_1(x) + \cdots ,$$

and ask whether this series converges to the function $f(x)$ in the L^2 sense. (This would be an infinite-dimensional version of a "spanning set".) If it does, then we will say that the set

$$\{1, c_1(x), s_1(x), c_2(x), s_2(x), \ldots\}$$

is *complete basis* for $L^2[0,1]$.

Theorem (Fatou, 1906). This set is a complete basis for $L^2[0,1]$. To be precise, let U_n be the subspace of $L^2[0,1]$ spanned by $\{1, c_1, s_1, \ldots, c_n, s_n\}$. From the previous section we know that the orthogonal projection of $f \in L^2[0,1]$ onto U_n is

$$f_n(x) := \langle f, 1\rangle + \langle f, c_1\rangle c_1(x) + \langle f, s_1\rangle s_1(x) + \cdots + \langle f, c_n\rangle c_n(x) + \langle f, s_n\rangle s_n(x).$$

Then we have $\|f - f_n\| \to 0$ in $L^2[0,1]$ as $n \to \infty$.

We omit the proof because it is quite involved. There is an interesting corollary of Fatou's result that is called *Parseval's identity*, which says that

$$\|f\|^2 = \langle f, 1\rangle^2 + \sum_{k\geq 1} \left(\langle f, c_k\rangle^2 + \langle f, s_k\rangle^2\right),$$

where the series converges in the usual sense of real numbers.

Proof. We saw in the previous section that the orthogonal projection f_n satisfies $\langle f - f_n, f_n\rangle = 0$, hence the Pythagorean theorem gives $\|f\|^2 = \|f - f_n\|^2 + \|f_n\|^2$. By using orthonormality we have

$$\|f_n\|^2 = \langle f_n, f_n\rangle = \langle f, 1\rangle^2 + \sum_{k=1}^{n} \left(\langle f, c_k\rangle^2 + \langle f, s_k\rangle^2\right).$$

Then use the fact that $\|f - f_n\| \to 0$ to show that $\|f_n\| \to \|f\|$ as $n \to \infty$. □

Example: Consider the square wave function

$$f(x) = \begin{cases} 1 & 0 \leq x < 1/2, \\ 0 & 1/2 \leq x \leq 1. \end{cases}$$


It is easy to check that $a_0 = \langle f(x), 1\rangle = 1/2$ and $a_n = \langle f(x), c_n\rangle = 0$ for all $n \geq 1$. Next, we compute

$$\begin{aligned} b_n &= \langle f(x), s_n\rangle \\ &= \sqrt{2}\int_0^1 f(x)\sin(2\pi n x)\,dx \\ &= \sqrt{2}\int_0^{1/2} \sin(2\pi n x)\,dx \\ &= \frac{\sqrt{2}}{2\pi n}\left[-\cos(2\pi n x)\right]_0^{1/2} \\ &= \frac{\sqrt{2}}{2\pi n}\left[-\cos(\pi n) + 1\right] \\ &= \frac{\sqrt{2}}{2\pi n}\left[-(-1)^n + 1\right] \\ &= \begin{cases} 0 & n \text{ even}, \\ \frac{\sqrt{2}}{\pi n} & n \text{ odd}. \end{cases} \end{aligned}$$

It follows that

$$f(x) = \frac{1}{2} + \frac{\sqrt{2}}{\pi}\sin(2\pi x) + \frac{\sqrt{2}}{3\pi}\sin(6\pi x) + \frac{\sqrt{2}}{5\pi}\sin(10\pi x) + \cdots,$$

with convergence in the L^2 sense. Here is a picture of 30th partial sum $f_{30}(x)$:



Finally, Parseval's identity gives the following interesting identity:

$$\begin{aligned}
\|f\|^2 &= \langle f,1\rangle^2 + \langle f,s_1\rangle^2 + \langle f,s_2\rangle^2 + \langle f,s_3\rangle^2 + \cdots \\
\int_0^1 f(x)^2\,dx &= \left(\frac{1}{2}\right)^2 + \left(\frac{\sqrt{2}}{\pi}\right)^2 + \left(\frac{\sqrt{2}}{3\pi}\right)^2 + \left(\frac{\sqrt{2}}{5\pi}\right)^2 + \cdots \\
\frac{1}{2} &= \frac{1}{4} + \frac{2}{\pi^2} + \frac{2}{3^2\pi^2} + \frac{2}{5^2\pi^2} + \cdots \\
\frac{1}{4} &= \frac{2}{\pi^2} + \frac{2}{3^2\pi^2} + \frac{2}{5^2\pi^2} + \cdots \\
\frac{1}{4} &= \frac{2}{\pi^2}\left(\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \cdots\right) \\
\frac{\pi^2}{8} &= \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \cdots
\end{aligned}$$

That's weird.

Remark: The previous series is closely related to the famous *Basel problem*. Pietro Mangoli in 1650 asked for an exact expression for the convergent sum

$$\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \cdots.$$

This problem was finally solved by Leonhard Euler in 1734, who showed that the sum is exactly $\pi^2/6$. The appearance of π was a big surprise. Exercise: Derive Euler's result by considering the Fourier series of the function $f(x) = x$ on the interval $[0, 1]$.

1.6 Inner Product Spaces Over $\mathbb{C}$

Now seems like a good time to involve complex numbers. I have a complaint with the American educational system, in that there is no course that reliably introduces complex numbers. The system will tell you that complex numbers are part of the pre-Calculus curriculum. However, the treatment is inadequate, and most math majors don't take pre-Calculus. Indeed, I believe it is possible for a student to graduate with a major in mathematics, having never seen a good introduction to complex numbers. As is traditional, I will give a quick review and pretend that you have seen this before, even if you haven't.

Complex Numbers. The complex numbers are defined as

$$\mathbb{C} = \{a + ib : a, b \in \mathbb{R}\},$$

where i is an abstract symbol satisfying $i^2 = -1$. Given a complex number $\alpha = a + ib$, we define its *absolute value* and *complex conjugate*:[24]

$$|\alpha| := \sqrt{a^2 + b^2},$$
$$\alpha^* := a - ib.$$

These satisfying the following properties.

(7) Properties of Complex Numbers. For all $\alpha, \beta \in \mathbb{C}$ we have

(a) $(\alpha + \beta)^* = \alpha^* + \beta^*$

(b) $(\alpha\beta)^* = \alpha^*\beta^*$

(c) $(\alpha^*)^* = \alpha$

(d) $\alpha = \alpha^* \iff \alpha \in \mathbb{R}$

(e) $|\alpha| \geq 0$ with $|\alpha| = 0$ if and only if $\alpha = 0$.

(f) $|\alpha| = \alpha^*\alpha$

(g) $|\alpha\beta| = |\alpha||\beta|$.

(h) If $\alpha \neq 0$ then $\alpha^{-1} = \alpha^*/|\alpha|^2$.

Many applications of linear algebra use complex instead of real scalars. All of the results from Sections 1.1 and 1.3 are still true for complex numbers,[25] but the definition of inner products needs a small modification. To see why this is necessary, consider the vector $\mathbf{v} = (1, 2i)$ in the vector space $\mathbb{C}^2$. The naive formula for length would give

$$\|\mathbf{v}\| = \sqrt{1^2 + (2i)^2} = \sqrt{1 - 2} = \sqrt{-1},$$

[24] I will use α^* instead of the traditional $\overline{\alpha}$ to avoid conflict with the whiteboard notation for vectors: $\vec{v}$.
[25] Indeed, all of these results hold for scalars in any field.

which is not acceptable. The purpose of norms is to measure distances, and distances must be non-negative real numbers. It turns out that there is a miraculously easy fix for this problem.

The Prototype: Hermitian Space. Consider the complex vector space

$$\mathbb{C}^n = \{\mathbf{v} = (v_1, \ldots, v_n) : v_1, \ldots, v_n \in \mathbb{C}\}.$$

If we want to define an inner product on $\mathbb{C}^n$, our first instinct is to consider the "dot product"

$$\mathbf{u} \bullet \mathbf{v} = u_1 v_1 + u_2 v_2 + \cdots + u_n v_n,$$

and then to define a norm by $\|\mathbf{v}\| := \sqrt{\mathbf{v} \bullet \mathbf{v}}$. But for this to make sense we need $\mathbf{v} \bullet \mathbf{v}$ to be a non-negative real number, and sometimes it it not.[26] The solution is to define the function

$$\langle \mathbf{u}, \mathbf{v} \rangle := u_1^* v_1 + u_2^* v_2 + \cdots + u_n^* v_n.$$

Then for any vector $\mathbf{v}$ we have

$$\langle \mathbf{v}, \mathbf{v} \rangle = v_1^* v_1 + v_2^* v_2 + \cdots + v_n^* v_n = |v_1|^2 + |v_2|^2 + \cdots + |v_n|^2,$$

which is always a non-negative real number. Furthermore, we have

$$\langle \mathbf{v}, \mathbf{v} \rangle = 0 \iff |v_i| = 0 \text{ for all } i \iff v_i = 0 \text{ for all } i \iff \mathbf{v} = \mathbf{0}.$$

Thus it makes sense to define $\|\mathbf{v}\| := \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}$. The major difference between this function and the naive dot product is that "symmetry" is replaced with "conjugate symmetry":

$$\begin{aligned} \langle \mathbf{v}, \mathbf{u} \rangle^* &= (v_1^* u_1 + \cdots + v_n^* u_n)^* \\ &= (v_1^*)^* u_1^* + \cdots + (v_n^*)^* u_n^* \\ &= v_1 u_1^* + \cdots + v_n u_n^* \\ &= u_1 v_1^* + \cdots + u_n v_n^* \\ &= \langle \mathbf{u}, \mathbf{v} \rangle. \end{aligned}$$

Here is the abstract concept.

(8) Axioms of Hermitian Inner Products. Let V be a vector space over $\mathbb{C}$, together with an algebraic operation

$$\mathbf{u}, \mathbf{v} \in V \rightsquigarrow \langle \mathbf{u}, \mathbf{v} \rangle \in \mathbb{C}.$$

We call this a *Hermitian inner product* if it satisfies the following axioms:

(a) $\langle \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{v}, \mathbf{u} \rangle^*$

(b) $\langle \mathbf{u}, \mathbf{v} + \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{u}, \mathbf{w} \rangle$[27]

[26] In fact, one can prove that **there does not exist a function** $\mathbb{C}^n \times \mathbb{C}^n \to \mathbb{C}$ satisfying all of the inner product axioms.

[27] By combining (8ab) we also have $\langle \mathbf{u} + \mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{w} \rangle + \langle \mathbf{v} + \mathbf{w} \rangle$.

(c) For all $\alpha \in \mathbb{C}$ we have $\langle \mathbf{u}, \alpha\mathbf{v}\rangle = \alpha\langle \mathbf{u}, \mathbf{v}\rangle$.[28]

(d) For all $\mathbf{v} \in V$, part (a) tells us that $\langle \mathbf{v}, \mathbf{v}\rangle \in \mathbb{R}$. Furthermore, we must have $\langle \mathbf{v}, \mathbf{v}\rangle \geq 0$ with $\langle \mathbf{v}, \mathbf{v}\rangle = 0$ if and only if $\mathbf{v} = \mathbf{0}$.

It follows from axioms (a) and (b) that

$$\begin{aligned}\langle \mathbf{u}+\mathbf{v}, \mathbf{w}\rangle &= \langle \mathbf{w}, \mathbf{u}+\mathbf{v}\rangle^* \\ &= (\langle \mathbf{w}, \mathbf{u}\rangle + \langle \mathbf{w}, \mathbf{v}\rangle)^* \\ &= \langle \mathbf{w}, \mathbf{u}\rangle^* + \langle \mathbf{w}, \mathbf{v}\rangle^* \\ &= \langle \mathbf{u}, \mathbf{w}\rangle + \langle \mathbf{v}, \mathbf{w}\rangle,\end{aligned}$$

just as before. However, it follows from axioms (a) and (c) that scalars in the first coordinate must be conjugated when pulling them outside:

$$\langle \alpha\mathbf{u}, \mathbf{v}\rangle = \langle \mathbf{v}, \alpha\mathbf{u}\rangle^* = (\alpha\langle \mathbf{v}, \mathbf{w}\rangle)^* = \alpha^*\langle \mathbf{v}, \mathbf{u}\rangle^* = \alpha^*\langle \mathbf{u}, \mathbf{v}\rangle.$$

Hermitian inner products are sometimes called *sesquilinear* (one-and-a-half times linear) because they are "linear" in the second coordinate,

$$\left\langle \mathbf{u}, \sum \beta_i \mathbf{u}_i \right\rangle = \sum \beta_i \langle \mathbf{u}, \mathbf{v}_i\rangle,$$

and conjugate-linear (i.e., half-linear?) in the first coordinate,

$$\left\langle \sum \alpha_i \mathbf{u}_i, \mathbf{v} \right\rangle = \sum \alpha_i^* \langle \mathbf{u}_i, \mathbf{v}\rangle.$$

Beware: The majority of books do it the other way: linear in the first coordinate and conjugate-linear in the second coordinate.

The whole point of the Hermitian inner product is that the function $\|\mathbf{v}\| := \sqrt{\langle \mathbf{v}, \mathbf{v}\rangle}$ satisfies the norm axioms. Again, the triangle inequality is the hard part. Luckily, the Cauchy-Schwarz inequality still holds exactly as before:

$$|\langle \mathbf{u}, \mathbf{v}\rangle| \leq \|\mathbf{u}\|\|\mathbf{v}\|,$$

where now $|\langle \mathbf{u}, \mathbf{v}\rangle|$ is the absolute value of the complex number $\langle \mathbf{u}, \mathbf{v}\rangle$. The proof is essentially the same. Then we obtain the triangle inequality as follows:

$$\begin{aligned}\|\mathbf{u}+\mathbf{v}\|^2 &= \langle \mathbf{u}+\mathbf{v}, \mathbf{u}+\mathbf{v}\rangle \\ &= \langle \mathbf{u}, \mathbf{u}\rangle + \langle \mathbf{u}, \mathbf{v}\rangle + \langle \mathbf{v}, \mathbf{u}\rangle + \langle \mathbf{v}, \mathbf{v}\rangle \\ &= \langle \mathbf{u}, \mathbf{u}\rangle + \langle \mathbf{u}, \mathbf{v}\rangle + \langle \mathbf{u}, \mathbf{v}\rangle^* + \langle \mathbf{v}, \mathbf{v}\rangle \\ &= \langle \mathbf{u}, \mathbf{u}\rangle + 2\,\mathrm{Re}(\langle \mathbf{u}, \mathbf{v}\rangle) + \langle \mathbf{v}, \mathbf{v}\rangle \\ &\leq \langle \mathbf{u}, \mathbf{u}\rangle + 2|\langle \mathbf{u}, \mathbf{v}\rangle| + \langle \mathbf{v}, \mathbf{v}\rangle && (*) \\ &\leq \langle \mathbf{u}, \mathbf{u}\rangle + 2\|\mathbf{u}\|\|\mathbf{v}\| + \langle \mathbf{v}, \mathbf{v}\rangle && \text{Cauchy-Schwarz}\end{aligned}$$

[28] By combining (8ac) we also have $\langle \alpha\mathbf{u}, \mathbf{v}\rangle = \alpha^*\langle \mathbf{u}, \mathbf{v}\rangle$.

$$= (\|\mathbf{u}\| + \|\mathbf{v}\|)^2.$$

In step $(*)$ we used the fact that for $\alpha = a + ib \in \mathbb{C}$ with real part $\mathrm{Re}(\alpha) = a$ we have

$$\mathrm{Re}(\alpha) = a \leq |a| \leq \sqrt{a^2 + b^2} = |\alpha|.$$

And here we used the fact that

$$\begin{aligned} a^2 &\leq a^2 + b^2 \\ \sqrt{a^2} &\leq \sqrt{a^2 + b^2} \\ |a| &\leq \sqrt{a^2 + b^2}. \end{aligned}$$

Quantum physics is the one of the big reasons for using complex Hermitian spaces, but the complex numbers also allow us to simplify many classical problems.

Example: Complex Fourier Series. Recall Euler's identities:

$$\begin{aligned} e^{i\theta} &= \cos\theta + i\sin\theta, \\ e^{-i\theta} &= \cos\theta - i\sin\theta, \\ \cos\theta &= (e^{i\theta} + e^{-i\theta})/2, \\ \sin\theta &= (e^{i\theta} - e^{-i\theta})/(2i). \end{aligned}$$

Suppose that we have a *real Fourier series*[29]

$$f(x) = a_0 + \sum_{n\geq 1} a_n \cos(2\pi n x) + \sum_{n\geq 1} b_n \sin(2\pi n x).$$

We can write this as a *complex Fourier series*

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{i2\pi n x}$$

by defining the complex coefficients:

$$c_n := \begin{cases} a_0 & n = 0, \\ (a_n - ib_n)/2 & n \geq 1, \\ (a_{-n} + ib_{-n})/2 & n \leq -1. \end{cases}$$

Why would we do this? Because the functions $e^{i2\pi n x}$ are easier to work with! Let's define the *complex L^2 space* $L^2[0,1]$ as the set of functions $[0,1] \to \mathbb{C}$ satisfying

$$\int_0^1 |f(x)|^2 \, dx < \infty.$$

[29] I'll absorb the $\sqrt{2}$ factors into the coefficients this time.

This space has a standard Hermitian inner product:

$$\langle f(x), g(x) \rangle = \int_0^1 f(x)^* g(x)\, dx.$$

It turns out that the functions $e^{i2\pi nx}$ for $n \in \mathbb{Z}$ are an orthonormal set. To see this, we first note that for all real numbers θ we have

$$(e^{i\theta})^* = (\cos\theta + i\sin\theta)^* = \cos\theta - i\sin\theta = \cos(-\theta) + i\sin(-\theta) = e^{-i\theta}.$$

Then we compute

$$\begin{aligned}
\langle e^{i2\pi mx}, e^{i2\pi nx} \rangle &= \int_0^1 (e^{i2\pi mx})^* e^{2\pi inx}\, dx \\
&= \int_0^1 e^{-i2\pi mx} e^{2\pi inx}\, dx \\
&= \int_0^1 e^{i2\pi(n-m)x}\, dx.
\end{aligned}$$

If $m = n$ then this gives

$$\langle e^{i2\pi nx}, e^{i2\pi nx} \rangle = \int_0^1 1\, dx = 1,$$

and if $m \neq n$ then we get

$$\begin{aligned}
\langle e^{i2\pi mx}, e^{i2\pi nx} \rangle &= \int_0^1 e^{i2\pi(n-m)x}\, dx \\
&= \frac{1}{i2\pi(n-m)} \left[e^{i2\pi(n-m)x} \right]_0^1 \\
&= \frac{1}{i2\pi(n-m)} \left[1 - 1 \right] \\
&= 0.
\end{aligned}$$

Note that this was much easier than messing around with trigonometric identities. It also means that we have a single formula for the complex Fourier coefficients:[30]

$$c_n = \langle e^{i2\pi nx}, f(x) \rangle = \int_0^1 e^{-i2\pi nx} f(x)\, dx.$$

It is easy to convert back to the real Fourier coefficients if desired.

[30] Taking the inner product in the other direction gives $\langle f(x), e^{i2\pi nx} \rangle = c_n^*$.

2 Linear Functions

2.1 Matrix Multiplication

In the last section we talked about individual vector spaces such as $\mathbb{R}^n$ and $L^2[0,1]$. Each of these has an inner product, hence it also has a vector norm and a metric. Now we discuss linear functions between different vector spaces. In the finite-dimensional case we can encode such functions as matrices. But matrix arithmetic does more than just encode linear functions; it is an extremely powerful language that gives out much more than we put in.

I assume you already know the definition of matrix multiplication. Here is a reminder.

Definition (Matrices and Matrix Multiplication). An $m \times n$ *matrix over* $\mathbb{R}$ is a rectangular array of real numbers with m rows and n columns. If A is a matrix we typically write $A = (a_{ij})$, where a_{ij} is the entry in the ith row and jth column:

$$A = \begin{pmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{m1} & \cdots & a_{mn} \end{pmatrix}.$$

We let $\mathbb{R}^{m\times n}$ denote the set of $m \times n$ matrices over $\mathbb{R}$. Given two matrices $A = (a_{ij}) \in \mathbb{R}^{\ell\times m}$ and $B = (b_{ij}) \in \mathbb{R}^{m\times n}$ (i.e., where the number of columns in A equals the number of rows in B), we define the *product matrix* $AB = (c_{ij}) \in \mathbb{R}^{\ell\times n}$ by the formula

$$\boxed{c_{ij} := \sum_{k=1}^{m} a_{ik} b_{kj}.}$$

It is important to remark that matrix multiplication is not commutative:

$$AB \neq BA \text{ in general.}$$

Indeed, one of the matrices AB an BA might not be defined, and if they are both defined then they might not have the same shape. Even in the case when AB and BA are both defined and have the same shape, there is no reason for them to be equal.

I could have postponed the definition of matrix multiplication until it emerged naturally from the theory. But, as I said, the topic of matrix arithmetic is more than the sum of its parts, so I wanted to explore the arithmetic properties of the definition before seeing the theory.

Row Times Column = Dot Product. Suppose that $\ell = n = 1$, so that

$$A = \begin{pmatrix} a_{11} & \cdots & a_{1m} \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} b_{11} \\ \vdots \\ b_{m1} \end{pmatrix}.$$

Then the matrix product AB has shape 1×1 (it is just a scalar) and corresponds to the dot product of vectors:

$$\begin{pmatrix} a_{11} & \cdots & a_{1m} \end{pmatrix} \begin{pmatrix} b_{11} \\ \vdots \\ b_{m1} \end{pmatrix} = a_{11}b_{11} + a_{12}b_{21} + \cdots + a_{1m}b_{m1}.$$

From now on we will identify vectors $\mathbf{v} = (v_1, \ldots, v_n) \in \mathbb{R}^n$ with **column vectors**:

$$\mathbf{v} = (v_1, \ldots, v_n) = \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix}.$$

In other words, we identify

$$\mathbb{R}^n = \mathbb{R}^{n \times 1}.$$

If we want to think of a vector $\mathbf{v} \in \mathbb{R}^n$ as a row vector then have to use the following operation.

Definition (Transposition). For any matrix $A = (a_{ij}) \in \mathbb{R}^{m \times n}$ we define the *transpose matrix* $A^T \in \mathbb{R}^{n \times m}$ by exchanging rows and columns:

$$A = \begin{pmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{m1} & \cdots & a_{mn} \end{pmatrix} \quad \mapsto \quad A^T := \begin{pmatrix} a_{11} & \cdots & a_{m1} \\ \vdots & & \vdots \\ a_{1n} & \cdots & a_{mn} \end{pmatrix}.$$

Thus the transpose of a column vector $\mathbf{v} \in \mathbb{R}^n$ is a row vector $\mathbf{v} \in \mathbb{R}^{1 \times n}$:

$$\mathbf{v}^T = \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix}^T = \begin{pmatrix} v_1 & \cdots & v_n \end{pmatrix}.$$

This allows us to express the dot product of any two vectors $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ in terms of matrix multiplication:

$$\mathbf{u}^T\mathbf{v} = \begin{pmatrix} u_1 & \cdots & u_n \end{pmatrix} \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix} u_1v_1 + \cdots + u_nv_n = \mathbf{u} \bullet \mathbf{v}.$$

We will see that this is quite useful.

Column Times Row $\neq$ Dot Product. A column times a row is not a scalar. For any vectors $\mathbf{u} \in \mathbb{R}^m$ and $\mathbf{v} \in \mathbb{R}^n$, the product $\mathbf{u}\mathbf{v}^T$ is the $m \times n$ matrix obtained from the pairwise products of all the entries of $\mathbf{u}$ and $\mathbf{v}$:[31]

$$\mathbf{u}\mathbf{v}^T = \begin{pmatrix} u_1 \\ \vdots \\ u_m \end{pmatrix} \begin{pmatrix} v_1 & \cdots & v_n \end{pmatrix} = \begin{pmatrix} u_1v_1 & \cdots & u_1v_n \\ \vdots & & \vdots \\ u_mv_1 & \cdots & u_mv_n \end{pmatrix}.$$

[31] Later we will call these *rank one matrices*.

Row times column and column times row are the two basic examples. In between there are many different ways to think about matrix multiplication. Here are the most important rules:

$$\begin{aligned}(ij \text{ entry of } AB) &= (i\text{th row of } A)(j\text{th col of } B),\\ (i\text{th row of } AB) &= (i\text{th row of } A)B,\\ (j\text{th col of } AB) &= A(j\text{th col of } B).\end{aligned}$$

If A has shape $\ell \times m$ and B has shape $m \times n$ then we also have

$$AB = \sum_{i=1}^{m} (i\text{th col of } A)(i\text{th row of } B).$$

where the right hand side is a sum of m matrices, each of shape $\ell \times n$. Each of these properties is useful for certain purposes, and each of them is an example of the following very general recursive property.

Theorem (Block Multiplication). Suppose that we partition two matrices into submatrices by inserting vertical and horizontal lines:

$$A = \left(\begin{array}{c|c|c} A_{11} & \cdots & A_{1m} \\ \hline \vdots & & \vdots \\ \hline A_{\ell 1} & \cdots & A_{\ell m} \end{array}\right) \quad \text{and} \quad B = \left(\begin{array}{c|c|c} B_{11} & \cdots & B_{1n} \\ \hline \vdots & & \vdots \\ \hline B_{m1} & \cdots & B_{mn}. \end{array}\right)$$

Let's say that each submatrix A_{ij} has shape $\lambda_i \times \mu_j$ and each B_{ij} has shape $\mu_i \times \nu_j$, so

$$\begin{aligned}\#(\text{rows of } A) &= \lambda_1 + \cdots + \lambda_\ell,\\ \#(\text{cols of } A) &= \mu_1 + \cdots + \mu_m,\\ \#(\text{rows of } B) &= \mu_1 + \cdots + \mu_m,\\ \#(\text{cols of } B) &= \nu_1 + \cdots + \nu_n.\end{aligned}$$

Then I claim that that the product matrix AB can be partitioned as

$$AB = \left(\begin{array}{c|c|c} C_{11} & \cdots & C_{1n} \\ \hline \vdots & & \vdots \\ \hline C_{\ell 1} & \cdots & C_{\ell n} \end{array}\right),$$

where the submatrix C_{ij} is given by

$$\boxed{C_{ij} = \sum_{k=1}^{m} A_{ik}B_{kj}.}$$

Note that $\#(\text{cols of } A_{ik}) = \mu_k = \#(\text{rows of } B_{kj})$ so that each matrix product $A_{ik}B_{kj}$ is defined and has shape $\ell_i \times \nu_j$. Thus C_{ij} is a sum of m matrices, each of shape $\lambda_i \times \nu_j$. In particular,

C_{ij} has shape $\lambda_i \times \mu_j$. Note that the standard formula for unpartitioned matrices corresponds to the case when each submatrix A_{ij} and B_{ij} has size 1×1.

I won't prove right this now because the notation is too messy.[32] Instead let's see some examples illustrating the few rules that we stated above. Let

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 1 & 0 \\ 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

Multiplying rows of A by columns of B gives

$$\left(\begin{array}{ccc} 1 & 1 & 1 \\ \hline 1 & 2 & 3 \end{array}\right) \left(\begin{array}{c|c} 1 & 0 \\ 1 & 1 \\ 0 & 1 \end{array}\right) = \left(\begin{array}{c|c} \begin{pmatrix} 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} & \begin{pmatrix} 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \\ \hline \begin{pmatrix} 1 & 2 & 3 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} & \begin{pmatrix} 1 & 2 & 3 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \end{array}\right) = \left(\begin{array}{c|c} 2 & 2 \\ \hline 3 & 5 \end{array}\right).$$

Multiplying rows of A by B gives

$$\left(\begin{array}{ccc} 1 & 1 & 1 \\ \hline 1 & 2 & 3 \end{array}\right) \begin{pmatrix} 1 & 0 \\ 1 & 1 \\ 0 & 1 \end{pmatrix} = \left(\begin{array}{c} \begin{pmatrix} 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 1 \\ 0 & 1 \end{pmatrix} \\ \hline \begin{pmatrix} 1 & 2 & 3 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 1 \\ 0 & 1 \end{pmatrix} \end{array}\right) = \left(\begin{array}{cc} 2 & 2 \\ \hline 3 & 5 \end{array}\right).$$

Multiplying A by columns of B gives

$$\begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \end{pmatrix} \left(\begin{array}{c|c} 1 & 0 \\ 1 & 1 \\ 0 & 1 \end{array}\right) = \left(\begin{array}{c|c} \begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} & \begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & 3 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix} \end{array}\right) = \left(\begin{array}{c|c} 2 & 2 \\ 3 & 5 \end{array}\right).$$

Finally, multiplying columns of A by rows of B gives

$$\begin{aligned} \left(\begin{array}{c|c|c} 1 & 1 & 1 \\ 1 & 2 & 3 \end{array}\right) \left(\begin{array}{c} 1 \quad 0 \\ \hline 1 \quad 1 \\ \hline 0 \quad 1 \end{array}\right) &= \begin{pmatrix} 1 \\ 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \end{pmatrix} + \begin{pmatrix} 1 \\ 2 \end{pmatrix} \begin{pmatrix} 1 & 1 \end{pmatrix} + \begin{pmatrix} 1 \\ 3 \end{pmatrix} \begin{pmatrix} 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} + \begin{pmatrix} 1 & 1 \\ 2 & 2 \end{pmatrix} + \begin{pmatrix} 0 & 1 \\ 0 & 3 \end{pmatrix} \\ &= \begin{pmatrix} 2 & 2 \\ 3 & 5 \end{pmatrix}. \end{aligned}$$

[32] Later it will follow from properties of linear functions between direct sums of vector spaces.

2.2 Linear Functions

Our ultimate goal is to hide all of the details of matrix arithmetic behind uppercase Roman letters. This lets us ignore irrelevant details to focus on higher level structure. The magic property that makes this work is the *associative property of matrix multiplication*.

Theorem (Associativity of Matrix Multiplication). Consider matrices A, B, C of shapes $\ell \times m$, $m \times n$ and $n \times p$, respectively. Note that the matrices $A(BC)$ and $(AB)C$ are both defined and have the same shape $\ell \times p$. It turns out that they are equal:

$$A(BC) = (AB)C.$$

This is not obvious from the definitions given above. A brute force proof is possible, but not enlightening. There is a much more conceptual explanation.

Definition (Linear Functions). Consider vector spaces U and V over $\mathbb{R}$ (or $\mathbb{C}$). A function $f : U \to V$ is called *linear* when it satisfies the following two properties:

- $f(\mathbf{u}_1 + \mathbf{u}_2) = f(\mathbf{u}_1) + f(\mathbf{u}_2)$ for all $\mathbf{u}_1, \mathbf{u}_2 \in U$,
- $f(\alpha \mathbf{u}) = \alpha f(\mathbf{u})$ for all $\mathbf{u} \in U$ and $\alpha \in \mathbb{R}$.

Setting $\alpha = 0$ in the second axiom shows that every linear function satisfies $f(\mathbf{0}) = \mathbf{0}$.[33] We can summarize the two axioms by saying that "a linear function preserves linear combinations":

$$f(\alpha_1 \mathbf{u}_1 + \cdots + \alpha_n \mathbf{u}_n) = \alpha_1 f(\mathbf{u}_1) + \cdots + \alpha_n f(\mathbf{u}_n).$$

As a teacher, it is a bit difficult to motivate the concept of linear functions. For now, let me just tell you that many natural functions are linear:

- Differention of functions is linear.
- Integration of functions is linear.
- Geometric operations such as rotation, reflection and projection are linear.
- An inner product $\langle -, - \rangle$ on V over $\mathbb{R}$ is *bilinear*. That is, for any $\mathbf{v} \in V$, each of the following two functions is linear:

$$\langle \mathbf{v}, - \rangle : V \to \mathbb{R} \quad \text{and} \quad \langle -, \mathbf{v} \rangle : V \to \mathbb{R}$$

- A Hermitian inner product $\langle -, - \rangle$ on V over $\mathbb{C}$ is sometimes called *sesquilinear* (one-and-a-half times linear). This means that for each fixed $\mathbf{v} \in V$, the function $V \to \mathbb{C}$ defined by $\mathbf{u} \mapsto \langle \mathbf{v}, \mathbf{u} \rangle$ is linear, but the function $V \to \mathbb{C}$ defined by $\mathbf{u} \mapsto \langle \mathbf{u}, \mathbf{v} \rangle$ is *conjugate linear*:

$$\langle \alpha_1 \mathbf{u}_1 + \cdots + \alpha_n \mathbf{u}_n, \mathbf{v} \rangle = \alpha_1^* \langle \mathbf{u}_1, \mathbf{v} \rangle + \cdots + \alpha_n^* \langle \mathbf{u}_n, \mathbf{v} \rangle.$$

[33] The first axiom also implies this, since $\mathbf{0} + \mathbf{0} = \mathbf{0}$ implies $f(\mathbf{0}) = f(\mathbf{0} + \mathbf{0}) = f(\mathbf{0}) + f(\mathbf{0})$ and then we can subtract $f(\mathbf{0})$ from both sides.

The big idea of this section is that linear functions and matrices are essentially the same concept. To express this cleanly, we let $\mathscr{L}(U,V)$ denote the set of all linear functions from U to V. If $\dim U = m$ and $\dim V = n$ then by choosing specific bases for U and V we will obtain a bijection with the set of $n \times m$ matrices:

$$\mathscr{L}(U,V) \quad \longleftrightarrow \quad \mathbb{R}^{n\times m}.$$

For now we use standard bases.[34] Thus we are looking for a bijection

$$\mathscr{L}(\mathbb{R}^m,\mathbb{R}^n) \quad \longleftrightarrow \quad \mathbb{R}^{n\times m}.$$

First we explain how to view a linear function $f:\mathbb{R}^m \to \mathbb{R}^n$ as a matrix. Let $\mathbf{e}_1,\dots,\mathbf{e}_m \in \mathbb{R}^m$ be the standard basis and consider any vector $\mathbf{x} = x_1\mathbf{e}_1 + \cdots + x_m\mathbf{e}_m \in \mathbb{R}^m$. By linearity of f we must have

$$\begin{aligned} f(\mathbf{x}) &= f(x_1\mathbf{e}_1 + \cdots + x_m\mathbf{e}_m) \\ &= x_1 f(\mathbf{e}_1) + \cdots + x_m f(\mathbf{e}_m). \end{aligned}$$

Recall that we view $\mathbf{x} \in \mathbb{R}^m$ and $f(\mathbf{e}_1),\dots,f(\mathbf{e}_m) \in \mathbb{R}^n$ as column vectors. Thus we can view $f(\mathbf{x})$ as a matrix product:

$$f(\mathbf{x}) = x_1 f(\mathbf{e}_1) + \cdots + x_m f(\mathbf{e}_m) = \begin{pmatrix} | & & | \\ f(\mathbf{e}_1) & \cdots & f(\mathbf{e}_m) \\ | & & | \end{pmatrix} \begin{pmatrix} x_1 \\ \vdots \\ x_m \end{pmatrix}.$$

This suggests that we should define the $n \times m$ matrix $[f] \in \mathbb{R}^{n\times m}$ by

$$[f] := \begin{pmatrix} | & & | \\ f(\mathbf{e}_1) & \cdots & f(\mathbf{e}_m) \\ | & & | \end{pmatrix},$$

so that

$$f(\mathbf{x}) = [f]\mathbf{x} \quad \text{ for all } \mathbf{x} \in \mathbb{R}^m.$$

Thus we have a function $f \mapsto [f]$ from the set of linear functions $\mathscr{L}(\mathbb{R}^m,\mathbb{R}^n)$ to the set of matrices $\mathbb{R}^{n\times m}$. I claim that this function is injective. Indeed, suppose we had another linear function $g:\mathbb{R}^m \to \mathbb{R}^n$ with $[f] = [g]$. We define two matrices to be equal when the entries are equal, so this implies that $f(\mathbf{e}_i) = g(\mathbf{e}_i)$ for every basis vector $\mathbf{e}_i$. But then from linearity we must have $f(\mathbf{x}) = g(\mathbf{x})$ for all $\mathbf{x} \in \mathbb{R}^m$ and hence $f = g$ as functions. Indeed, for any $\mathbf{x} = x_1\mathbf{e}_1 + \cdots + x_m\mathbf{e}_m \in \mathbb{R}^m$, the linearity of f and g imply that

$$\begin{aligned} f(\mathbf{x}) &= x_1 f(\mathbf{e}_1) + \cdots + x_m f(\mathbf{e}_m) \\ &= x_1 g(\mathbf{e}_1) + \cdots + x_m g(\mathbf{e}_m) = g(\mathbf{x}). \end{aligned}$$

We summarize this proof with a slogan:

[34] To discuss arbitrary bases right now would hide the important ideas under too much notation. We will return to this topic later when we discuss "diagonalization".

A linear function is uniquely determined by its values on a basis.

It only remains to check that the function $f \mapsto [f]$ is surjective. That is, for a given matrix $A \in \mathbb{R}^{n\times m}$ we must show that there exists some linear function $f : \mathbb{R}^m \to \mathbb{R}^n$ satisfying $[f] = A$. For this purpose we define the function $\ell_A(\mathbf{x}) := A\mathbf{x}$ by matrix multiplication. If $\mathbf{a}_1, \ldots, \mathbf{a}_m \in \mathbb{R}^n$ are the column vectors of A, we recall from the definition of matrix multiplication that $A\mathbf{x}$ is a linear combination of the columns of A:

$$A\mathbf{x} = A\begin{pmatrix} x_1 \\ \vdots \\ x_m \end{pmatrix} = x_1\mathbf{a}_1 + \cdots + x_m\mathbf{a}_m.$$

It follows easily from this that ℓ_A is a linear function, and that $\ell_A(\mathbf{e}_j) = \mathbf{a}_j$ for all j, so that $[\ell_A] = A$ as desired. Here is a summary.

Theorem (Linear Functions = Matrices). Given a linear function $f : \mathbb{R}^m \to \mathbb{R}^n$ we let $[f] \in \mathbb{R}^{n\times m}$ be the matrix whose jth column is $f(\mathbf{e}_j)$, and given a matrix $A \in \mathbb{R}^{n\times m}$ we define the function $\ell_A : \mathbb{R}^m \to \mathbb{R}^n$ by matrix multiplication: $\ell_A(\mathbf{x}) := A\mathbf{x}$. The assignments $f \mapsto [f]$ and $A \mapsto \ell_A$ are inverse bijections between the set of linear functions $\mathscr{L}(\mathbb{R}^m, \mathbb{R}^n)$ and the set of matrices $\mathbb{R}^{n\times m}$:

$$\begin{array}{rcl} \mathscr{L}(\mathbb{R}^m, \mathbb{R}^n) & \longleftrightarrow & \mathbb{R}^{n\times m} \\ f & \mapsto & [f] \\ \ell_A & \leftarrow\!\shortmid & A. \end{array}$$

To say that the assignments are inverse means that $\ell_{[f]} = f$, which says that $f(\mathbf{x}) = [f]\mathbf{x}$ for all $\mathbf{x} \in \mathbb{R}^m$, and $[\ell_A] = A$, which says that $A\mathbf{e}_j$ is the jth column vector of A.

Remark: Recall that we identify vectors with one-column matrices: $\mathbb{R}^n = \mathbb{R}^{n\times 1}$. This means that each vector $\mathbf{x} \in \mathbb{C}^n$ should correspond to a linear function $\ell_{\mathbf{x}} : \mathbb{R}^1 \to \mathbb{R}^n$. Indeed, this is just the function that sends the scalar $t \in \mathbb{R}$ to the vector $t\mathbf{x} \in \mathbb{R}^n$, which we can view as a parametrization of a line.

So far this is only slightly interesting. It becomes much more interesting when we consider composition of linear functions. Let $f : \mathbb{R}^m \to \mathbb{R}^n$ and $g : \mathbb{R}^\ell \to \mathbb{R}^m$ be linear functions and consider the composite function $f \circ g : \mathbb{R}^\ell \to \mathbb{R}^n$ as in the following diagram:

$$\mathbb{R}^\ell \xrightarrow{\ g\ } \mathbb{R}^m \xrightarrow{\ f\ } \mathbb{R}^n, \qquad \mathbb{R}^\ell \xrightarrow{\ f\circ g\ } \mathbb{R}^n$$

We observe that the composite function is also linear, since for any vectors $\mathbf{v}_i \in \mathbb{R}^\ell$ and scalars $c_i \in \mathbb{R}$ we have

$$(f \circ g)\left(\sum c_i\mathbf{v}_i\right) = f\left(g\left(\sum c_i\mathbf{v}_i\right)\right)$$

$$
\begin{aligned}
&= f\left(\sum c_i g(\mathbf{v}_i)\right) && \text{linearity of } g \\
&= \sum c_i f(g(\mathbf{v}_i)) && \text{linearity of } f \\
&= \sum c_i (f \circ g)(\mathbf{v}_i).
\end{aligned}
$$

Thus we have three matrices:

$$
\begin{aligned}
[f] &\text{ has shape } m \times n, \\
[g] &\text{ has shape } \ell \times m, \\
[f \circ g] &\text{ has shape } \ell \times n.
\end{aligned}
$$

We observe that the product matrix $[f][g]$ is defined and has the same shape as the matrix $[f \circ g]$. It turns out that they are equal.

Theorem (Matrix Multiplication = Composition of Functions). Let $f : \mathbb{R}^m \to \mathbb{R}^n$ and $g : \mathbb{R}^\ell \to \mathbb{R}^m$ be linear functions. Then the composite function $f \circ g : \mathbb{R}^\ell \to \mathbb{R}^n$ is linear, and the corresponding three matrices are related by

$$[f \circ g] = [f][g],$$

where the operation on the right is matrix multiplication.

Proof. The proof will use the following property of matrix multiplication:

$$(j\text{th col of } AB) = A(j\text{th col of } B).$$

Our goal is to show that $[f \circ g]$ and $[f][g]$ have the same columns. From the definition of the matrix $[f \circ g]$ we have

$$(j\text{th col of } [f \circ g]) = (f \circ g)(\mathbf{e}_j) = f(g(\mathbf{e}_j)).$$

On the other hand, from the above property of matrix multiplication we have

$$(j\text{th col of } [f][g]) = [f](j\text{th col of } [g]) = [f]g(\mathbf{e}_j) = f(g(\mathbf{e}_j)).$$

□

Remark: It is worth meditating on this proof. When you understand it then you can say that you really understand the concept of matrix multiplication. In fact, it is possible to **define** matrix multiplication via this theorem.

Example (The Identity Matrix). Another important example of a linear function is the *identity function* $\mathrm{id} : \mathbb{R}^n \to \mathbb{R}^n$ defined by $\mathrm{id}(\mathbf{x}) = \mathbf{x}$. The corresponding $n \times n$ matrix is

$$
[\mathrm{id}] = \begin{pmatrix} | & | & & | \\ \mathrm{id}(\mathbf{e}_1) & \mathrm{id}(\mathbf{e}_2) & \cdots & \mathrm{id}(\mathbf{e}_n) \\ | & | & & | \end{pmatrix} = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & 1 \end{pmatrix}.
$$

Since this matrix contains so many zeroes, it is conventional to leave these entries blank. We define the $n \times n$ *identity matrix* as


$$I_n := \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & \ddots & \\ & & & 1 \end{pmatrix}.$$

For any $m \times n$ matrix $A \in \mathbb{R}^{n \times m}$ we have $I_n A = A$. Indeed, we know that $A = [f]$ for some linear function $f : \mathbb{R}^m \to \mathbb{R}^n$. Since $\mathrm{id} \circ f = f$ we have

$$I_n A = [\mathrm{id}][f] = [\mathrm{id} \circ f] = [f] = A.$$

Similarly, we have $AI_m = A$. Of course, these identities can also be proved directly from the definition of matrix multiplication.

Before moving on to some examples, we pause to give the *correct* proof that matrix multiplication is associative. Consider any three functions $f : \mathbb{R}^m \to \mathbb{R}^n$, $g : \mathbb{R}^\ell \to \mathbb{R}^m$ and $h : \mathbb{R}^k \to \mathbb{R}^\ell$, as in the following diagram:



From the definition of functional composition, we have

$$(f \circ (g \circ h))(\mathbf{x}) = f((g \circ h)(\mathbf{x})) = f(g(h(\mathbf{x}))$$

and

$$((f \circ g) \circ h))(\mathbf{x}) = (f \circ g)(h(\mathbf{x})) = f(g(h(\mathbf{x}))$$

for all $\mathbf{x} \in \mathbb{R}^k$, which means that $f \circ (g \circ h)$ and $(f \circ g) \circ h$ are the same function. In other words, the composition of functions is naturally associative. The same property is inherited by matrix multiplication.

Proof of Associativity. Consider matrices A, B, C of shapes $m \times n$, $\ell \times m$ and $k \times \ell$, respectively. We know from an earlier theorem that there exist linear functions $f : \mathbb{R}^m \to \mathbb{R}^n$, $g : \mathbb{R}^\ell \to \mathbb{R}^m$ and $h : \mathbb{R}^k \to \mathbb{R}^\ell$ satisfying $A = [f]$, $B = [g]$ and $C = [h]$. Then from the previous theorem and from the associativity of composition, we have

$$\begin{aligned} A(BC) &= [f]([g][h]) \\ &= [f][g \circ h] \\ &= [f \circ (g \circ h)] \\ &= [(f \circ g) \circ h] \end{aligned}$$

$$
\begin{aligned}
&= [f \circ g][h] \\
&= ([f][g])[h] \\
&= (AB)C.
\end{aligned}
$$

□

We will see in this course that the associative property of matrix multiplication is shockingly useful. We end this section by considering the question of invertibility. We say that a function $f : U \to V$ between sets U and V is invertible if it is injective and surjective. Equivalently, f is invertible if there exists a function $g : V \to U$ satisfying $f \circ g = \mathrm{id}_V$ and $g \circ f = \mathrm{id}_U$. If such a function g exists, then it is unique. Indeed, suppose we had another function $h : V \to U$ satisfying $f \circ h = \mathrm{id}_V$ and $h \circ f = \mathrm{id}_U$. Then we must have

$$g = \mathrm{id}_U \circ g = (h \circ f) \circ g = h \circ (f \circ g) = h \circ \mathrm{id}_V = h.$$

If it exists, we denote the unique inverse function by f^{-1}.

Now suppose that U and V are vector spaces, and $f : U \to V$ is a linear function. If the inverse function $f^{-1} : V \to U$ exists, then it must also be linear. To see this, consider any vectors $\mathbf{v}_1, \mathbf{v}_2 \in V$ and scalars $\alpha, \beta \in \mathbb{R}$. Let $\mathbf{u}_1 := f^{-1}(\mathbf{v}_1)$ and $\mathbf{u}_2 := f^{-1}(\mathbf{v}_2)$. By linearity of f we have

$$\alpha\mathbf{v}_1 + \beta\mathbf{v}_2 = \alpha f(\mathbf{u}_1) + \beta f(\mathbf{u}_2) = f(\alpha\mathbf{u}_1 + \beta\mathbf{u}_2).$$

Then apply f^{-1} to both sides to get

$$
\begin{aligned}
f^{-1}(\alpha\mathbf{v}_1 + \beta\mathbf{v}_2) &= f^{-1}(f(\alpha\mathbf{u}_1 + \beta\mathbf{u}_2)) \\
&= \alpha\mathbf{u}_1 + \beta\mathbf{u}_2 \\
&= \alpha f^{-1}(\mathbf{u}_1) + \beta f^{-1}(\mathbf{u}_2).
\end{aligned}
$$

Now let $U = \mathbb{R}^m$ and $V = \mathbb{R}^n$, and consider the corresponding matrices $[f] \in \mathbb{R}^{n \times m}$ and $[f^{-1}] \in \mathbb{R}^{m \times n}$. From the previous results, we have

$$[f][f^{-1}] = I_m \quad \text{ and } [f^{-1}][f] = I_n.$$

In this case we will say that the matrix $[f^{-1}]$ is the inverse of the matrix $[f]$.

Definition (Inverse Matrices). Let A be an $m \times n$ matrix. Any $n \times m$ matrix B satisfying

$$AB = I_m$$

is called a *right inverse* of A. Any $n \times m$ matrix C satisfying

$$CA = I_n$$

is called a *left inverse* of A. Left and right inverses, if they exist, need not be unique. However, suppose that A has **both** a right inverse B **and** a left inverse C. Then we must have

$$B = I_n B = (CA)B = C(AB) = CI_m = C.$$

In this case $B = C$ is the unique *two-sided inverse* of A, and we write

$$A^{-1} = B = C.$$

When A has a two-sided inverse we say that A is *invertible*. In order to forestall potential confusion, let me immediately mention two key facts:

- If A is invertible then A must be square.
- If A and B are square of the same size, then $AB = I$ implies $BA = I$.

These facts important to know but **their proofs are surprisingly subtle**. We will derive the full proofs from the "rank-nullity theorem" in the next chapter.

2.3 Matrix Analysis

We will see that matrices are extremely useful. There are two main reasons for this:

- Sometimes a complicated situation can be summarized with a single matrix equation.
- Sometimes matrix equations can be solved purely symbolically, without mentioning the entries of the matrices.

In this section we develop some language for working with matrices without mentioning their entries. First we note that the set $\mathbb{R}^{m\times n}$ of $m \times n$ matrices is actually a vector space. For any matrices $A, B \in \mathbb{R}^{m\times n}$ and scalar $\alpha \in \mathbb{R}$, we define matrices $A + B \in \mathbb{R}^{m\times n}$ and $\alpha A \in \mathbb{R}^{m\times n}$ "componentwise":

$$\begin{aligned}(ij \text{ entry of } A + B) &= (ij \text{ entry of } A) + (ij \text{ entry of } B),\\ (ij \text{ entry of } \alpha A) &= \alpha(ij \text{ entry of } A).\end{aligned}$$

The zero vector of this space is the $m \times n$ *zero matrix*:

$$O_{m\times n} = \begin{pmatrix} 0 & \cdots & 0 \\ \vdots & & \vdots \\ 0 & \cdots & 0 \end{pmatrix}.$$

We will simply write O when the shape of the zero matrix is understood . It is easy to verify the vector space properties:

$$\begin{aligned} A + B &= B + A,\\ A + (B + C) &= (A + B) + C,\\ A + O &= O + A = A,\\ 1A &= A,\\ 0A &= O,\\ \alpha(\beta A) &= (\alpha\beta)A, \end{aligned}$$

$$(\alpha+\beta)A = \alpha A + \beta A,$$
$$\alpha(A+B) = \alpha A + \alpha B.$$

Furthermore, there is a *standard basis* of matrices E_{ij} with $1 \leq i \leq m$ and $1 \leq j \leq n$, with the entry 1 in the ij position and all other entries equal to zero:

$$E_{ij} = \begin{array}{c} \\ i \end{array} \begin{array}{c} \begin{array}{c} j \end{array} \\ \begin{pmatrix} \vdots \\ \cdots\ 1 \\ \\ \end{pmatrix} \end{array}$$

(When a matrix contains many zero entries we will simply leave them blank.) Since there are mn such basis matrices, it follows that

$$\dim \mathbb{R}^{m\times n} = mn.$$

Next we have the all-important operation of matrix multiplication. For any three integers ℓ, m, n we can view this as a function from the set $\mathbb{R}^{\ell\times m} \times \mathbb{R}^{m\times n}$ to the set $\mathbb{R}^{\ell\times n}$.

$$\begin{array}{ccc} \mathbb{R}^{\ell\times m} \times \mathbb{R}^{m\times n} & \to & \mathbb{R}^{\ell\times n} \\ (A,B) & \mapsto & AB. \end{array}$$

In terms of the vector space structure on matrices, we say that multiplication is "bilinear" because it satisfies the following distributive rules:

$$A(\beta B + \gamma C) = \beta AB + \gamma AC,$$
$$(\alpha A + \beta B)C = \alpha AC + \beta BC.$$

For each n there is also an identity matrix I_n, as discussed in the previous section. Multiplication interacts with the zero matrices and identity matrices by

$$AO = O,$$
$$OA = O,$$
$$AI = A,$$
$$IA = A,$$

whenever these products are defined. We don't label the shape of the matrices unless necessary.

Next we discuss the symbolic properties of inner products. By default we will work with complex numbers because real inner product spaces are just a special case of complex Hermitian spaces.[35] Here is the motivating question:

[35] Many textbooks prioritize linear algebra over abstract fields, but then they have to relegate inner products to later chapters, which I think is backwards.

How does the Hermitian structure on $\mathbb{C}^n$ interact with linear functions?

This question leads to the following concept.

Theorem (Adjoint of a Linear Operator). Consider the vector spaces $\mathbb{C}^m$ and $\mathbb{C}^n$ with their natural Hermitian inner products, and consider any linear function $f : \mathbb{C}^m \to \mathbb{C}^n$. Then there exists a unique linear function $f^* : \mathbb{C}^n \to \mathbb{C}^m$ satisfying the property

$$\langle f(\mathbf{x}), \mathbf{y} \rangle = \langle \mathbf{x}, f^*(\mathbf{y}) \rangle \quad \text{for all } \mathbf{x} \in \mathbb{C}^m \text{ and } \mathbf{y} \in \mathbb{C}^n.$$

We call f^* the *adjoint of* f. If $A = (a_{ij}) = [f] \in \mathbb{C}^{n \times m}$ is the matrix corresponding to f and $A^* = (b_{ij}) = [f^*] \in \mathbb{C}^{m \times n}$ is the matrix corresponding to f^*, then we have $b_{ij} = a^*_{ji}$. In other words, A^* is the *conjugate transpose* matrix:

$$\begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1m} \\ a_{21} & a_{22} & \cdots & a_{2m} \\ \vdots & \vdots & & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nm} \end{pmatrix}^* = \begin{pmatrix} a^*_{11} & a^*_{21} & \cdots & a^*_{n1} \\ a^*_{12} & a^*_{22} & \cdots & a^*_{n2} \\ \vdots & \vdots & & \vdots \\ a^*_{1m} & a^*_{2m} & \cdots & a^*_{nm} \end{pmatrix}.$$

In the case of real matrices, everything is the same except there is no conjugation. In that case you will often see the notation $A^* = A^T$ for the *transpose* matrix. We won't use this notation much because it is unnecessary.

Before giving the proof, we make a few remarks. If we treat 1×1 matrices as scalars, then the adjoint is just the complex conjugate. This is the reason that I use α^* for complex conjugation instead of the more common $\overline{\alpha}$. The concept of the conjugate transpose matrix allows us to express the inner product on $\mathbb{C}^n$ in terms of matrix multiplication:

$$\langle \mathbf{u}, \mathbf{v} \rangle = u_1^* v_1 + \cdots + u_n^* v_n = \begin{pmatrix} u_1^* & \cdots & u_n^* \end{pmatrix} \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix} = \mathbf{u}^* \mathbf{v}.$$

We will also use the identity $(A\mathbf{x})^* = \mathbf{x}^* A^*$ for vectors $\mathbf{x}$, which can be checked directly from the definitions of matrix multiplication and conjugate transpose.

Proof. Let $A = [f] \in \mathbb{C}^{n \times m}$, let $A^* \in \mathbb{C}^{m \times n}$ be the conjugate transpose matrix, and let $g : \mathbb{C}^n \to \mathbb{C}^m$ be the corresponding linear function, so that $[g] = A^*$. We will show that g satisfies the defining property of the adjoint:

$$\langle f(\mathbf{x}), \mathbf{y} \rangle = \langle \mathbf{x}, g(\mathbf{y}) \rangle \quad \text{for all } \mathbf{x} \in \mathbb{C}^m \text{ and } \mathbf{y} \in \mathbb{C}^n.$$

Indeed, since $f(\mathbf{x}) = A\mathbf{x}$ for all $\mathbf{x} \in \mathbb{C}^m$ and $g(\mathbf{y}) = A^*\mathbf{y}$ for all $\mathbf{y} \in \mathbb{C}^n$, we have[36]

$$\langle f(\mathbf{x}), \mathbf{y} \rangle = \langle A\mathbf{x}, \mathbf{y} \rangle$$

[36]The associativity step $(\mathbf{x}^* A^*)\mathbf{y} = \mathbf{x}^*(A^*\mathbf{y})$ is doing real work here. It is hiding the regrouping of a double summation. This is an example of the miracle of associativity.

$$
\begin{aligned}
&= (A\mathbf{x})^*\mathbf{y} \\
&= (\mathbf{x}^*A^*)\mathbf{y} \\
&= \mathbf{x}^*(A^*\mathbf{y}) \\
&= \langle \mathbf{x}, A^*\mathbf{y}\rangle \\
&= \langle \mathbf{x}, g(\mathbf{y})\rangle.
\end{aligned}
$$

It only remains to show uniqueness. Suppose we have another linear function $h : \mathbb{C}^n \to \mathbb{C}^m$ satisfying $\langle f(\mathbf{x}), \mathbf{y}\rangle = \langle \mathbf{x}, h(\mathbf{y})\rangle$ for all $\mathbf{x} \in \mathbb{C}^m$ and $\mathbf{y} \in \mathbb{C}^n$. Then we have

$$
\begin{aligned}
\langle \mathbf{x}, g(\mathbf{y})\rangle &= \langle \mathbf{x}, h(\mathbf{y})\rangle \\
\langle \mathbf{x}, g(\mathbf{y})\rangle - \langle \mathbf{x}, h(\mathbf{y})\rangle &= 0 \\
\langle \mathbf{x}, g(\mathbf{y}) - h(\mathbf{y})\rangle &= 0.
\end{aligned}
$$

Since this holds for all $\mathbf{x} \in \mathbb{C}^m$ we may choose $\mathbf{x} = g(\mathbf{y}) - h(\mathbf{y})$, so that

$$\langle g(\mathbf{y}) - h(\mathbf{y}), g(\mathbf{y}) - h(\mathbf{y})\rangle = 0.$$

This implies that $g(\mathbf{y}) - h(\mathbf{y}) = 0$ and hence $g(\mathbf{y}) = h(\mathbf{y})$. Finally, since this holds for all $\mathbf{y} \in \mathbb{C}^n$ we conclude that $g = h$. □

Remark: The concept of adjoint operators extends to infinite dimensional inner product spaces, but, in general, not every operator has an adjoint. If $f : U \to V$ is a linear operator between complete[37] inner product spaces, then f has an adjoint if and only if f is continuous with respect to the induced norms on U and V.

We summarize the algebraic properties of the conjugate transpose:

$$
\begin{aligned}
(A^*)^* &= A, \\
(A+B)^* &= A^* + B^* \\
(\alpha A)^* &= \alpha^* A^*, \\
(AB)^* &= B^* A^*.
\end{aligned}
$$

Each of these identities is immediate except for the last one. One can give a brute force proof from the definition of matrix multiplication. For a conceptual proof, let $A = [f]$ and $B = [g]$ for some linear functions $f : \mathbb{C}^m \to \mathbb{C}^n$ and $g : \mathbb{C}^{\ell\times m}$. For all $\mathbf{x} \in \mathbb{C}^\ell$ and $\mathbf{y} \in \mathbb{C}^n$ we have

$$
\begin{aligned}
\langle (f\circ g)(\mathbf{x}), \mathbf{y}\rangle &= \langle f(g(\mathbf{x})), \mathbf{y}\rangle \\
&= \langle g(\mathbf{x}), f^*(\mathbf{y})\rangle \\
&= \langle \mathbf{x}, g^*(f^*(\mathbf{y})\rangle \\
&= \langle \mathbf{x}, (g^*\circ f^*)(\mathbf{y})\rangle,
\end{aligned}
$$

[37] Complete means that every Cauchy sequence converges.

which implies that $(f \circ g)^* = g^* \circ f^*$ by uniqueness of the adjoint. Then the same identity holds for the corresponding matrices:

$$(AB)^* = ([f][g])^* = [f \circ g]^* = [(f \circ g)^*] = [g^* \circ f^*] = [g^*][f^*] = B^*A^*.$$

Now we observe that there is a natural inner product on the vector space of $m \times n$ matrices. It is a direct generalization of the Hermitian inner product on $\mathbb{C}^m = \mathbb{C}^{m\times 1}$.

Definition (Inner Product of Matrices). For any two matrices $A, B \in \mathbb{C}^{m\times n}$ of the same shape, we define

$$\langle A, B\rangle := \sum_{i,j} a_{ij}^* b_{ij}.$$

For column vectors $\mathbf{u}, \mathbf{v} \in \mathbb{C}^{m\times 1}$ we observe that this agrees with the usual inner product.

$$\langle \mathbf{u}, \mathbf{v}\rangle = \mathbf{u}^*\mathbf{v}.$$

To see why this is an inner product, we just need to identify the vector space $\mathbb{C}^{m\times n}$ of $m \times n$ matrices with the vector space $\mathbb{C}^{mn}$ of $mn \times 1$ column vectors. This can be done in many ways, for example by reading across each row like a book. Note that this construction respects addition and scalar products,[38] so the inner product properties on $\mathbb{C}^{m\times n}$ follow from the same properties on $\mathbb{C}^{mn\times 1}$.

Remark: The definition can also be expressed as

$$\langle A, B\rangle = \operatorname{tr}(A^*B),$$

where we define the *trace* of a square matrix $M = (m_{ij})$ as the sum of its diagonal entries:[39]

$$\operatorname{tr}(M) := \sum_i m_{ii}.$$

The trace function can also be defined for operators on some infinite dimensional spaces, in which case $\langle f, g\rangle = \operatorname{tr}(f^* \circ g)$ is called the *Hilbert-Schmidt* inner product.

Now we have an obvious way to measure the size of matrices.

Definition (The Frobenius Norm). For any matrix $A \in \mathbb{C}^{m\times n}$ we define

$$\|A\|_F := \sqrt{\langle A, A\rangle} = \sum_{i,j} a_{ij}^* a_{ij} = \sum_{i,j} |a_{ij}|^2.$$

Note that for any column vector $\mathbf{x}$ we have $\|\mathbf{x}\|_F = \|\mathbf{x}\|$. Since the Frobenius norm is induced from an inner product, it automatically satisfies the norm axioms:

- $\|A\|_F \geq 0$,

[38] Technically, it is an "isomorphism" of vector spaces $\mathbb{C}^{m\times n} \cong \mathbb{C}^{mn\times 1}$.

[39] We will have much more to say about this concept later.

- $\|A\|_F = 0$ implies $A = O$,
- $\|\alpha A\|_F = |\alpha|\|A\|_F$,
- $\|A + B\|_F \leq \|A\|_F + \|B\|_F$.

But this concept is not very useful unless it also has something to do with matrix multiplication. It turns out that

$$\|AB\|_F \leq \|A\|_F \|B\|_F$$

for all matrices A, B such that AB is defined. You will prove this on the homework by first proving the special case $\|A\mathbf{x}\| \leq \|A\|_F\|\mathbf{x}\|$ for all vectors $\mathbf{x}$. Here is the general concept.

Definition (Matrix Norm). A *matrix norm* is a family of norm functions on the vector spaces $\mathbb{C}^{m\times n}$, with the extra property that

$$\|AB\| \leq \|A\|\|B\|,$$

whenever the product matrix AB is defined.

Matrix norms allow us to measure distance between matrices and to define "convergence" for sequences of matrices. To be precise, we say that a sequence of matrices $X_1, X_2, \ldots \in \mathbb{C}^{m\times n}$ converges to a matrix $Y \in \mathbb{C}^{m\times n}$, with respect to a given matrix norm $\|-\|$,[40] if the sequence of scalars $\|X_k - Y\|$ converges to zero. As a first application, we note that for any square matrix A satisfying $\|A\| < 1$, the powers of A converge to the zero matrix:

$$\|A\| < 1 \quad \Longrightarrow \quad A^k \to O.$$

Indeed, by repeatedly applying the sub-multiplicative property, we have $\|A^k\| \leq \|A\|^k$, hence

$$\|A^k - O\| = \|A^k\| \leq \|A\|^k,$$

which converges to the zero scalar if $\|A\| < 1$. Here is a more interesting example.

Example: Geometric Series. Let $\|-\|$ be a matrix norm and let A be a square matrix satisfying $\|A\| < 1$. Then the infinite series $I + A + A^2 + \cdots$ converges to some matrix M with the property that $M(I - A) = (I - A)M$. In other words:

$$\|A\| < 1 \quad \Longrightarrow \quad I + A + A^2 + \cdots = (I - A)^{-1}.$$

Here is a rough sketch of the proof. Let $r := \|A\|$ and consider the partial sums $S_k = I + A + \cdots + A^k$. For $m < n$ we have

$$\|S_n - S_m\| = \left\| \sum_{k=m+1}^{n} A^k \right\| \leq \sum_{k=m+1}^{n} \|A\|^k = r^{m+1} \cdot \frac{1 - r^{n-m}}{1 - r} \leq r^{m+1} \cdot \frac{1}{1 - r}.$$

[40] It turns out that any two norms on a finite dimensional vector space are "topologically equivalent", so it doesn't matter which one we use. Infinite dimensional vector spaces are more complicated.


Since $r < 1$, we can make the right side arbitrarily small by taking m sufficiently large. In other words, the sequence of partial sums $S_1, S_2, \ldots$ is a Cauchy sequence. Since the complex numbers are complete, this implies that S_k converges to some matrix M. Now observe that $S_k(I - A) = I - A^{k+1}$. By our previous remark, the right side converges to I, and by continuity of matrix multiplication, the left side converges to $M(I - A)$. Similarly, the equation $(I - A)S_k = I - A^{k+1}$ leads to $(I - A)M = I$. □

Note that this result holds for any matrix norm. Apart from the Frobenius norm, there is another norm I should mention.

Definition (The Operator Norm). Given a matrix $A \in \mathbb{C}^{m \times n}$, we define the its *operator norm* as the supremum of the length of the vector $A\mathbf{u} \in \mathbb{C}^m$ over all unit vectors $\mathbf{u} \in \mathbb{C}^n$:

$$\|A\|_{\text{op}} := \sup_{\|\mathbf{u}\|=1} \|A\mathbf{u}\|.$$

In the case of real numbers, the image of the unit sphere is an ellipsoid, and the operator norm measures the longest axis of this ellipsoid:



Let's prove that the operator norm is a matrix norm. It turns out that it does not come from an inner product, so we have to prove all of the properties from scratch.

First note that $\|A\|_{\text{op}} \geq 0$ because $\|A\mathbf{u}\| \geq 0$ for all unit vectors $\mathbf{u}$. Next, suppose that $\|A\|_{\text{op}} = 0$. This implies that $\|A\mathbf{u}\| = 0$, and hence $A\mathbf{u} = \mathbf{0}$, for all unit vectors $\mathbf{u}$. But every vector $\mathbf{x}$ can be expressed as $\mathbf{x} = \|\mathbf{x}\|\mathbf{u}$ where $\mathbf{u}$ is a unit vector, hence $A\mathbf{x} = \|\mathbf{x}\|A\mathbf{u} = \mathbf{0}$, so A is the zero matrix. Next, we observe that $\|\alpha A\mathbf{u}\| = |\alpha|\|A\mathbf{u}\|$ for any scalar α and unit vector $\mathbf{u}$, hence

$$\|\alpha A\|_{\text{op}} = \sup_{\|\mathbf{u}\|=1} \|\alpha A\mathbf{u}\| = |\alpha| \sup_{\|\mathbf{u}\|=1} \|A\mathbf{u}\| = |\alpha|\|A\|_{\text{op}}.$$

To prove the triangle inequality, we use the triangle inequality for the vector norm. For any unit vector $\mathbf{u}$ we have $\|A\|_{\text{op}} \leq \|A\mathbf{u}\|$ and $\|B\|_{\text{op}} \leq \|B\mathbf{u}\|$, hence

$$\|(A+B)\mathbf{u}\| = \|A\mathbf{u} + B\mathbf{u}\| \leq \|A\mathbf{u}\| + \|B\mathbf{u}\| \leq \|A\|_{\text{op}} + \|B\|_{\text{op}}.$$

Then since $\|(A+B)\mathbf{u}\| \leq \|A\|_{\text{op}} + \|B\|_{\text{op}}$ for any unit vector $\mathbf{u}$, the supremum $\|A+B\|_{\text{op}}$ is also less than or equal to $\|A\|_{\text{op}} + \|B\|_{\text{op}}$. To prove sub-multiplicativity, we first show that

$$\|A\mathbf{x}\| \leq \|A\|_{\text{op}}\|\mathbf{x}\|$$

for any vector $\mathbf{x}$. Indeed, if $\mathbf{x} = \mathbf{0}$ then the inequality holds trivially. If $\mathbf{x} \neq \mathbf{0}$ then consider the unit vector $\mathbf{u} = \frac{1}{\|\mathbf{x}\|}\mathbf{x}$. Then by definition we have

$$\begin{aligned} \|A\mathbf{u}\| &\leq \|A\|_{\text{op}} \\ \left\|A\frac{1}{\|\mathbf{x}\|}\mathbf{x}\right\| &\leq \|A\|_{\text{op}} \\ \frac{1}{\|\mathbf{x}\|}\|A\mathbf{x}\| &\leq \|A\|_{\text{op}} \\ \|A\mathbf{x}\| &\leq \|A\|_{\text{op}}\|\mathbf{x}\|. \end{aligned}$$

Finally, for any unit vector $\mathbf{u}$ we have

$$\|AB\mathbf{u}\| \leq \|A\|_{\text{op}}\|B\mathbf{u}\| \leq \|A\|_{\text{op}}\|B\|_{\text{op}},$$

and hence $\|AB\|_{\text{op}} \leq \|A\|_{\text{op}}\|B\|_{\text{op}}$. This completes the proof that the operator norm is a matrix norm. It is worth mentioning that the operator norm is generally smaller than the Frobenius norm:

$$\|A\|_{\text{op}} \leq \|A\|_F.$$

Indeed, for any unit vector $\mathbf{u}$ we have $\|A\mathbf{u}\| \leq \|A\|_F\|\mathbf{u}\| = \|A\|_F$, and hence $\|A\|_{\text{op}} \leq \|A\|_F$. For example, the operator norm of the identity matrix is $\|I_n\|_{\text{op}} = 1$, but the Frobenius norm is $\|I_n\|_F = \sqrt{n}$.

We end this section by stating the symbolic properties of matrix inversion:

- If A^{-1} exists then $(A^*)^{-1}$ exists, and $(A^*)^{-1} = (A^{-1})^*$.
- If A^{-1}, B^{-1} and AB exist, then $(AB)^{-1}$ exists, and $(AB)^{-1} = B^{-1}A^{-1}$.

For the first statement we need to show that $A^*(A^{-1})^* = I$ and $(A^{-1})^*A^* = I$. Indeed,[41]

$$A^*(A^{-1})^* = (A^{-1}A)^* = I^* = I,$$

and a similar argument shows that $(A^{-1})^*A^* = I$. For the second statement we need to show that $(AB)(B^{-1}A^{-1}) = I$ and $(B^{-1}A^{-1})AB = I$. This follows from the associativity of matrix multiplication:

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AIA^{-1} = AA^{-1} = I.$$


The other direction is similar.

[41] Recall that $A^*B^* = (BA)^*$.

2.4 Rotations, Reflections, Projections

In this section we consider the most important examples of 2×2 matrices.

Rotations. Consider the function $R_t : \mathbb{R}^2 \to \mathbb{R}^2$ that rotates each point by angle t, counterclockwise around the origin. This function is linear because it sends the origin to itself and it sends parallelograms to parallelograms. To determine the corresponding matrix we only need to rotate the standard basis vectors:



Since no confusion will result, I will use the notation R_t for the function and for the corresponding matrix. Thus we have

$$R_t = \begin{pmatrix} R_t \begin{pmatrix} 1 \\ 0 \end{pmatrix} & R_t \begin{pmatrix} 0 \\ 1 \end{pmatrix} \end{pmatrix} = \begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix}.$$

Once we have the matrix we can use this to rotate a general point:

$$R_t \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \cos t - y \sin t \\ x \sin t + y \cos t \end{pmatrix}.$$

It would be much harder to solve this problem without the theory of matrices. Next we consider the composition of two rotations. Thinking in terms of functions, it is clear that $R_s R_t = R_{s+t} = R_t R_s$, since rotating first by one angle and then by the other angle is the same as rotating once by the sum of the two angles. On the other hand, since matrix multiplication is the same as functional composition, we obtain the following matrix identity, which is equivalent

to the angle sum trigonometric identities:

$$\begin{pmatrix} \cos s & -\sin s \\ \sin s & \cos s \end{pmatrix} \begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix} = \begin{pmatrix} \cos(s+t) & -\sin(s+t) \\ \sin(s+t) & \cos(s+t) \end{pmatrix}.$$

Note that rotation clockwise by angle t is the same as rotation counterclockwise by angle $-t$. Thus the functions R_t and R_{-t} are inverses:

$$R_t R_{-t} = R_{-t} R_t = R_0 = I.$$

Note that rotation by angle zero is just the identity function. It is interesting to observe that


$$\begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix}^{-1} = (R_t)^{-1} = R_{-t} = \begin{pmatrix} \cos(-t) & -\sin(-t) \\ \sin(-t) & \cos(-t) \end{pmatrix} = \begin{pmatrix} \cos t & \sin t \\ -\sin t & \cos t \end{pmatrix} = (R_t)^T.$$

We will see below the matrices satisfying $A^{-1} = A^T$ are called *orthogonal matrices*. Finally, let me remark that the determinant of a rotation matrix is always 1:

$$\det(R_t) = \det \begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix} = \cos^2 t + \sin^2 t = 1.$$

We will discuss the general theory of determinants later.

Reflections. Let $F_t : \mathbb{R}^2 \to \mathbb{R}^2$ be the function that reflects each point across the line that makes angle $t/2$ from the positive x-axis. Again, this is a linear function because it sends the origin to itself and sends parallelograms to parallelograms. To determine the corresponding matrix we reflect the standard basis vectors:



Thus we obtain the matrix

$$F_t = \begin{pmatrix} F_t \begin{pmatrix} 1 \\ 0 \end{pmatrix} & F_t \begin{pmatrix} 0 \\ 1 \end{pmatrix} \end{pmatrix} = \begin{pmatrix} \cos t & \sin t \\ \sin t & -\cos t \end{pmatrix}.$$

The composition of two reflections in two different lines turns out to be a rotation:

$$\begin{aligned} F_s F_t &= \begin{pmatrix} \cos s & \sin s \\ \sin s & -\cos s \end{pmatrix} \begin{pmatrix} \cos t & \sin t \\ \sin t & -\cos t \end{pmatrix} \\ &= \begin{pmatrix} \cos s \cos t + \sin s \sin t & \cos s \sin t - \sin s \cos t \\ \sin s \cos t - \cos s \sin t & \sin s \sin t + \cos s \cos t \end{pmatrix} \\ &= \begin{pmatrix} \cos(s-t) & -\sin(s-t) \\ \sin(s-t) & \cos(s-t) \end{pmatrix} \\ &= R_{s-t}. \end{aligned}$$

This would be more difficult to see geometrically. In particular, we find that reflection matrices do not commute in general:

$$F_s F_t = R_{s-t} \neq R_{t-s} = F_t F_s \quad \text{unless angles } s-t \text{ and } t-s \text{ are equal.}$$

Taking $s = t$ shows that the composition of a reflection with itself is the identity matrix:

$$F_t^2 = F_t F_t = R_{t-t} = R_0 = I.$$

In other words, reflecting in the same line twice is the same thing as doing nothing. This implies that each reflection matrix F_t is equal to its own inverse:

$$(F_t)^{-1} = F_t.$$


It also happens that $(F_t)^T = F_t$, so F_t is another example of an orthogonal matrix. Finally, let me remark that the determinant of any reflection matrix is -1:

$$\det(F_t) = \det \begin{pmatrix} \cos t & \sin t \\ \sin t & -\cos t \end{pmatrix} = -\cos^2 t - \sin^2 t = -1.$$

Projections. Consider the following matrix:

$$P_t = \begin{pmatrix} \cos^2 t & \cos t \sin t \\ \cos t \sin t & \sin^2 t \end{pmatrix}.$$

As with any 2×2 matrix, this defines a linear function $\mathbb{R}^2 \to \mathbb{R}^2$. We will show that this is the orthogonal projection onto the line generated by the vector $\mathbf{a} = (\cos t, \sin t)$. In fact, we will prove a more general statement. Let $\mathbf{a} \in \mathbb{R}^n$ be a nonzero vector in n-dimensional Euclidean space and let $P_{\mathbf{a}}$ be the orthogonal projection onto the line generated by $\mathbf{a}$. Based on the explicit formulas in the previous chapter we know that this function is linear, and for a given point $\mathbf{x}$, the point $P_{\mathbf{a}}\mathbf{x}$ is uniquely characterized by the fact that $P_{\mathbf{a}}\mathbf{x}$ is in the line and $P_{\mathbf{a}}\mathbf{x} - \mathbf{x}$ is orthogonal to the line generated by $\mathbf{a}$:



We can express these two facts algebraically:

- $P_{\mathbf{a}}\mathbf{x} = \alpha\mathbf{a}$ for some $\alpha \in \mathbb{R}$,
- $\mathbf{a}^T(P_{\mathbf{a}}\mathbf{x} - \mathbf{x}) = \langle \mathbf{a}, P_{\mathbf{a}}\mathbf{x} - \mathbf{x}\rangle = 0$.

Combining these two facts gives

$$\begin{aligned} \mathbf{a}^T(P_{\mathbf{a}}\mathbf{x} - \mathbf{x}) &= 0 \\ \mathbf{a}^T(\alpha\mathbf{a} - \mathbf{x}) &= 0 \\ \alpha\mathbf{a}^T\mathbf{a} - \mathbf{a}^T\mathbf{x} &= 0 \\ \alpha &= \mathbf{a}^T\mathbf{x}/\mathbf{a}^T\mathbf{a} \\ \alpha &= \mathbf{a}^T\mathbf{x}/\|\mathbf{a}\|^2. \end{aligned}$$

Hence the projection of $\mathbf{x}$ is given by

$$P_{\mathbf{a}}\mathbf{x} = \alpha\mathbf{a} = \underbrace{\left(\frac{\mathbf{a}^T\mathbf{x}}{\|\mathbf{a}\|^2}\right)}_{\text{scalar}} \underbrace{\mathbf{a}}_{\text{vector}}.$$

Note that this agrees with the abstract formula from the previous chapter:

$$P_{\mathbf{a}}\mathbf{x} = \frac{\langle \mathbf{a}, \mathbf{x}\rangle}{\langle \mathbf{a}, \mathbf{a}\rangle}\,\mathbf{a}.$$

To find a formula for the $n \times n$ projection matrix $P_{\mathbf{a}}$ we simply rearrange using the fact that matrix multiplication is associative:

$$P_{\mathbf{a}}\mathbf{x} = \underbrace{\left(\frac{\mathbf{a}^T\mathbf{x}}{\|\mathbf{a}\|^2}\right)}_{\text{scalar}} \underbrace{\mathbf{a}}_{\text{vector}}$$

$$= \frac{1}{\|\mathbf{a}\|^2}\mathbf{a}(\mathbf{a}^T\mathbf{x}) \qquad \text{scalars commute with matrices}$$

$$= \underbrace{\frac{1}{\|\mathbf{a}\|^2}(\mathbf{a}\mathbf{a}^T)}_{n \times n \text{ matrix}} \underbrace{\mathbf{x}}_{\text{vector}}. \qquad \text{associativity}$$

Since this identity holds for any $\mathbf{x}$, we conclude that the projection matrix is given by

$$P_{\mathbf{a}} = \frac{1}{\|\mathbf{a}\|^2}\,\mathbf{a}\mathbf{a}^T.$$

If $\mathbf{a} = \mathbf{u}$ is a unit vector then the formula is particularly simple:

$$P_{\mathbf{u}} = \mathbf{u}\mathbf{u}^T = \text{the projection onto the line in } \mathbb{R}^n \text{ spanned by unit vector } \mathbf{u}.$$


Now we go back to two dimensions. Consider the line in $\mathbb{R}^2$ that makes angle t counterclockwise from the positive x-axis. This line is generated by the unit vector $\mathbf{u} = (\cos t, \sin t)$. Hence the matrix that projects onto this line is

$$P_t = \mathbf{u}\mathbf{u}^T = \begin{pmatrix}\cos t \\ \sin t\end{pmatrix}\begin{pmatrix}\cos t & \sin t\end{pmatrix} = \begin{pmatrix}\cos^2 t & \cos t \sin t \\ \cos t \sin t & \sin^2 t\end{pmatrix}.$$

The image of a general point $\mathbf{x} = (x, y)$ under this projection is

$$P_t \begin{pmatrix}x \\ y\end{pmatrix} = \begin{pmatrix}\cos^2 t & \cos t \sin t \\ \cos t \sin t & \sin^2 t\end{pmatrix}\begin{pmatrix}x \\ y\end{pmatrix} = \begin{pmatrix}x\cos^2 t + y \cos t \sin t \\ x \cos t \sin t + y \sin^2 t\end{pmatrix}.$$

Here is a picture:



Note that this projection is **not invertible**. To see this, let's consider the point $(-\sin t, \cos t)$. This point gets projected to the origin:

$$\begin{aligned} P_t \begin{pmatrix} -\sin t \\ \cos t \end{pmatrix} &= \begin{pmatrix} \cos^2 t & \cos t \sin t \\ \cos t \sin t & \sin^2 t \end{pmatrix} \begin{pmatrix} -\sin t \\ \cos t \end{pmatrix} \\ &= \begin{pmatrix} -\cos^2 t \sin t + \cos^2 t \sin t \\ -\cos t \sin^2 t + \cos t \sin^2 t \end{pmatrix} \\ &= \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \end{aligned}$$

But we also have $P_t \mathbf{0} = \mathbf{0}$. If P_t had an inverse matrix $(P_t)^{-1}$ then this would imply that

$$\begin{aligned} P_t \begin{pmatrix} -\sin t \\ \cos t \end{pmatrix} &= P_t \begin{pmatrix} 0 \\ 0 \end{pmatrix} \\ (P_t)^{-1} P_t \begin{pmatrix} -\sin t \\ \cos t \end{pmatrix} &= (P_t)^{-1} P_t \begin{pmatrix} 0 \\ 0 \end{pmatrix} \\ \begin{pmatrix} -\sin t \\ \cos t \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \end{aligned}$$

which is a contradiction. More generally, let $P_{\mathbf{u}} = \mathbf{u}\mathbf{u}^T$ be the matrix that projects onto the line in $\mathbb{R}^n$ generated by some unit vector $\mathbf{u}$ and let $\mathbf{x} \in \mathbb{R}^n$ be any nonzero vector that is perpendicular to $\mathbf{u}$, so that $\mathbf{u}^T \mathbf{x} = 0$. Then we have

$$P_{\mathbf{u}} \mathbf{x} = (\mathbf{u}\mathbf{u}^T)\mathbf{x} = \mathbf{u}(\mathbf{u}^T \mathbf{x}) = \mathbf{u}(0) = \mathbf{0}.$$

If $P_{\mathbf{u}}$ had an inverse then this would imply that $\mathbf{x} = I\mathbf{x} = (P_{\mathbf{u}})^{-1} P_{\mathbf{u}} \mathbf{x} = (P_{\mathbf{u}})^{-1} \mathbf{0} = \mathbf{0}$, which contradicts the fact that $\mathbf{x}$ is nonzero.[42] Finally, let me note that the matrix P_t has determinant zero:

$$\det P_t = \det \begin{pmatrix} \cos^2 t & \cos t \sin t \\ \cos t \sin t & \sin^2 t \end{pmatrix} = \cos^2 t \sin^2 t - \cos^2 t \sin^2 t = 0.$$

Later we will see that a square matrix A is invertible if and only if $\det A \neq 0$.

2.5 Unitary and Orthogonal Matrices

One of the most important classes of matrices are those that have orthonormal columns. There is a nice way to express this in terms of matrix multiplication. Consider a complex matrix $U : \mathbb{C}^m \to \mathbb{C}^n$. Then we have

$$U \text{ has orthonormal columns} \quad \Longleftrightarrow \quad U^* U = I_n.$$

Indeed, let $\mathbf{u}_1, \ldots, \mathbf{u}_n \in \mathbb{C}^m$ denote the column vectors of U, so that $\mathbf{u}_1^*, \ldots, \mathbf{u}_n^*$ are the row vectors of U^*. From the properties of matrix multiplication we have

$$(ij \text{ entry of } U^*U) = (i\text{-th row of } U^*)(j\text{-th column of } U) = \mathbf{u}_i^* \mathbf{u}_j = \langle \mathbf{u}_i, \mathbf{u}_j \rangle.$$

[42] The general principle is that a matrix that sends some nonzero vector to zero is not invertible. If the matrix is square we will later see that the converse is also true.

Let δ_{ij} be the Kronecker delta. Then we have

$$\begin{aligned} U^*U = I_n &\iff (ij \text{ entry of } U^*U) = \delta_{ij} \\ &\iff \langle \mathbf{u}_i, \mathbf{u}_j \rangle = \delta_{ij} \\ &\iff \mathbf{u}_1, \ldots, \mathbf{u}_n \text{ are orthonormal.} \end{aligned}$$

(In the context of real numbers, one can use the notation $U^T = U^*$.) Geometrically, these matrices correspond to linear functions that preserve distances and angles.

Theorem (Linear Isometries). Consider a matrix $U \in \mathbb{C}^{m \times n}$. The following are equivalent:

(1) $U^*U = I_n$,

(2) $\langle U\mathbf{x}, U\mathbf{y} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle$ for all $\mathbf{x}, \mathbf{y} \in \mathbb{C}^n$,

(3) $\|U\mathbf{x}\| = \|\mathbf{x}\|$ for all $\mathbf{x} \in \mathbb{C}^n$,

(4) $\|U\mathbf{x} - U\mathbf{y}\| = \|\mathbf{x} - \mathbf{y}\|$ for all $\mathbf{x}, \mathbf{y} \in \mathbb{C}^n$.

Part (4) says that U preserves distances between points, which is the meaning of "isometry".

The hardest part of the proof is to show that (3) implies (2). For that we will use the idea of "polarization", which allows us to recover the inner product from its norm.

Lemma (Polarization). In complex Hermitian space we have

$$\Re\langle \mathbf{x}, \mathbf{y} \rangle = \frac{1}{4} \left(\|\mathbf{x} + \mathbf{y}\|^2 - \|\mathbf{x} - \mathbf{y}\|^2 \right).$$

In real Euclidean space we have $\Re\langle \mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle$, so that

$$\langle \mathbf{x}, \mathbf{y} \rangle = \frac{1}{4} \left(\|\mathbf{x} + \mathbf{y}\|^2 - \|\mathbf{x} - \mathbf{y}\|^2 \right).$$

Proof (Polarization). We have

$$\|\mathbf{x} + \mathbf{y}\|^2 = \langle \mathbf{x} + \mathbf{y}, \mathbf{x} + \mathbf{y} \rangle = \langle \mathbf{x}, \mathbf{x} \rangle + \langle \mathbf{y}, \mathbf{y} \rangle + \langle \mathbf{x}, \mathbf{y} \rangle + \langle \mathbf{y}, \mathbf{x} \rangle$$

and

$$\|\mathbf{x} - \mathbf{y}\|^2 = \langle \mathbf{x} - \mathbf{y}, \mathbf{x} - \mathbf{y} \rangle = \langle \mathbf{x}, \mathbf{x} \rangle + \langle \mathbf{y}, \mathbf{y} \rangle - \langle \mathbf{x}, \mathbf{y} \rangle - \langle \mathbf{y}, \mathbf{x} \rangle.$$

Subtracting the two equations gives

$$\|\mathbf{x} + \mathbf{y}\|^2 - \|\mathbf{x} - \mathbf{y}\|^2 = 2(\langle \mathbf{x}, \mathbf{y} \rangle + \langle \mathbf{y}, \mathbf{x} \rangle) = 4\Re\langle \mathbf{x}, \mathbf{y} \rangle.$$

The last equation holds because $\langle \mathbf{x}, \mathbf{y} \rangle$ and $\langle \mathbf{y}, \mathbf{x} \rangle$ are complex conjugates, and $\alpha + \alpha^* = 2\Re\alpha$ for any complex number $\alpha \in \mathbb{C}$. □

Proof (Linear Isometries). First we will prove

$$(1) \Leftrightarrow (2) \Rightarrow (3) \Leftrightarrow (4),$$

and then we will prove the harder implication (3) $\Rightarrow$ (2). (1)$\Rightarrow$(2): Suppose $U^*U = I$. Then

$$\langle U\mathbf{x}, U\mathbf{y}\rangle = (U\mathbf{x})^*(U\mathbf{y}) = \mathbf{x}^*U^*U\mathbf{y} = \mathbf{x}^*I\mathbf{y} = \mathbf{x}^*\mathbf{y} = \langle \mathbf{x}, \mathbf{y}\rangle.$$

(2)$\Rightarrow$(1): Suppose that $\langle U\mathbf{x}, U\mathbf{y}\rangle = \langle \mathbf{x}, \mathbf{y}\rangle$ for all $\mathbf{x}, \mathbf{y} \in \mathbb{C}^n$. In matrix language this says that $\mathbf{x}^*U^*U\mathbf{y} = \mathbf{x}^*\mathbf{y}$. Letting $\mathbf{x} = \mathbf{e}_i$ and $\mathbf{y} = \mathbf{e}_j$ be standard basis vectors shows that

$$(ij \text{ entry of } U^*U) = \mathbf{e}_i^*U^*U\mathbf{e}_j = \mathbf{e}_i^*\mathbf{e}_j = \begin{cases} 1 & i = j, \\ 0 & i \neq j, \end{cases}$$

and hence $U^*U = I$. (2)$\Rightarrow$(3): Suppose that $\langle U\mathbf{x}, U\mathbf{y}\rangle = \langle \mathbf{x}, \mathbf{y}\rangle$ for all $\mathbf{x}, \mathbf{y} \in \mathbb{C}^n$. Taking $\mathbf{y} = \mathbf{x}$ gives

$$\|U\mathbf{x}\|^2 = \langle U\mathbf{x}, U\mathbf{x}\rangle = \langle \mathbf{x}, \mathbf{x}\rangle = \|x\|^2,$$

and then taking square roots gives the result. (3)$\Rightarrow$(4): Suppose that $\|U\mathbf{z}\| = \|\mathbf{z}\|$ for all $\mathbf{z} \in \mathbb{C}^n$. Then taking $\mathbf{z} = \mathbf{x} - \mathbf{y}$ gives

$$\|U\mathbf{x} - U\mathbf{y}\| = \|U(\mathbf{x} - \mathbf{y})\| = \|U\mathbf{z}\| = \|\mathbf{z}\| = \|\mathbf{x} - \mathbf{y}\|.$$

(4)$\Rightarrow$(3): Suppose $\|U\mathbf{x} - U\mathbf{y}\| = \|\mathbf{x} - \mathbf{y}\|$ for all $\mathbf{x}, \mathbf{y} \in \mathbb{C}^n$. Then taking $\mathbf{y} = \mathbf{0}$ gives

$$\|U\mathbf{x}\| = \|U\mathbf{x} - \mathbf{0}\| = \|U\mathbf{x} - U\mathbf{0}\| = \|\mathbf{x} - \mathbf{0}\| = \|\mathbf{x}\|.$$

Now we prove the harder implication (3) $\Rightarrow$ (2). Suppose that $\|U\mathbf{x}\| = \|\mathbf{x}\|$ for all $\mathbf{x} \in \mathbb{C}^n$. Then for all $\mathbf{x}, \mathbf{y} \in \mathbb{C}^n$, the polarization identity gives

$$\begin{aligned} \Re\langle U\mathbf{x}, U\mathbf{y}\rangle &= \frac{1}{4}\left(\|U\mathbf{x} + U\mathbf{y}\|^2 - \|U\mathbf{x} - U\mathbf{y}\|^2\right) \\ &= \frac{1}{4}\left(\|U(\mathbf{x} + \mathbf{y})\|^2 - \|U(\mathbf{x} - \mathbf{y})\|^2\right) \\ &= \frac{1}{4}\left(\|\mathbf{x} + \mathbf{y}\|^2 - \|\mathbf{x} - \mathbf{y}\|^2\right) \\ &= \Re\langle \mathbf{x}, \mathbf{y}\rangle. \end{aligned}$$

In order to complete the proof that $\langle U\mathbf{x}, U\mathbf{y}\rangle = \langle \mathbf{x}, \mathbf{y}\rangle$, we also need to show that the imaginary parts satisfy $\Im\langle U\mathbf{x}, U\mathbf{y}\rangle = \Im\langle \mathbf{x}, \mathbf{y}\rangle$. For this we will use the fact that $\langle i\mathbf{x}, \mathbf{y}\rangle = -i\langle \mathbf{x}, \mathbf{y}\rangle$, so that $\Im\langle \mathbf{x}, \mathbf{y}\rangle = \Re\langle i\mathbf{x}, \mathbf{y}\rangle$. Then for all $\mathbf{x}, \mathbf{y} \in \mathbb{C}^n$ we have

$$\Im\langle U\mathbf{x}, U\mathbf{y}\rangle = \Re\langle iU\mathbf{x}, U\mathbf{y}\rangle = \Re\langle U(i\mathbf{x}), U\mathbf{y}\rangle = \Re\langle i\mathbf{x}, \mathbf{y}\rangle = \Im\langle \mathbf{x}, \mathbf{y}\rangle.$$

□

There is a similar theorem for matrices $U \in \mathbb{C}^{m\times n}$ satisfying $UU^* = I_m$:

$$U \text{ has orthonormal rows} \quad \Longleftrightarrow \quad UU^* = I_m$$

$$\begin{aligned} &\iff && \langle U^*\mathbf{x}, U^*\mathbf{y}\rangle = \langle \mathbf{x}, \mathbf{y}\rangle \text{ for all } \mathbf{x}, \mathbf{y} \in \mathbb{C}^m \\ &\iff && \|U^*\mathbf{x}\| = \|\mathbf{x}\| \text{ for all } \mathbf{x} \in \mathbb{C}^m \\ &\iff && \|U^*\mathbf{x} - U^*\mathbf{y}\| = \|\mathbf{x} - \mathbf{y}\| \text{ for all } \mathbf{x}, \mathbf{y} \in \mathbb{C}^m. \end{aligned}$$

In this case we say that $U^* : \mathbb{C}^m \to \mathbb{C}^n$ is a linear isometry.

Definition (Unitary and Orthogonal Matrices). We say that a matrix U with complex entries is *unitary* if both U and U^* are isometries. That is, if

$$U^*U = I \quad \text{and} \quad UU^* = I.$$

A unitary matrix with real entries is called an *orthogonal* matrix.

Remark: To avoid confusion, let me remind you that if matrices A and B satisfy $AB = I$ and $BA = I$, then A and B must be square. Furthermore, for square matrices A and B we have $AB = I$ if and only if $BA = I$. (Both of these facts will follow from the fundamental theorem in the next chapter.) So we could have defined a unitary matrix as a square matrix U satisfying $U^*U = I$.

We note that unitary matrices preserve the Frobenius and operator norms. That is, for any matrix $A \in \mathbb{C}^{m\times n}$ and unitary matrices $U \in \mathbb{C}^{m\times m}, V \in \mathbb{C}^{n\times n}$, we have

$$\|UAV\|_F = \|A\|_F \quad \text{and} \quad \|UAV\|_{\text{op}} = \|A\|_{\text{op}}.$$

For the Frobenius norm, recall that $\|A^*\|_F = \|A\|_F = \sqrt{\text{tr}(A^*A)}$. For any unitary matrix U this implies that

$$\|UA\| = \sqrt{\text{tr}((UA)^*UA)} = \sqrt{\text{tr}(A^*U^*UA)} = \sqrt{\text{tr}(A^*A)} = \|A\|_F.$$

Then for any unitary V we have

$$\|AV\|_F = \|(AV)^*\|_F = \|V^*A^*\|_F = \|A^*\|_F = \|A\|_F.$$

For the operator norm, recall that $\|A\|_{\text{op}}$ is the supremum of $\|A\mathbf{u}\|$ over unit vectors $\mathbf{u}$. If U is unitary then $\|UA\mathbf{u}\| = \|A\mathbf{u}\|$, so the supremum is the same. If V is unitary we note that $\|\mathbf{u}\| = 1$ implies $\|V\mathbf{u}\| = 1$, and since V is invertible (indeed, $V^{-1} = V^*$), $V\mathbf{u}$ ranges over all unit vectors as $\mathbf{u}$ ranges over all unit vectors. So the supremum of $\|AV\mathbf{u}\|$ as $\mathbf{u}$ ranges over unit vectors equals the supremum of $\|A(V\mathbf{u})\|$ as $V\mathbf{u}$ ranges over unit vectors, which is $\|A\|_{\text{op}}$.

What happens in higher dimensions?
For the remainder of this section, we discuss the relationship between real orthogonal matrices and the geometry of Euclidean space.

Theorem (2×2 Orthogonal Matrices). Any real 2×2 orthogonal matrix is either a rotation or a reflection.

Proof. Let A be a 2×2 matrix with columns $\mathbf{a}_1, \mathbf{a}_2 \in \mathbb{R}^2$. If $A^T A = I$ then the vectors $\mathbf{a}_1, \mathbf{a}_2$ are orthonormal. Since $\|\mathbf{a}_1\| = 1$ we can write $\mathbf{a}_1 = (\cos t, \sin t)$ for some unique angle $0 \leq t < 2\pi$. Then since $\|\mathbf{a}_2\| = 1$ and $\mathbf{a}_1^T \mathbf{a}_2 = 0$ we musty have $\mathbf{a}_2 = (-\sin t, \cos t)$, in which case $A = R_t$ is a rotation matrix, or $\mathbf{a}_2 = (\sin t, -\cos t)$, in which case $A = F_t$ is a reflection matrix.[43] □

State and prove the isometry theorem, using the polarization identity as a lemma.

Define reflections $F_{\mathbf{u}}$ and state their relation to projections $P_{\mathbf{u}}$. State and prove the Cartan-Dieudonné theorem.

Cartan-Dieudonné Let $A \in \mathbb{R}^{n \times n}$ satisfy $A^T A = I$. We will show that there exist

Let $A^T A = I$ be orthogonal (square) and let $U \subseteq \mathbb{R}^n$ be a subspace. If A fixes U then A stabilizes $U^\perp$. Proof: Let $\mathbf{x} \in U^\perp$. Then for any $\mathbf{u} \in U$ we have $0 = \mathbf{x}^T \mathbf{u} = \mathbf{x}^T A^T A \mathbf{u} = (A\mathbf{x})^T (A\mathbf{u}) = (A\mathbf{x})^T \mathbf{u}$. □ Let U_k be the span of $\mathbf{e}_1, \ldots, \mathbf{e}_k$. We will construct reflections $F_1, \ldots, F_n$ such that $A_k := F_k \cdots F_1 A$ fixes U_k, so that $A_n = I$. Suppose done for $A_{k-1} = F_{k-1} \cdots F_1 A$. If $\mathbf{v}_k := A_{k-1}\mathbf{e}_k - \mathbf{e}_k \neq \mathbf{0}$, let $\mathbf{u}_k = \mathbf{v}_k / \|\mathbf{v}_k\|$ so that $F_k := F_{\mathbf{u}_k}$ exchanges $A_{k-1}\mathbf{e}_k$ and $\mathbf{e}_k$, so $A_k \mathbf{e}_k = \mathbf{e}_k$. If $\mathbf{v}_k = \mathbf{0}$ let $F_k = I$. Since A_{k-1} is orthogonal and fixes U_{k-1}, it preserves $U_{k-1}^\perp$. Since $\mathbf{e}_k \in U_{k-1}^\perp$, we also have $A_{k-1}\mathbf{e}_k \in U_{k-1}^\perp$, hence $\mathbf{v}_k$ and $\mathbf{u}_k \in U_{k-1}^\perp$. But then F_k fixes U_{k-1} because $F_k \mathbf{x} = \mathbf{x} - 2\mathbf{u}_k \mathbf{u}_k^T \mathbf{x} = \mathbf{x}$ for any $\mathbf{u}_k^T \mathbf{x} = 0$. Since A_{k-1} and F_k fix U_{k-1}, so does A_k. Since A_k also fixes $\mathbf{e}_k$, A_k fixes U_k. Done.

3 Subspaces Associated to Linear Functions

3.1 The Rank-Nullity Theorem

In the last section we discussed purely symbolic properties of matrix inversion. Recall: Let A be an $m \times n$ matrix. An $n \times m$ matrix B is called a *right inverse of* A when $AB = I_m$ and an $m \times n$ matrix C is called a *left inverse of* A when $CA = I_n$. If A has both a right inverse B and a left inverse C then the two must be equal because

$$B = I_n B = (CA)B = C(AB) = CI_m = C.$$

In this case we say that $A^{-1} = B = C$ is the unique *two-sided inverse of* A. Any matrix having a two-sided inverse is called *invertible*. We also proved the following basic facts: If A^{-1} exists then $(A^*)^{-1}$ exists and is equal to $(A^{-1})^*$. If A^{-1}, B^{-1} and AB exist then $(AB)^{-1}$ exists and is equal to $B^{-1}A^{-1}$. Then I mentioned "two subtle theorems", that we were not able to prove at the time:

(1) If a matrix A is invertible then it must be square.

(2) If A and B are square of the same size, then $AB = I$ implies $BA = I$.

We will prove these results in this section, using a fundamental result called the "rank-nullity theorem". The idea is that we must ascend to a higher level of abstraction, and consider the dimensions of various subspaces associated to linear functions.

[43] The matrices R_t and F_t were discussed in the previous section.

Definition (Kernel and Image). Consider a linear function $f : U \to V$ between vector spaces. We define the *kernel* and the *image* of f as follows:

$$\begin{aligned} ker(f) &:= \{\text{the set of } \mathbf{u} \in U \text{ such that } f(\mathbf{u}) = \mathbf{0}\}, \\ im(f) &:= \{\text{the set of } \mathbf{v} \in V \text{ such that } \mathbf{v} = f(\mathbf{u}) \text{ for some } \mathbf{u} \in V\}. \end{aligned}$$

We note that $ker(f) \subseteq U$ and $im(f) \subseteq V$ are subspaces. To see that $ker(f) \subseteq U$ is a subspace, consider any vectors $\mathbf{u}_i \in ker(f)$ and scalars α_i. Then from the linearity of f we have

$$f\left(\sum_i \alpha_i \mathbf{u}_i\right) = \sum_i \alpha_i f(\mathbf{u}_i) = \sum_i \alpha_i \mathbf{0} = \mathbf{0},$$

and hence $\sum_i \alpha_i \mathbf{u}_i \in ker(f)$. To see that $im(f) \subseteq V$ is a subspace, consider any vectors $\mathbf{v}_i \in im(f)$ and scalars β_i. From the definition of the image there exist some vectors $\mathbf{u}_i \in U$ satisfying $f(\mathbf{u}_i) = \mathbf{v}_i$. Then from the linearity of f we have

$$\sum_i \beta_i \mathbf{v}_i = \sum_i \beta_i f(\mathbf{u}_i) = f\left(\sum_i \beta_i \mathbf{u}_i\right),$$

which is an element of $im(f)$ because $\sum_i \beta_i \mathbf{u}_i$ is an element of U.

Questions of invertibility can be rephrased in terms of kernel and image. Recall that a function is invertible if and only if it is injective and surjective, and that the inverse of a linear function, if it exists, is necessarily linear. For a linear function $f : U \to V$ we see that

$$f \text{ is surjective} \quad \iff \quad im(f) = V.$$

Indeed, this is true by definition. The characterization of injectivity in terms of kernels actually needs a proof, so we state this as a lemma.

Lemma (Injective $\Leftrightarrow$ Trivial Kernel). Let $f : U \to V$ be a linear function. Then

$$f \text{ is injective} \quad \iff \quad ker(f) = \{\mathbf{0}\}.$$

Proof. First suppose that f is injective. To show that $ker(f) = \{\mathbf{0}\}$, consider any element $\mathbf{u} \in ker(f)$, so that $f(\mathbf{u}) = \mathbf{0}$. Since f is linear, we also have $f(\mathbf{0}) = \mathbf{0}$. Then since $f(\mathbf{u}) = f(\mathbf{0})$, the injectivity of f implies $\mathbf{u} = \mathbf{0}$. For the other direction, suppose that $ker(f) = \{\mathbf{0}\}$. To show that f is injective, suppose that $f(\mathbf{u}_1) = f(\mathbf{u}_2)$. Then we have

$$\begin{aligned} f(\mathbf{u}_1) &= f(\mathbf{u}_2) && \\ f(\mathbf{u}_1) - f(\mathbf{u}_2) &= \mathbf{0} && \\ f(\mathbf{u}_1 - \mathbf{u}_2) &= \mathbf{0} && \text{linearity of } f \\ \mathbf{u}_1 - \mathbf{u}_2 &= \mathbf{0} && ker(f) = \{\mathbf{0}\} \\ \mathbf{u}_1 &= \mathbf{u}_2. && \end{aligned}$$

Hence f is injective. $\square$

Thus we have shown for a linear function $f : U \to V$ that

$$f \text{ is invertible} \iff im(f) = V \text{ and } ker(f) = \{\mathbf{0}\}.$$

If U and V are finite-dimensional then we can rephrase this in terms of dimension. Recall that for a subspace $V' \subseteq V$ we have $V' = V$ if and only if $\dim V' = \dim V$, and for a subspace $U' \subseteq U$ we have $U' = \{\mathbf{0}\}$ if and only if $\dim U' = 0$. Thus we have

$$f \text{ is invertible} \iff \dim im(f) = \dim V \text{ and } \dim ker(f) = 0.$$

This becomes particularly useful because of the following theorem relating the dimensions of the image and kernel. The image is sometimes called the "range" and its dimension is called the "rank" of f. The kernel is sometimes called the "nullspace" and its dimension is called the "nullity" of f. Hence the name of the theorem.

Theorem (Rank-Nullity Theorem). Let $f : U \to V$ be a linear function. If U is finite-dimensional, then $im(f)$ and $ker(f)$ are finite dimensional, and we have

$$\boxed{\dim im(f) + \dim ker(f) = \dim U.}$$

Proof. Let $\dim U = n$. Since any subspace of a finite-dimensional space is also finite-dimensional, we have $\dim ker(f) = k$ for some $0 \leq k \leq n$. Let $\mathbf{u}_1, \ldots, \mathbf{u}_k$ be a basis for $ker(f)$ and extend this to a basis $\mathbf{u}_1, \ldots, \mathbf{u}_k, \mathbf{u}_{k+1}, \ldots, \mathbf{u}_n$ for U. We will show that $f(\mathbf{u}_{k+1}), \ldots, f(\mathbf{u}_n)$ is a basis for $im(f)$, and hence

$$\dim im(f) = n - k = \dim U - \dim ker(f).$$

To show that it is a spanning set, consider any $\mathbf{v} \in im(f)$. By definition of the image, we have $\mathbf{v} = f(\mathbf{u})$ for some $\mathbf{u} \in U$. Write $\mathbf{u} = \alpha_1\mathbf{u}_1 + \cdots + \alpha_n\mathbf{u}_n$. Then by linearity of f we have

$$\begin{aligned}
\mathbf{v} &= f(\mathbf{u}) \\
&= \alpha_1 f(\mathbf{u}_1) + \cdots + \alpha_n f(\mathbf{u}_n) \\
&= \alpha_1\mathbf{0} + \cdots + \alpha_k\mathbf{0} + \alpha_{k+1} f(\mathbf{u}_{k+1}) + \cdots + \alpha_n f(\mathbf{u}_n) \\
&= \alpha_{k+1} f(\mathbf{u}_{k+1}) + \cdots + \alpha_n f(\mathbf{u}_n),
\end{aligned}$$

and we conclude that $\mathbf{v}$ is in the span of $f(\mathbf{u}_{k+1}), \ldots, f(\mathbf{u}_n)$. To show that the set is independent, suppose that we have

$$\alpha_{k+1} f(\mathbf{u}_{k+1}) + \cdots + \alpha_n f(\mathbf{u}_n) = \mathbf{0}$$

for some scalars $\alpha_{k+1}, \ldots, \alpha_n$. By linearity, this implies

$$f(\alpha_{k+1}\mathbf{u}_{k+1} + \cdots + \alpha_n\mathbf{u}_n) = \alpha_{k+1} f(\mathbf{u}_{k+1}) + \cdots + \alpha_n f(\mathbf{u}_n) = \mathbf{0},$$

and hence $\alpha_{k+1}\mathbf{u}_{k+1} + \cdots + \alpha_n\mathbf{u}_n \in ker(f)$. Since $\mathbf{u}_1, \ldots, \mathbf{u}_k$ is a basis for $ker(f)$ we can write

$$\alpha_{k+1}\mathbf{u}_{k+1} + \cdots + \alpha_n\mathbf{u}_n = \beta_1\mathbf{u}_1 + \cdots + \beta_k\mathbf{u}_k$$

for some scalars $\beta_1, \ldots, \beta_k$. But then we have

$$-\beta_1\mathbf{u}_1 - \cdots - \beta_k\mathbf{u}_k + \alpha_{k+1}\mathbf{u}_{k+1} + \cdots + \alpha_n\mathbf{u}_n = \mathbf{0}.$$

Finally, the independence of the vectors $\mathbf{u}_1, \ldots, \mathbf{u}_n$ implies that all of the coefficients are zero. In particular, $\alpha_{k+1} = \cdots = \alpha_n = 0$. □

Here are some immediate, interesting consequences.[44]

Corollary. Let $f : U \to V$ be a linear function between finite-dimensional vector spaces.

- If f is injective then $\dim U \leq \dim V$.
- If f is surjective then $\dim U \geq \dim V$.
- If f is bijective then $\dim U = \dim V$.
- If $\dim U = \dim V$ then

$$f \text{ is injective} \iff f \text{ is surjective} \iff f \text{ is bijective.}$$

Proof. Suppose f is injective. From the above result this implies $ker(f) = \{\mathbf{0}\}$, and hence $\dim ker(f) = 0$. Then from rank-nullity we have $\dim U = \dim im(f)$. And since $im(f)$ is a subspace of V, we have $\dim im(f) \leq \dim V$. Hence $\dim U \leq \dim V$. Next suppose that f is surjective, so that $im(f) = V$. Then from rank-nullity we have $\dim V + \dim ker(f) = \dim U$, which implies that $\dim U \geq \dim V$. If f is bijective then it is injective and surjective. From the previous two results this implies $\dim U \leq \dim V$ and $\dim U \geq \dim V$, hence $\dim U = \dim V$. Finally, suppose that $\dim U = \dim V$. Then we have

$$\begin{aligned}
f \text{ is injective} &\iff ker(f) = \{\mathbf{0}\} \\
&\iff \dim ker(f) = 0 \\
&\iff \dim im(f) = \dim U && \text{rank-nullity} \\
&\iff \dim im(f) = \dim V && \dim U = \dim V \\
&\iff im(f) = V \\
&\iff f \text{ is surjective.}
\end{aligned}$$

□

We now have enough tools to prove the results (1) and (2) from the beginning of this section.

[44] These results are analogues of results about injective, surjective and bijective functions between finite sets, where dimension plays the role of cardinality.

Theorem (Invertible ⇒ Square). Consider a matrix $A \in \mathbb{C}^{m\times n}$, and suppose there exists an inverse matrix $A^{-1} : \mathbb{C}^{n\times m}$, satisfying $AA^{-1} = I_m$ and $A^{-1}A = I_n$. Then

$$m = n.$$

Proof. Consider the linear function $\ell_A : \mathbb{C}^n \to \mathbb{C}^m$ defined by $\ell_A(\mathbf{x}) := A\mathbf{x}$. This function is invertible, with inverse defined by $\ell_A^{-1}(\mathbf{x}) := A^{-1}\mathbf{x}$. Then since ℓ_A is a bijective linear function between $\mathbb{C}^m$ and $\mathbb{C}^n$, it follows from the previous corollary that

$$m = \dim \mathbb{C}^m = \dim \mathbb{C}^n = n.$$

□

Theorem (One-Sided Inverse ⇒ Two-Sided Inverse). Let $A, B \in \mathbb{C}^{n\times n}$ be square matrices of the same size. Then

$$AB = I_n \quad \Longleftrightarrow \quad BA = I_n.$$

Proof. Suppose that $AB = I_n$ and consider the linear function $\ell_A : \mathbb{C}^n \to \mathbb{C}^n$ defined by $\ell_A(\mathbf{x}) = A\mathbf{x}$. This function is surjective, since for any $\mathbf{y} \in \mathbb{C}^n$ we have $\mathbf{y} = I_n\mathbf{y} = (AB)\mathbf{y} = A(B\mathbf{y}) = \ell_A(B\mathbf{y})$. Since ℓ_A is a surjective linear function between vector spaces of the same dimension, we conclude that ℓ_A is also injective. We will use this property to show that $BA\mathbf{x} = \mathbf{x}$ for all vectors $\mathbf{x} \in \mathbb{C}^n$, and hence $BA = I_n$.[45] Indeed, for all $\mathbf{x} \in \mathbb{C}^n$ we have

$$\ell_A(BA\mathbf{x}) = A(BA\mathbf{x}) = (AB)A\mathbf{x} = I_nA\mathbf{x} = A\mathbf{x} = \ell_A(\mathbf{x}),$$

and hence $BA\mathbf{x} = \mathbf{x}$ by injectivity of ℓ_A. □

I think that proof was a bit too tricky, so I will give another, more conceptual proof, based on the following general principle.

Theorem (Left and Right Invertibility). Let $f : U \to V$ be a linear function. We say that f is *left-invertible* if there exists a linear function $g : V \to U$ satisfying $g \circ f = \mathrm{id}_U$, i.e., satisfying $g(f(\mathbf{u})) = \mathbf{u}$ for all $\mathbf{u}$. We say that f is *right-invertible* if there exists a linear function $h : V \to U$ satisfying $f \circ h = \mathrm{id}_V$, i.e., satisfying $f(h(\mathbf{v})) = \mathbf{v}$ for all $\mathbf{v}$. We note that

$$\begin{aligned} f \text{ is left-invertible} &\quad \Longleftrightarrow \quad f \text{ is injective,} \\ f \text{ is right-invertible} &\quad \Longleftrightarrow \quad f \text{ is surjective.} \end{aligned}$$

The theorem is true for arbitrary vector spaces, but the proof involves the axiom of choice when U and V are infinite-dimensional. Therefore we will assume for concreteness that U and V are finite dimensional, say $\dim U = m$ and $\dim V = n$. The left-to-right implications

[45] Let $\mathbf{x} = \mathbf{e}_j$ to show that the jth columns of BA and I_n are the same for all j.

are easy. The right-to-left implications depend on arbitrary choices. We will use the general principle that we can create a linear function by specifying arbitrary values on a basis.

Proof. First, assume that f has a left inverse: $g \circ f = \mathrm{id}_U$. To show that f is injective, suppose that $f(\mathbf{u}_1) = f(\mathbf{u}_2)$ for some $\mathbf{u}_1, \mathbf{u}_2 \in U$. Then applying g on the left gives

$$\begin{aligned} f(\mathbf{u}_1) &= f(\mathbf{u}_2) \\ g(f(\mathbf{u}_1)) &= g(f(\mathbf{u}_2)) \\ \mathbf{u}_1 &= \mathbf{u}_2. \end{aligned}$$

Conversely, suppose that f is injective. Our goal is to find a function $g : V \to U$ satisfying $g(f(\mathbf{u})) = \mathbf{u}$ for all $\mathbf{u}$. To construct this g we begin with a basis $\mathbf{u}_1, \ldots, \mathbf{u}_m$ for U. Consider the images $\mathbf{v}_1 := f(\mathbf{u}_1), \ldots, \mathbf{v}_m := f(\mathbf{u}_m) \in V$. The injectivity of f implies that the $\mathbf{v}_i$ are distinct. I claim that they are linearly independent. Indeed, suppose that $\sum \alpha_i \mathbf{v}_i = \mathbf{0}$ for some scalars α_i. Then by linearity and injectivity of f, we have

$$\begin{aligned} \alpha_1 \mathbf{v}_1 + \cdots + \alpha_m \mathbf{v}_m &= \mathbf{0} \\ \alpha_1 f(\mathbf{u}_1) + \cdots + \alpha_m f(\mathbf{u}_m) &= \mathbf{0} \\ f(\alpha_1 \mathbf{u}_1 + \cdots + \alpha_m \mathbf{u}_m) &= \mathbf{0} \\ f(\alpha_1 \mathbf{u}_1 + \cdots + \alpha_m \mathbf{u}_m) &= f(\mathbf{0}) \\ \alpha_1 \mathbf{u}_1 + \cdots + \alpha_m \mathbf{u}_m &= \mathbf{0}. \end{aligned}$$

Since the $\mathbf{u}_i$ are independent, this implies that $\alpha_i = 0$ for all i. We can extend the independent set $\mathbf{v}_1, \ldots, \mathbf{v}_m$ to a basis $\mathbf{v}_1, \ldots, \mathbf{v}_m, \mathbf{v}_{m+1}, \ldots, \mathbf{v}_n$ for V. Finally, we define the linear function $g : V \to U$ on this basis by setting

$$g(\mathbf{v}_i) := \begin{cases} \mathbf{u}_i & 1 \le i \le m, \\ \mathbf{0} & m < i \le n. \end{cases}$$

To verify that $g \circ f = \mathrm{id}_U$, consider any $\mathbf{u} = \sum \alpha_i \mathbf{u}_i \in U$. Then

$$\begin{aligned} g(f(\mathbf{u})) &= g(f(\alpha_1 \mathbf{u}_1 + \cdots + \alpha_m \mathbf{u}_m)) \\ &= g(\alpha_1 f(\mathbf{u}_1) + \cdots + \alpha_m f(\mathbf{u}_m)) \\ &= g(\alpha_1 \mathbf{v}_1 + \cdots + \alpha_m \mathbf{v}_m) && \text{definition of } \mathbf{v}_i \\ &= \alpha_1 g(\mathbf{v}_1) + \cdots + \alpha_m g(\mathbf{v}_m) \\ &= \alpha_1 \mathbf{u}_1 + \cdots + \alpha_m \mathbf{u}_m && \text{definition of } g \\ &= \mathbf{u}. \end{aligned}$$

For the second equivalence in the theorem, suppose that f has a right inverse: $f \circ h = \mathrm{id}_V$. By definition this means that $f(h(\mathbf{v})) = \mathbf{v}$ for any $\mathbf{v} \in V$. But this immediately implies that f is surjective, since any $\mathbf{v}$ has the form $f(\mathbf{u})$, where $\mathbf{u} = h(\mathbf{v})$. Conversely, suppose that f is surjective. Our goal is to find a linear function $h : V \to U$ satisfying $f(h(\mathbf{v})) = \mathbf{v}$ for all $\mathbf{v}$. To do this, we choose any basis $\mathbf{v}_1, \ldots, \mathbf{v}_n$ for V. Since f is surjective, there exist

vectors $\mathbf{u}_i$ satisfying $f(\mathbf{u}_i) = \mathbf{v}_i$ for all i. Define the linear function $h : V \to U$ by specifying the values $h(\mathbf{v}_i) := \mathbf{u}_i$ on the basis $\mathbf{v}_i$. To verify that $f \circ h = \mathrm{id}_V$, consider any vector $\mathbf{v} = \beta_1\mathbf{v}_1 + \cdots + \beta_n\mathbf{v}_n \in V$. Then we have

$$\begin{aligned} f(h(\mathbf{v})) &= f(h(\beta_1\mathbf{v}_1 + \cdots + \beta_n\mathbf{v}_n)) \\ &= f(\beta_1 h(\mathbf{v}_1) + \cdots + \beta_n h(\mathbf{v}_n)) \\ &= f(\beta_1\mathbf{u}_1 + \cdots + \beta_n\mathbf{u}_n) \\ &= \beta_1 f(\mathbf{u}_1) + \cdots + \beta_n f(\mathbf{u}_n) \\ &= \beta_1\mathbf{v}_1 + \cdots + \beta_n\mathbf{v}_n \\ &= \mathbf{v}. \end{aligned}$$

□

Remark: The left and right inverses constructed in this proof are **not unique**. The construction of g depends on the extension of the set $\mathbf{v}_1, \ldots, \mathbf{v}_m$ to a basis of V, and the construction of h depends on the choices of the pre-images $\mathbf{u}_1, \ldots, \mathbf{u}_n$ for the basis $\mathbf{v}_1, \ldots, \mathbf{v}_n$.[46]

Based on this theorem, we can give a more conceptual proof that $AB = I$ implies $BA = I$ for square matrices.

Another Proof (One-Sided Inverse ⇒ Two-Sided Inverse). Let A be square. Then

$$\begin{aligned} AB = I \quad &\Longrightarrow \quad A \text{ is right-invertible} \\ &\Longleftrightarrow \quad A \text{ is surjective} \\ &\Longleftrightarrow \quad A \text{ is injective} \qquad\qquad \text{rank-nullity, } A \text{ is square} \\ &\Longleftrightarrow \quad A \text{ is left-invertible} \\ &\Longleftrightarrow \quad CA = I \text{ for some } C. \end{aligned}$$

But then we have

$$C = CI = C(AB) = (CA)B = IB = B.$$

Hence $BA = I$. □

Here is a surprising corollary.

Corollary (Orthonormal Columns ⇔ Orthonormal Rows). Consider a square matrix $A \in \mathbb{C}^{n\times n}$. Then we have the following equivalences:

$$\begin{aligned} A \text{ has orthonormal columns} \quad &\Longleftrightarrow \quad A^*A = I \\ &\Longleftrightarrow \quad AA^* = I \\ &\Longleftrightarrow \quad A \text{ has orthonormal rows,} \end{aligned}$$

[46] One can show that the set of left-inverses is $m(n-m)$-dimensional (where $m \le n$ because f is injective), and the set of right-inverses is $n(m-n)$-dimensional (where $m \ge n$ because f is surjective). These sets are not vector spaces, but they are "affine subspaces" of the vector space $\mathscr{L}(V, U)$ of linear functions from V to U.

where the middle equivalence comes from the previous theorem. The other two equivalences come from Section ?.

Before moving on, I should introduce the terminology of "isomorphism".

Definition (Isomorphism of Vector Spaces). Let U and V be vector spaces. If there exists a bijective linear map $f : U \to V$ then we will say that U and V are *isomorphic*, and we will write

$$U \cong V.$$

In this case we say that f is an *isomorphism* between U and V. Isomorphism of finite-dimensional vector spaces is particulrly simple.

Theorem (Isomorphic $\Leftrightarrow$ Same Dimension). Let U and V be finite-dimensional vector spaces. Then

$$U \cong V \quad \iff \quad \dim U = \dim V.$$

Proof. If $U \cong V$ then there exists a bijective linear function $f : U \to V$, so we have $\dim U = \dim V$ from the corollary of the rank-nullity theorem. Conversely, suppose that $\dim U = \dim V = n$. In this case we need to find a bijective linear function $f : U \to V$. There are many choices. Let $\mathbf{u}_1, \ldots, \mathbf{u}_n \in U$ and $\mathbf{v}_1, \ldots, \mathbf{v}_n \in V$ be arbitrary bases for U and V, and define the linear function $f : U \to V$ by specifying the values $f(\mathbf{u}_i) := \mathbf{v}_i$ on the basis $\mathbf{u}_1, \ldots, \mathbf{u}_n$. Is it easy to check that this is a bijection. □

This result justifies our preference for Cartesian coordinates, because any n-dimensional vector space over $\mathbb{R}$ is isomorphic to $\mathbb{R}^n$. To be precise, let V be any n-dimensional vector space over $\mathbb{R}$ and let $\mathbf{v}_1, \ldots, \mathbf{v}_n \in V$ be any basis. Then we obtain an isomorphism $V \cong \mathbb{R}^n$ by sending $\mathbf{v}_1, \ldots, \mathbf{v}_n \in V$ to the standard basis vectors $\mathbf{e}_1, \ldots, \mathbf{e}_n \in \mathbb{R}^n$:

$$a_1\mathbf{v}_1 + \cdots + a_n\mathbf{v}_n \mapsto a_1\mathbf{e}_1 + \cdots + a_n\mathbf{e}_n = \begin{pmatrix} a_1 \\ \vdots \\ a_n \end{pmatrix}.$$

3.2 Subspaces Associated to Matrices

In this section we relate the results of the previous section to the rows and columns of matrices. Recall that to any matrix $A \in \mathbb{C}^{m \times n}$ we associate the linear function $\ell_A : \mathbb{C}^n \to \mathbb{C}^m$ defined by matrix multiplication: $\ell_A(\mathbf{x}) := A\mathbf{x}$. The image and kernel of ℓ_A have special names. The kernel is called the *nullspace* of A:

$$\mathcal{N}(A) := ker(\ell_A) = \{\mathbf{x} \in \mathbb{C}^n : A\mathbf{x} = \mathbf{0}\}.$$

The image is called the *column space* of A:

$$\mathcal{C}(A) := im(\ell_A) = \{A\mathbf{x} : \mathbf{x} \in \mathbb{C}^n\}.$$

The name indicates the fact that this is the subspace of $\mathbb{C}^m$ spanned by the column vectors of A. To see this, let $\mathbf{a}_1, \ldots, \mathbf{a}_n \in \mathbb{C}^m$ denote the columns of A. Then for any vector $\mathbf{x} = (x_1, \ldots, x_n) \in \mathbb{C}^n$ we recall that

$$A\mathbf{x} = \begin{pmatrix} | & & | \\ \mathbf{a}_1 & \cdots & \mathbf{a}_n \\ | & & | \end{pmatrix} \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} = x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \cdots + x_n\mathbf{a}_n.$$

Here is the matrix version of the rank-nullity theorem.

Theorem (Rank-Nullity for Matrices). Let $A \in \mathbb{C}^{m\times n}$ be an $m \times n$ matrix with column space $\mathcal{C}(A) \subseteq \mathbb{C}^m$ and nullspace $\mathcal{N}(A) \subseteq \mathbb{C}^n$. Then

$$\dim \mathcal{C}(A) + \dim \mathcal{N}(A) = n = \#\text{ columns of } A.$$

Proof: Consider the linear function $\ell_A : \mathbb{C}^n \to \mathbb{C}^m$. The rank-nullity theorem tells us that

$$\dim im(\ell_A) + \dim ker(\ell_A) = \dim \mathbb{C}^n = n.$$

□

What about the rows of A? Suppose we are working over the real numbers, so the rows of A are the columns of the transpose matrix A^T. Then we define the *row space*

$$\mathcal{R}(A) := \mathcal{C}(A^T).$$

Over the complex numbers we prefer to work with the conjugate row space

$$\mathcal{R}(A) := \mathcal{C}(A^*),$$

i.e., the span of the complex conjugates of the rows of A.[47] Here is the key observation.

Theorem (Row Space $\perp$ Nullspace). Let $A \in \mathbb{C}^{m\times n}$. We observe that the nullspace $\mathcal{N}(A)$ and the row space $\mathcal{R}(A)$ are both subspaces of $\mathbb{C}^n$. In fact, they are orthogonal complements:

$$\mathcal{N}(A) = \mathcal{R}(A)^\perp.$$

Then it follows from Section ? that

$$\dim \mathcal{R}(A) + \dim \mathcal{N}(A) = \dim \mathbb{C}^n = n = \#\text{ columns of } A.$$

Proof. We want to show that $\mathcal{N}(A) = \mathcal{C}(A^*)^\perp$. Recall that $\mathcal{N}(A) \subseteq \mathbb{C}^n$ consists of vectors $\mathbf{x} \in \mathbb{C}^n$ satisfying $A\mathbf{x} = \mathbf{0}$, and $\mathcal{C}(A^*) \subseteq \mathbb{C}^n$ consists of vectors of the form $A^*\mathbf{y}$ with $\mathbf{y} \in \mathbb{C}^m$. Furthermore, note that $\langle A\mathbf{x}, \mathbf{y}\rangle = \langle \mathbf{x}, A^*\mathbf{y}\rangle$. Thus we have

$$\mathbf{x} \in \mathcal{N}(A) \quad \Longleftrightarrow \quad A\mathbf{x} = \mathbf{0}$$

[47] The distinction is not really important because $\dim \mathcal{C}(A^T) = \dim \mathcal{C}(A^*)$, where A^T is the usual (non-conjugate) transpose applied to a complex matrix. However, we prefer **never** to refer to the non-conjugate transpose of a complex matrix.

$$\begin{aligned}
&\iff \quad \langle A\mathbf{x}, \mathbf{y}\rangle = 0 \text{ for all } \mathbf{y} \in \mathbb{C}^m \\
&\iff \quad \langle \mathbf{x}, A^*\mathbf{y}\rangle = 0 \text{ for all } \mathbf{y} \in \mathbb{C}^m \\
&\iff \quad \langle \mathbf{x}, \mathbf{z}\rangle = 0 \text{ for all } \mathbf{z} \in \mathcal{C}(A^*) \\
&\iff \quad \mathbf{x} \in \mathcal{C}(A^*)^\perp.
\end{aligned}$$

□

This is not as hard as it looks. If $\mathbf{b}_1^*, \ldots, \mathbf{b}_m^*$ are the row vectors of A then the entries of the vector $A\mathbf{x}$ can be expressed in terms of inner products:

$$A\mathbf{x} = \begin{pmatrix} - & \mathbf{b}_1^* & - \\ & \vdots & \\ - & \mathbf{b}_m^* & - \end{pmatrix} \mathbf{x} = \begin{pmatrix} \mathbf{b}_1^*\mathbf{x} \\ \vdots \\ \mathbf{b}_m^*\mathbf{x} \end{pmatrix} = \begin{pmatrix} \langle \mathbf{b}_1, \mathbf{x}\rangle \\ \vdots \\ \langle \mathbf{b}_m, \mathbf{x}\rangle \end{pmatrix}.$$

Thus we have $A\mathbf{x} = \mathbf{0}$ if and only if $\langle \mathbf{b}_i, \mathbf{x}\rangle = 0$ for all i, i.e., if and only if $\mathbf{x}$ is orthogonal to every (conjugate) row of A.

Combining the previous two equalities gives another surprising result.

Theorem (Existence of Rank). For any matrix A we have

$$\boxed{\dim \mathcal{R}(A) = \dim \mathcal{C}(A).}$$

That is, the dimension of the row space equals the dimension of the column space. We call this common dimension the *rank* of the matrix A, and we denote it by

$$\operatorname{rank}(A) := \dim \mathcal{R}(A) = \dim \mathcal{C}(A).$$

Suppose $A \in \mathbb{C}^{m\times n}$ with rank r. Since $\mathcal{C}(A)$ is a subspace of $\mathbb{C}^m$ we have $r \leq m$ and since $\mathbb{R}(A)$ is a subspace of $\mathbb{C}^n$, we have $r \leq n$. In summary:

$$r \leq \min\{m, n\}.$$

Note that $r = m$ if and only if the rows are independent, while $r = n$ if and only if the columns are independent. Indeed, if n vectors span a vector space of dimension n, then these vectors must be independent because ... I should probably state the two-out-of-three theorem somewhere.[48]

We can incorporate this theorem into our list of equivalent conditions for invertibility.

Corollary (Conditions for Matrix Invertibility). Let $A \in \mathbb{C}^{m\times n}$. Then

$$A \text{ has a right inverse} \quad \iff \quad A \text{ is surjective}$$

[48] Let $\dim V = n$ and consider a subset $S \subseteq V$. Then any two of the following three statements imply the third: (1) $\#S = n$, (2) S is spanning, (3) S is independent.

$$\begin{aligned} &\iff && \mathcal{C}(A) = \mathbb{C}^m \\ &\iff && \dim \mathcal{C}(A) = m \\ &\iff && \dim \mathcal{R}(A) = m \\ &\iff && \text{the rows of } A \text{ are independent.} \end{aligned}$$

And

$$\begin{aligned} A \text{ has a left inverse} &\iff && A \text{ is injective} \\ &\iff && \mathcal{N}(A) = \{\mathbf{0}\} \\ &\iff && \text{the columns of } A \text{ are independent} \\ &\iff && \dim \mathcal{C}(A) = n \\ &\iff && \dim \mathcal{R}(A) = n \\ &\iff && \mathcal{R}(A) = \mathbb{C}^n. \end{aligned}$$

Proof. The only part that we haven't mentioned yet is the equivalence

$$\mathcal{N}(A) = \{\mathbf{0}\} \quad \iff \quad \text{the columns of } A \text{ are independent.}$$

This is true almost by definition. Indeed, let $\mathbf{a}_1, \ldots, \mathbf{a}_n \in \mathbb{C}^m$ be the column vectors of A and let $\mathbf{x} = (x_1, \ldots, x_n) \in \mathbb{C}^n$ be an arbitrary vector. We note that

$$\mathbf{x} \in \mathcal{N}(A) \quad \iff \quad A\mathbf{x} = \mathbf{0} \quad \iff \quad x_1\mathbf{a}_1 + \cdots + x_n\mathbf{a}_n = \mathbf{0}.$$

First suppose that $\mathcal{N}(A) = \{\mathbf{0}\}$. Then we have

$$\begin{aligned} x_1\mathbf{a}_1 + \cdots + x_n\mathbf{a}_n = \mathbf{0} &\implies && \mathbf{x} \in \mathcal{N}(A) \\ &\implies && \mathbf{x} = \mathbf{0} \\ &\implies && x_1 = x_2 = \cdots = x_n = 0. \end{aligned}$$

Conversely, suppose that $\mathbf{a}_1, \ldots, \mathbf{a}_n$ are independent. Then

$$\begin{aligned} \mathbf{x} \in \mathcal{N}(A) &\implies && x_1\mathbf{a}_1 + \cdots + x_n\mathbf{a}_n = \mathbf{0} \\ &\implies && x_1 = x_2 = \cdots = x_n = 0 \\ &\implies && \mathbf{x} = \mathbf{0}. \end{aligned}$$

□

3.3 Linear Systems and Affine Subspaces

I assume you have some familiarity with linear systems of equations, which is the main topic of Linear Algebra I. In this section we will go deeper into the topic. Recall that a system of m linear equations in n unknowns has the form

$$\left\{ \begin{array}{ccccccc} a_{11}x_1 & + & \cdots & + & a_{1n}x_n & = & b_1, \\ \vdots & & & & \vdots & \vdots & \vdots \\ a_{m1}x_1 & + & \cdots & + & a_{mn}x_n & = & b_m, \end{array} \right.$$

which can be expressed as a single matrix equation:

$$\begin{pmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{m1} & \cdots & a_{mn} \end{pmatrix} \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} b_1 \\ \vdots \\ b_m \end{pmatrix}.$$

More abstractly, we just write $A\mathbf{x} = \mathbf{b}$. Given a matrix of coefficients $A \in \mathbb{R}^{m\times n}$ and a vector of constants $\mathbf{b} \in \mathbb{R}^m$, the goal is to solve for the vector of unknowns $\mathbf{x} \in \mathbb{R}^n$.

First of all, how many solutions do we expect? Note that the system $A\mathbf{x} = \mathbf{b}$ has a solution if and only if $\mathbf{b}$ is in the column space $\mathcal{C}(A)$. If a solution exists, then there may be infinitely many solutions. In fact, the set of solutions forms an "affine subspace" that is parallel to the nullspace $\mathcal{N}(A)$.

Theorem (Affine Subspace). Let V be a vector space and consider a nonempty subset $S \subseteq V$. We say that S is an *affine subspace* of V if it is closed under *affine linear combinations*:

$$\mathbf{x}_1, \ldots, \mathbf{x}_k \in S \text{ and } \alpha_1 + \cdots + \alpha_k = 1 \quad \Longrightarrow \quad \sum \alpha_i \mathbf{x}_i \in S.$$

In particular, given two points $\mathbf{x}, \mathbf{y}$ in S, the entire line $(1-t)\mathbf{x} + t\mathbf{y}$ must be in S. We define the *underlying subspace* as the set of differences

$$S_0 := \{\mathbf{x} - \mathbf{y} : \mathbf{x}, \mathbf{y} \in S\},$$

which is an actual subspace of V.[49] For any specific point $\mathbf{x}' \in S$ we can write

$$S = S_0 + \mathbf{x}' := \{\mathbf{x}_0 + \mathbf{x}' : \mathbf{x}_0 \in S_0\},$$

and we say that S is a *translate* of S_0. We define the dimension as follows:

$$\dim S := \dim S_0.$$

Remark: An intersection of affine subspaces is either an affine subspace or the empty set. For this reason, some authors regard the empty set as an affine subspace.[50]

Proof. We need to show that $S_0 \subseteq V$ is a subspace and that $S = S_0 + \mathbf{x}'$ for any $\mathbf{x}' \in S$. Since S is nonempty, we can choose some $\mathbf{x}' \in S$. Now consider any vectors $\mathbf{v}_i = \mathbf{x}_i - \mathbf{y}_i$ in S_0, with $\mathbf{x}_i, \mathbf{y}_i \in S$. Then for any scalars α_i we have

$$\begin{aligned} \sum \alpha_i \mathbf{v}_i &= \sum \alpha_i (\mathbf{x}_i - \mathbf{y}_i) \\ &= \left(\mathbf{x}' + \sum \alpha_i \mathbf{x}_i - \sum \alpha_i \mathbf{y}_i\right) - \mathbf{x}'. \end{aligned}$$

[49] Sometimes we call actual subspaces "linear subspaces", to distinguish them from "affine subspaces".

[50] And some authors say that $\dim \emptyset = -1$, but we will have no need for this convention.

Since the coefficients in the parenthesized expression sum to 1, we have

$$\mathbf{y}' := \mathbf{x}' + \sum \alpha_i \mathbf{x}_i - \sum \alpha_i \mathbf{y}_i \in S,$$

and hence $\sum \alpha_i \mathbf{v}_i = \mathbf{y}' - \mathbf{x}' \in S_0$. Thus we have shown that $S_0 \subseteq V$ is a subspace. Next we show that $S = S_0 + \mathbf{x}'$ for any $\mathbf{x}' \in S$. So fix $\mathbf{x}' \in S$. For the first direction, consider any $\mathbf{x} \in S$. Since $\mathbf{x} - \mathbf{x}' \in S_0$, we have $\mathbf{x} = (\mathbf{x} - \mathbf{x}') + \mathbf{x}' \in S_0 + \mathbf{x}'$. For the other direction, consider any $\mathbf{v} + \mathbf{x}' \in S_0 + \mathbf{x}'$. Since $\mathbf{v} \in S_0$ we can write $\mathbf{v} = \mathbf{x} - \mathbf{y}$ with $\mathbf{x}, \mathbf{y} \in S$. But then $\mathbf{v} + \mathbf{x}' = \mathbf{x} - \mathbf{y} + \mathbf{x}' \in S$ because the coefficients $1, -1, 1$ sum to 1. □

Theorem (Solutions of Linear Systems). Given a matrix $A \in \mathbb{C}^{m \times n}$ and vector $\mathbf{b}$ consider

$$S = \{\mathbf{x} : A\mathbf{x} = \mathbf{b}\}.$$

If $\mathbf{b} \notin \mathcal{C}(A)$, then S is empty. If $\mathbf{b} \in \mathcal{C}(A)$, then S is an affine subspace of $\mathbb{C}^n$ with underlying subspace equal to the nullspace $S_0 = \mathcal{N}(A)$.

Proof. We need to show that S is closed under affine combinations. Consider any solutions $\mathbf{x}_i \in S$ and scalars α_i satisfying $\sum \alpha_i = 1$. Then we have

$$A \sum \alpha_i \mathbf{x}_i = \sum \alpha_i A\mathbf{x}_i = \sum \alpha_i \mathbf{b} = \left(\sum \alpha_i\right) \mathbf{b} = 1\mathbf{b} = \mathbf{b},$$

and hence $\sum \alpha_i \mathbf{x}_i \in S$. To show that the underlying subspace is the nullspace, consider any $\mathbf{x} - \mathbf{y} \in S_0$, with $\mathbf{x}, \mathbf{y} \in S$, so that $A\mathbf{x} = \mathbf{b}$ and $A\mathbf{y} = \mathbf{b}$. Then we have

$$A(\mathbf{x} - \mathbf{y}) = A\mathbf{x} - A\mathbf{y} = \mathbf{b} - \mathbf{b} = \mathbf{0},$$

and hence $\mathbf{x} - \mathbf{y} \in \mathcal{N}(A)$. Conversely, consider any $\mathbf{x} \in \mathcal{N}(A)$. We have assumed that S is non-empty, so there exists some $\mathbf{x}'$ satisfying $A\mathbf{x}' = \mathbf{b}$. Then $A(\mathbf{x}' + \mathbf{x}) = A\mathbf{x}' + A\mathbf{x} = \mathbf{b} + \mathbf{0} = \mathbf{b}$, and hence $\mathbf{x}' + \mathbf{x} \in S$. It follows that $\mathbf{x} = (\mathbf{x}' + \mathbf{x}) - \mathbf{x}' \in S_0$. □

It follows from this that if $\mathbf{x}' \in \mathbb{C}^n$ is any one particular solution to the equation $A\mathbf{x} = \mathbf{b}$, then the general solution can be expressed as


$$S = \mathcal{N}(A) + \mathbf{x}' = \{\mathbf{x}_0 + \mathbf{x}' : A\mathbf{x}_0 = \mathbf{0}\}.$$

In more colloquial terms:[51]

$$\text{(general solution)} = \text{(one particular solution)} + \text{(general } \textit{homogeneous} \text{ solution)}.$$

Here is a picture where the nullspace is a 2-dimensional plane living in $\mathbb{R}^n$, so the general solution is also a 2-dimensional plane:

[51] The system $A\mathbf{x} = \mathbf{0}$ is sometimes called the *homogeneous system* associated to $A\mathbf{x} = \mathbf{b}$.



There is also a "dual" point of view, which is expressed in terms of hyperplanes.

Definition (Linear and Affine Hyperplanes). Let V be a finite-dimensional inner product space. A *affine hyperplane* is an affine subspace $H \subseteq V$ satisfying

$$\dim H = \dim V - 1.$$

We say that H is a *linear hyperplane* if it is also a linear subspace, i.e., if it passes through the origin. Any linear hyperplane can be expressed as

$$H = \mathbf{u}^\perp := \{\mathbf{x} \in V : \langle \mathbf{u}, \mathbf{x} \rangle = 0\}$$

for some unit vector $\mathbf{u} \in V$. Any affine hyperplane can be expressed as

$$H_{\mathbf{u},k} := \{\mathbf{x} \in V : \langle \mathbf{u}, \mathbf{x} \rangle = k\},$$

for some unit vector $\mathbf{u} \in V$ and scalar k. The linear hyperplane underlying $H_{\mathbf{u},k}$ is $H_{\mathbf{u},0} = \mathbf{u}^\perp$.

Proof. Let H be an affine hyperplane with underlying linear hyperplane H_0. Since $\dim H = \dim H_0 = \dim V - 1$, we have $\dim H_0^\perp = 1$. This means that $H_0^\perp = \text{Span}\{\mathbf{u}\}$ for some unit vector $\mathbf{u} \in V$. But then $H_0 = (H_0^\perp)^\perp = \text{Span}\{\mathbf{u}\}^\perp$. If $\mathbf{x}' \in H$ is any particular point, it follows that $H = H_0 + \mathbf{x}'$. Set $k := \langle \mathbf{u}, \mathbf{x}' \rangle$. Then for any other point $\mathbf{x} \in H$ we have $\mathbf{x} - \mathbf{x}' \in H_0$, hence

$$\begin{aligned} \langle \mathbf{u}, \mathbf{x} - \mathbf{x}' \rangle &= 0 \\ \langle \mathbf{u}, \mathbf{x} \rangle &= \langle \mathbf{u}, \mathbf{x}' \rangle \\ \langle \mathbf{u}, \mathbf{x} \rangle &= k. \end{aligned}$$

It follows that $H \subseteq H_{\mathbf{u},k}$. Conversely, suppose $\mathbf{x} \in H_{\mathbf{u},k}$, so that $\langle \mathbf{u}, \mathbf{x} \rangle = k$. It follows from the previous argument that $\langle \mathbf{u}, \mathbf{x} - \mathbf{x}' \rangle = 0$, so that $\mathbf{x} - \mathbf{x}' \in H_0$ and hence $\mathbf{x} \in H$. $\square$

We can also view an affine hyperplane as a linear equation:

$$a_1x_1 + \cdots + a_nx_n = b.$$

If $\mathbf{a} = (a_1, \ldots, a_n) \in \mathbb{R}^n$, then the set of solutions to this equation has the form

$$\{\mathbf{x} : \langle \mathbf{a}, \mathbf{x} \rangle = b\} = \{\mathbf{x} : \langle \mathbf{u}, \mathbf{x} \rangle = k\} = H_{\mathbf{u},k},$$

where $\mathbf{u} = \mathbf{a}/\|\mathbf{a}\|$ is a unit vector and $k = b/\|\mathbf{a}\|$. Thus we can view the set of solutions of a system of m linear equations in n unknowns as the intersection of m affine hyperplanes in n-dimensional space. In particular, this shows that the set of solutions is an affine subspace.[52]

3.4 Matrix Elimination

We now discuss an algorithm for solving a linear system of the form $A\mathbf{x} = \mathbf{b}$. If a solution $A\mathbf{x}' = \mathbf{b}$ exists, we recall that the general solution is

$$\{\mathbf{x} : A\mathbf{x} = \mathbf{b}\} = \{\mathbf{x} : A\mathbf{x} = \mathbf{0}\} + \mathbf{x}' = \mathcal{N}(A) + \mathbf{x}'.$$

So there are two problems to solve:

- Find a specific solution, or prove that none exists.
- Find an explicit description (e.g., a basis) for the nullspace $\mathcal{N}(A)$.

In practice, we solve these problems with the method of elimination. Suppose that the matrix E has a left inverse $E'E = I$. Then the systems $A\mathbf{x} = \mathbf{b}$ and $(EA)\mathbf{x} = (E\mathbf{b})$ have exactly the same set of solutions. Indeed, if $A\mathbf{x} = \mathbf{b}$ then multiplying both sides on the left by E gives $EA\mathbf{x} = E\mathbf{b}$. On the other hand, suppose that $EA\mathbf{x} = E\mathbf{b}$. Then multiplying both sides on the left by E' gives

$$\begin{aligned} EA\mathbf{x} &= E\mathbf{b} \\ E'EA\mathbf{x} &= E'E\mathbf{b} \\ A\mathbf{x} &= \mathbf{b}. \end{aligned}$$

The goal is to choose the matrix E so that the matrix EA becomes as simple as possible.

Theorem. Consider any matrices A, E, F so that EAF exists. Suppose that E has a left inverse, $E'E = I$, and F has a right inverse, $FF' = I$. Then

$$\begin{aligned} \mathcal{C}(AF) &= \mathcal{C}(A), \\ \mathcal{R}(EA) &= \mathcal{R}(A), \\ \dim \mathcal{C}(EA) &= \dim \mathcal{C}(A), \\ \dim \mathcal{R}(AF) &= \dim \mathcal{R}(A). \end{aligned}$$

[52] The intersection of affine subspaces is an affine subspace.

Proof. Recall that the column space $\mathcal{C}(B)$ is the set of all vectors B(some vector). To show that $\mathcal{C}(AF) \subseteq \mathcal{C}(A)$, consider any element $(AF)\mathbf{x} \in \mathcal{C}(AF)$. But then $(AF)\mathbf{x} = A(F\mathbf{x}) = A$(some vector) $\in \mathcal{C}(A)$, hence $\mathcal{C}(AF) \subseteq \mathcal{C}(A)$. Note that this argument holds for arbitrary matrices A and F. To show that $\mathcal{C}(A) \subseteq \mathcal{C}(AF)$, let F have a right inverse: $FF' = I$. Then

$$\mathcal{C}(A) = \mathcal{C}(AI) = \mathcal{C}(AFF') \subseteq \mathcal{C}(AF).$$

The last inclusion is just $\mathcal{C}(BC) \subseteq \mathcal{C}(B)$ with $B = AF$ and $C = F'$. Now recall the definition of the row space: $\mathcal{R}(A) := \mathcal{C}(A^*)$. If E has a left inverse E', then

$$\begin{aligned} E'E &= I \\ (E'E)^* &= I^* \\ E^*(E')^* &= I, \end{aligned}$$

so that E^* has a right inverse. It follows that

$$\mathcal{R}(EA) = \mathcal{C}((EA)^*) = \mathcal{C}(A^*E^*) = \mathcal{C}(A^*) = \mathcal{R}(A).$$

For the dimension statements, we note that $\mathcal{C}(EA)$ is not equal to $\mathcal{C}(A)$. Indeed, they are not even subspaces of the same space. But there is a natural linear function $\varphi : \mathcal{C}(A) \to \mathcal{C}(EA)$ defined by sending $\mathbf{y} \mapsto E\mathbf{y}$. I claim that φ is an isomorphism. First note that φ is well-defined, i.e., that $\mathbf{y} \in \mathcal{C}(A)$ implies $\varphi(\mathbf{y}) \in \mathcal{C}(EA)$. To see this, note that $\mathbf{y} \in \mathcal{C}(A)$ implies $\mathbf{y} = A\mathbf{x}$ for some $\mathbf{x}$. Then $\varphi(\mathbf{y}) = E\mathbf{y} = EA\mathbf{x} \in \mathcal{C}(EA)$. To see that φ is surjective, consider any $EA\mathbf{x} \in \mathcal{C}(EA)$ and observe that $EA\mathbf{x} = \varphi(A\mathbf{x})$. To see that φ is injective, suppose that $\varphi(\mathbf{y}_1) = \varphi(\mathbf{y}_2)$, so that $E\mathbf{y}_1 = E\mathbf{y}_2$. Then multiply both sides on the left by E' to get $\mathbf{y}_1 = \mathbf{y}_2$. We have shown that φ is an isomorphism, and hence $\dim \mathcal{C}(EA) = \dim \mathcal{C}(A)$. The corresponding statement for row spaces follows exactly as before:

$$\begin{aligned} \dim \mathcal{R}(AF) &= \dim \mathcal{C}((AF)^*) \\ &= \dim \mathcal{C}(F^*A^*) \\ &= \dim \mathcal{C}(A^*) \\ &= \dim \mathcal{R}(A). \end{aligned}$$

□

A particularly important observation is that $\mathcal{R}(EA) = \mathcal{R}(A)$ implies $\mathcal{N}(EA) = \mathcal{N}(A)$ because

$$\mathcal{N}(EA) = \mathcal{R}(EA)^\perp = \mathcal{R}(A)^\perp = \mathcal{N}(A).$$

Now we discuss how to find an appropriate left-invertible matrix E so that EA is as simple as possible. In practice, E will be a square invertible matrix, and we will construct it from certain "elementary matrices".

Definition (Elementary Matrices). We define three families of square matrices:

- For any index i and **nonzero** scalar λ we define[53]

$$D_i(\lambda) = \begin{pmatrix} 1 & & & & \\ & 1 & & & \\ & & \lambda & & \\ & & & 1 & \\ & & & & 1 \end{pmatrix} = I + (\lambda - 1)E_{ii}.$$

  The main diagonal entries are 1, except for the ii entry, which is λ. The off-diagonal entries are all zero.

- For any indices $i \neq j$ and any scalar λ we define

$$L_{ij}(\lambda) = \begin{pmatrix} 1 & & & & \\ & 1 & \cdots & \lambda & \\ & & 1 & \vdots & \\ & & & 1 & \\ & & & & 1 \end{pmatrix} = I + \lambda E_{ij}.$$

  The main diagonal entries are 1. The only other nonzero entry is λ in the ij position.

- For any indices $i \neq j$ we define

$$T_{ij} = \begin{pmatrix} 1 & & & & \\ & 0 & \cdots & 1 & \\ & \vdots & 1 & \vdots & \\ & 1 & \cdots & 0 & \\ & & & & 1 \end{pmatrix} = I - E_{ii} - E_{jj} + E_{ij} + E_{ji}.$$

  The main diagonal entries are 1 except for zeros in the ii and jj positions. The off-diagonal entries are zero except for 1 in the ij and ji positions.

The purpose of elementary matrices is to perform elementary row and column operations.

Lemma (Elementary Row and Column Operations).

- The function $A \rightsquigarrow D_i(\lambda)A$ scales the ith row of A by λ.
- The function $A \rightsquigarrow T_{ij}A$ swaps the ith and jth rows of A.
- The function $A \rightsquigarrow L_{ij}(\lambda)A$ adds $\lambda(j$-th row) to the ith row of A.
- The function $A \rightsquigarrow AD_i(\lambda)$ scales the ith column of A by λ.
- The function $A \rightsquigarrow AT_{ij}$ swaps the ith and jth column of A.

[53] Recall that E_{ij} is the matrix with a 1 in the i, j position and zeroes elsewhere.

- The function $A \rightsquigarrow AL_{ij}(\lambda)$ adds $\lambda(i\text{-th column})$ to the jth column of A.

Based on these interpretations, it is easy to see that elementary matrices are invertible, and their inverses are also elementary matrices:

$$D_i(\lambda)^{-1} = D_i(1/\lambda)$$
$$L_{ij}(\lambda)^{-1} = L_{ij}(-\lambda)$$
$$T_{ij}^{-1} = T_{ij}.$$

Theorem (Reduced Row-Echelon Form). We say that two matrices $A, B \in \mathbb{C}^{m\times n}$ are *row-equivalent*[54] if there exists a sequence of elementary matrices $E_1, \ldots, E_k$ such that

$$B = E_k \cdots E_2 E_1 A.$$

We say that a matrix R is in *reduced row-echelon form* (RREF) if

- Every nonzero row has leftmost nonzero entry equal to 1. This entry is called a *pivot*.
- The pivots follow a staircase ("echelon") shape. That is, if $i_1 < i_2$ then the pivot in row i_2 occurs to the right of the pivot in row i_1.
- The zero rows (i.e., the non-pivot rows) occur at the bottom of the matrix.
- Each pivot is the only nonzero entry in its column.[55]

A matrix in RREF looks like this, where the empty space below the staircase represents zeros, and the asterisks denote arbitrary scalars:

$$\begin{pmatrix} 1 & * & * & 0 & * & 0 & * \\ & & & 1 & * & 0 & * \\ & & & & & 1 & * \\ & & & & & & \end{pmatrix}$$

The theorem says that every matrix A is row-equivalent to a **unique** matrix R in reduced row-echelon form, and we write

$$R = \mathrm{RREF}(A).$$

The details of the proof are a bit annoying, so we proceed in casual fashion.

Casual Proof of Existence. The proof of existence is an algorithm, called *Gaussian elimination*. The idea is to multiply A on the left by a sequence of elementary matrices $E_1, E_2, \ldots,$ corresponding to elementary row operations, until the RREF is reached. Here is a casual description of the algorithm:

[54] Later we will see that this definition of row-equivalence is equivalent to the statement that $\mathcal{R}(A) = \mathcal{R}(B)$, which is a more natural condition.

[55] This condition is the "reduced" part of "reduced row-echelon form".

(1) Find the leftmost nonzero column. Apply T_{ij} to place a nonzero entry at the top of this column. Apply $D_i(\lambda)$ to turn this entry into a 1, which becomes a new pivot. Apply $L_{ij}(\lambda)$ to turn all entries below the pivot into zeroes.

(2) Repeat step (1) on the submatrix below and to the right of the previous pivot. This process eventually stops, leaving the original matrix in "row-echelon form".

(3) To reach the "reduced row-echelon form", apply $L_{ij}(\lambda)$ to turn all entries above the pivots into zeroes.

The proof of uniqueness is based on an important observation, which we isolate as a lemma.

Lemma (Row Operations Preserve Column Relations). Let $A \in \mathbb{C}^{m\times n}$ have column vectors $\mathbf{a}_1, \ldots, \mathbf{a}_n \in \mathbb{C}^m$. We can think of elements of the nullspace $\mathbf{x} = (x_1, \ldots, x_n) \in \mathcal{N}(A)$ as *linear relations* among the columns, because

$$A\mathbf{x} = \mathbf{0} \quad \Longleftrightarrow \quad x_1\mathbf{a}_1 + \cdots + x_n\mathbf{a}_n = \mathbf{0}.$$

If E is an elementary matrix (hence invertible), we showed previously that $\mathcal{N}(EA) = \mathcal{N}(A)$. This implies that the columns of EA (which are $E\mathbf{a}_1, \ldots, E\mathbf{a}_n$) satisfy exactly the same linear relations as the columns of A.

Casual Proof of Uniqueness. Suppose A is row equivalent to matrices R and S, both in RREF. By the previous lemma, the matrices R and S satisfy exactly the same column relations. For a matrix in RREF, the pivot columns are precisely those columns that are not linear combinations of previous columns. Hence R and S have the same pivot columns. Finally, each non-pivot column is a unique linear combination of pivot columns, and its entries are the coefficients of this linear combination. But this linear combination is the same in R and S, so R and S also have the same non-pivot columns. □

Remark: We could have used the the RREF theorem as an independent path to the rank-nullity theorem and related dimension theorems. Let $R = \mathrm{RREF}(A)$. Since $R = EA$ for some invertible matrix E (the product of the elementary matrices corresponding to the steps in Gaussian elimination), we have $\mathcal{R}(A) = \mathcal{R}(EA) = \mathcal{R}(R)$. Since the pivot rows of R are clearly a basis for the row space $\mathcal{R}(R)$, we obtain

$$\dim \mathcal{R}(A) = \#\text{ pivot rows in } \mathrm{RREF}(A).$$

On the other hand, we also know that $\dim \mathcal{C}(A) = \dim \mathcal{C}(EA) = \dim \mathcal{C}(R)$.[56] Since the pivot columns of R are clearly a basis for the column space $\mathcal{C}(R)$, we have

$$\dim \mathcal{C}(A) = \#\text{ pivot columns in } \mathrm{RREF}(A).$$

But there is a bijection between pivot rows and pivot columns provided by the pivots, hence

$$\dim \mathcal{R}(A) = \#\text{ pivot rows in } \mathrm{RREF}(A)$$

[56] We showed in a previous lemma that the vector spaces $\mathcal{C}(A)$ and $\mathcal{C}(EA)$ are isomorphic.

$$
\begin{aligned}
&= \#\text{ pivots in RREF}(A) \\
&= \#\text{ pivot columns in RREF}(A) \\
&= \dim \mathcal{C}(A).
\end{aligned}
$$

This is the path pursued in certain elementary textbooks, for example in Gilbert Strang's *Introduction to Linear Algebra*.

Now we discuss applications of the RREF to solving linear systems. As a first remark, consider any block matrix of the form $(A|B)$. Then block multiplication tells us that

$$E(A|B) = (EA|EB).$$

If E is an elementary matrix then this corresponds to the fact that performing a row operation on the matrix $(A|B)$ performs the same row operation on the blocks A and B simultaneously. Because Gaussian elimination works from left to right, it is not difficult to check that

$$\text{RREF}(A|B) = (\text{RREF}(A)|B'),$$

for some matrix B'. There is a nice application of this to the computation of inverse matrices.

Theorem (Computation of Matrix Inverses). Let $A \in \mathbb{C}^{n\times n}$ be a square matrix. Then

$$A^{-1} \text{ exists} \quad \iff \quad \text{RREF}(A) = I,$$

in which case we have

$$\text{RREF}(A|I) = (I|A^{-1}).$$

Proof. Let $R = \text{RREF}(A)$. Since $R = EA$ for some invertible matrix E, we know that $\mathcal{R}(R) = \mathcal{R}(A)$. Furthermore, we know that $\dim \mathcal{R}(A)$ is the number of pivots in R. Now recall that A^{-1} exists if and only if $\dim \mathcal{R}(A) = n$. But the only $n \times n$ matrix in RREF with n pivots is the identity matrix. Thus the equivalence is established. Now consider the augmented matrix $(A|I) \in \mathbb{C}^{n\times 2n}$. If A^{-1} exists then $\text{RREF}(A) = I$ and we have

$$\text{RREF}(A|I) = (\text{RREF}(A)|B) = (I|B),$$

for some matrix $B \in \mathbb{C}^{n\times n}$. The RREF is obtained by multiplying on the left by an invertible matrix E (which is the product of the elementary matrices corresponding to the steps of the algorithm), hence we have

$$E(A|I) = \text{RREF}(A|I) = (I|B).$$

Since $E(A|I) = (EA|E)$, this implies that $EA = I$ and $E = B$, hence $B = A^{-1}$. □

For example, let's compute the inverse of the matrix

$$A = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & 0 \\ 1 & 0 & 0 \end{pmatrix}.$$

Here are the steps of Gaussian elimination, with the corresponding elementary matrices labeled:

$$\left(\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 1 & 2 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 1 \end{array}\right)$$

$$\xrightarrow{L_{21}(-1)} \left(\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 1 \end{array}\right)$$

$$\xrightarrow{L_{31}(-1)} \left(\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & -1 & -1 & -1 & 0 & 1 \end{array}\right)$$

$$\xrightarrow{L_{32}(1)} \left(\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & -2 & -2 & 1 & 1 \end{array}\right)$$

$$\xrightarrow{D_3(-1/2)} \left(\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & 1 & -1/2 & -1/2 \end{array}\right)$$

$$\xrightarrow{L_{23}(1)} \left(\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1/2 & -1/2 \\ 0 & 0 & 1 & 1 & -1/2 & -1/2 \end{array}\right)$$

$$\xrightarrow{L_{13}(-1)} \left(\begin{array}{ccc|ccc} 1 & 1 & 0 & 0 & 1/2 & 1/2 \\ 0 & 1 & 0 & 0 & 1/2 & -1/2 \\ 0 & 0 & 1 & 1 & -1/2 & -1/2 \end{array}\right)$$

$$\xrightarrow{L_{12}(-1)} \left(\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 1/2 & -1/2 \\ 0 & 0 & 1 & 1 & -1/2 & -1/2 \end{array}\right).$$

It follows that

$$\begin{pmatrix} 1 & 1 & 1 \\ 1 & 2 & 0 \\ 1 & 0 & 0 \end{pmatrix}^{-1} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1/2 & -1/2 \\ 1 & -1/2 & -1/2 \end{pmatrix}$$

Here is an interesting corollary.

Corollary. If A^{-1} exists then A can be expressed as a product of elementary matrices. Indeed, if A^{-1} exists then we know that $\mathrm{RREF}(A) = I$. If the steps of the algorithm correspond to the elementary matrices $E_1, E_2, \ldots, E_k$, then this means that

$$E_k \cdots E_2 E_1 A = I$$
$$A = E_1^{-1} E_2^{-1} \cdots E_k^{-1}.$$

Since the inverse of an elementary matrix is also elementary, we conclude that A is a product of elementary matrices. In the previous example, we have

$$A = L_{21}(-1)^{-1} L_{31}(-1)^{-1} L_{32}(1)^{-1} D_3(-1/2)^{-1} L_{23}(1)^{-1} L_{13}(-1)^{-1} L_{12}(-1)^{-1}$$

$$= L_{21}(1)L_{31}(1)L_{32}(-1)D_3(-2)L_{23}(-1)L_{13}(1)L_{12}(1).$$

If A^{-1} does not exist, then A cannot be a product of elementary matrices, because elementary matrices are invertible.

What happens if we try to invert a non-invertible matrix? Consider the matrix

$$A = \begin{pmatrix} 1 & 3 & 8 \\ 1 & 2 & 6 \\ 0 & 1 & 2 \end{pmatrix}.$$

The RREF of the augmented matrix is

$$\left(\begin{array}{ccc|ccc} 1 & 3 & 8 & 1 & 0 & 0 \\ 1 & 2 & 6 & 0 & 1 & 0 \\ 0 & 1 & 2 & 0 & 0 & 1 \end{array}\right) \xrightarrow{\text{RREF}} \left(\begin{array}{ccc|ccc} 1 & 0 & 2 & 0 & 1 & -2 \\ 0 & 1 & 2 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & -1 & -1 \end{array}\right).$$

This computation implies that

$$\text{RREF}(A) = \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix},$$

which implies that the inverse A^{-1} does not exist.

Next, we show how to use the RREF to solve a linear system $A\mathbf{x} = \mathbf{b}$ with $A \in \mathbb{C}^{m\times n}$ and $\mathbf{b} \in \mathbb{C}^m$. Recall that the general solution has the form $\mathcal{N}(A)+\mathbf{x}'$, where $\mathbf{x}'$ is any one particular solution. First we show how to compute the nullspace using the RREF. Since $\text{RREF}(A) = EA$ for some invertible matrix (the product of the elementary matrices corresponding to the steps of the algorithm), we know that

$$\mathcal{N}(A) = \mathcal{N}(\text{RREF}(A)).$$

It is easy to read off the nullspace of a matrix in reduced row-echelon form. Consider our previous example:

$$A = \begin{pmatrix} 1 & 3 & 8 \\ 1 & 2 & 6 \\ 0 & 1 & 2 \end{pmatrix} \quad \text{and} \quad \text{RREF}(A) = \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix}.$$

The nullspace consists of vectors $\mathbf{x} = (x, y, z)$ satisfying

$$\begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \implies \begin{cases} x + 0 + 2z = 0, \\ 0 + y + 2z = 0, \\ 0 + 0 + 0 = 0. \end{cases}$$

In general, we know that the dimension of the nullspace is equal to the number of non-pivot columns. We let the variables corresponding to these columns be *free variables*. In this example, we let $t := z$ be a free variable. Then the solution is

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -2z \\ -2z \\ z \end{pmatrix} = t \begin{pmatrix} -2 \\ -2 \\ 1 \end{pmatrix}.$$

In other words, the nullspace is the line generated by $(-2, -2, 1)$. Now consider the non-homogeneous system

$$\left\{ \begin{array}{rcrcrcl} x & + & 3y & + & 8z & = & 2, \\ x & + & 2y & + & 6z & = & 1, \\ 0 & + & y & + & 2z & = & 1, \end{array} \right.$$

which can be expressed in matrix notation as

$$\begin{pmatrix} 1 & 3 & 8 \\ 1 & 2 & 6 \\ 0 & 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \\ 1 \end{pmatrix},$$

$$A\mathbf{x} = \mathbf{b}.$$

If a solution $A\mathbf{x}' = \mathbf{b}$ exists, then the general solution will be

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} + t \begin{pmatrix} -2 \\ -2 \\ 1 \end{pmatrix}.$$

We can use the RREF to find a solution $\mathbf{x}'$, or prove that none exists. To do this, we form the augmented matrix $(A|\mathbf{b})$ and then compute the RREF, which has the form

$$\mathrm{RREF}(A|\mathbf{b}) = (\mathrm{RREF}(A)|\mathbf{b}'),$$

for some vector $\mathbf{b}'$. In our case, we have

$$\left(\begin{array}{ccc|c} 1 & 3 & 8 & 2 \\ 1 & 2 & 6 & 1 \\ 0 & 1 & 2 & 1 \end{array} \right) \xrightarrow{\text{RREF}} \left(\begin{array}{ccc|c} 1 & 0 & 2 & -1 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & 0 & 0 \end{array} \right).$$

Let E be the matrix that converts $(A|\mathbf{b})$ into RREF, so that $\mathrm{RREF}(A|\mathbf{b}) = E(A|\mathbf{b}) = (EA|E\mathbf{b})$. Since E is invertible, we recall that the systems $A\mathbf{x} = \mathbf{b}$ and $EA\mathbf{x} = E\mathbf{b}$ have the same set of solutions. In our example,

$$\left\{ \begin{array}{rcrcrcl} x & + & 3y & + & 8z & = & 2 \\ x & + & 2y & + & 6z & = & 1 \\ 0 & + & y & + & 2z & = & 1 \end{array} \right\} \iff \left\{ \begin{array}{rcrcrcl} x & + & 0 & + & 2z & = & -1 \\ 0 & + & y & + & 2z & = & 1 \\ 0 & + & 0 & + & 0 & = & 0 \end{array} \right\}.$$

By setting $z = 0$ in the right hand system, we observe that $\mathbf{x}' = (-1, 1, 0)$ is a solution, hence the general solution is

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -1 \\ 1 \\ 0 \end{pmatrix} + t \begin{pmatrix} -2 \\ -2 \\ 1 \end{pmatrix}.$$

Changing the right hand side to $\mathbf{b} = (1, 0, 0)$ gives

$$\left(\begin{array}{ccc|c} 1 & 3 & 8 & 1 \\ 1 & 2 & 6 & 0 \\ 0 & 1 & 2 & 0 \end{array}\right) \xrightarrow{\text{RREF}} \left(\begin{array}{ccc|c} 1 & 0 & 2 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 1 \end{array}\right),$$

which leads to the equivalence

$$\left\{\begin{array}{rcrcrcl} x & + & 3y & + & 8z & = & 1 \\ x & + & 2y & + & 6z & = & 0 \\ 0 & + & y & + & 2z & = & 0 \end{array}\right\} \iff \left\{\begin{array}{rcrcrcl} x & + & 0 & + & 2z & = & 0 \\ 0 & + & y & + & 2z & = & 0 \\ 0 & + & 0 & + & 0 & = & 1 \end{array}\right\}.$$

The presence of the equation "$0 = 1$" indicates that this system has no solution. In general, the system $A\mathbf{x} = \mathbf{b}$ has a solution if and only if the final column of $\text{RREF}(A|\mathbf{b})$ is not a pivot column. We can use this as a test for "subspace membership":

$$\begin{aligned} \mathbf{b} \in \mathcal{C}(A) &\iff A\mathbf{x} = \mathbf{b} \text{ has a solution} \\ &\iff \text{the last column of } \text{RREF}(A|\mathbf{b}) \text{ is not a pivot column.} \end{aligned}$$

3.5 Least Squares Approximation

In Section 1.4 we discussed a continuous version of "least squares approximation". Now we discuss the discrete version, and its relation to matrix algebra.

Definition (Least Squares Solutions). Consider a matrix $A \in \mathbb{C}^{m\times n}$ and a vector $\mathbf{b} \in \mathbb{C}^m$. If $\mathbf{b}$ is not in the column space of A then the system $A\mathbf{x} = \mathbf{b}$ has no solution $\mathbf{x} \in \mathbb{C}^n$. In this case we will settle for an approximate solution. A vector $\mathbf{x} \in \mathbb{C}^n$ is called a *least squares solution* of $A\mathbf{x} = \mathbf{b}$ if the length of $A\mathbf{x} - \mathbf{b}$ is as small as possible, i.e., if

$$\|A\mathbf{x} - \mathbf{b}\| \leq \|A\mathbf{y} - \mathbf{b}\| \text{ for all } \mathbf{y} \in \mathbb{C}^n.$$

We already have the tools to deal with this.

Theorem (Least Squares From Orthogonal Projection). Consider a matrix $A \in \mathbb{C}^{m\times n}$ and a vector $\mathbf{b} \in \mathbb{C}^m$. Let $P_{\mathcal{C}(A)}\mathbf{b}$ be the orthogonal projection of $\mathbf{b}$ onto the column space $\mathcal{C}(A)$. Then the least squares solutions of $A\mathbf{x} = \mathbf{b}$ are the exact solutions of the equation

$$A\mathbf{x} = P_{\mathcal{C}(A)}\mathbf{b}.$$

A solution $\mathbf{x}$ always exists because $P_{\mathcal{C}(A)}\mathbf{b}$ is in $\mathcal{C}(A)$, but the solution might not be unique.

Proof. Recall from Section 1.4 that the orthogonal projection of a point $\mathbf{b} \in \mathbb{C}^m$ onto a subspace $U \subseteq \mathbb{C}^m$ is the **unique** point $\mathbf{p} \in U$ satisfying

$$\|\mathbf{p} - \mathbf{b}\| \leq \|\mathbf{q} - \mathbf{b}\| \text{ for all } \mathbf{q} \in U.$$

If $U = \mathcal{C}(A) = \{A\mathbf{x} : \mathbf{x} \in \mathbb{C}^n\}$ is the column space of A, then we can write $\mathbf{p} = A\mathbf{x}$ and $\mathbf{q} = A\mathbf{y}$ for some $\mathbf{x}, \mathbf{y} \in \mathbb{C}^n$ (possibly not unique). Thus the inequality becomes

$$\|A\mathbf{x} - \mathbf{b}\| \leq \|A\mathbf{y} - \mathbf{b}\| \text{ for all } \mathbf{y} \in \mathbb{C}^n.$$

In other words, $\mathbf{x}$ is a least squares solution if and only if $A\mathbf{x} = \mathbf{p}$. □

We remark that the least squares solution of $A\mathbf{x} = \mathbf{b}$ is unique if and only if A has independent columns. Indeed, recall that A has independent columns if and only A has some left inverse, say $BA = I$. If $\mathbf{x}, \mathbf{x}' \in \mathbb{C}^n$ are two least squares solutions, so that

$$A\mathbf{x} = P_{\mathcal{C}(A)}\mathbf{b} = A\mathbf{x}',$$

then multiplying by B on the left gives $\mathbf{x} = \mathbf{x}'$. Conversely, if A has dependent columns, then there exists a nonzero vector $\mathbf{v} \in \mathbb{C}^n$ satisfying $A\mathbf{v} = \mathbf{0}$. If $\mathbf{x}$ is a least-square solution, then so is $\mathbf{x} + \mathbf{v}$:

$$A\mathbf{x} = P_{\mathcal{C}(A)}\mathbf{b} \quad \Longrightarrow \quad A(\mathbf{x} + \mathbf{v}) = A\mathbf{x} + A\mathbf{v} = A\mathbf{x} + \mathbf{0} = P_{\mathcal{C}(A)}\mathbf{b}.$$

Hence the least squares solution is not unique.

Now we discuss explicit formulas and methods of computation. The following lemma is crucial.

Lemma (The Matrices A^*A and AA^*). For any matrix $A \in \mathbb{C}^{m\times n}$, we have

$$\operatorname{rank}(A^*A) = \operatorname{rank}(A) = \operatorname{rank}(A^*) = \operatorname{rank}(AA^*).$$

Proof. First we show that[57]

$$\mathcal{N}(A^*A) = \mathcal{N}(A).$$

One direction is easy: If $A\mathbf{x} = \mathbf{0}$ then $(A^*A)\mathbf{x} = A^*(A\mathbf{x}) = A^*\mathbf{0} = \mathbf{0}$, hence $\mathcal{N}(A) \subseteq \mathcal{N}(A^*A)$. For the other direction, suppose that $A^*A\mathbf{x} = \mathbf{0}$. Then we have

$$\|A\mathbf{x}\|^2 = (A\mathbf{x})^*(A\mathbf{x}) = \mathbf{x}^*(A^*A\mathbf{x}) = \mathbf{x}^*\mathbf{0} = 0,$$

which implies $A\mathbf{x} = \mathbf{0}$. Hence $\mathcal{N}(A^*A) \subseteq \mathcal{N}(A)$. For any matrix, we recall from the rank-nullity theorem that

$$\dim \mathcal{N}(B) + \operatorname{rank}(B) = \#\text{ columns of } B.$$

Since A^*A and A have the same number of columns, and the same nullspace, it follows that they have the same rank. Thus we have shown that $\operatorname{rank}(A^*A) = \operatorname{rank}(A)$. Then by replacing A with A^* we obtain $\operatorname{rank}(A^*) = \operatorname{rank}(A^{**}A^*) = \operatorname{rank}(AA^*)$. The middle equality comes from the fact that $\mathcal{N}(A^*) = \mathcal{C}(A)^\perp$.[58] Then rank-nullity gives

$$\operatorname{rank}(A^*) = m - \dim \mathcal{N}(A^*)$$

[57] We have already seen that $\mathcal{N}(EA) = \mathcal{N}(A)$ when E has a left inverse, but A^* need not have a left inverse.
[58] This is equivalent to the statement $\mathcal{N}(A) = \mathcal{R}(A)^\perp$ by replacing A with A^*.

$$
\begin{aligned}
&= m - \dim \mathcal{C}(A)^{\perp} \\
&= m - (m - \dim \mathcal{C}(A)) \\
&= \dim \mathcal{C}(A) \\
&= \operatorname{rank}(A).
\end{aligned}
$$

□

For any matrix $A \in \mathbb{C}^{m\times n}$, the matrices $A^*A \in \mathbb{C}^{n\times n}$ and $AA^* \in \mathbb{C}^{m\times m}$ are always square, so they are potentially invertible. It follows from the previous lemma that

$$
\begin{array}{rcl}
(A^*A)^{-1} \text{ exists} & \iff & \operatorname{rank}(A^*A) = n \\
& \iff & \operatorname{rank}(A) = n \\
& \iff & A \text{ has independent columns} \\
& \iff & A \text{ has a left inverse} \\
& & \text{etc.}
\end{array}
$$

In fact, if $(A^*A)^{-1}$ exists, then the matrix $(A^*A)^{-1}A^*$ itself is a left inverse of A:

$$
[(A^*A)^{-1}A^*]A = (A^*A)^{-1}(A^*A) = I.
$$

Similarly, we have

$$
\begin{array}{rcl}
(AA^*)^{-1} \text{ exists} & \iff & \operatorname{rank}(AA^*) = m \\
& \iff & \operatorname{rank}(A^*) = m \\
& \iff & A \text{ has independent rows} \\
& \iff & A \text{ has a right inverse} \\
& & \text{etc.},
\end{array}
$$

in which case $A^*(AA^*)^{-1}$ is one particular right inverse of A.


Theorem (Least Squares with Independent Columns). Let $\mathbf{a}_1, \ldots, \mathbf{a}_n \in \mathbb{C}^m$ be the columns of a matrix $A \in \mathbb{C}^{m\times n}$. If the columns are independent, then the matrix of the projection onto the column space is

$$
P_{\mathcal{C}(A)} = A(A^*A)^{-1}A^*,
$$

and for any $\mathbf{b} \in \mathbb{C}^m$, the unique least squares solution to the system $A\mathbf{x} = \mathbf{b}$ is

$$
\mathbf{x} = (A^*A)^{-1}A^*\mathbf{b}.
$$

Proof. The projection $P_{\mathcal{C}(A)}\mathbf{b}$ has the form $A\mathbf{x}$ because it is in $\mathcal{C}(A)$. Geometrically, the vector $A\mathbf{x} - \mathbf{b}$ is in $\mathcal{C}(A)^{\perp}$, which means that $A\mathbf{x} - \mathbf{b}$ is orthogonal to every column of A, as in the following picture:



We can express this situation with the single matrix equation $A^*(A\mathbf{x}-\mathbf{b})=\mathbf{0}$. Then we solve this to obtain

$$\begin{aligned} A^*(A\mathbf{x}-\mathbf{b}) &= \mathbf{0} \\ A^*A\mathbf{x}-A^*\mathbf{b} &= \mathbf{0} \\ A^*A\mathbf{x} &= A^*\mathbf{b}. \end{aligned}$$

So far, this holds for any matrix A. If A has independent columns, then $(A^*A)^{-1}$ exists and we may continue as follows:

$$\begin{aligned} A^*A\mathbf{x} &= A^*\mathbf{b} \\ \mathbf{x} &= (A^*A)^{-1}A^*\mathbf{b}. \end{aligned}$$

This gives the unique least squares solution. To obtain the projection matrix we multiply on the left by A:

$$\begin{aligned} \mathbf{x} &= (A^*A)^{-1}A^*\mathbf{b} \\ A\mathbf{x} &= A(A^*A)^{-1}A^*\mathbf{b} \\ P_{\mathcal{C}(A)}\mathbf{b} &= A(A^*A)^{-1}A^*\mathbf{b}. \end{aligned}$$


Since this holds for any $\mathbf{b}\in\mathbb{C}^m$, we conclude that $P_{\mathcal{C}(A)}=A(A^*A)^{-1}A^*$. □

The classical application of least squares is to curve fitting. Indeed, this is the purpose for which Gauss invented the method.[59] For example, suppose that we have a collection of n data points in the x,y-plane:


$$(x_1,y_1),(x_2,y_2),\ldots,(x_n,y_n).$$

We would like to find the line of the form $y=a+bx$ that is the "best fit" for these points:


[59] He used it to fit the elliptical orbit of the dwarf planet Ceres to a collection of observed data points.



There are different ways one might interpret the word "best". The most obvious definition might be to minimize the orthogonal distances[60] from the points to the line:



This idea is called *total least squares*, or *orthogonal least squares*. It is a hard non-linear problem, which we will solve after discussing the singular value decomposition. In statistics this problem is called *principal component analysis*. It is much easier to minimize the sum of squares of the vertical distances:



This problem is called *ordinary least squares*, or just least squares regression. Here's how we solve it. We start by being optimistic and assuming that all of the data points fit perfectly on

[60] Typically we want to minimize the sum of squared distances.

the line, which leads to a system of n linear equations in the two unknowns a and b:

$$\begin{cases} a + bx_1 = y_1, \\ a + bx_2 = y_2, \\ \qquad\vdots \\ a + bx_n = y_n, \end{cases}$$

It is an unfortunate feature of curve fitting problems that the roles of variables and constants get switched around, so instead of a system looking like $A\mathbf{x} = \mathbf{b}$ we get a system looking like $X\mathbf{a} = \mathbf{y}$. In our case we have

$$X\mathbf{a} = \mathbf{y}$$
$$\begin{pmatrix} 1 & x_1 \\ \vdots & \vdots \\ 1 & x_n \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} y_1 \\ \vdots \\ y_n \end{pmatrix}.$$

This system almost certainly does not have a solution, since any three or more points almost certainly do not fit perfectly on a straight line. Hence we will apply the method of least squares. If the data points do not all have the same x value, then the two columns of X are are independent and we get a unique solution:[61]


$$\begin{aligned} X\mathbf{a} &= \mathbf{y} \\ X^T X\mathbf{a} &= X^T\mathbf{y} \\ \mathbf{a} &= (X^T X)^{-1} X\mathbf{y}. \end{aligned}$$

Recall that this "least squares solution" minimizes the length $\|X\mathbf{a}-\mathbf{y}\|$, hence it also minimizes the squared length $\|X\mathbf{a} - \mathbf{y}\|^2$. In terms of the data points, this becomes

$$\|X\mathbf{a} - \mathbf{y}\|^2 = \left\| \begin{pmatrix} a + bx_i - y_i \\ \vdots \\ a + bx_n - y_n \end{pmatrix} \right\|^2 = \sum (a + bx_i - y_i)^2,$$

which is, indeed, the sum of the squared vertical errors:

[61] Here we are dealing with real numbers, so we use the transpose notation X^T instead of the adjoint notation X^*.



To be explicit, the normal equation has the following form, which you might recognize:

$$X^T X\mathbf{a} = X^T\mathbf{y}$$

$$\begin{pmatrix} 1 & \cdots & 1 \\ x_1 & \cdots & x_n \end{pmatrix} \begin{pmatrix} 1 & x_1 \\ \vdots & \vdots \\ 1 & x_n \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 1 & \cdots & 1 \\ x_1 & \cdots & x_n \end{pmatrix} \begin{pmatrix} y_1 \\ \vdots \\ y_n \end{pmatrix}$$

$$\begin{pmatrix} n & \sum x_i \\ \sum x_i & \sum x_i^2 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} \sum y_i \\ \sum x_i y_i \end{pmatrix},$$

which is equivalent to the linear system

$$\begin{cases} an & + & b\sum x_i & = & \sum y_i, \\ a\sum x_i & + & b\sum x_i^2 & = & \sum x_i y_i. \end{cases}$$

This is the form usually presented in introductory statistics courses, when the students don't know linear algebra.

However, the linear algebra formulation is much more powerful because it generalizes easily. For example, we can fit our data to polynomial curve of degree d:

$$y = a_0 + a_1 x + \cdots + \cdots + a_d x^d.$$

Assuming optimistically that all n data points lie on this curve gives a system of n linear equations in the $d+1$ unknown coefficients $a_0, \ldots, a_d$:

$$X\mathbf{a} = \mathbf{y}$$

$$\begin{pmatrix} 1 & x_1 & x_1^2 & \cdots & x_1^d \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & x_n & x_n^2 & \cdots & x_n^d \end{pmatrix} \begin{pmatrix} a_0 \\ \vdots \\ a_d \end{pmatrix} = \begin{pmatrix} y_1 \\ \vdots \\ y_n \end{pmatrix}.$$

Then the least squares solution (which minimizes the sum of squares of the vertical errors) is given by the normal equation $X^T X\mathbf{a} = X^T\mathbf{y}$. This equation is much harder to obtain using calculus, and the explicit formulas for the entries of the matrix $X^T X$ are not so nice.[62]

For completeness, we discuss the details of least squares solutions to $A\mathbf{x} = \mathbf{b}$ when the matrix A does not have independent columns, and hence the least squares solution is not unique. Nevertheless, there is always a unique least squares solution with minimum length. The amazing fact is that this minimum length solution is determined by a matrix.

Theorem (Existence of the Pseudoinverse). Consider any matrix $A \in \mathbb{C}^{m\times n}$. There exists a matrix $A^+ \in \mathbb{C}^{n\times m}$, called the *pseudoinverse of* A with the following property: For any vector $\mathbf{b} \in \mathbb{C}^m$, the unique least squares solution of minimum length equals the matrix product $A^+\mathbf{b}$. Schematically:

$$A\mathbf{x} = \mathbf{b} \quad \xrightarrow{\text{minimum length least squares solution}} \quad \hat{\mathbf{x}} = A^+\mathbf{b}.$$

Remark: This implies $O^+ = O^*$ for any shape of zero matrix.

Proof. First we prove the existence of a unique least squares solution of minimum length. By a previous theorem, we know that the least squares solutions to $A\mathbf{x} = \mathbf{b}$ are the exact solutions to $A\mathbf{x} = P_{\mathcal{C}(A)}\mathbf{b}$, where $P_{\mathcal{C}(A)}$ is the orthogonal projection onto the column space. Denote the set of least squares solutions by

$$S_{A,\mathbf{b}} = \{\mathbf{x} : A\mathbf{x} = P_{\mathcal{C}(A)}\mathbf{b}\}.$$

If $\mathbf{x}' \in S_{A,\mathbf{b}}$ is any one solution, we know from the previous section that $S_{A,\mathbf{b}} = \mathbf{x}' + \mathcal{N}(A)$. We need to prove that there exists a unique element $\hat{\mathbf{x}} \in S_{A,\mathbf{b}}$ such $\|\hat{\mathbf{x}}\| \leq \|\mathbf{x}' + \mathbf{y}\|$ for all $\mathbf{y} \in \mathcal{N}(A)$. From the distance-minimizing property of orthogonal projection, we have

$$\|\mathbf{x}' - P_{\mathcal{N}(A)}\mathbf{x}'\| \leq \|\mathbf{x}' - \mathbf{z}\| \text{ for all } \mathbf{z} \in \mathcal{N}(A),$$

with equality if and only if $\mathbf{z} = P_{\mathcal{N}(A)}\mathbf{x}'$. If $\mathbf{y} = -\mathbf{z}$ then since $\mathcal{N}(A)$ is a subspace we observe that $\mathbf{y}$ ranges over all of $\mathcal{N}(A)$ as $\mathbf{z}$ does, hence

$$\|\mathbf{x}' - P_{\mathcal{N}(A)}\mathbf{x}'\| \leq \|\mathbf{x}' + \mathbf{y}\| \text{ for all } \mathbf{y} \in \mathcal{N}(A),$$

with equality if and only if $\mathbf{y} = -P_{\mathcal{N}(A)}\mathbf{x}'$. We conclude that $\hat{\mathbf{x}} = \mathbf{x}' - P_{\mathcal{N}(A)}\mathbf{x}'$ is the desired unique element of $S_{A,\mathbf{b}}$ of minimum norm. Since $\mathcal{N}(A)$ and $\mathcal{R}(A)$ are orthogonal complements, their projection matrices satisfy $P_{\mathcal{N}(A)} + P_{\mathcal{R}(A)} = I$, hence we can also write

$$\hat{\mathbf{x}} = P_{\mathcal{R}(A)}\mathbf{x}'.$$

I claim we can also characterize $\hat{\mathbf{x}}$ as the unique least squares solution in the row space of A:

$$S_{A,\mathbf{b}} \cap \mathcal{R}(A) = \{\hat{\mathbf{x}}\},$$

[62]Remark: One can show that this matrix X has independent columns if and only if at least $d+1$ of the numbers $x_1, \ldots, x_n$ are distinct.

which shows that $\hat{\mathbf{x}}$ is independent of the choice of $\mathbf{x}'$. Indeed, since $P_{\mathcal{R}(A)}\mathbf{x}' \in \mathcal{R}(A)$ we already know that $\hat{\mathbf{x}} \in S_{A,\mathbf{b}} \cap \mathcal{R}(A)$. Conversely, consider any $\mathbf{x} \in S_{A,\mathbf{b}} \cap \mathcal{R}(A)$. Since $\mathbf{x}, \hat{\mathbf{x}} \in S_{A,\mathbf{b}}$ we have $\mathbf{x} - \hat{\mathbf{x}} \in \mathcal{N}(A)$ and since $\mathbf{x}, \hat{\mathbf{x}} \in \mathcal{R}(A)$ we have $\mathbf{x} - \hat{\mathbf{x}} \in \mathcal{R}(A)$. But $\mathcal{N}(A) \cap \mathcal{R}(A) = \{\mathbf{0}\}$, which implies that $\mathbf{x} - \hat{\mathbf{x}} = \mathbf{0}$, hence $\mathbf{x} = \hat{\mathbf{x}}$.

Finally, we show that for a given matrix A, the map sending $\mathbf{b} \mapsto \hat{\mathbf{x}}$ is linear, hence we can represent it by a matrix, which we denote by $\hat{\mathbf{x}} = A^+\mathbf{b}$. To see this, consider any vectors $\mathbf{b}_1, \mathbf{b}_2 \in \mathbb{C}^m$ with images $\hat{\mathbf{x}}_1, \hat{\mathbf{x}}_2$. By definition, this means that

$$A\hat{\mathbf{x}}_1 = P_{\mathcal{C}(A)}\mathbf{b}_1, \quad A\hat{\mathbf{x}}_2 = P_{\mathcal{C}(A)}\mathbf{b}_2, \quad \hat{\mathbf{x}}_1, \hat{\mathbf{x}}_2 \in \mathcal{R}(A).$$

For any scalars $\alpha, \beta \in \mathbb{C}$, we need to show that $\alpha\mathbf{b}_1 + \beta\mathbf{b}_2$ gets sent to $\alpha\hat{\mathbf{x}}_1 + \beta\hat{\mathbf{x}}_2$, i.e., that

$$A(\alpha\hat{\mathbf{x}}_1 + \beta\hat{\mathbf{x}}_2) = P_{\mathcal{C}(A)}(\alpha\mathbf{b}_1 + \beta\mathbf{b}_2) \quad \text{and} \quad \alpha\hat{\mathbf{x}}_1 + \beta\hat{\mathbf{x}}_2 \in \mathcal{R}(A).$$

But the first statement is true because matrix multiplication is linear, and the second statement is true because $\mathcal{R}(A)$ is a subspace. □

This theorem proves the existence of A^+ but doesn't tell us how to compute it. Here is a beautiful, explicit formula.

Definition (Rank Factorization). Let $A \in \mathbb{C}^{m\times n}$ and consider a factorization

$$A = CR,$$

where C has independent columns and R has independent rows. If A has rank r then we must have $C \in \mathbb{C}^{m\times r}$ and $R \in \mathbb{C}^{r\times n}$. Indeed, since C has independent columns, we know that $\dim \mathcal{R}(A) = \dim \mathcal{R}(R)$, and since R has independent columns, we know that $\dim \mathcal{C}(A) = \dim \mathcal{C}(C)$. Any such factorization $A = CR$ is called a *rank factorization*.

Rank factorizations always exist. Indeed, we can choose any r independent columns of A and let these be the columns of C. Then let R encode the expressions for the columns of A in terms of the columns of C. In our previous example, we can take the first and second columns of A, so that

$$A = \begin{pmatrix} 1 & 3 & 8 \\ 1 & 2 & 6 \\ 0 & 1 & 2 \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 3 \\ 1 & 2 \\ 0 & 1 \end{pmatrix}}_{C} \underbrace{\begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 2 \end{pmatrix}}_{R}.$$

Note that the resulting R necessarily has independent rows because it contains an $r \times r$ identity submatrix.[63]

Theorem (Computation of the Pseudoinverse). Consider a nonzero matrix $A \in \mathbb{C}^{m\times n}$ and let $A = CR$ be a rank factorization. Note that $(C^*C)^{-1}$ and $(RR^*)^{-1}$ exist because C has independent columns and R has independent rows. Then the pseudoinverse of A is

$$A^+ = R^*(RR^*)^{-1}(C^*C)^{-1}C^*.$$

[63] Remark: If C consists of the pivot columns of A, then R is just the RREF of A with the zero rows deleted. Thus one can easily read off a rank factorization of A from $\mathrm{RREF}(A)$.

Note that this product is independent of the choice of rank factorization.

Proof. Let $B = R^*(RR^*)^{-1}(C^*C)^{-1}C^*$, and consider any vector $\mathbf{b}$. From the existence theorem, we know that $A^+\mathbf{b}$ is uniquely characterized by the properties

$$A(A^+\mathbf{b}) = P_{\mathcal{C}(A)}\mathbf{b} \quad \text{and} \quad A^+\mathbf{b} \in \mathcal{R}(A).$$

We will show that the matrix $B\mathbf{b}$ has these same properties, and hence $A^+\mathbf{b} = B\mathbf{b}$. Since this holds for all $\mathbf{b}$, it will follow that $A^+ = B$. To prove the properties, first note that $\mathcal{C}(A) = \mathcal{C}(C)$ because R has independent rows. Then since C has independent columns, we have $P_{\mathcal{C}(C)} = C(C^*C)^{-1}C^*$, and hence

$$\begin{aligned} AB &= [CR][R^*(RR^*)^{-1}(C^*C)^{-1}C^*] \\ &= C(RR^*)(RR^*)^{-1}(C^*C)^{-1}C^* \\ &= C(C^*C)^{-1}C^* \\ &= P_{\mathcal{C}(C)} \\ &= P_{\mathcal{C}(A)}. \end{aligned}$$

It follows that $A(B\mathbf{b}) = P_{\mathcal{C}(A)}\mathbf{b}$, as desired. For the other condition, note that

$$B\mathbf{b} \in \mathcal{C}(R^*) = \mathcal{R}(R) = \mathcal{R}(CR) = \mathcal{R}(A).$$

The equality $\mathcal{R}(R) = \mathcal{R}(CR)$ holds because C has independent columns. □

In our previous example, one may compute that

$$\begin{aligned} A^+ &= R^*(RR^*)^{-1}(C^*C)^{-1}C^* \\ &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ 2 & 2 \end{pmatrix} \begin{pmatrix} 5 & 4 \\ 4 & 5 \end{pmatrix}^{-1} \begin{pmatrix} 2 & 5 \\ 5 & 14 \end{pmatrix}^{-1} \begin{pmatrix} 1 & 1 & 0 \\ 3 & 2 & 1 \end{pmatrix} \\ &= \frac{1}{9} \begin{pmatrix} -3 & 8 & -11 \\ 3 & -7 & 10 \\ 0 & 2 & -2 \end{pmatrix}. \end{aligned}$$

Recall from the previous section that the following system of equations has no solution:

$$A\mathbf{x} = \begin{pmatrix} 1 & 3 & 8 \\ 1 & 2 & 6 \\ 0 & 1 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \mathbf{b}$$

Since the columns of A are not independent, there are infinitely many least squares solutions. The unique least squares solution of minimum length is

$$\hat{\mathbf{x}} = A^+\mathbf{b} = \frac{1}{9} \begin{pmatrix} -3 & 8 & -11 \\ 3 & -7 & 10 \\ 0 & 2 & -2 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} -1/3 \\ 1/3 \\ 0 \end{pmatrix}.$$

We end by listing some interesting properties of the pseudoinverse. The proofs are fun exercises with the explicit formula $A^+ = R^*(RR^*)^{-1}(C^*C)^{-1}C^*$.

Theorem (Properties of the Pseudoinverse). Consider any nonzero matrix $A \in \mathbb{C}^{m\times n}$.

- $(A^+)^+ = A$
- $(A^*)^+ = (A^+)^*$
- If A has independent columns then $A^+ = (A^*A)^{-1}A^*$.
- If A has independent rows then $A^+ = A^*(AA^*)^{-1}$.
- If A is invertible then $A^+ = A^{-1}$.

Proof. Choose a rank factorization $A = CR$. Let $D := R^*(RR^*)^{-1}$ and $S := (C^*C)^{-1}C^*$, so that $A^+ = DS$. Since D has independent columns and S has independent rows, we have

$$(A^+)^+ = S^*(SS^*)^{-1}(D^*D)^{-1}D^*.$$

To simplify this formula, note that

$$\begin{aligned} SS^* &= (C^*C)^{-1}C^*[(C^*C)^{-1}C^*]^* \\ &= (C^*C)^{-1}C^*[C((C^*C)^{-1})^*] \\ &= (C^*C)^{-1}C^*[C((C^*C)^*)^{-1}] \\ &= (C^*C)^{-1}C^*[C(C^*C)^{-1}] \\ &= (C^*C)^{-1}(C^*C)(C^*C)^{-1} \\ &= (C^*C)^{-1}, \end{aligned}$$

and a similar computation shows that $D^*D = (RR^*)^{-1}$. One can also check that $S^* = C(C^*C)^{-1}$ and $D^* = (RR^*)^{-1}R$, hence we obtain

$$\begin{aligned} (A^+)^+ &= S^*(SS^*)^{-1}(D^*D)^{-1}D^* \\ &= S^*(C^*C)(RR^*)D^* \\ &= C(C^*C)^{-1}(C^*C)(RR^*)(RR^*)^{-1}R \\ &= CR \\ &= A. \end{aligned}$$

To show that $(A^*)^+ = (A^+)^{-1}$, we first note that $A^* = (CR)^* = R^*C^*$. Since R^* has independent columns and C^* has independent rows, this implies that

$$(A^*)^+ = C(C^*C)^{-1}(RR^*)^{-1}R = [R^*(RR^*)^{-1}(C^*C)^{-1}C^*]^* = (A^+)^*.$$

Next suppose that A has independent columns, so $r = n$. In this case we can write $A = AI_n$, where I_n has independent rows, hence

$$A^+ = I^*(II^*)^{-1}(A^*A)^{-1}A^* = (A^*A)^{-1}A^*.$$

A similar argument shows that if A has independent rows, then $A^+ = A^*(AA^*)^{-1}$. Finally, if A is invertible then we can use the independent columns formula to obtain

$$A^+ = (A^*A)^{-1}A^* = A^{-1}(A^*)^{-1}A^* = A^{-1}.$$

□

Remark: In general, we have $(AB)^+ \neq B^+A^+$.

4 Linear and Bilinear Forms

4.1 Linear Forms

Let V be a vector space over $\mathbb{R}$ (or $\mathbb{C}$). A linear function

$$\varphi : V \to \mathbb{R}$$

is called a *linear form*. If V is an infinite dimensional space of functions such as L^2 then a linear form is usually called a *linear functional*.

Linear forms on $\mathbb{R}^n$ are particularly simple. Let $\varphi : \mathbb{R}^n \to \mathbb{R}$ be a linear form and for each basis vector $\mathbf{e}_i$ define the scalar

$$b_i := \varphi(\mathbf{e}_i).$$

Then for any vector $\mathbf{x} = x_1\mathbf{e}_1 + \cdots + x_n\mathbf{e}_n \in \mathbb{R}^n$ we have

$$\begin{aligned}\varphi(\mathbf{x}) &= \varphi(x_1\mathbf{e}_1 + \cdots + x_n\mathbf{e}_n) \\ &= x_1\varphi(\mathbf{e}_1) + \cdots + x_n\varphi(\mathbf{e}_n) \\ &= x_1b_1 + \cdots + x_nb_n.\end{aligned}$$

If we write $\mathbf{b} = (b_1, \ldots, b_n)$ then this becomes

$$\varphi(\mathbf{x}) = \mathbf{b}^T\mathbf{x}.$$

We will denote the function $\mathbf{x} \mapsto \mathbf{b}^T\mathbf{x}$ by $\varphi_{\mathbf{b}} : \mathbb{R}^n \to \mathbb{R}$. Thus we obtain a bijection between vectors and linear forms:

$$\begin{array}{rcl}\mathbb{R}^n & \to & \text{linear forms on } \mathbb{R}^n \\ \mathbf{b} & \mapsto & \varphi_{\mathbf{b}}.\end{array}$$

Indeed, the function $\varphi_{\mathbf{b}}(\mathbf{x}) = \mathbf{b}^T\mathbf{x}$ is linear, and we have just seen that every linear function $\varphi : \mathbb{R}^n \to \mathbb{R}$ is equal to $\varphi_{\mathbf{b}}$ for some $\mathbf{b} \in \mathbb{R}^n$.

More abstractly, let V be an inner product space over $\mathbb{R}$ (or a Hermitian space over $\mathbb{C}$). Then for any vector $\mathbf{u} \in V$ we can define a linear form

$$\varphi_{\mathbf{u}}(\mathbf{v}) := \langle \mathbf{u}, \mathbf{v} \rangle.$$

Again, this gives a map[64] from V to the set of linear forms on V:

$$\begin{array}{rcl} V & \to & \text{linear forms on } V \\ \mathbf{u} & \mapsto & \varphi_{\mathbf{u}}. \end{array}$$

But this need not be a bijection in general. To investigate this, suppose that vectors $\mathbf{u}_1, \mathbf{u}_2 \in V$ correspond to the same functional, so that for all $\mathbf{v} \in V$ we have

$$\begin{aligned} \varphi_{\mathbf{u}_1}(\mathbf{v}) &= \varphi_{\mathbf{u}_2}(\mathbf{v}) \\ \langle \mathbf{u}_1, \mathbf{v} \rangle &= \langle \mathbf{u}_2, \mathbf{v} \rangle \\ \langle \mathbf{u}_1, \mathbf{v} \rangle - \langle \mathbf{u}_2, \mathbf{v} \rangle &= 0 \\ \langle \mathbf{u}_1 - \mathbf{u}_2, \mathbf{v} \rangle &= 0. \end{aligned}$$

Since this applies to any $\mathbf{v}$ we can take $\mathbf{v} = \mathbf{u}_1 - \mathbf{u}_2$ to obtain

$$\langle \mathbf{u}_1 - \mathbf{u}_2, \mathbf{u}_1 - \mathbf{u}_2 \rangle = 0.$$

But it is an axiom of (Hermitian) inner products that $\langle \mathbf{x}, \mathbf{x} \rangle = 0$ implies $\mathbf{x} = \mathbf{0}$, hence we must have

$$\begin{aligned} \mathbf{u}_1 - \mathbf{u}_2 &= 0 \\ \mathbf{u}_1 &= \mathbf{u}_2. \end{aligned}$$

This shows that the map from V to the set of linear forms on V is always injective. However, it is not necessarily surjective. This is the subject of the Riesz Representation Theorem.

Theorem (Riesz Representation). Let V be a *Hilbert space*. This means that V is an inner product space over $\mathbb{R}$ (or a Hermitian space over $\mathbb{C}$), and that Cauchy sequences with respect to the norm $\| - \| = \sqrt{\langle -, - \rangle}$ converge.[65] Let $\varphi : V \to \mathbb{R}$ be a linear functional. Then

$$\varphi = \varphi_{\mathbf{u}} \text{ for some } \mathbf{u} \in V \quad \iff \quad \varphi \text{ is continuous with respect to } \| - \|.$$

If V is finite dimensional then every linear functional is continuous. If V is infinite dimensional then there exist **discontinuous** functionals, but they are often ignored.

Let me introduce a some jargon. Given a vector space V over $\mathbb{R}$ (or $\mathbb{C}$) we define its *dual space* as the set of linear forms:[66]

$$\begin{aligned} V^\vee &= \text{the dual space} \\ &= \{\text{all linear forms } V \to \mathbb{R}\}. \end{aligned}$$

[64] So many different words for "function". The purpose is to avoid confusion when discussing many different kinds of functions at the same time.

[65] Recall: We say that $\mathbf{v}_1, \mathbf{v}_2, \ldots$ is a Cauchy sequence if for all $k \geq \ell \geq N$ we have $\|\mathbf{v}_k - \mathbf{v}_\ell\| \to 0$ as $N \to \infty$.

[66] It is more common to write V^* for the dual space, but I am already using that notation for the conjugate transpose.

As the name suggests, the set $V^\vee$ is also a vector space over $\mathbb{R}$. For a given list of forms $\varphi_i : V \to \mathbb{R}$ and scalars $a_i \in \mathbb{F}$ we define the form $\sum a_i\varphi_i : V \to \mathbb{R}$ "pointwise":

$$\left(\sum a_i\varphi_i\right)(\mathbf{v}) := \sum a_i\varphi_i(\mathbf{v}) \text{ for all } \mathbf{v} \in V.$$

I claim that this definition makes the map $V \to V^\vee$ into a linear map. To see this, let's give the map a name. Let Φ denote the map that sends the vector $\mathbf{u} \in V$ to the form $\varphi_\mathbf{u} \in V^\vee$:

$$\begin{array}{rccc} \Phi : V & \to & V^\vee \\ \mathbf{u} & \mapsto & \varphi_\mathbf{u} \end{array}$$

Then for any linear combination of vectors $\sum a_i\mathbf{u}_i \in V$, I claim that

$$\Phi\left(\sum a_i\mathbf{u}_i\right) = \sum a_i\Phi(\mathbf{u}_i),$$

where each side of the equation is a linear form. To show that two forms are equal we must show that they define the same function $V \to \mathbb{R}$. So consider any vector $\mathbf{v} \in V$. Then since $\Phi(\mathbf{u})$ is just another name for $\varphi_\mathbf{u}$, we have

$$\begin{aligned} \left[\Phi\left(\sum a_i\mathbf{u}_i\right)\right](\mathbf{v}) &= \varphi_{\sum a_i\mathbf{u}_i}(\mathbf{v}) \\ &= \left\langle \sum a_i\mathbf{u}_i, \mathbf{v} \right\rangle \\ &= \sum a_i\langle \mathbf{u}_i, \mathbf{v}\rangle \\ &= \sum a_i\varphi_{\mathbf{u}_i}(\mathbf{v}) \\ &= \left[\sum a_i\Phi(\mathbf{u}_i)\right](\mathbf{v}). \end{aligned}$$

Thus $\Phi : V \to V^\vee$ is an injective linear map, and if V is finite dimensional then it is also surjective, hence it is an isomorphism $V \cong V^\vee$. When V is infinite dimensional then Φ is **not surjective**, however it is common to restrict the definition of $V^\vee$ as follows:

$$V^\vee = \{\text{the set of } \textbf{continuous} \text{ linear functionals } V \to \mathbb{R}\}.$$

Then from the Riesz Reprentation Theorem we will still have $V \cong V^\vee$.

Another piece of jargon is the Dirac *bra-ket notation* from quantum physics. To motivate this, consider the isomorphism between $\mathbb{R}^n$ and its dual:

$$\begin{array}{ccc} \mathbb{R}^n & \cong & (\mathbb{R}^n)^\vee \\ \mathbf{b} & \leftrightarrow & \varphi_\mathbf{b}, \end{array}$$

where the form $\varphi_\mathbf{b} : \mathbb{R}^n \to \mathbb{R}$ corresponding to the column vector $\mathbf{b}$ is defined by $\varphi_\mathbf{b}(\mathbf{x}) = \mathbf{b}^T\mathbf{x}$. But every linear function corresponds to a matrix, and the linear function $\varphi_\mathbf{b} : \mathbb{R}^n \to \mathbb{R}$ corresponds to the $1 \times n$ row vector $\mathbf{b}^T$. In the language of Chapter 2, we have

$$[\varphi_\mathbf{b}] = \mathbf{b}^T.$$

Thus it makes sense to identify the dual space $(\mathbb{R}^n)^\vee$ with the space of row vectors, and the isomorphism $\mathbb{R}^n \cong (\mathbb{R}^n)^\vee$ with transposition:[67]

$$\begin{array}{ccc} \mathbb{R}^n & \cong & (\mathbb{R}^n)^\vee \\ \mathbf{b} & \leftrightarrow & \mathbf{b}^T. \end{array}$$

For infinite dimensional spaces we can no longer use matrices. However, if V is an infinite dimensional Hilbert space of functions, such as $L^2(\mathbb{C})$, and $V^\vee$ is its dual space of **continuous** functionals, Dirac introduced the following notation:

$$\begin{array}{ccc} V & \cong & V^\vee \\ |f\rangle & \leftrightarrow & \langle f|. \end{array}$$

This notation is compatible with the inner product notation $\langle -, - \rangle$ since, by definition, the functional $\langle f| \in V^\vee$ acts on the vector $|g\rangle \in V$ by

$$\langle f| \text{ acting on } |g\rangle = \langle f, g\rangle.$$

Hence in the physics notation the inner product is written as $\langle f|g\rangle$.

4.2 Bilinear Forms

Let V be a vector space over $\mathbb{R}$ (or $\mathbb{C}$). A *bilinear form* is a function

$$\varphi : V \times V \to \mathbb{R}$$

that is linear in each coordinate:

- $\varphi(\mathbf{u}, \sum a_i \mathbf{v}_i) = \sum a_i \varphi(\mathbf{u}, \mathbf{v}_i)$,
- $\varphi(\sum a_i \mathbf{u}_i, \mathbf{v}) = \sum a_i \varphi(\mathbf{u}_i, \mathbf{v})$.

Remark: Over $\mathbb{C}$ we want one of the coordinates to be conjugate linear. In this course I have picked the first coordinate:

$$\varphi(\sum a_i \mathbf{u}_i, \mathbf{v}) = \sum a_i^* \varphi(\mathbf{u}_i, \mathbf{v}).$$

In this case we say that φ is *sesquilinear* (one-and-a-half times linear) instead of bilinear. For example, an inner product is a bilinear function and a Hermitian inner product is a sesquilinear function.

As with linear forms, we begin with the case of Euclidean space. Let $\varphi : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$ be a bilinear form, and for any two basis vectors $\mathbf{e}_i, \mathbf{e}_j \in \mathbb{R}^n$ define the scalar

$$b_{ij} := \varphi(\mathbf{e}_i, \mathbf{e}_j).$$

[67] Another piece of jargon: Sometimes the elements of $(\mathbb{R}^n)^\vee$ are called *co-vectors*.

Then for any vectors $\mathbf{x} = x_1\mathbf{e}_1 + \cdots + x_n\mathbf{e}_n$ and $\mathbf{y} = y_1\mathbf{e}_1 + \cdots + y_n\mathbf{e}_n$ we have

$$\begin{aligned}\varphi(\mathbf{x},\mathbf{y}) &= \varphi(x_1\mathbf{e}_1 + \cdots + x_n\mathbf{e}_n, y_1\mathbf{e}_1 + \cdots + y_n\mathbf{e}_n)\\ &= \sum x_i y_j \varphi(\mathbf{e}_i,\mathbf{e}_j)\\ &= \sum x_i y_i b_{ij}.\end{aligned}$$

If we let B be the $n \times n$ matrix with ij entry b_{ij} then this becomes

$$\varphi(\mathbf{x},\mathbf{y}) = \mathbf{x}^T B\mathbf{y} = \begin{pmatrix} x_1 & \cdots & x_n \end{pmatrix} \begin{pmatrix} b_{11} & \cdots & b_{1n} \\ \vdots & & \vdots \\ b_{n1} & \cdots & b_{nn} \end{pmatrix} \begin{pmatrix} y_1 \\ \vdots \\ y_n \end{pmatrix}.$$

Exercise: Verify this. Conversely, for any $n \times n$ matrix B we can define a bilinear form φ_B by

$$\varphi_B(\mathbf{x},\mathbf{y}) := \mathbf{x}^T B\mathbf{y}.$$

If B has ij entry b_{ij} then it follows that

$$\varphi_B(\mathbf{e}_i,\mathbf{e}_j) = \mathbf{e}_i^T B\mathbf{e}_j = (i\text{th row of } B)\mathbf{e}_j = b_{ij}.$$

Hence for any $n \times n$ matrices B and C we have

$$\begin{aligned}\varphi_B = \varphi_C \quad &\Longrightarrow \quad \varphi_B(\mathbf{x},\mathbf{y}) = \varphi_C(\mathbf{x},\mathbf{y}) \text{ for all } \mathbf{x},\mathbf{y} \in \mathbb{R}^n\\ &\Longrightarrow \quad \varphi_B(\mathbf{e}_i,\mathbf{e}_j) = \varphi_C(\mathbf{e}_i,\mathbf{e}_j) \text{ for all } i,j\\ &\Longrightarrow \quad b_{ij} = c_{ij} \text{ for all } i,j\\ &\Longrightarrow \quad B = C.\end{aligned}$$

In summary, we obtain a bijection between $n \times n$ matrices and bilinear forms:

$$\begin{aligned}\text{square matrices } \mathbb{R}^{n\times n} \quad &\leftrightarrow \quad \text{bilinear forms } \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}\\ B \quad &\leftrightarrow \quad \varphi_B.\end{aligned}$$

We can also view this as an isomorphism of vector spaces, since bilinear forms can be added and multiplied by scalars, as can any kind of functions with values in $\mathbb{R}$. The following result compares properties of the form φ_B to properties of the matrix B.

Theorem (Properties of Bilinear Forms). Let B be an $n \times n$ matrix over $\mathbb{R}$ (or $\mathbb{C}$) and consider the bilinear (or sesquilinear) form φ_B defined by[68]

$$\varphi_B(\mathbf{x},\mathbf{y}) = \mathbf{x}^T B\mathbf{y} \text{ over } \mathbb{R} \qquad \text{or} \qquad \varphi_B(\mathbf{x},\mathbf{y}) = \mathbf{x}^* B\mathbf{y} \text{ over } \mathbb{C}.$$

[68]Recall: For any matrix A with complex entries, A^* denotes the conjugate transpose matrix. If $\mathbf{x}$ is a column vector then $\mathbf{x}^*$ is a row vector.

(a) **Symmetric.** We have

$$\begin{aligned} \varphi_B(\mathbf{x},\mathbf{y}) = \varphi_B(\mathbf{y},\mathbf{x}) \text{ for all } \mathbf{x},\mathbf{y} \quad &\Longleftrightarrow \quad B^T = B, \\ \varphi_B(\mathbf{x},\mathbf{y})^* = \varphi_B(\mathbf{y},\mathbf{x}) \text{ for all } \mathbf{x},\mathbf{y} \quad &\Longleftrightarrow \quad B^* = B. \end{aligned}$$

In the first case we say that the form φ_B and the matrix B are *symmetric*. In second case we say they are *Hermitian*.

(b) **Positive Semi-Definite.** We have

$$\varphi_B(\mathbf{x},\mathbf{x}) \geq 0 \text{ for all } \mathbf{x} \quad \Longleftrightarrow \quad B = A^T A \text{ (or } B = A^* A) \text{ for some matrix } A.$$

In this case the form φ_B and the matrix B are called *positive semi-definite*.[69]

(c) **Positive Definite.** Let φ_B be positive semi-definite, so that $B = A^T A$ (or $B = A^* A$) as in part (b). Then we have

$$\varphi_B(\mathbf{x},\mathbf{x}) = 0 \text{ implies } x = \mathbf{0} \quad \Longleftrightarrow \quad \text{the matrix } A \text{ has independent columns.}$$

In this case the form φ_B and the matrix B are called *positive definite*.

(d) **Negative.** If $B = -A^T A$ (or $B = -A^* A$) for some matrix A then we have

$$\varphi_B(\mathbf{x},\mathbf{x}) \leq 0 \text{ for all } x,$$

in which case we say that φ_B and B are *negative semi-definite*. If, in addition, the matrix A has independent columns then

$$\varphi_B(\mathbf{x},\mathbf{x}) = 0 \text{ implies } \mathbf{x} = \mathbf{0},$$

in which case we say that φ_B and B are *negative definite*.

(e) **Indefinite.** If B is not of the form $\pm A^T A$ (or $\pm A^* A$) for some matrix A, then there exist points $\mathbf{x}$ and $\mathbf{y}$ such that

$$\varphi_B(\mathbf{x},\mathbf{x}) > 0 \quad \text{and} \quad \varphi_B(\mathbf{y},\mathbf{y}) < 0.$$

In this case we say that φ_B and B are *indefinite*.

Example: The identity matrix I corresponds to the standard dot product $\varphi_I(\mathbf{x},\mathbf{y}) = \mathbf{x}^T\mathbf{y}$ on $\mathbb{R}^n$ and the standard Hermitian product $\varphi_I(\mathbf{x},\mathbf{y}) = \mathbf{x}^*\mathbf{y}$ on $\mathbb{C}^n$, both of which are positive definite. Indeed, we can write $I = I^T I$, where I has independent columns.

Remark: Many problems in applied mathematics seek to minimize an expression of the form $\mathbf{x}^T B\mathbf{x}$ (or $\mathbf{x}^* B\mathbf{x}$). If we know that $B = A^T A$ (or $B = A^* A$) for some matrix A with independent columns then we are guaranteed that a unique minimum exists. Indeed, from part (b) we know that $\mathbf{x}^T B\mathbf{x} \geq 0$ for all $\mathbf{x}$ and from part (c) we know that $\mathbf{x}^T B\mathbf{x} > 0$ for all $\mathbf{x} \neq \mathbf{0}$.

[69] Some books use the alternate term *non-positive definite*.

Proof. We only prove the complex versions, since the real versions are just a special case. Furthermore, we will only prove one direction of (b) and (c). The other directions are harder and we will prove them after discussing the Spectral Theorem.

(a): If b_{ij} is the ij entry of the matrix B then we have seen that $\varphi_B(\mathbf{e}_i, \mathbf{e}_j) = b_{ij}$ where $\mathbf{e}_i$ and $\mathbf{e}_j$ are standard basis vectors. Suppose that $\varphi_B(\mathbf{x}, \mathbf{y})^* = \varphi_B(\mathbf{y}, \mathbf{x})$ for all $\mathbf{x}, \mathbf{y} \in \mathbb{C}^n$, then in particular we must have

$$b_{ij}^* = \varphi_B(\mathbf{e}_i, \mathbf{e}_j)^* = \varphi_B(\mathbf{e}_j, \mathbf{e}_j) = b_{ij},$$

and hence $B^* = B$. Conversely, suppose that $B^* = B$. Then for all $\mathbf{x}, \mathbf{y} \in \mathbb{C}^n$ we have

$$\begin{aligned}
\varphi_B(\mathbf{x}, \mathbf{y})^* &= (\mathbf{x}^* B \mathbf{y})^* \\
&= \mathbf{y}^* B^* (\mathbf{x}^*)^* \\
&= \mathbf{y}^* B \mathbf{x} \\
&= \varphi_B(\mathbf{y}, \mathbf{x}).
\end{aligned}$$

(b): Suppose that $B = A^* A$ for some matrix A, and let $\|\mathbf{v}\| = \sqrt{\mathbf{v}^* \mathbf{v}}$ be the standard Hermitian norm on $\mathbb{C}^n$. Then for all $\mathbf{x} \in \mathbb{C}^n$ we have

$$\begin{aligned}
\varphi_B(\mathbf{x}, \mathbf{x}) &= \mathbf{x}^* B \mathbf{x} \\
&= \mathbf{x}^* A^* A \mathbf{x} \\
&= (A\mathbf{x})^* (A\mathbf{x}) \\
&= \|A\mathbf{x}\|^2 \geq 0.
\end{aligned}$$

(c): Continuing from (b), suppose that $\varphi_B(\mathbf{x}, \mathbf{x}) = 0$, so that $\|A\mathbf{x}\|^2 = 0$. This implies that $A\mathbf{x} = \mathbf{0}$ because of properties of the standard Hermitian norm.[70] But if A has independent columns then this implies that $\mathbf{x} = \mathbf{0}$. There are many ways to see this. One method uses the fact that $(A^T A)^{-1}$ exists to get

$$\begin{aligned}
A\mathbf{x} &= \mathbf{0} \\
A^T A\mathbf{x} &= A^T \mathbf{0} \\
A^T A\mathbf{x} &= \mathbf{0} \\
\mathbf{x} &= (A^T A)^{-1} \mathbf{0} \\
\mathbf{x} &= \mathbf{0}.
\end{aligned}$$

(d): This follows from (b) and (c), and the fact that

$$\varphi_{-B}(\mathbf{x}, \mathbf{x}) = \mathbf{x}^T (-B)\mathbf{x} = -\mathbf{x}^T B \mathbf{x} = -\varphi_B(\mathbf{x}, \mathbf{x}).$$

[70] Recall that $\|\mathbf{v}\|^2 = |v_1|^2 + \cdots + |v_n|^2$, so that $\|\mathbf{v}\| = 0$ if and only if $|v_i| = 0$ (and hence $v_i = 0$) for all i.

(e): This follows from (b), (c) and (d). □

As with linear forms, it is also possible to define bilinear (sesquilinear) forms on infinite dimensional vector spaces. Let V be any Hermitian inner product space over $\mathbb{C}$ and let $B : V \to V$ be any linear operator.[71] Then we can define a function $\varphi_B : V \times V \to \mathbb{C}$ by

$$\varphi_B(\mathbf{x}, \mathbf{y}) = \langle \mathbf{x}, B\mathbf{y} \rangle.$$

In the finite dimensional case this corresponds to $\langle \mathbf{x}, B\mathbf{y} \rangle = \mathbf{x}^* B\mathbf{y}$, where B is a matrix. If B^* is the conjugate transpose matrix, then we observe that

$$\langle B^*\mathbf{x}, \mathbf{y} \rangle = (B^*\mathbf{x})^*\mathbf{y} = \mathbf{x}^*(B^*)^*\mathbf{y} = \mathbf{x}^* B\mathbf{y} = \langle \mathbf{x}, B\mathbf{y} \rangle.$$

This computation suggests a way to define a "conjugate transpose operator" $B^* : V \to V$, even when V is infinite dimensional. The definition is really a theorem.

Theorem (Adjoint Operators). Let V be a complex Hilbert space and consider a linear operator $B : V \to V$. If B is **continuous** with respect to the standard norm $\| - \| = \sqrt{\langle -, - \rangle}$ then there exists a unique linear operator $B^* : V \to V$, which is also continuous, satisfying

$$\langle B^*\mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{u}, B\mathbf{v} \rangle \text{ for all } \mathbf{u}, \mathbf{v} \in V.$$

The operator B^* is called the *adjoint* of B.[72]

These ideas are particularly important in quantum mechanics. In the standard statistical interpretation, a nonzero vector in Hilbert space $\psi \in V$ corresponds to the *state* of a quantum system. An operator $Q : V \to V$ satisfyiing $Q^* = Q$ corresponds to an *observable quantity*. The outcome of a measurement is random but the *expected value* of quantity Q on state ψ is

$$\langle \psi, Q\psi \rangle \quad \text{or } \langle \psi | Q | \psi \rangle \text{ in Dirac notation.}$$

Those who study quantum mechanics will notice that it is mostly linear algebra, but the notation is different and the vectors and operators are sometimes just pretend.[73]

4.3 Quadratic Forms

Let V be a vector space over $\mathbb{R}$. Given a bilinear form $\varphi : V \times V \to \mathbb{R}$ we define the corresponding *quadratic form* $Q : V \to \mathbb{R}$ by

$$Q(\mathbf{x}) := \varphi(\mathbf{x}, \mathbf{x}).$$

[71] Yet another fancy word that just means "function".

[72] An operator is continuous if and only if it is bounded

[73] Indeed, we have seen that the "functions" $\delta(x)$ and $e^{2\pi i x}$ are treated as elements of $L^2(\mathbb{C})$, even though $e^{2\pi i x}$ is not square integrable and $\delta(x)$ doesn't really exist. Furthermore, the theorem on adjoints applies to **continuous** operators, but many operators of interest in quantum mechanics, such as position and momentum, are not continuous.

In the case of Euclidean space $V = \mathbb{R}^n$ suppose that $\varphi(\mathbf{x}, \mathbf{y}) = \varphi_B(\mathbf{x}, \mathbf{y}) = \mathbf{x}^T B\mathbf{y}$ for a square matrix B. Then the corresponding quadratic form is

$$Q_B(\mathbf{x}) = \mathbf{x}^T B\mathbf{x}.$$

Quadratic forms give a relationship between polynomials of degree 2 and linear algebra. For example, consider a polynomial in two variables:

$$f(x, y) = 2 + x - y + 3x^2 + 2xy + 4y^2.$$

We can express this in terms of linear algebra as follows:

$$f(x, y) = 2 + \begin{pmatrix} 1 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} x & y \end{pmatrix} \begin{pmatrix} 3 & 2 \\ 0 & 4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}.$$

Indeed, for any 2×2 matrix B we observe that

$$\begin{aligned} \mathbf{x}^T B\mathbf{x} &= \begin{pmatrix} x & y \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \\ &= \begin{pmatrix} x & y \end{pmatrix} \begin{pmatrix} ax + by \\ cx + dy \end{pmatrix} \\ &= x(ax + by) + y(cx + dy) \\ &= ax^2 + bxy + cyx + dy^2 \\ &= ax^2 + (b + c)xy + dy^2. \end{aligned}$$

This formula shows that the choice of b and c is not unique. It is common to choose $b = c$ so that the corresponding matrix B is symmetric. Thus we can express any polynomial $\alpha x^2 + \beta xy + \gamma y^2$ in terms of a symmetric matrix:

$$\alpha x^2 + \beta xy + \gamma y^2 = \begin{pmatrix} x & y \end{pmatrix} \begin{pmatrix} \alpha & \beta/2 \\ \beta/2 & \gamma \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}.$$

And we can rewrite the polynomial $f(x, y)$ above using a symmetric matrix:

$$f(x, y) = 2 + \begin{pmatrix} 1 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} x & y \end{pmatrix} \begin{pmatrix} 3 & 1 \\ 1 & 4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}.$$

More generally, let $\mathbf{x} = (x_1, \ldots, x_n)$ be a vector of n unknowns. Then any polynomial $f(\mathbf{x}) = f(x_1, \ldots, x_n)$ of degree 2 has a unique expression of the form

$$f(\mathbf{x}) = b + \mathbf{b}^T\mathbf{x} + \mathbf{x}^T B\mathbf{x},$$

where b is a scalar, $\mathbf{b}^T$ is a row vector and B is a symmetric matrix. For example, in the case $n = 3$ it is common to write $\mathbf{x} = (x, y, z)$ instead of $\mathbf{x} = (x_1, x_2, x_3)$. Then we have

$$f(x, y, z) = b + b_1x + b_2y + b_3z + b_{11}x^2 + b_{22}y^2 + b_{33}z^2 + b_{12}xy + b_{13}xz + b_{23}yz$$


$$= b + \begin{pmatrix} b_1 & b_2 & b_3 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} + \begin{pmatrix} x & y & z \end{pmatrix} \begin{pmatrix} b_{11} & b_{12}/2 & b_{13}/2 \\ b_{12}/2 & b_{22} & b_{23}/2 \\ b_{13}/2 & b_{23}/2 & b_{33} \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}$$
$$= b + \mathbf{b}^T\mathbf{x} + \mathbf{x}^T B\mathbf{x}.$$

Thus the degree zero terms correspond to a scalar b, the degree 1 terms correspond to a vector $\mathbf{b}^T$,[74] and the degree 2 terms correspond to a matrix B. To describe higher degree polynomials we would need cubes of numbers, hypercubes of numbers, etc. Such objects are called "tensors" and they are more difficult to work with. Luckily, degree 2 polynomials are sufficient for most applications.[75]

Here are three simplest examples of quadratic forms. Let

$$B = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \text{ so that } Q_B(\mathbf{x}) = \mathbf{x}^T B\mathbf{x} = x^2 + y^2.$$

The graph of $Q_B(x, y)$ in $\mathbb{R}^3$ looks like a paraboloid with a unique minimum at $(0, 0)$:




Indeed, this matrix is positive definite because it can be factored as $B = I^T I$, where I is the identity matrix, which has independent columns. Next, let

$$B = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \text{ so that } Q_B(\mathbf{x}) = \mathbf{x}^T B\mathbf{x} = x^2.$$

The graph of $Q_B(x, y)$ in $\mathbb{R}^3$ is a parabolic cylinder:

[74] It doesn't matter whether we write the degree 1 terms as $\mathbf{b}^T\mathbf{x}$ or $\mathbf{x}^T\mathbf{b}$. I am simply following the convention from Section 1.1, where linear forms correspond to row vectors.

[75] It is a curious fact that most physical laws can be expressed in terms of first and second derivatives. Higher derivatives are almost never useful.




This time the minimum is not unique, since $Q_B(0, y) = 0$ for any value of y. Indeed, this matrix can be factored as

$$B = A^T A = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \end{pmatrix},$$

where the matrix A does **not** have independent columns. Finally, let

$$B = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \text{ so that } Q_B(\mathbf{x}) = \mathbf{x}^T B \mathbf{x} = x^2 - y^2.$$

This time the graph of $Q_B(x, y)$ in $\mathbb{R}^3$ is a saddle:



Since Q_B takes both positive and negative values, it follows from the previous section that B cannot be factored as $B = A^T A$ for any matrix A, although this is a bit hard to see directly.

In the next chapter we will prove the Spectral Theorem, which makes the analysis of quadratic forms much easier. As a preview, we will prove the following results. Let B be a square matrix satisfying $B^T = B$. Then:

- The eigenvalues of B are real.
- B is positive semi-definite if and only if all eigenvalues are ≥ 0.
- B is positive definite if and only if all eigenvalues are > 0.
- B if indefinite if and only if there exist both positive and negative eigenvalues.

4.4 Multivariable Taylor Expansion

From calculus we are familiar with the idea of a Taylor series. Suppose that a function $f : \mathbb{R} \to \mathbb{R}$ is differentiable k times at the point $p \in \mathbb{R}$. Then for small values of x we have

$$f(p+x) = f(p) + f'(p)x + \frac{1}{2}f''(p)x^2 + \cdots + \frac{1}{k!}f^{(k)}(p)x^k + \text{higher terms},$$

where the higher terms are vanishingly small.[76]

The concept of Taylor series can be generalized to higher dimensions using a little bit of linear algebra. Consider a real valued function $f : \mathbb{R}^n \to \mathbb{R}$ written as

$$f(\mathbf{x}) = f(x_1, \ldots, x_n),$$

where $\mathbf{x} \in \mathbb{R}^n$ is the input vector. We will denote first partial derivatives by

$$f_i = \frac{\partial}{\partial x_i} f,$$

and second partial derivatives by

$$f_{ij} = \frac{\partial}{\partial x_j}\frac{\partial}{\partial x_i} f.$$

Note that f_i and f_{ij} are themselves functions from $\mathbb{R}^n$ to $\mathbb{R}$. Suppose that the first and second partials exist and are continuous at some point $\mathbf{p} \in \mathbb{R}^n$. Then Clairaut's theorem tells us that

$$f_{ij}(\mathbf{p}) = f_{ji}(\mathbf{p}) \text{ for all } i, j.$$

Furthermore, we define the gradient vector at $\mathbf{p}$:

$$(\nabla f)_{\mathbf{p}} = \begin{pmatrix} f_1(\mathbf{p}) \\ \vdots \\ f_n(\mathbf{p}) \end{pmatrix}$$

[76] The exact nature of the higher terms will not concern us; we don't do analysis in this course.

and the Hessian matrix at $\mathbf{p}$:

$$(Hf)_{\mathbf{p}} = \begin{pmatrix} f_{11}(\mathbf{p}) & \cdots & f_{1n}(\mathbf{p}) \\ \vdots & & \vdots \\ f_{n1}(\mathbf{p}) & \cdots & f_{nn}(\mathbf{p}) \end{pmatrix}.$$

Note that the Hessian matrix is symmetric. Then for small vectors $\mathbf{x} \in \mathbb{R}^n$, the multivariable Taylor series tells us that

$$f(\mathbf{p}+\mathbf{x}) = f(\mathbf{p}) + (\nabla f)_p^T \mathbf{x} + \frac{1}{2}\mathbf{x}^T (Hf)_{\mathbf{p}} \mathbf{x} + \text{higher terms},$$

where the higher terms are vanishingly small. Note the relationship to linear and bilinear forms. The linear part of the Taylor series is a linear form

$$\mathbf{x} \mapsto (\nabla f)_{\mathbf{p}}^T \mathbf{x} = f_1(\mathbf{p})x_1 + f_2(\mathbf{p})x_2 + \cdots + f_n(\mathbf{p})x_n,$$

and the quadratic part of the Taylor series is a quadratic form

$$\mathbf{x} \mapsto \frac{1}{2}\mathbf{x}^T (Hf)_{\mathbf{p}} \mathbf{x} = \frac{1}{2} \sum f_{ij}(\mathbf{p}) x_i x_j.$$

Higher terms of the Taylor series can be described by multilinear forms, but, as I said, these don't come up much in applications.

For example, consider again the polynomial function, with $(x_1, x_2) = (x, y)$:

$$f(x,y) = 2 + x - y + 3x^2 + 2xy + 4y^2.$$

We compute the first and second partial derivatives:

$$\begin{aligned} f_1 &= 1 + 6x + 2y, \\ f_2 &= -1 + 2x + 8y, \\ f_{11} &= 6, \\ f_{12} &= 2, \\ f_{21} &= 2, \\ f_{22} &= 8. \end{aligned}$$

This gives the following gradient vector and Hessian matrix:

$$\nabla f = \begin{pmatrix} 1 + 6x + 2y \\ -1 + 2x + 8y \end{pmatrix} \quad \text{and} \quad Hf = \begin{pmatrix} 6 & 2 \\ 2 & 4 \end{pmatrix}.$$

The Taylor expansion at $\mathbf{p} = (0,0)$ is

$$f(0+x, 0+y) = f(0,0) + (\nabla f)_{(0,0)}^T \mathbf{x} + \frac{1}{2}\mathbf{x}^T (Hf)_{(0,0)} \mathbf{x}$$

$$= 2 + \begin{pmatrix} 1 & -1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \frac{1}{2} \begin{pmatrix} x & y \end{pmatrix} \begin{pmatrix} 6 & 2 \\ 2 & 8 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix},$$


which we already computed in the previous section. The Taylor expansion at $\mathbf{p} = (1, 1)$ is

$$\begin{aligned} f(1+x, 1+y) &= f(1,1) + (\nabla f)^T_{(1,1)}\mathbf{x} + \frac{1}{2}\mathbf{x}^T (Hf)_{(1,1)}\mathbf{x} \\ &= 11 + \begin{pmatrix} 9 & 9 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \frac{1}{2} \begin{pmatrix} x & y \end{pmatrix} \begin{pmatrix} 6 & 2 \\ 2 & 8 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}. \end{aligned}$$

And the Taylor expansion at $\mathbf{p} = \left(\frac{-10}{44}, \frac{8}{44}\right)$ is

$$\begin{aligned} f\left(\frac{-10}{44} + x, \frac{8}{44} + y\right) &= f\left(\frac{-10}{44}, \frac{8}{44}\right) + (\nabla f)^T_{\left(\frac{-10}{44}, \frac{8}{44}\right)}\mathbf{x} + \frac{1}{2}\mathbf{x}^T (Hf)_{\left(\frac{-10}{44}, \frac{8}{44}\right)}\mathbf{x} \\ &= \frac{79}{44} + \begin{pmatrix} 0 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \frac{1}{2} \begin{pmatrix} x & y \end{pmatrix} \begin{pmatrix} 6 & 2 \\ 2 & 8 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \\ &= \frac{79}{44} + \begin{pmatrix} x & y \end{pmatrix} \begin{pmatrix} 3 & 1 \\ 1 & 4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}. \end{aligned}$$

Note that $\mathbf{p} = \left(\frac{-10}{44}, \frac{8}{44}\right)$ is a *critical point* of f, since the gradient vector vanishes: $(\nabla f)_{\mathbf{p}} = \mathbf{0}$. Recall that $(\nabla f)_{\mathbf{p}}$ is the direction of greatest increase of f near the point $\mathbf{p}$. If $(\nabla f)_{\mathbf{p}} = \mathbf{0}$ then the function is in equilibrium because it can't decide which way is "up". Here is a picture:



A multivariable function $f : \mathbb{R}^n \to \mathbb{R}$ near a critical point $\mathbf{p}$ is approximately a quadratic form:

$$f(\mathbf{p}+\mathbf{x}) = f(\mathbf{p}) + \frac{1}{2}\mathbf{x}^T(Hf)_{\mathbf{p}}\mathbf{x} + \text{higher terms.}$$

Thus we have the following facts, which are sometimes called the *multivariable second derivative test*. Assume that $(\nabla f)_{\mathbf{p}} = \mathbf{0}$. Then:

- f has a local minimum at $\mathbf{p}$ if and only if $(Hf)_{\mathbf{p}}$ is positive definite.
- f has a local maximum at $\mathbf{p}$ if and only if $(Hf)_{\mathbf{p}}$ is negative definite.

Indeed, if $(Hf)_{\mathbf{p}}$ is positive definite then we have

$$\mathbf{x}^T(Hf)_{\mathbf{p}}\mathbf{x} \geq 0 \text{ for all } \mathbf{x}, \text{ and } \mathbf{x}^T(Hf)_{\mathbf{p}}\mathbf{x} = 0 \text{ if and only if } \mathbf{x} = \mathbf{0}.$$

so that[77]

$$f(\mathbf{p}+\mathbf{x}) \geq f(\mathbf{p}) \text{ for all } \mathbf{x}, \text{ and } f(\mathbf{p}+\mathbf{x}) = f(\mathbf{p}) \text{ if and only if } \mathbf{x} = \mathbf{0},$$

If $(Hf)_{\mathbf{p}}$ is positive (or negative) semi-definite then there is a local minimum (or maximum) in some directions, but in some directions the function is constant. Otherwise, if $(Hf)_{\mathbf{p}}$ is indefinite then there exist small $\mathbf{x}$ and $\mathbf{y}$ such that $f(\mathbf{p}+\mathbf{x}) > f(\mathbf{p})$ and $f(\mathbf{p}+\mathbf{y}) < f(\mathbf{p})$. Geometrically, this is a higher dimensional saddle point.

In the previous example, it happens that

$$\text{the matrix } B = \begin{pmatrix} 3 & 1 \\ 1 & 4 \end{pmatrix} \text{ is positive definite,}$$

so the function $f(x,y) = f(x,y) = 2 + x - y + 3x^2 + 2xy + 4y^2$ has a local minimum at $\mathbf{p} = \left(\frac{-10}{44}, \frac{8}{44}\right)$. To verify that B is positive definite, I computed the eigenvalues $7+\sqrt{5}$ and $7-\sqrt{5}$, which are both positive. Later I will show you how to find a matrix A with independent columns such that $B = A^TA$. Such a matrix is not unique; here is one example, called the *Cholesky decomposition*:

$$A = \begin{pmatrix} \sqrt{3} & \sqrt{3}/3 \\ 0 & \sqrt{33}/3 \end{pmatrix}.$$

Check:

$$A^TA = \begin{pmatrix} \sqrt{3} & 0 \\ \sqrt{3}/3 & \sqrt{33}/3 \end{pmatrix}\begin{pmatrix} \sqrt{3} & \sqrt{3}/3 \\ 0 & \sqrt{33}/3 \end{pmatrix} = \begin{pmatrix} 3 & 1 \\ 1 & 4 \end{pmatrix}.$$

[77] Remember, the higher order terms are vanishingly small, so they don't affect the inequality.

5 Determinants

5.1 Multilinear Forms

We have studied linear and bilinear forms. Now we discuss the general situation. Let V be a vector space over $\mathbb{R}$, and recall the notation for Cartesian product:

$$V^k := V \times V \times \cdots \times V = \{(\mathbf{x}_1, \ldots, \mathbf{x}_k) : \mathbf{x}_i \in \mathbb{R}^n \text{ for all } i\}.$$

A *multilinear k-form* is a function

$$\varphi : V^k \to \mathbb{R}$$

that is linear in each input. In other words, for any index i we have

$$\varphi\left(\mathbf{v}_1, \ldots, \mathbf{v}_{i-1}, \sum a_j \mathbf{u}_j, \mathbf{v}_{i+1}, \ldots, \mathbf{v}_k\right) = \sum a_i \varphi(\mathbf{v}_1, \ldots, \mathbf{v}_{i-1}, \mathbf{u}_j, \mathbf{v}_{i+1}, \ldots, \mathbf{v}_k).$$

(This time we don't bother with Hermitian forms, since it's not clear where to put the complex conjugates.) Just as with linear and bilinear forms, k-forms can be added and multiplied by scalars. That is, given k-forms φ, ψ and scalar a, we define the k-form $\varphi + a\psi$ by

$$(\varphi + a\psi)(\mathbf{v}_1, \ldots, \mathbf{v}_k) = \varphi(\mathbf{v}_1, \ldots, \mathbf{v}_k) + a\psi(\mathbf{v}_1, \ldots, \mathbf{v}_k).$$

Thus we obtain a vector space of multilinear k-forms:[78]

$$\mathcal{T}^k(V) = \{\text{multilinear } k\text{-forms } \varphi : V^k \to \mathbb{R}\}.$$

In the case $k = 1$ we also use the notation of the dual space

$$V^\vee = \mathcal{T}^1(V) = \{\text{linear forms } V \to \mathbb{R}\}.$$

For example, consider Euclidean space $V = \mathbb{R}^n$. In the previous section we proved that $\mathcal{T}^1(\mathbb{R}^n)$ is isomorphic to the vector space of row vectors:

$$\mathcal{T}^1(\mathbb{R}^n) \cong \{1 \times n \text{ row vectors}\} = \mathbb{R}^{1\times n},$$

and hence

$$\dim \mathcal{T}^1(\mathbb{R}^n) = n.$$

We also proved that $\mathcal{T}^2(\mathbb{R}^n)$ is isomorphic to the vector space of $n \times n$ matrices:

$$\mathcal{T}^2(\mathbb{R}^n) \cong \{n \times n \text{ matrices}\} = \mathbb{R}^{n\times n},$$

and hence[79]

$$\dim \mathcal{T}^1(\mathbb{R}^n) = n^2.$$

[78] The letter $\mathcal{T}$ is for "tensor".

[79] An $n \times n$ matrix is uniquely determined by its n^2 entries. More formally, let E_{ij} the $n \times n$ matrix with 1 in the ij position and zeros elsewhere. Then the set of matrices E_{ij} with $1 \le i, j \le n$ is a basis for $\mathbb{R}^{n\times n}$. More generally, one can show that $\mathbb{R}^{m\times n}$ has dimension mn.

More generally, I claim that

$$\dim \mathcal{T}^k(\mathbb{R}^n) = n^k.$$

In order to prove this we will construct a "standard basis" for $\mathcal{T}^k(\mathbb{R}^n)$.

Theorem (The Dual Standard Basis). Let $\mathbf{e}_1, \ldots, \mathbf{e}_n$ be the standard basis for $\mathbb{R}^n$. Now we will construct a corresponding "standard basis" for the dual space $(\mathbb{R}^n)^\vee = \mathcal{T}^1(\mathbb{R}^n)$. For all $1 \leq i \leq n$, let $\varepsilon_i : \mathbb{R}^n \to \mathbb{R}$ be the linear form defined by picking out the ith coordinate:

$$\varepsilon_i(\mathbf{x}) = \varepsilon_i \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} = x_i.$$

To see that this ε_i is linear, consider any linear combination $\sum a_j \mathbf{x}_j \in \mathbb{R}^n$, where x_{ij} is the ith entry of the vector $\mathbf{x}_j \in \mathbb{R}^n$. Then we have

$$\varepsilon_i \left(\sum a_j \mathbf{x}_j \right) = \varepsilon_i \begin{pmatrix} \sum a_j x_{1j} \\ \vdots \\ \sum a_j x_{nj} \end{pmatrix} = \sum a_j x_{ij} = \sum a_j \varepsilon_i(\mathbf{x}_j).$$

In the previous section we showed that every linear form $\varphi : \mathbb{R}^n \to \mathbb{R}$ can be expressed as $\varphi(\mathbf{x}) = \mathbf{b}^T \mathbf{x}$ for some unique vector $\mathbf{b} = (b_1, \ldots, b_n)$. Equivalently, each linear form φ can be expressed as

$$\varphi = b_1 \varepsilon_1 + \cdots + b_n \varepsilon_n,$$

for some unique scalars $b_1, \ldots, b_n$. This shows that $\varepsilon_1, \ldots, \varepsilon_2$ is indeed a basis for $(\mathbb{R}^n)^\vee$. In terms of matrices, note that

$$\varepsilon_i(\mathbf{x}) = \varepsilon_i \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} = x_i = \begin{pmatrix} 0 & \cdots & 0 & 1 & 0 & \cdots & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix},$$

which shows that the linear function ε_i corresponds to a standard row vector:

$$[\varepsilon_i] = \begin{pmatrix} 0 & \cdots & 0 & 1 & 0 & \cdots & 0 \end{pmatrix}.$$

Finally, we say that the bases $\mathbf{e}_1, \ldots, \mathbf{e}_n \in \mathbb{R}^n$ and $\varepsilon_1, \ldots, \varepsilon_n \in (\mathbb{R}^n)^\vee$ are "dual" because

$$\varepsilon_i(\mathbf{e}_j) = \begin{cases} 1 & i = j, \\ 0 & i \neq j. \end{cases}$$

If we were only going to talk about row vectors and column vectors then this level of abstraction is completely unnecessary. However, it becomes necessary when we talk about k-forms.

Tensor Product of Forms. Let V be a vector space over $\mathbb{R}$. Consider a k-form $\varphi : V^k \to \mathbb{R}$ and an ℓ-form $\psi : V^\ell \to \mathbb{R}$. Then the *tensor product* $\varphi \otimes \psi$ is a $(k+\ell)$-form defined as follows:

$$(\varphi \otimes \psi)(\mathbf{v}_1, \ldots, \mathbf{v}_k, \mathbf{v}_{k+1}, \ldots, \mathbf{v}_{k+\ell}) := \varphi(\mathbf{v}_1, \ldots, \mathbf{v}_k) \cdot \psi(\mathbf{v}_{k+1}, \ldots, \mathbf{v}_{k+\ell}).$$

It is straightforward to check that this function is linear, and hence $\varphi \otimes \psi \in \mathcal{T}^{k+\ell}(V)$. One can also check that the tensor product is associative, hence if φ, ψ, ω are k, ℓ, m-forms, respectively, then we obtain a $(k+\ell+m)$-form:

$$\varphi \otimes \psi \otimes \omega = (\varphi \otimes \psi) \otimes \omega = \varphi \otimes (\psi \otimes \omega).$$

For example, for any standard 1-forms ε_i and ε_j we obtain a 2 form $\varepsilon_i \otimes \varepsilon_j$ defined as follows:

$$(\varepsilon_i \otimes \varepsilon_j)(\mathbf{v}_1, \mathbf{v}_2) = \varepsilon_i(\mathbf{v}_1) \cdot \varepsilon_j(\mathbf{v}_2).$$

And for any standard 1-forms $\varepsilon_i, \varepsilon_j, \varepsilon_k$ we obtain a 3-form $\varepsilon_i \otimes \varepsilon_j \otimes \varepsilon_k$ by

$$(\varepsilon_i \otimes \varepsilon_j \otimes \varepsilon_k)(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3) = \varepsilon_i(\mathbf{v}_1) \cdot \varepsilon_j(\mathbf{v}_2) \cdot \varepsilon_k(\mathbf{v}_3).$$

To be more explicit let's consider an example with $V = \mathbb{R}^3$. Then we have

$$(\varepsilon_1 \otimes \varepsilon_2) \left(\begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}, \begin{pmatrix} 2 \\ 3 \\ 4 \end{pmatrix} \right) = \varepsilon_1 \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} \cdot \varepsilon_2 \begin{pmatrix} 2 \\ 3 \\ 4 \end{pmatrix} = (1)(3) = 3$$

and

$$(\varepsilon_2 \otimes \varepsilon_1) \left(\begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}, \begin{pmatrix} 2 \\ 3 \\ 4 \end{pmatrix} \right) = \varepsilon_2 \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix} \cdot \varepsilon_1 \begin{pmatrix} 2 \\ 3 \\ 4 \end{pmatrix} = (-1)(2) = -2,$$

which shows that $\varepsilon_1 \otimes \varepsilon_2$ and $\varepsilon_2 \otimes \varepsilon_1$ define different bilinear functions. In other words, we see that the **tensor product is not commutative**.

Theorem (The Standard Basis of k-Forms). Let $\mathbf{e}_1, \ldots, \mathbf{e}_n$ be the standard basis of $\mathbb{R}^n$ and let $\varepsilon_1, \ldots, \varepsilon_n$ be the dual standard basis of $\mathcal{T}^1(\mathbb{R}^n)$. Then I claim that the following set is a basis for the vector space of k-forms:

$$\left\{ \varepsilon_{i_1} \otimes \varepsilon_{i_2} \otimes \cdots \otimes \varepsilon_{i_k} : i_1, i_2, \ldots, i_k \in \{1, 2, \ldots, n\} \right\}.$$

Note that this basis contains n^k elements, and hence

$$\dim \mathcal{T}^k(\mathbb{R}^n) = n^k.$$

We won't bother to prove this since we have already proved the cases $k = 1$ and $k = 2$ in the previous section. The general proof is similar, but with more horrible notation. To see how this works, we will repeat our proof for $k = 2$ in the new language. Let B be an $n \times n$ matrix with ij entry b_{ij} and consider the 2-form

$$\varphi_B(\mathbf{x}, \mathbf{y}) = \mathbf{x}^T B \mathbf{y}.$$

Note that for any basis vectors $\mathbf{e}_i, \mathbf{e}_j$ we have

$$\varphi_B(\mathbf{e}_i, \mathbf{e}_j) = \mathbf{e}_i^T B \mathbf{e}_j = b_{ij}.$$

Furthermore, for any vectors $\mathbf{x} = (x_1, \ldots, x_n)$ and $\mathbf{y} = (y_1, \ldots, y_n)$ we have

$$\varphi_B(\mathbf{x}, \mathbf{y}) = \mathbf{x}^T B \mathbf{y} = \sum b_{ij} x_i y_j.$$

On the other hand, since $(\varepsilon_i \otimes \varepsilon_j)(\mathbf{x}, \mathbf{y}) = x_i y_j$, we can express this as

$$\varphi_B(\mathbf{x}, \mathbf{y}) = \mathbf{x}^T B y = \sum b_{ij} x_i y_j = \sum b_{ij} (\varepsilon_i \otimes \varepsilon_j)(\mathbf{x}, \mathbf{y}) = \left(\sum b_{ij} (\varepsilon_i \otimes \varepsilon_j) \right) (\mathbf{x}, \mathbf{y}),$$

and hence

$$\varphi_B = \sum b_{ij} (\varepsilon_i \otimes \varepsilon_j).$$

More generally, any 3-form $\varphi \in \mathcal{T}^3(\mathbb{R}^n)$ corresponds to an $n \times n \times n$ cube of numbers b_{ijk}:

$$\varphi = \sum b_{ijk} (\varepsilon_i \otimes \varepsilon_j \otimes \varepsilon_k).$$

These are some kind of "higher dimensional matrices", but they are much harder to work with. In this course we will focus only on very special kinds of k-forms.

Symmetric and Alternating k-Forms. We say that a k-form $\varphi \in \mathcal{T}^k(V)$ is *symmetric* if switching any two inputs leaves the output unchanged. For example, if φ is symmetric then

$$\varphi(\mathbf{v}_2, \mathbf{v}_1, \mathbf{v}_3, \ldots, \mathbf{v}_k) = \varphi(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \ldots, \mathbf{v}_k).$$

We say that a k-form φ is *alternating* if switching any two inputs multiplies the output by -1. For example, if φ is alternating then

$$\varphi(\mathbf{v}_2, \mathbf{v}_1, \mathbf{v}_3, \ldots, \mathbf{v}_k) = -\varphi(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \ldots, \mathbf{v}_k).$$

To be more explicit, let's consider $V = \mathbb{R}^3$. I claim that the 2-form $\varphi = \varepsilon_1 \otimes \varepsilon_2 + \varepsilon_2 \otimes \varepsilon_1$ is symmetric. Indeed, for any vectors $\mathbf{x}, \mathbf{y} \in \mathbb{R}^3$ we observe that

$$\begin{aligned}
\varphi(\mathbf{x}, \mathbf{y}) &= (\varepsilon_1 \otimes \varepsilon_2 + \varepsilon_2 \otimes \varepsilon_1) \left(\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}, \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} \right) \\
&= (\varepsilon_1 \otimes \varepsilon_2) \left(\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}, \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} \right) + (\varepsilon_2 \otimes \varepsilon_1) \left(\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}, \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} \right) \\
&= \varepsilon_1 \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \cdot \varepsilon_2 \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} + \varepsilon_2 \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \cdot \varepsilon_1 \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} \\
&= x_1 y_2 + x_2 y_1
\end{aligned}$$

is equal to

$$\varphi(\mathbf{y}, \mathbf{x}) = (\varepsilon_1 \otimes \varepsilon_2 + \varepsilon_2 \otimes \varepsilon_1) \left(\begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}, \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \right)$$

$$
\begin{aligned}
&= (\varepsilon_1 \otimes \varepsilon_2)\left(\begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}, \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}\right) + (\varepsilon_2 \otimes \varepsilon_1)\left(\begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}, \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}\right) \\
&= \varepsilon_1 \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} \cdot \varepsilon_2 \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} + \varepsilon_2 \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} \cdot \varepsilon_1 \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \\
&= y_1 x_2 + y_2 x_1 \\
&= x_1 y_2 + x_2 y_1.
\end{aligned}
$$

On the other hand, the 2-form $\psi = \varepsilon_1 \otimes \varepsilon_2 - \varepsilon_2 \otimes \varepsilon_1$ is alternating since

$$\psi(\mathbf{x}, \mathbf{y}) = (\varepsilon_1 \otimes \varepsilon_2 - \varepsilon_2 \otimes \varepsilon_1)(\mathbf{x}, \mathbf{y}) = x_1 y_2 - x_2 y_1$$

and

$$\psi(\mathbf{y}, \mathbf{x}) = (\varepsilon_1 \otimes \varepsilon_2 - \varepsilon_2 \otimes \varepsilon_1)(\mathbf{y}, \mathbf{x}) = y_1 x_2 - y_2 x_1 = -(x_1 y_2 - x_2 y_1) = -\psi(\mathbf{x}, \mathbf{y}).$$

Since the sum of symmetric forms is symmetric, and the sum of alternating forms is alternating, we can define the following vector spaces[80]

$$
\begin{aligned}
\mathcal{S}^k(V) &= \text{the space of symmetric } k\text{-forms } V^k \to \mathbb{R}, \\
\mathcal{A}^k(V) &= \text{the space of alternating } k\text{-forms } V^k \to \mathbb{R}.
\end{aligned}
$$

For small k and n, it is not too hard to write down a basis for $\mathcal{S}^k(\mathbb{R}^n)$ in terms of the standard basis of $\mathcal{T}^k(\mathbb{R}^n)$. To save space, let's write

$$\varepsilon_{ij} = \varepsilon_i \otimes \varepsilon_j, \quad \varepsilon_{ijk} = \varepsilon_i \otimes \varepsilon_j \otimes \varepsilon_k, \quad \text{etc.}$$

Then, for example, we have

$$
\begin{aligned}
\mathcal{S}^1(\mathbb{R}^2) &= \operatorname{Span}\{\varepsilon_1, \varepsilon_2\}, \\
\mathcal{S}^2(\mathbb{R}^2) &= \operatorname{Span}\{\varepsilon_{11}, \varepsilon_{12} + \varepsilon_{21}, \varepsilon_{22}\}, \\
\mathcal{S}^3(\mathbb{R}^2) &= \operatorname{Span}\{\varepsilon_{111}, \varepsilon_{112} + \varepsilon_{121} + \varepsilon_{211}, \varepsilon_{122} + \varepsilon_{212} + \varepsilon_{211}, \varepsilon_{222}\},
\end{aligned}
$$

and

$$
\begin{aligned}
\mathcal{S}^1(\mathbb{R}^3) &= \operatorname{Span}\{\varepsilon_1, \varepsilon_2, \varepsilon_3\}, \\
\mathcal{S}^2(\mathbb{R}^3) &= \operatorname{Span}\{\varepsilon_{11}, \varepsilon_{22}, \varepsilon_{33}, \varepsilon_{12} + \varepsilon_{21}, \varepsilon_{13} + \varepsilon_{31}, \varepsilon_{23} + \varepsilon_{32}\}, \\
\mathcal{S}^3(\mathbb{R}^3) &= \operatorname{Span}\{\varepsilon_{111}, \varepsilon_{222}, \varepsilon_{333}, \\
&\qquad \varepsilon_{112} + \varepsilon_{121} + \varepsilon_{211}, \varepsilon_{113} + \varepsilon_{131} + \varepsilon_{311}, \varepsilon_{223} + \varepsilon_{232} + \varepsilon_{322}, \\
&\qquad \varepsilon_{221} + \varepsilon_{212} + \varepsilon_{122}, \varepsilon_{331} + \varepsilon_{313} + \varepsilon_{133}, \varepsilon_{332} + \varepsilon_{323} + \varepsilon_{233},
\end{aligned}
$$

[80] Alternating forms are also called *anti-symmetric*. In advanced calculus, a *differential form* is an alternating k-form whose coefficients can change from point to point. More precisely, a differential form on a k-dimensional manifold assigns an alternating k-form to the tangent space at each point.

$$\varepsilon_{123} + \varepsilon_{132} + \varepsilon_{213} + \varepsilon_{231} + \varepsilon_{312} + \varepsilon_{321}\}.$$

In particular, we have

$$\dim \mathcal{S}^1(\mathbb{R}^3) = 2, \quad \dim \mathcal{S}^2(\mathbb{R}^3) = 6, \quad \dim \mathcal{S}^3(\mathbb{R}^3) = 10.$$

Maybe you can see a pattern here. In general, one can use a combinatorial argument[81] to show that

$$\dim \mathcal{S}^k(\mathbb{R}^n) = \binom{n+k-1}{k}.$$

Let's test this on the special case $k = 2$. Recall from the previous section that a symmetric bilinear form is the same thing as a symmetric $n \times n$ matrix, hence $\mathcal{S}^2(\mathbb{R}^n)$ can be identified with the space of symmetric $n \times n$ matrices. A symmetric matrix is uniquely determined by the n diagonal elements and the $n(n-1)/2$ elements above the diagonal. (We don't need to specify the entries below the diagonal because they are equal to the above-diagonal elements.) Hence we must have

$$\dim \mathcal{S}^2(\mathbb{R}^n) = n + \frac{n(n-1)}{2} = \frac{2n + n(n-1)}{2} = \frac{n^2+n}{2} = \frac{(n+1)n}{2},$$

which agrees with the formula

$$\binom{n+2-1}{2} = \binom{n+1}{2} = \frac{(n+1)n}{2}.$$

It is trickier to find a basis for the space of alternating k-forms. Here are some small examples:

$$\begin{aligned}
\mathcal{A}^1(\mathbb{R}^2) &= \text{Span}\{\varepsilon_1, \varepsilon_2\}, \\
\mathcal{A}^2(\mathbb{R}^2) &= \text{Span}\{\varepsilon_{12} - \varepsilon_{21}\}, \\
\mathcal{A}^k(\mathbb{R}^2) &= \{0\} \text{ for } k > 2, \\
\mathcal{A}^1(\mathbb{R}^3) &= \text{Span}\{\varepsilon_1, \varepsilon_2, \varepsilon_3\}, \\
\mathcal{A}^2(\mathbb{R}^3) &= \text{Span}\{\varepsilon_{12} - \varepsilon_{21}, \varepsilon_{13} - \varepsilon_{31}, \varepsilon_{23} - \varepsilon_{32}\}, \\
\mathcal{A}^3(\mathbb{R}^3) &= \text{Span}\{\varepsilon_{123} + \varepsilon_{231} + \varepsilon_{312} - \varepsilon_{132} - \varepsilon_{213} - \varepsilon_{321}\}, \\
\mathcal{A}^k(\mathbb{R}^3) &= \{0\} \text{ for } k > 3.
\end{aligned}$$

You will prove on the homework that $\dim \mathcal{A}^k(\mathbb{R}^n) = 0$ for all $k > n$. That is, if $k > n$ then any alternating k-form on $\mathbb{R}^n$ must be the zero function that sends any k-tuple of vectors in $\mathbb{R}^n$ to zero. For $0 \le k \le n$ I claim that[82]

$$\dim \mathcal{A}^k(\mathbb{R}^n) = \binom{n}{k}.$$

[81] There is one basis element of $\mathcal{S}^k(\mathbb{R}^n)$ for each weakly increasing sequence $1 \le i_1 \le i_2 \le \cdots \le i_k \le n$ of k numbers between 1 and n. Such a weakly increasing sequence can be encoded as a word of length $n+k-1$ containing k "stars" and $n-1$ "bars". For example, the word $**|*||***$ corresponds to $1 \le 1 \le 2 \le 4 \le 4 \le 4$. Such a word has length $k + (n-1) = n + k - 1$. The number of such words is $\binom{n+k-1}{k}$ since from $n+k-1$ possible positions, we must choose k positions to place the stars.

[82] The definition of "alternating" doesn't really apply to 0-forms and 1-forms. However, it is convenient to define $\mathcal{A}^0 := \mathcal{T}^0 := \{0\}$ and $\mathcal{A}^1 := \mathcal{T}^1$, so the dimension formula is still correct when $k = 0$ and $k = 1$.

We won't prove this theorem in general, but we will prove the special case when $k = n$:

$$\dim \mathcal{A}^n(\mathbb{R}^n) = \binom{n}{n} = 1.$$

In other words, there exists a unique (up to scalar multiplication) alternating n-form on $\mathbb{R}^n$. At the risk of spoiling the surprise, I will tell you right now that this unique form is called the *determinant*.

According to the examples listed above, we have

$$\begin{aligned}\mathcal{A}^2(\mathbb{R}^2) &= \operatorname{Span}\{\varepsilon_{12} - \varepsilon_{21}\},\\ \mathcal{A}^3(\mathbb{R}^3) &= \operatorname{Span}\{\varepsilon_{123} + \varepsilon_{231} + \varepsilon_{312} - \varepsilon_{132} - \varepsilon_{213} - \varepsilon_{321}\}.\end{aligned}$$

Recall that $\varepsilon_{12} - \varepsilon_{21}$ represents the 2-form $\varepsilon_1 \otimes \varepsilon_2 - \varepsilon_2 \otimes \varepsilon_1$, which we have already discussed. When applied to two vectors $\mathbf{x} = (x_1, x_2)$ and $\mathbf{y} = (y_1, y_2)$ in $\mathbb{R}^2$ it gives

$$(\varepsilon_{12} - \varepsilon_{21})(\mathbf{x}, \mathbf{y}) = \varepsilon_{12}(\mathbf{x}, \mathbf{y}) - \varepsilon_{21}(\mathbf{x}, \mathbf{y}) = x_1 y_2 - x_2 y_1.$$

In general, if $\varphi \in \mathcal{T}^k(\mathbb{R}^n)$ is a k-form on $\mathbb{R}^n$ and if A is a $n \times k$ matrix with columns $\mathbf{a}_1, \ldots, \mathbf{a}_k \in \mathbb{R}^n$, it is convenient to define

$$\varphi(A) := \varphi(\mathbf{a}_1, \ldots, \mathbf{a}_k).$$

Thus for any 2×2 matrix we have

$$(\varepsilon_{12} - \varepsilon_{21})(A) = (\varepsilon_{12} - \varepsilon_{21}) \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix} = \varepsilon_{12} \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix} - \varepsilon_{21} \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix} = a_1 b_2 - a_2 b_1,$$

and for any 3×3 matrix

$$A = \begin{pmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{pmatrix}$$

we have

$$\begin{aligned}&(\varepsilon_{123} + \varepsilon_{231} + \varepsilon_{312} - \varepsilon_{132} - \varepsilon_{213} - \varepsilon_{321})(A)\\ &\quad = \varepsilon_{123}(A) + \varepsilon_{231}(A) + \varepsilon_{312}(A) - \varepsilon_{132}(A) - \varepsilon_{231}(A) - \varepsilon_{321}(A),\\ &\quad = a_1 b_2 c_3 + a_2 b_3 c_1 + a_3 b_1 c_2 - a_1 b_3 c_2 - a_2 b_3 c_1 - a_3 b_2 c_1.\end{aligned}$$

You may recognize these formulas from your previous linear algebra course. But where do they come from? And how do we know that there are no other alternating 2-forms on $\mathbb{R}^2$ and no other alternating 3-forms on $\mathbb{R}^3$?

5.2 Uniqueness of the Determinant

As we have seen, the formula for the determinant of a 3×3 matrix is rather complicated. I could give a general formula right now, but it is actually more useful to work with the **properties** of the determinant. Explicit formulas for the determinant are messy, but the properties of the determinant are easy to describe.

As before, we will think of a k-form $\varphi \in \mathcal{T}^k(\mathbb{R}^n)$ as a function sending $n \times k$ matrices to scalars. That is, for any matrix A with columns $\mathbf{a}_1, \ldots, \mathbf{a}_k \in \mathbb{R}^n$ we will write

$$\varphi(A) := \varphi(\mathbf{a}_1, \ldots, \mathbf{a}_k).$$

This function is "multilinear in the columns of A". For example, consider some $n \times 3$ matrices

$$A = \left(\begin{array}{c|c|c} \mathbf{u} & \mathbf{v} & \mathbf{a} \end{array} \right), \quad B = \left(\begin{array}{c|c|c} \mathbf{u} & \mathbf{v} & \mathbf{b} \end{array} \right), \quad C = \left(\begin{array}{c|c|c} \mathbf{u} & \mathbf{v} & \mathbf{a} + \lambda \mathbf{b} \end{array} \right),$$

with $\mathbf{u}, \mathbf{v}, \mathbf{a}, \mathbf{b} \in \mathbb{R}^n$ and $\lambda \in \mathbb{R}$. Then for any 3-form $\varphi \in \mathcal{T}^3(\mathbb{R}^3)$ we have

$$\varphi(C) = \varphi(A) + \lambda \cdot \varphi(B).$$

Warning: Multilinear functions are not linear. For example, consider any bilinear function $\varphi \in \mathcal{T}^2(\mathbb{R}^n)$, and consider any two $n \times 2$ matrices

$$A = \left(\begin{array}{c|c} \mathbf{a}_1 & \mathbf{a}_2 \end{array} \right) \text{ and } B = \left(\begin{array}{c|c} \mathbf{b}_1 & \mathbf{b}_2 \end{array} \right), \text{ hence } A + B = \left(\begin{array}{c|c} \mathbf{a}_1 + \mathbf{b}_1 & \mathbf{a}_2 + \mathbf{b}_2 \end{array} \right).$$

Then we have

$$\begin{aligned} \varphi(A + B) &= \varphi(\mathbf{a}_1 + \mathbf{b}_1, \mathbf{a}_2 + \mathbf{b}_2) \\ &= \varphi(\mathbf{a}_1, \mathbf{a}_2) + \varphi(\mathbf{b}_1, \mathbf{b}_2) + \varphi(\mathbf{a}_1, \mathbf{b}_2) + \varphi(\mathbf{b}_1, \mathbf{a}_2) \\ &= \varphi(A) + \varphi(B) + \varphi(\mathbf{a}_1, \mathbf{b}_2) + \varphi(\mathbf{b}_1, \mathbf{a}_2), \end{aligned}$$

which is **not** equal to $\varphi(A) + \varphi(B)$.[83]

As mentioned in the previous section, there exists a unique (up to scalar multiplication) alternating n-form on $\mathbb{R}^n$, which can be interpreted as the determinant of an $n \times n$ matrix. In this section we will prove that there is no more than one such function, so that

$$\dim \mathcal{A}^n(\mathbb{R}^n) \leq 1,$$

and in the next section we will show that there is at least one such function, so that

$$\dim \mathcal{A}^n(\mathbb{R}^n) \geq 1.$$

Theorem (Uniqueness of the Determinant). Let φ be a function sending $n \times n$ matrices to scalars. We say that φ is a *determinant function* if it satisfies the following three properties:

[83] For the same reason, we will have $\det(A + B) \neq \det(A) + \det(B)$.

(1) *Multilinear.* The function φ is linear in each individual column.

(2) *Alternating.* If A' is obtained from A by swapping two columns, then $\varphi(A') = -\varphi(A)$.

(3) *Normalized.* The function φ sends the identity matrix I_n to 1.

In other words, a determinant function is an alternating n-form $\varphi \in \mathcal{A}^n(\mathbb{R}^n)$ that is appropriately normalized so that

$$\varphi(I_n) = \varphi(\mathbf{e}_1, \ldots, \mathbf{e}_n) = 1.$$

I claim that

there is at most one determinant function.

In order to streamline the proof I will isolate several lemmas, which have independent interest.

Lemma A. Let φ be a determinant function. If A has a repeated column then

$$\varphi(A) = 0.$$

Proof. Suppose that the ith and jth columns are equal and let A' be the matrix obtained from A by switching the ith and jth columns. On the one hand we have $A' = A$. On the other hand, property (2) tells us that

$$\begin{aligned} \varphi(A') &= -\varphi(A) \\ \varphi(A) &= -\varphi(A) \\ 2\varphi(A) &= 0 \\ \varphi(A) &= 0. \end{aligned}$$

□

Lemma B. Let φ be a determinant function. If A has dependent columns then

$$\varphi(A) = 0.$$

Proof. Let A have columns $\mathbf{a}_1, \ldots, \mathbf{a}_n \in \mathbb{R}^n$. If these columns are dependent then there exists some i such that $\mathbf{a}_i$ can be expressed as a linear combination of the other columns. Without loss of generality,[84] suppose that $i = 1$, so we can write

$$\mathbf{a}_1 = b_1\mathbf{a}_2 + \cdots + b_n\mathbf{a}_n,$$

for some scalars $b_2, \ldots, b_n$. Now let $\hat{A}_1(\mathbf{a}_j)$ denote the matrix A with the first column replaced by $\mathbf{a}_j$. From property (1) we have

$$\varphi(A) = b_1 \cdot \varphi(\hat{A}_1(\mathbf{a}_2)) + \cdots + b_n \cdot \varphi(\hat{A}_1(\mathbf{a}_n)).$$

[84] By applying property (2) we can swap the 1st and ith columns, which does not affect whether the determinant is zero or nonzero.

But each matrix $\hat{A}_1(\mathbf{a}_j)$ with $j \neq 1$ has a repeated column, so from Lemma A we must have

$$\begin{aligned}\varphi(A) &= b_1 \cdot \varphi(\hat{A}_1(\mathbf{a}_2)) + \cdots + b_n \cdot \varphi(\hat{A}_1(\mathbf{a}_n)) \\ &= b_1 \cdot 0 + b_2 \cdot 0 + \cdots + b_n \cdot 0 \\ &= 0.\end{aligned}$$

□

The next lemma refers to the elementary matrices, which we discussed in the previous chapter:

$$D_i(\lambda) = \begin{pmatrix} 1 & & & & & \\ & 1 & & & & \\ & & \lambda & & & \\ & & & 1 & & \\ & & & & 1 & \\ & & & & & 1 \end{pmatrix},$$

$$L_{ij}(\lambda) = \begin{pmatrix} 1 & & & & & \\ & 1 & \cdots & \lambda & & \\ & & 1 & \vdots & & \\ & & & 1 & & \\ & & & & 1 & \\ & & & & & 1 \end{pmatrix},$$

$$T_{ij} = \begin{pmatrix} 1 & & & & & \\ & 0 & \cdots & 1 & & \\ & \vdots & 1 & \vdots & & \\ & 1 & \cdots & 0 & & \\ & & & & 1 & \\ & & & & & 1 \end{pmatrix}.$$

Lemma C. Let φ be a determinant function. Then for any square matrix A we have

$$\begin{aligned}\varphi(AD_i(\lambda)) &= \lambda \cdot \varphi(A), \\ \varphi(AL_{ij}(\lambda)) &= \varphi(A), \\ \varphi(AT_{ij}) &= -\varphi(A).\end{aligned}$$

Proof. First, note that $AD_i(\lambda)$ has the same columns as A except that the ith column has been scaled by λ, hence $\varphi(AD_i(\lambda)) = \lambda \cdot \varphi(A)$ follows from property (1). Next, note that AT_{ij} is obtained from A by switching columns i and j, hence the identity $\varphi(AT_{ij}) = -\varphi(A)$ is just a restatement of (2). Finally, note that kth column of $AL_{ij}(\lambda)$ is equal to the kth column of A, except in the case $k = j$, in which case

$$(j\text{th column of } AL_{ij}(\lambda)) = (j\text{th column of } A) + \lambda \cdot (i\text{th column of } A).$$

To simplify notation, let $\mathbf{a}_1, \ldots, \mathbf{a}_n$ be the columns of A and let $\hat{A}_j(\mathbf{v})$ denote the matrix A with the jth column replaced by vector $\mathbf{v}$. Then from property (1) we have

$$\varphi(AL_{ij}(\lambda)) = \varphi(A) + \lambda \cdot \varphi(\hat{A}_j(\mathbf{a}_i)).$$

But the matrix $\hat{A}_j(\mathbf{a}_i)$ has a repeated column, so it follows from Lemma A that

$$\varphi(AL_{ij}(\lambda)) = \varphi(A) + \lambda \cdot 0 = \varphi(A).$$

□

Lemma D. Let φ be a determinant function. Then we have

$$\varphi(D_i(\lambda)) = \lambda, \quad \varphi(L_{ij}(\lambda)) = 1, \quad \varphi(T_{ij}) = -1.$$

Proof. Taking $A = I$ in Lemma C and using property (3) gives[85]

$$\begin{aligned} \varphi(D_i(\lambda)) &= \varphi(ID_i(\lambda)) = \lambda \cdot \varphi(I) = \lambda, \\ \varphi(L_{ij}(\lambda)) &= \varphi(IL_{ij}(\lambda)) = \varphi(I) = 1, \\ \varphi(T_{ij}) &= \varphi(IT_{ij}) = -\varphi(I) = -1. \end{aligned}$$

□

Lemma E. Let φ be a determinant function. For elementary matrices $E_1, \ldots, E_k$ we have

$$\varphi(E_1 E_2 \cdots E_k) = \varphi(E_1)\varphi(E_2) \cdots \varphi(E_k).$$

Proof. By applying Lemma D, we can rephrase Lemma C as saying that

$$\varphi(AE) = \varphi(A)\varphi(E) \text{ for any elementary matrix } E.$$

If $E_1, \ldots, E_k$ are elementary matrices, then it follows by induction that

$$\begin{aligned} \varphi(E_1 \cdots E_k) &= \varphi(E_1 \cdots E_{k-1})\varphi(E_k) \\ &= \varphi(E_1) \cdots \varphi(E_{k-1})\varphi(E_k). \end{aligned}$$

□

Proof of the Theorem. Let δ_1 and δ_2 be any two determinant functions. Our goal is to show that $\delta_1 = \delta_2$. If A is not invertible then the columns of A are dependent and it follows from Lemma B that $\delta_1(A) = 0 = \delta_2(A)$. So let us suppose that A is invertible. In this case we can apply column operations to reduce A to the identity matrix:

$$AE_1E_2 \cdots E_k = I.$$

Since elementary matrices are invertible, this becomes

$$A = E_k^{-1} \cdots E_1^{-1}.$$

[85] This is our first and only use of property (3).

If E is elementary then E^{-1} is also elementary, so Lemma D implies that $\delta_1(E^{-1}) = \delta_2(E^{-1})$. Finally, by Lemma E we have

$$\begin{aligned}\delta_1(A) &= \delta_1(E_k^{-1} \cdots E_1^{-1}) \\ &= \delta_1(E_k^{-1}) \cdots \delta_1(E_1^{-1}) \\ &= \delta_2(E_k^{-1}) \cdots \delta_2(E_1^{-1}) \\ &= \delta_2(E_k^{-1} \cdots E_1^{-1}) \\ &= \delta_2(A).\end{aligned}$$

□

Thus we have proved that there exists at most one determinant function. From this point on, we will use the notation $\det(A)$ to refer to this function.

We end this section by giving a new criterion for invertibility of square matrices.

Theorem. For any square matrix A we have

$$A \text{ is invertible} \quad \Longleftrightarrow \quad \det(A) \neq 0.$$

Proof. If A is not invertible then A has dependent columns and it follows from Lemma B that $\det(A) = 0$. Conversely, suppose that A is invertible. In the previous chapter we showed that a square matrix is invertible if and only if its Reduced Row Echelon Form is an identity matrix, so that

$$E_k \cdots E_2 E_1 A = I$$

for some elementary matrices $E_1, \ldots, E_k$. From Lemma E it follows that

$$\begin{aligned}A &= E_1^{-1} E_2^{-1} \cdots E_k^{-1} \\ \det(A) &= \det(E_1^{-1}) \det(E_2^{-1}) \cdots \det(E_k^{-1}) \neq 0.\end{aligned}$$

□

Note that we only use elementary matrices $D_i(\lambda)$ with $\lambda \neq 0$ so that $\det(E) \neq 0$ for every elementary matrix E.

5.3 Algebraic Properties of the Determinant

In the previous section we studied the application of determinant functions to elementary matrices, and we used this to prove that there exists at most one determinant function. In this section we will apply the same lemmas to prove some interesting algebraic properties of determinants. Only in the next section will we finally prove that determinants exist!

Theorem. For any square matrices A and B we have

(a) $\det(A^T) = \det(A)$,

(b) $\det(AB) = \det(A)\det(B)$,

(c) $\det(A^{-1}) = 1/\det(A)$.

Proof. (a): Note that A^T is invertible if and only if A is invertible, hence $\det(A^T) = 0$ if and only if $\det(A) = 0$. If $\det(A) \neq 0$ then A is invertible and we can write

$$A = E_1 \cdots E_k$$

for some elementary matrices $E_1, \ldots, E_k$. Note that the transpose E^T of an elementary matrix E is also elementary, and from Lemma C we have $\det(E^T) = \det(E)$. It follows that

$$\begin{aligned} A^T &= E_k^T \cdots E_1^T \\ \det(A^T) &= \det(E_k^T \cdots E_1^T) \\ &= \det(E_k^T) \cdots \det(E_1^T) \\ &= \det(E_k) \cdots \det(E_1) \\ &= \det(E_1) \cdots \det(E_k) \\ &= \det(E_1 \cdots E_k) \\ &= \det(A). \end{aligned}$$

(b): Note that AB is invertible if and only if both of A and B are invertible, so that $\det(AB) = 0$ if and only if $\det(A)\det(B) = 0$. If $\det(A) \neq 0$ and $\det(B) \neq 0$ then A and B are both invertible, hence we can write

$$\begin{aligned} A &= E_1 \cdots E_k, \\ B &= F_1 \cdots F_\ell, \end{aligned}$$

for some elementary matrices $E_1, \ldots, E_k$ and $F_1, \ldots, F_\ell$. It follows that

$$\begin{aligned} \det(AB) &= \det(E_1 \cdots E_k F_1 \cdots F_\ell) \\ &= \det(E_1) \cdots \det(E_k)\det(F_1) \cdots \det(F_\ell) \\ &= [\det(E_1) \cdots \det(E_k)][\det(F_1) \cdots \det(F_\ell)] \\ &= \det(E_1 \cdots E_k)\det(F_1 \cdots F_\ell) \\ &= \det(A)\det(B). \end{aligned}$$

(c): If A is invertible then $\det(A) \neq 0$ and from (b) we obtain

$$\begin{aligned} A^{-1}A &= I \\ \det(A^{-1}A) &= \det(I) \\ \det(A^{-1})\det(A) &= 1 \\ \det(A^{-1}) &= 1/\det(A). \end{aligned}$$

□

As you see, the elementary matrices are quite useful.

5.4 Formulas for the Determinant

I hope you have developed an appreciation for the remarkable properties of determinants. In this section I will prove that determinants actually exist, and in the next section I will finally tell you what determinants "really are". I guess I could have told you that first, but it didn't fit the narrative.

There are several equivalent ways to define the determinant of an $n \times n$ matrix. If A is not invertible then we must have $\det(A) = 0$, so let us suppose that A is invertible. In this case we can perform row (or column) operations to transform A into the identity matrix, which allows us to write A as a product of elementary matrices:

$$A = E_1 \cdots E_k.$$

Then from Lemma E in Section 2.2 we must have

$$\det(A) = \det(E_1) \cdots \det(E_k),$$

where the determinants of elementary matrices are trivial to compute. You might think we could use this formula to **define** the determinant, but the factorization of A into elementary matrices is not unique, and it's not clear that we wouldn't get different values of $\det(A)$ from different factorizations of A. Essentially this has to do with the uniqueness of the RREF, but I don't want to prove this. Instead I'll just give an example computation.

Computing the Determinant by Elimination. Consider again the matrix

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 1 & 1 & 1 \\ 2 & 4 & 1 \end{pmatrix}.$$

First we perform down-elimination steps to put A in upper triangular form:

$$L_{31}(-2)L_{21}(-1)A = \begin{pmatrix} 1 & 2 & 3 \\ 0 & -1 & -2 \\ 0 & 0 & -5 \end{pmatrix}. \qquad (*)$$

Then we scale the rows to turn the pivots into ones:

$$D_3(-1/5)D_2(-1)L_{31}(-2)L_{21}(-1)A = \begin{pmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{pmatrix}.$$

Then we perform up-elimination to obtain an identity matrix:

$$L_{12}(-2)L_{13}(-3)L_{23}(-1)D_3(-1/5)D_2(-1)L_{31}(-2)L_{21}(-1)A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Taking the elementary matrices to the other side gives

$$\begin{aligned} A &= L_{21}(-1)^{-1}L_{31}(-2)^{-1}D_2(-1)^{-1}D_3(-1/5)^{-1}L_{23}(-1)^{-1}L_{13}(-3)^{-1}L_{12}(-2)^{-1} \\ &= L_{21}(1)L_{31}(2)D_2(-1)D_3(-5)L_{23}(1)L_{13}(3)L_{12}(2), \end{aligned}$$

and taking the determinant of each side gives

$$\begin{aligned} \det(A) &= 1 \cdot 1 \cdot (-1) \cdot (-5) \cdot 1 \cdot 1 \cdot 1 \\ &= 5. \end{aligned}$$

Note that this is the product of the pivot entries in step $(*)$. Hence we could have stopped there. In general, if no row transpositions are required, then the determinant is just the product of the diagonal entries after down-elimination.

Next I will give the traditional definition of the determinant, which expresses it as an "alternating sum" over permutations. After that I will give a recursive formula, which is more useful.

Permutation Definition of the Determinant. Let S_n denote the set of *permutations*, i.e., the set of bijective functions $\{1, \ldots, n\} \to \{1, \ldots, n\}$. It is convenient to express a permutation by listing the sequence of values:

$$\sigma = (\sigma(1), \sigma(2), \ldots, \sigma(n)).$$

Each permutation $\sigma \in S_n$ has a well-defined *sign*, or *parity*:

$$\operatorname{sgn}(\sigma) \in \{1, -1\}.$$

Essentially this tells us the number of swaps necessary to obtain the list $(\sigma(1), \ldots, \sigma(n))$ from the list $(1, \ldots, n)$, or vice versa. The number of swaps is not unique, but it turns out that it is always even, or always odd. For example, we can get from $(1, 2, 3)$ to $(3, 2, 1)$ using 3 swaps:

$$(1, 2, 3) \to (2, 1, 3) \to (2, 3, 1) \to (3, 2, 1),$$

Or we can get there using 5 swaps:

$$(1, 2, 3) \to (1, 3, 2) \to (3, 1, 2) \to (2, 1, 3) \to (2, 3, 1) \to (3, 2, 1).$$

But we could never get there using an even number of swaps.[86] Since we can get from $(1, 2, 3)$ to $(3, 2, 1)$ using only odd numbers of swaps, we define

$$\operatorname{sgn}(3, 2, 1) = -1.$$

[86] It is a bit tricky to prove this so we won't bother. It fits better in a course on "group theory".

Of the $n!$ permutations in S_n, it turns out that exactly half are "even" and half are "odd". For example, here is the sign table for S_3:

σ	$\operatorname{sgn}(\sigma)$
$(1,2,3)$	$+1$
$(2,3,1)$	$+1$
$(3,1,2)$	$+1$
$(1,3,2)$	-1
$(2,1,3)$	-1
$(3,2,1)$	-1

Finally, recall the standard basis of k-forms:

$$\varepsilon_{i_1} \otimes \varepsilon_{i_2} \otimes \cdots \otimes \varepsilon_{i_k} \text{ for all } i_1, \ldots, i_k \in \{1, \ldots, n\}.$$

Then we define the determinant function $\det \in \mathcal{A}^n(\mathbb{R}^n)$ as follows:

$$\boxed{\det = \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) \cdot \varepsilon_{\sigma(1)} \otimes \varepsilon_{\sigma(2)} \otimes \cdots \otimes \varepsilon_{\sigma(n)}.}$$

For example, when $n = 3$, the above table of signs gives

$$\begin{aligned} \det = {} & \varepsilon_1 \otimes \varepsilon_2 \otimes \varepsilon_3 + \varepsilon_2 \otimes \varepsilon_3 \otimes \varepsilon_1 + \varepsilon_3 \otimes \varepsilon_1 \otimes \varepsilon_2 \\ & - \varepsilon_1 \otimes \varepsilon_3 \otimes \varepsilon_2 - \varepsilon_2 \otimes \varepsilon_1 \otimes \varepsilon_3 - \varepsilon_3 \otimes \varepsilon_2 \otimes \varepsilon_1. \end{aligned}$$

Equivalently, if A is an $n \times n$ matrix with ij entry a_{ij} then we define

$$\boxed{\det(A) = \sum_{\sigma \in S_n} \operatorname{sgn}(\sigma) \cdot a_{\sigma(1),1} a_{\sigma(2),2} \cdots a_{\sigma(n),n}.}$$

One can check that this function satisfies the three properties of a determinant function, but to do so requires a more thorough study of permutations than we have time for.

Laplace Expansion. The permutation definition of the determinant is explicit but it's mostly useless. Another, recursive, definition called *Laplace expansion* or *expansion by cofactors* has many applications.

For any $n \times n$ matrix A we let $\hat{A}_{ij}$ denote the $(n-1) \times (n-1)$ matrix obtained from A by deleting the ith row and the jth column. To **expand along the ith row**, we fix some i and then compute

$$\boxed{\det(A) = \sum_j (-1)^{i+j} a_{ij} \det(\hat{A}_{ij}).}$$

To **expand along the jth column** we fix some j and compute

$$\boxed{\det(A) = \sum_i (-1)^{i+j} a_{ij} \det(\hat{A}_{ij}).}$$

One must check that these formulas agree with the permutation definition of the determinant. Alternatively, one could prove that these formulas obey the three rules for determinant functions. But I'm not going to do this. Instead I will just give some examples.

First we compute a general 3×3 determinant by expanding along the second row:

$$\begin{aligned}\det \begin{pmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{pmatrix} &= -a_2 \cdot \det \begin{pmatrix} b_1 & c_1 \\ b_3 & c_3 \end{pmatrix} + b_2 \cdot \det \begin{pmatrix} a_1 & c_1 \\ a_3 & c_3 \end{pmatrix} - c_2 \cdot \det \begin{pmatrix} a_1 & b_1 \\ a_3 & b_3 \end{pmatrix} \\ &= -a_2(b_1c_3 - b_3c_1) + b_2(a_1c_3 - a_3c_1) - c_2(a_1b_3 - a_3b_1) \\ &= a_1b_2c_3 + a_2b_3c_1 + a_3b_1c_2 - a_1b_3c_2 - a_2b_1c_3 - a_3b_2c_1.\end{aligned}$$

Next we expand a specific our favorite matrix along the second column:

$$\begin{aligned}\det \begin{pmatrix} 1 & 2 & 3 \\ 1 & 1 & 1 \\ 2 & 4 & 1 \end{pmatrix} &= -2 \cdot \det \begin{pmatrix} 1 & 1 \\ 2 & 1 \end{pmatrix} + 1 \cdot \det \begin{pmatrix} 1 & 3 \\ 2 & 1 \end{pmatrix} - 4 \cdot \det \begin{pmatrix} 1 & 3 \\ 1 & 1 \end{pmatrix} \\ &= -2(1-2) + 1(1-6) - 4(1-3) \\ &= -2(-1) + 1(-5) - 4(-2) \\ &= 2 - 5 + 8 \\ &= 5.\end{aligned}$$

And also along the first row:

$$\begin{aligned}\det \begin{pmatrix} 1 & 2 & 3 \\ 1 & 1 & 1 \\ 2 & 4 & 1 \end{pmatrix} &= 1 \cdot \det \begin{pmatrix} 1 & 1 \\ 4 & 1 \end{pmatrix} - 2 \cdot \det \begin{pmatrix} 1 & 1 \\ 2 & 1 \end{pmatrix} + 3 \cdot \det \begin{pmatrix} 1 & 1 \\ 2 & 4 \end{pmatrix} \\ &= 1(1-4) - 2(1-2) + 3(4-2) \\ &= 1(-3) - 2(-1) + 3(2) \\ &= -3 + 2 + 6 \\ &= 5.\end{aligned}$$

5.5 Cramer's Rule (Optional)

While we're on the subject, there is a famous trick relating determinants to solutions of linear systems. Let A be a square $n \times n$ matrix and consider the linear system

$$A\mathbf{x} = \mathbf{b}$$

$$\begin{pmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{n1} & \cdots & a_{nn} \end{pmatrix} \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix}.$$

Assume that A is invertible, so the system has a unique solution $\mathbf{x} = (x_1, \ldots, x_n)$. Then the ith coordinate of the solution is given by

$$\boxed{x_i = \frac{\det(\hat{A}_i(\mathbf{b}))}{\det(A)},}$$

where $\hat{A}_i(\mathbf{b})$ is the matrix obtained from A by replacing its ith column with $\mathbf{b}$:

$$\hat{A}_i(\mathbf{b}) = \left(\begin{array}{c|c|c|c|c|c|c} \mathbf{a}_1 & \cdots & \mathbf{a}_{i-1} & \mathbf{b} & \mathbf{a}_{i+1} & \cdots & \mathbf{a}_n \end{array} \right).$$

Proof. Consider the matrix

$$X_i := \hat{I}_i(\mathbf{x}) = \left(\begin{array}{c|c|c|c|c|c|c} \mathbf{e}_1 & \cdots & \mathbf{e}_{i-1} & \mathbf{x} & \mathbf{e}_{i+1} & \cdots & \mathbf{e}_n \end{array} \right) = \begin{pmatrix} 1 & & x_1 & & \\ & 1 & \vdots & & \\ & & x_i & & \\ & & \vdots & 1 & \\ & & x_n & & 1 \end{pmatrix}.$$

By Laplace expansion along the ith column, we observe that[87]

$$\det(X_i) = (-1)^{i+i} x_i \det(I_{n-1}) = x_i.$$

Next we observe that

$$\begin{aligned} AX_i &= A \left(\begin{array}{c|c|c|c|c|c|c} \mathbf{e}_1 & \cdots & \mathbf{e}_{i-1} & \mathbf{x} & \mathbf{e}_{i+1} & \cdots & \mathbf{e}_n \end{array} \right) \\ &= \left(\begin{array}{c|c|c|c|c|c|c} A\mathbf{e}_1 & \cdots & A\mathbf{e}_{i-1} & A\mathbf{x} & A\mathbf{e}_{i+1} & \cdots & A\mathbf{e}_n \end{array} \right) \\ &= \left(\begin{array}{c|c|c|c|c|c|c} \mathbf{a}_1 & \cdots & \mathbf{a}_{i-1} & \mathbf{b} & \mathbf{a}_{i+1} & \cdots & \mathbf{a}_n \end{array} \right) \\ &= \hat{A}_i(\mathbf{b}), \end{aligned}$$

and hence

$$\begin{aligned} AX_i &= \hat{A}_i(\mathbf{b}) \\ \det(A)\det(X_i) &= \det(\hat{A}_i(\mathbf{b})) \\ \det(X_i) &= \det(\hat{A}_i(\mathbf{b}))/\det(A) \\ x_i &= \det(\hat{A}_i(\mathbf{b}))/\det(A). \end{aligned}$$

□

For example, let A be the 3×3 matrix from the previous section and consider the linear system

$$\begin{pmatrix} 1 & 2 & 3 \\ 1 & 1 & 1 \\ 2 & 4 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}.$$

[87] The matrix obtained by deleting the ith row and column of X_i is the $(n-1)\times(n-1)$ identity matrix I_{n-1}. Every other $(n-1)\times(n-1)$ matrix in the expansion has a row (also a column) of zeros, hence its determinant is zero.

Then we have

$$x_1 = \det\begin{pmatrix}1&2&3\\0&1&1\\0&4&1\end{pmatrix} \Big/ \det\begin{pmatrix}1&2&3\\1&1&1\\2&4&1\end{pmatrix} = \frac{-3}{5},$$

$$x_2 = \det\begin{pmatrix}1&1&3\\1&0&1\\2&0&1\end{pmatrix} \Big/ \det\begin{pmatrix}1&2&3\\1&1&1\\2&4&1\end{pmatrix} = \frac{1}{5},$$

$$x_3 = \det\begin{pmatrix}1&2&1\\1&1&0\\2&4&0\end{pmatrix} \Big/ \det\begin{pmatrix}1&2&3\\1&1&1\\2&4&1\end{pmatrix} = \frac{2}{5}.$$

Cramer's Rule is useful when we want to pick out a specific coordinate of the solution. We can use this idea to give an explicit formula for the entries of an inverse matrix. Let A be an invertible $n \times n$ square matrix and let $X = \begin{pmatrix} \mathbf{x}_1 \,\big|\, \cdots \,\big|\, \mathbf{x}_n \end{pmatrix}$ be a matrix whose columns $\mathbf{x}_1, \ldots, \mathbf{x}_n \in \mathbb{R}^n$ are unknown vectors. If X is the inverse of A then we must have

$$\begin{aligned} AX &= I \\ A\begin{pmatrix} \mathbf{x}_1 \,\big|\, \cdots \,\big|\, \mathbf{x}_n \end{pmatrix} &= \begin{pmatrix} \mathbf{e}_1 \,\big|\, \cdots \,\big|\, \mathbf{e}_n \end{pmatrix} \\ \begin{pmatrix} A\mathbf{x}_1 \,\big|\, \cdots \,\big|\, A\mathbf{x}_n \end{pmatrix} &= \begin{pmatrix} \mathbf{e}_1 \,\big|\, \cdots \,\big|\, \mathbf{e}_n \end{pmatrix}, \end{aligned}$$

which is equivalent to n matrix equations: $A\mathbf{x}_i = \mathbf{e}_i$ for each i. Let x_{ij} be the ij entry of the unknown matrix X, which is the ith entry of the jth column vector $\mathbf{x}_j$. Then Cramer's Rule says that

$$\begin{aligned} x_{ij} &= i\text{th coordinate of } \mathbf{x}_j \\ &= i\text{th coordinate of the solution to } A\mathbf{x}_j = \mathbf{e}_j \\ &= \det(\hat{A}_i(\mathbf{e}_j))/\det(A), \end{aligned}$$

where $\hat{A}_i(\mathbf{e}_j)$ is the matrix obtained from A by replacing its ith column with $\mathbf{e}_j$. By Laplace expansion along the ith column we have

$$\det(\hat{A}_i(\mathbf{e}_j)) = (-1)^{i+j}\det(\hat{A}_{ji}),$$

where $\hat{A}_{ji}$ is the $(n-1) \times (n-1)$ matrix obtained from A by deleting the jth row and ith column. If $\det(A) \neq 0$ then we conclude that

$$\boxed{(ij \text{ entry of } A^{-1}) = \frac{1}{\det(A)}(-1)^{i+j}\det(\hat{A}_{ji}).}$$

Warning: Note that the positions of i and j are switched in this formula![88]

[88] I have forgotten this many times.

For example, suppose that

$$AX = I,$$
$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Then we have

$$\begin{aligned} x_{11} &= (-1)^{1+1} \det(\hat{A}_{11})/\det(A) = a_{22}/\det(A), \\ x_{12} &= (-1)^{1+2} \det(\hat{A}_{21})/\det(A) = -a_{12}/\det(A), \\ x_{21} &= (-1)^{2+1} \det(\hat{A}_{12})/\det(A) = -a_{21}/\det(A), \\ x_{22} &= (-1)^{2+2} \det(\hat{A}_{22})/\det(A) = a_{11}/\det(A), \end{aligned}$$


which is just the usual formula for the inverse of a 2×2 matrix:

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}.$$

5.6 Geometric Interpretation

My bias is that algebra is based on geometry,[89] hence for me the "true meaning" of the determinant is its geometric interpretation.

Consider two vectors in the plane, $\mathbf{u}, \mathbf{v} \in \mathbb{R}^2$ with angle θ between them. The area of the parallelogram they generate is $\|\mathbf{u}\|\|\mathbf{v}\||\sin\theta|$.[90] Indeed, in the following picture the red parallelogram and the blue rectangle have the same area:



[89] And geometry is based on physics. I believe that physics is the true foundation of mathematics, not axiomatic set theory.

[90] The absolute value accounts for negative angles.

On the other hand, we can interpret this area as a determinant. Let A be the 2×2 matrix with columns $\mathbf{u}$ and $\mathbf{v}$:

$$A = \left(\begin{array}{c|c} \mathbf{u} & \mathbf{v} \end{array} \right).$$

I claim that the area of the parallelogram equals (the absolute value of) the determinant of A. To prove this we use a clever trick. First we observe that

$$\sqrt{\det(A^T A)} = \sqrt{\det(A^T)\det(A)} = \sqrt{\det(A)\det(A)} = \sqrt{\det(A)^2} = |\det(A)|.$$


But the determinant of $A^T A$ can also be computed as follows:

$$\begin{aligned} A^T A &= \left(\begin{array}{c} \mathbf{u}^T \\ \hline \mathbf{v}^T \end{array} \right) \left(\begin{array}{c|c} \mathbf{u} & \mathbf{v} \end{array} \right) \\ A^T A &= \begin{pmatrix} \|\mathbf{u}\|^2 & \mathbf{u} \bullet \mathbf{v} \\ \mathbf{u} \bullet \mathbf{v} & \|\mathbf{v}\|^2 \end{pmatrix} \\ \det(A^T A) &= \|\mathbf{u}\|^2 \|\mathbf{v}\|^2 - (\mathbf{u} \bullet \mathbf{v})^2 \\ &= \|\mathbf{u}\|^2 \|\mathbf{v}\|^2 - (\|\mathbf{u}\| \|\mathbf{v}\| \cos\theta)^2 \\ &= \|\mathbf{u}\|^2 \|\mathbf{v}\|^2 (1 - \cos^2\theta) \\ &= \|\mathbf{u}\|^2 \|\mathbf{v}\|^2 \sin^2\theta. \end{aligned}$$

So we conclude that

$$|\det(A)| = \sqrt{\det(A^T A)} = \sqrt{\|\mathbf{u}\|^2 \|\mathbf{v}\|^2 \sin^2\theta} = \|\mathbf{u}\| \|\mathbf{v}\| |\sin\theta|.$$

This trick is much more important than it looks. Suppose now that our parallelogram lives in n-dimensional space, generated by vectors $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ with angle θ:




For geometric reasons, the area of the parallelogram is still $\|\mathbf{u}\|\|\mathbf{v}\||\sin\theta|$, but now the $n \times 2$ matrix $A = \begin{pmatrix} \mathbf{u} \mid \mathbf{v} \end{pmatrix}$ is **not square**, so $\det(A)$ is not defined. However, the matrix A^TA is still square, so we may still consider $\det(A^TA)$, and the same calculation as above shows that

$$\sqrt{\det(A^TA)} = \|\mathbf{u}\|\|\mathbf{v}\||\sin\theta|.$$

In general, we have the following theorem.

Theorem (Geometric Interpretation of the Determinant). Let A be an $n \times k$ matrix with columns $\mathbf{a}_1, \ldots, \mathbf{a}_k \in \mathbb{R}^n$, which generate a *$k$-parallelotope* living in n-dimensional space:



Let $\text{Vol}_k(A)$ denote the volume of this k-parallelotope, measured within the k-dimensional subspace that it spans. We call this the *k-volume* of the k-parallelotope. Then we have[91]

$$\boxed{\text{Vol}_k(A) = \sqrt{\det(A^T A)}.}$$

If $k = n$, then we are measuring the full n-dimensional volume of an n-parallelotope in $\mathbb{R}^n$. In this case the matrix A is square, and we obtain

$$\boxed{\text{Vol}_n(A) = |\det(A)|.}$$

Note that we already proved this theorem in the case $k = 2$. The proof of the general case proceeds in four steps:

(1) For $n \times n$ matrices A we have $\text{Vol}_n(A) = |\det(A)|$.

(2) For $n \times n$ matrices A we have $|\det(A)| = \sqrt{\det(A^T A)}$.

(3) It follows from (1) and (2) that the n-volume of an n-parallelotope in $\mathbb{R}^n$ depends only on the lengths and angles between its generating vectors.

(4) Hence we also have $\text{Vol}_k(A) = \sqrt{\det(A^T A)}$, even when $k \neq n$.

[91] This volume can very well be zero, which happens when the columns of A are not independent. In this case, the k-parallelotope generated by $\mathbf{a}_1, \ldots, \mathbf{a}_k$ is "flat", i.e., it lives in a smaller-dimensional subspace of $\mathbb{R}^n$. For example, a 3-parallelogram generated by dependent vectors is actually some kind of 2-dimensional hexagon. I guess there is a recursive formula for the lower-dimensional volume but I don't want to work it out.

The hardest part is (1), which we will prove below. The proof of (2) is a simple calculation, which was given above. For the proof of (3) let A be $n \times n$. We observe that the ij entry of the $n \times n$ matrix $A^T A$ is

$$\mathbf{a}_i^T \mathbf{a}_j = \mathbf{a}_i \bullet \mathbf{a}_j = \|\mathbf{a}_i\| \|\mathbf{a}_j\| \cos \theta_{ij},$$

where θ_{ij} is the angle between $\mathbf{a}_i$ and $\mathbf{a}_j$. Since from (1) we have

$$\mathrm{Vol}_n(A) = |\det(A)| = \sqrt{\det(A^T A)},$$

and since the entries of $A^T A$ only depend on the lengths $\|\mathbf{a}_i\|$ and angles θ_{ij}, it follows that the volume $\mathrm{Vol}_n(A)$ only depends on the lengths and angles. But now suppose that A is $k \times n$ with columns $\mathbf{a}_1, \ldots, \mathbf{a}_k \in \mathbb{R}^n$. In this case the ij entry of $A^T A$ is still given by

$$\mathbf{a}_i^T \mathbf{a}_j = \mathbf{a}_i \bullet \mathbf{a}_j = \|\mathbf{a}_i\| \|\mathbf{a}_j\| \cos \theta_{ij},$$

hence $\det(A^T A)$ has exactly the same formula in terms of $\|\mathbf{a}_i\|$ and θ_{ij} as it does when A is a $k \times k$ square matrix. Then from the square case we conclude that

$$\mathrm{Vol}_k(A) = \text{some formula involving the lengths } \|\mathbf{a}_i\| \text{ and angles } \theta_{ij} = \sqrt{\det(A^T A)}.$$

This completes the proof, except for part (1). □

Before diving into the proof of (1), we consider the case $k = 3$. The technical name for a 3-parallelogram is a *parallelepiped*.

Volume of a Parallelepiped. Let A be an $n \times 3$ matrix with columns $\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3 \in \mathbb{R}^n$, and let θ_{ij} be the angle between vectors $\mathbf{a}_i$ and $\mathbf{a}_j$, which can be computed via the dot product:

$$\theta_{ij} = \arccos\left(\frac{\mathbf{a}_i \bullet \mathbf{a}_j}{\|\mathbf{a}_i\| \|\mathbf{a}_j\|}\right).$$

Then the volume (i.e., the 3-volume) of the parallelepiped generated by $\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_3$ is given by

$$\begin{aligned}
\mathrm{Vol}_3(A)^2 &= \det(A^T A) \\
&= \det \begin{pmatrix} \|\mathbf{a}_1\|^2 & \mathbf{a}_1 \bullet \mathbf{a}_2 & \mathbf{a}_1 \bullet \mathbf{a}_3 \\ \mathbf{a}_1 \bullet \mathbf{a}_2 & \|\mathbf{a}_2\|^2 & \mathbf{a}_2 \bullet \mathbf{a}_3 \\ \mathbf{a}_1 \bullet \mathbf{a}_3 & \mathbf{a}_2 \bullet \mathbf{a}_3 & \|\mathbf{a}_3\|^2 \end{pmatrix} \\
&= \det \begin{pmatrix} \|\mathbf{a}_1\|^2 & \|\mathbf{a}_1\| \|\mathbf{a}_2\| \cos\theta_{12} & \|\mathbf{a}_1\| \|\mathbf{a}_3\| \cos\theta_{13} \\ \|\mathbf{a}_1\| \|\mathbf{a}_2\| \cos\theta_{12} & \|\mathbf{a}_2\|^2 & \|\mathbf{a}_2\| \|\mathbf{a}_3\| \cos\theta_{23} \\ \|\mathbf{a}_1\| \|\mathbf{a}_3\| \cos\theta_{13} & \|\mathbf{a}_2\| \|\mathbf{a}_3\| \cos\theta_{23} & \|\mathbf{a}_3\|^2 \end{pmatrix},
\end{aligned}$$

which after some simplification becomes

$$\mathrm{Vol}_3(A) = \|\mathbf{a}_1\| \|\mathbf{a}_2\| \|\mathbf{a}_3\| \sqrt{(1 + 2\cos\theta_{12}\cos\theta_{13}\cos\theta_{23} - (\cos^2\theta_{12} + \cos^2\theta_{13} + \cos^2\theta_{23}))}.$$

This formula is much more difficult to derive without determinants.[92]

Proof of (1). For an $n \times n$ matrix A we need to prove that

$$|\det(A)| = \mathrm{Vol}_n(A).$$

Actually, we will prove that

$$\det(A) = \pm \mathrm{Vol}_n(A),$$


where the sign depends on the ordering of the columns, and is not relevant to the geometry. Thus we will show that the determinant can be interpreted as a "signed volume".[93] According to Section 2.2, we only need to show that the function Vol_n from $(\mathbb{R}^n)^n$ to $\mathbb{R}$ satisfies the three rules of a determinant function:

- *Multilinear.* The function $\mathrm{Vol}_n(A)$ is linear in each individual column of A.
- *Alternating.* If A' is obtained from A by switching two columns, then
$$\mathrm{Vol}_n(A') = -\mathrm{Vol}_n(A).$$
- *Normalized.* We have $\mathrm{Vol}_n(I_n) = 1$.

The third property is part of the **definition** of volume. It just says that the unit n-cube has n-volume 1. And we can just assume that the second property is true, since we don't care about the sign of the volume. Thus we only need to show that Vol_n is multilinear.

There is a subtle difficulty here, since to prove a theorem about volume, one must have a formal definition of volume, which we don't. In fact, the most common formal definition of volume is based the determinant! But any proof using this formalization would be circular.

Instead of developing a rigorous "measure theory",[94] we will proceed intuitively. It is intuitively obvious that scaling one of the columns scales the volume by the same amount. For example, doubling one side of a parallelogram doubles the area:




[92] If $n = 3$ then we can also express the volume in terms of the cross product, but doing so breaks the symmetry, and the cross product doesn't generalize to higher dimensions.

[93] This should be familiar from Calculus, since the area under a curve is actually a "signed area", with regions below the x-axis having "negative area". See the next section.

[94] Measure theory is the term for the modern, rigorous, theory of integration.

Thus we only need to show that Vol_n preserves addition in each column. In the case of parallelograms, we need to show that the areas of the parallelograms generated by $\mathbf{u}, \mathbf{w}$ and $\mathbf{v}, \mathbf{w}$ add to the area of the parallelogram generated by $\mathbf{u}+\mathbf{v}$ and $\mathbf{w}$. For example, in the following picture we need to show that the areas of the red and green parallelograms add to the area of the blue parallelogram:



The proof uses the dotted line, which is parallel to $\mathbf{w}$. This line divides the blue parallelogram into two pieces, which have the same areas as the red and green parallelograms. This follows because parallelograms with the same base and height have the same area.

In higher dimensions the scaling argument is still plausible but the addition argument is harder to visualize. Instead of trying to generalize the above picture, we will base our argument on a general geometric principle called Cavalieri's Principle, which we take as an axiom.[95]

Cavalieri's Principle. An *n-prism* in $\mathbb{R}^n$ has the following form. Let $V \subseteq \mathbb{R}^n$ be an $(n-1)$-dimensional subspace. For any subset $S \subseteq V$ and for any vector $\mathbf{a} \in \mathbb{R}^n$ that is **not** in V, we define the "prism over S generated by $\mathbf{a}$":

$$\mathrm{Prism}_S(\mathbf{a}) = \{\mathbf{p} + t\mathbf{a} : \mathbf{p} \in S \text{ and } 0 \leq t \leq 1\}.$$

Then Cavalieri's principle says that


$$\mathrm{Vol}_n(\mathrm{Prism}_S(\mathbf{a})) = \mathrm{Vol}_n(\mathrm{Prism}_S(\mathbf{a}+\mathbf{v})) \quad \text{for any vector } \mathbf{v} \in V.$$

More colloquially:

two prisms with the same base and the same height have the same volume.

Here is a picture:

[95] This principle is often taken as an axiom, for example when deriving the volume of a sphere in $\mathbb{R}^3$ without calculus.



For any $n \times n$ matrix A we will show that applying an elementary matrix of the form $L_{ij}(\lambda)$ to A does not change the volume of the n-parallelotope:

$$\text{Vol}_n(AL_{ij}(\lambda)) = \text{Vol}_n(A).$$

To be precise, let A have columns $\mathbf{a}_1, \ldots, \mathbf{a}_n \in \mathbb{R}^n$ and let S_j be the $(n-1)$-parallelogram living in the $(n-1)$-dimensional subspace $V \subseteq \mathbb{R}^n$ generated by the vectors $\mathbf{a}_1, \ldots, \mathbf{a}_n$, except for $\mathbf{a}_j$. We can view the n-parallelotope generated by A as $\text{Prism}_{S_j}(\mathbf{a}_j)$. If A' is obtained from A by replacing column $\mathbf{a}_j$ by itself plus any vector $\mathbf{v} \in V$, then Cavalieri's principle says

$$\text{Vol}_n(A') = \text{Vol}_n(\text{Prism}_{S_j}(\mathbf{a}_j + \mathbf{v})) = \text{Vol}_n(\text{Prism}_{S_j}(\mathbf{a}_j)) = \text{Vol}_n(A).$$

We are interested in the special case when $\mathbf{v} = \lambda \mathbf{a}_i$ for some $i \neq j$, in which case $A' = AL_{ij}(\lambda)$.

And that's enough about that.

5.7 Application to Calculus

In the previous section we showed that the a determinant can be viewed as the n-volume of an n-parallelotope living in $\mathbb{R}^n$. Now we apply this idea to volumes of arbitrary shapes.

Scaling Factor. Consider an $n \times n$ matrix A with columns $\mathbf{a}_1, \ldots, \mathbf{a}_n \in \mathbb{R}^n$. We can think of A as the linear function $\mathbb{R}^n \to \mathbb{R}^n$ that sends $\mathbf{x} \mapsto A\mathbf{x}$. Hence A sends the unit n-cube generated by the standard basis vectors $\mathbf{e}_1, \ldots, \mathbf{e}_n$ to the n-parallelotope generated by the vectors $A\mathbf{e}_i = \mathbf{a}_i$. Since the unit n-cube has volume 1 (by definition), we see that

$$\begin{aligned} \text{Vol}_n(A) &= |\det(A)| \\ \text{Vol}_n(A) &= |\det(A)| \cdot 1 \\ \text{Vol}_n(n\text{-parallelotope generated by } \mathbf{a}_1, \ldots, \mathbf{a}_n) &= |\det(A)| \cdot \text{Vol}_n(\text{unit } n\text{-cube}). \end{aligned}$$

More generally, consider an $n \times n$ matrix B with columns $\mathbf{b}_1, \ldots, \mathbf{b}_n \in \mathbb{R}^n$. Then A sends the n-parallelotope generated by $\mathbf{b}_1, \ldots, \mathbf{b}_n$ to the n-parallelotope generated by $A\mathbf{b}_1, \ldots, A\mathbf{b}_n$, which are the columns of AB. Hence we have

$$\begin{aligned}
&\text{Vol}_n(\text{image of the parallelotope } \mathbf{b}_1, \ldots, \mathbf{b}_n \text{ under the function } A) \\
&\qquad = \text{Vol}_n(\text{parallelotope generated by } A\mathbf{b}_1, \ldots, A\mathbf{b}_n) \\
&\qquad = \text{Vol}_n(A\mathbf{b}_1, \ldots, A\mathbf{b}_n) \\
&\qquad = \text{Vol}_n(AB) \\
&\qquad = |\det(AB)| \\
&\qquad = |\det(A)\det(B)| \\
&\qquad = |\det(A)| \cdot |\det(B)| \\
&\qquad = |\det(A)| \cdot \text{Vol}_n(B) \\
&\qquad = |\det(A)| \cdot \text{Vol}_n(\text{parallelotope } \mathbf{b}_1, \ldots, \mathbf{b}_n).
\end{aligned}$$

For example, the unit n-cube corresponds to the identity matrix $B = I_n$. It is also worth mentioning the case when $A = \lambda I_n$ for some scalar λ, so that A is the function that dilates $\mathbb{R}^n$ by a factor of λ. In this case we have[96]

$$\det(\lambda I_n) = \lambda^n,$$

so the function A scales volumes in $\mathbb{R}^n$ by a factor of λ^n. Indeed, if you double the side length of a cube in $\mathbb{R}^3$ then its volume gets multiplied by $8 = 2^3$.

We can think of a square matrix A in two ways. If we think of it as a collection of numbers then the determinant is the (signed) volume of the parallelotope generated by the columns. On the other hand, if we think of A as a linear function $\mathbb{R}^n \to \mathbb{R}^n$ then we should think of $|\det(A)|$ as a "volume scaling factor". Indeed, we have shown that applying A to any parallelotope in $\mathbb{R}^n$ scales its volume by $|\det(A)|$. I claim that the same idea holds for arbitrary[97] subsets of $\mathbb{R}^n$. To be precise, for any subset $S \subseteq \mathbb{R}^n$ we define the image set

$$A(S) := \{\text{the set of points } A\mathbf{p} \text{ for all } \mathbf{p} \in S\}.$$


In this case I claim that

$$\boxed{\text{Vol}_n(A(S)) = |\det(A)| \cdot \text{Vol}_n(S).}$$

The idea of the proof is that any reasonable subset of $\mathbb{R}^n$ can be approximated as a union of tiny parallelotopes. To simplify the discussion we will use tiny cubes. Suppose that the set $S \subseteq \mathbb{R}^n$ is a union of tiny cubes. Then the image $A(S) \subseteq \mathbb{R}^n$ is a union of tiny parallelotopes, each of whose volume has been scaled by $|\det(A)|$. But the total volume is just the sum of the volumes of the tiny pieces. Hence the total volume is also scaled by $|\det(A)|$. Here is a picture:

[96] This follows from multilinearity. Multiplying one column by λ multiplies the determinant λ. Multiplying each of the n columns by λ multiplies the determinant by λ^n.

[97] Arbitrary "measurable" subsets. The real numbers are wild enough that they admit pathological examples such as "sets whose volume cannot be defined". I am happy to ignore such things.



Thinking of determinants as volume scaling factors of linear functions gives an intuitive explanation for the identity $\det(AB) = \det(A)\det(B)$. Indeed, for any subset $S \subseteq \mathbb{R}^n$ and for any $n \times n$ matrices A, B we have

$$\mathrm{Vol}_n((AB)(S)) = |\det(AB)| \cdot \mathrm{Vol}_n(S),$$

but also

$$\begin{aligned}\mathrm{Vol}_n((AB)(S)) &= \mathrm{Vol}_n(A(B(S)) \\ &= |\det(A)| \cdot \mathrm{Vol}_n(B(S)) \\ &= |\det(A)| \cdot |\det(B)| \cdot \mathrm{Vol}_n(S),\end{aligned}$$

which implies that $|\det(AB)| = |\det(A)| \cdot |\det(B)|$. (The sign is a bit trickier to handle.) This idea also gives meaning to the determinant of an abstract linear function $f : V \to V$, independent of choosing a basis for V.

Linear Approximation. We have seen that a linear function $A : \mathbb{R}^n \to \mathbb{R}^n$ scales the n-volume of an arbitrary shape $S \subseteq \mathbb{R}^n$ by a factor of $|\det(A)|$. In this section we will generalize from linear to **non-linear functions**.

A general function $\mathbf{r} : \mathbb{R}^m \to \mathbb{R}^n$ has the form

$$\mathbf{r}(x_1, \ldots, x_n) = \mathbf{r}(\mathbf{x}) = (\mathbf{r}_1(\mathbf{x}), \ldots, \mathbf{r}_m(\mathbf{x})),$$

where each component function $\mathbf{r}_i(x_1, \ldots, x_n)$ sends $\mathbb{R}^n \to \mathbb{R}$. Suppose that each $\mathbf{r}_i$ has continuous partial derivatives near some point $\mathbf{p} \in \mathbb{R}^n$, and consider the Taylor expansion:

$$\mathbf{r}_i(\mathbf{p} + \mathbf{x}) = \mathbf{r}_i(\mathbf{p}) + (\nabla \mathbf{r}_i)_{\mathbf{p}}^T \mathbf{x} + \text{higher terms},$$

where the higher terms are small when $\mathbf{x}$ is close to $\mathbf{0}$. Then we collect the components into a column vector:

$$\begin{aligned}
\mathbf{r}(\mathbf{p}+\mathbf{x}) &= \begin{pmatrix} \mathbf{r}_1(\mathbf{p}+\mathbf{x}) \\ \vdots \\ \mathbf{r}_m(\mathbf{p}+\mathbf{x}) \end{pmatrix} \\
&\approx \begin{pmatrix} \mathbf{r}_1(\mathbf{p}) + (\nabla \mathbf{r}_1)_{\mathbf{p}}^T \mathbf{x} \\ \vdots \\ \mathbf{r}_m(\mathbf{p}) + (\nabla \mathbf{r}_m)_{\mathbf{p}}^T \mathbf{x} \end{pmatrix} \\
&= \begin{pmatrix} \mathbf{r}_1(\mathbf{p}) \\ \vdots \\ \mathbf{r}_m(\mathbf{p}) \end{pmatrix} + \begin{pmatrix} (\nabla \mathbf{r}_1)_{\mathbf{p}}^T \mathbf{x} \\ \vdots \\ (\nabla \mathbf{r}_m)_{\mathbf{p}}^T \mathbf{x} \end{pmatrix} \\
&= \mathbf{r}(\mathbf{p}) + \begin{pmatrix} (\nabla \mathbf{r}_1)_{\mathbf{p}}^T \\ \vdots \\ (\nabla \mathbf{r}_m)_{\mathbf{p}}^T \end{pmatrix} \mathbf{x} \\
&= \mathbf{r}(\mathbf{p}) + \begin{pmatrix} \frac{\partial \mathbf{r}_1}{\partial x_1}(\mathbf{p}) & \cdots & \frac{\partial \mathbf{r}_1}{\partial x_n}(\mathbf{p}) \\ \vdots & & \vdots \\ \frac{\partial \mathbf{r}_m}{\partial x_1}(\mathbf{p}) & \cdots & \frac{\partial \mathbf{r}_m}{\partial x_n}(\mathbf{p}) \end{pmatrix} \mathbf{x}.
\end{aligned}$$

The $m \times n$ matrix of partial derivatives of the components of $\mathbf{r}$ is called the *Jacobian matrix*:

$$J\mathbf{r} := \begin{pmatrix} \partial \mathbf{r}_1/\partial x_1 & \cdots & \partial \mathbf{r}_1/\partial x_n \\ \vdots & & \vdots \\ \partial \mathbf{r}_m/\partial x_1 & \cdots & \partial \mathbf{r}_m/\partial x_n \end{pmatrix}.$$

This matrix plays the role of the "linear part" of the multi-multivariable Taylor expansion:

$$\mathbf{r}(\mathbf{p}+\mathbf{x}) = \mathbf{r}(\mathbf{p}) + (J\mathbf{r})_{\mathbf{p}}\mathbf{x} + \text{higher terms}.$$

In summary, suppose that a **possibly non-linear function** $\mathbf{r} : \mathbb{R}^n \to \mathbb{R}^m$ behaves nicely near a point $\mathbf{p} \in \mathbb{R}^n$. Then near this point the function $\mathbf{r}$ is approximately equal to the **linear function** corresponding to the $m \times n$ Jacobian matrix $(J\mathbf{r})_{\mathbf{p}}$. If $\mathbf{r}$ happens to be linear, corresponding to an $m \times n$ matrix A, then one can check that $(J\mathbf{r})_{\mathbf{p}} = A$ for any point $\mathbf{p}$. If $\mathbf{r}$ is non-linear then the matrix $(J\mathbf{r})_{\mathbf{p}}$ changes from point to point.

Application to Integration. In the previous sections we showed the following:

- If a function $\mathbf{r} : \mathbb{R}^m \to \mathbb{R}^n$ has continuous partial derivatives near a point $\mathbf{p} \in \mathbb{R}^n$ then we can approximate $\mathbf{r}$ near $\mathbf{p}$ by an $m \times n$ matrix $(J\mathbf{r})_{\mathbf{p}}$.
- A linear function $A : \mathbb{R}^n \to \mathbb{R}^n$ scales volume by the factor $|\det(A)|$.

Combining these ideas gives us a method to compute the volumes of parametrized shapes in $\mathbb{R}^n$. Before showing some examples, I will state the general theorem.

Theorem (Volume of a k-dimensional submanifold of $\mathbb{R}^n$). We wish to compute the k-volume of a k-dimensional subset $T \subseteq \mathbb{R}^n$. To do this, we look for a *parametrization function* $\mathbf{r} : \mathbb{R}^k \to \mathbb{R}^n$ whose image is T. Suppose that $\mathbf{r}$ sends the subset $S \subseteq \mathbb{R}^k$ to the subset $T \subseteq \mathbb{R}^n$. Then we can compute[98] the k-volume of T by integrating a suitable "volume stretch factor" over the region $S \subseteq \mathbb{R}^k$ using standard Euclidean coordinates:

$$\boxed{\mathrm{Vol}_k(T) = \mathrm{Vol}_k(\mathbf{r}(S)) = \int_{\mathbf{p} \in S} \sqrt{\det((J\mathbf{r})_{\mathbf{p}}^T (J\mathbf{r})_{\mathbf{p}})} \cdot d\mathbf{p}.}$$

Remark: We require the shapes S, T and the function $\mathbf{r}$ to be sufficiently nice. This involves several technical conditions that I am happy to ignore. Basically, S and T should be reasonably smooth, and $\mathbf{r}$ should parametrize T without any overlaps or kinks.

Proof. A tiny cube at the point $\mathbf{p} \in S$ has a tiny volume $d\mathbf{p}$. The function $\mathbf{r}$ is approximately linear at $\mathbf{p}$, given by the $n \times k$ matrix $(J\mathbf{r})_{\mathbf{p}}$. This matrix sends the tiny cube at the point $\mathbf{p}$ to a tiny k-parallelotope at the point $\mathbf{r}(\mathbf{p})$. For any small shape near $\mathbf{p}$, the linear function $(J\mathbf{r})_{\mathbf{p}}$ scales its volume by a factor of[99]

$$\sqrt{\det((J\mathbf{r})_{\mathbf{p}}^T (J_{\mathbf{r}}))}.$$

Hence the volume of the tiny k-parallelotope at the point $\mathbf{r}(\mathbf{p})$ is

$$\sqrt{\det((J\mathbf{r})_{\mathbf{p}}^T (J_{\mathbf{r}}))} \cdot (\text{volume of the tiny cube}) = \sqrt{\det((J\mathbf{r})_{\mathbf{p}}^T (J_{\mathbf{r}}))} \cdot d\mathbf{p}.$$

Then we just add up all these tiny volumes to get the k-volume of T. □

To end this section, I will illustrate how this result unifies several formulas from Calculus III.

Example: Arc Length. Let $\mathbf{r} : \mathbb{R} \to \mathbb{R}^n$ be a parametrized path in $\mathbb{R}^n$. Usually we think of the parameter as time, and we write $\mathbf{r}(t) = (x_1(t), x_2(t), \ldots, x_n(t))$. The Jacobian matrix at time t is just the velocity vector:

$$J\mathbf{r}(t) = \begin{pmatrix} \partial x_1/\partial t \\ \vdots \\ \partial x_n/\partial t \end{pmatrix} = \mathbf{r}'(t).$$

In this case, $(J\mathbf{r})^T (J\mathbf{r})$ is just a scalar, and the 1-volume (i.e., length) scaling factor is just the speed of the parametrization:

$$(J\mathbf{r})^T (J\mathbf{r}) = \mathbf{r}'(t)^T \mathbf{r}(t)$$

[98] In fact, this formula is often used as the **definition** of volume.

[99] We only proved this in the case $k = n$, when $(J_{\mathbf{r}})_{\mathbf{p}}$ is a square matrix and the scaling factor reduces to $|\det((J\mathbf{r})_{\mathbf{p}})|$. The general case follows by the same argument as in 2.6.

$$(J\mathbf{r})^T(J\mathbf{r}) = \|\mathbf{r}'(t)\|^2$$
$$\det((J\mathbf{r})^T(J\mathbf{r})) = \|\mathbf{r}'(t)\|^2$$
$$\sqrt{\det((J\mathbf{r})^T(J\mathbf{r}))} = \|\mathbf{r}'(t)\|.$$

Then the theorem tells us that the arc length of the curve is just the integral of the speed:


$$(\text{length of the curve } \mathbf{r}(t) \text{ between times } t = a \text{ and } t = b) = \int_a^b \|\mathbf{r}'(t)\|\, dt.$$

Of course this makes sense because distance is the time integral of speed.

Example: Surface Area. Let $\mathbf{r} : \mathbb{R}^2 \to \mathbb{R}^n$ be a parametrization for a 2-dimensional surface $T \subseteq \mathbb{R}^n$. It is common to write $\mathbf{r}(u, v) = (x_1(u, v), \ldots, x_n(u, v))$, where each coordinate x_i is a function from $\mathbb{R}^2$ to $\mathbb{R}$. The Jacobian matrix is

$$J\mathbf{r} = \begin{pmatrix} \partial x_1/\partial u & \partial x_1/\partial v \\ \vdots & \vdots \\ \partial x_n/\partial u & \partial x_n/\partial v \end{pmatrix} = \begin{pmatrix} | & | \\ \mathbf{r}_u & \mathbf{r}_v \\ | & | \end{pmatrix},$$

where $\mathbf{r}_u$ and $\mathbf{r}_v$ are the "velocity vectors" of $\mathbf{r}$ in the u and v directions[100]



[100] If one of u or v is fixed then you can think of the other as time.

In this case the 2-volume (i.e., area) scaling factor is the area of the parallelogram generated by $\mathbf{r}_u$ and $\mathbf{r}_v$:

$$\begin{aligned}
(J\mathbf{r})^T(J\mathbf{r}) &= \begin{pmatrix} - & \mathbf{r}_u & - \\ - & \mathbf{r}_v & - \end{pmatrix} \begin{pmatrix} | & | \\ \mathbf{r}_u & \mathbf{r}_v \\ | & | \end{pmatrix} \\
&= \begin{pmatrix} \|\mathbf{r}_u\|^2 & \mathbf{r}_u \bullet \mathbf{r}_v \\ \mathbf{r}_u \bullet \mathbf{r}_v & \|\mathbf{r}_v\|^2 \end{pmatrix} \\
\det((J\mathbf{r})^T(J\mathbf{r})) &= \|\mathbf{r}_u\|^2\|\mathbf{r}_v\|^2 - (\mathbf{r}_u \bullet \mathbf{r}_v)^2 \\
&= \|\mathbf{r}_u\|^2\|\mathbf{r}_v\|^2 - (|\mathbf{r}_u\|\|\mathbf{r}_v\| \cos\theta_{uv})^2 \\
&= \|\mathbf{r}_u\|^2\|\mathbf{r}_v\|^2(1 - \cos^2\theta_{uv}) \\
&= \|\mathbf{r}_u\|^2\|\mathbf{r}_v\|^2 \sin^2\theta_{uv} \\
\sqrt{\det((J\mathbf{r})^T(J\mathbf{r}))} &= \|\mathbf{r}_u\|\|\mathbf{r}_v\|\,|\sin\theta_{uv}|,
\end{aligned}$$

where θ_{uv} is the angle between the velocity vectors $\mathbf{r}_u$ and $\mathbf{r}_v$. In the special case of a surface in $\mathbb{R}^3$, we can also describe this area as the length of the cross product vector:


$$\|\mathbf{r}_u \times \mathbf{r}_v\| = \|\mathbf{r}_u\|\|\mathbf{r}_v\|\,|\sin\theta_{uv}|.$$

To compute the area of the surface, we add up all of the areas of tiny parallelograms:

$$(\text{area of the surface } T \subseteq \mathbb{R}^n) = \int \sqrt{\|\mathbf{r}_u\|^2\|\mathbf{r}_v\|^2 - (\mathbf{r}_u \bullet \mathbf{r}_v)^2} \cdot du dv.$$

Example: Change of Coordinates. A parametrization of an n-dimensional shape in n-dimensional space is sometimes viewed as a "change of coordinates" $\mathbf{r} : \mathbb{R}^n \to \mathbb{R}^n$. For example, take the parametrization of $\mathbb{R}^2$ by polar coordinates:

$$\mathbf{r}(r,\theta) = \begin{pmatrix} x(r,\theta) \\ y(r,\theta) \end{pmatrix} = \begin{pmatrix} r\cos\theta \\ r\sin\theta \end{pmatrix}.$$



The Jacobian stretch factor at the point (r, θ) is

$$\begin{aligned} J\mathbf{r} &= \begin{pmatrix} \partial x/\partial r & \partial x/\partial \theta \\ \partial y/\partial r & \partial y/\partial \theta \end{pmatrix} \\ &= \begin{pmatrix} \cos\theta & -r\sin\theta \\ \sin\theta & r\cos\theta \end{pmatrix} \\ \sqrt{\det((J\mathbf{r})^T(J\mathbf{r}))} &= |\det(J\mathbf{r})| \\ &= |r\cos^2\theta + r\sin^2\theta| \\ &= |r|. \end{aligned}$$

Hence the area of a region T in the x, y-plane, which is parametrized by a region S in the r, θ-plane is given by[101]

$$\int_S r \cdot dr d\theta.$$

Since a change of coordinates maps a space into itself, changes of coordinates can be composed. Suppose we have functions $\mathbf{r} : \mathbb{R}^n \to \mathbb{R}^n$ and $\mathbf{s} : \mathbb{R}^n \to \mathbb{R}^n$, with composition $\mathbf{r} \circ \mathbf{s} : \mathbb{R}^n \to \mathbb{R}^n$. The multi-multivariable version of the chain rule says that the Jacobian matrix of the composition $\mathbf{r} \circ \mathbf{s}$ is equal to the product of the Jacobian matrices of $\mathbf{r}$ and $\mathbf{s}$. That is, for any point $\mathbf{p} \in \mathbb{R}^n$ we have

$$(J(\mathbf{r} \circ \mathbf{s}))_{\mathbf{p}} = (J\mathbf{r})_{\mathbf{s}(\mathbf{p})} \cdot (J\mathbf{s})_{\mathbf{p}}.$$

[101] In order to ensure the "niceness" of the parametrization, we will take $r \geq 0$ (so that $|r| = r$) and $0 \leq \theta < 2\pi$.

Hence the Jacobian scaling factors multiply:

$$|\det((J(\mathbf{r}\circ\mathbf{s}))_{\mathbf{p}})| = |\det((J\mathbf{r})_{\mathbf{s}(\mathbf{p})})|\cdot|\det((J\mathbf{s})_{\mathbf{p}})|.$$

Observe that the notation is getting complicated. Indeed, the subject of differential geometry is known for its impenetrable notation. Since no two authors can understand each other, they often invent their own personal notations. Einstein's notation is the most popular among physicists.

6 Eigenvalues and Eigenvectors

6.1 A Motivating Example

In order to motivate the concepts of eigenvalues and eigenvectors I will develop one specific example in detail. Some of the steps will seem miraculous, and only make sense later when we discuss the general theory.

Our example comes from the theory of "Markov chains". Consider the following matrix:

$$A = \begin{pmatrix} .8 & .3 \\ .2 & .7 \end{pmatrix}.$$

This matrix has the special property that each of its columns sums to 1. In matrix notation:[102]


$$\begin{pmatrix} 1 & 1 \end{pmatrix} A = \begin{pmatrix} 1 & 1 \end{pmatrix} \begin{pmatrix} .8 & .3 \\ .2 & .7 \end{pmatrix} = \begin{pmatrix} .8+.2 & .3+.7 \end{pmatrix} = \begin{pmatrix} 1 & 1 \end{pmatrix}.$$

By induction, this implies that every power of A has columns that sum to 1:

$$\begin{aligned} \begin{pmatrix} 1 & 1 \end{pmatrix} A^n &= \begin{pmatrix} 1 & 1 \end{pmatrix} (AA^{n-1}) \\ &= (\begin{pmatrix} 1 & 1 \end{pmatrix} A)A^{n-1} \\ &= \begin{pmatrix} 1 & 1 \end{pmatrix} A^{n-1} \\ &\vdots \\ &= \begin{pmatrix} 1 & 1 \end{pmatrix}. \end{aligned}$$

Such a matrix is called a *Markov matrix* or a *stochastic matrix*. We can interpret the matrix entries as probabilities. Suppose that a certain particle can be in one of two states. At each discrete time step, the particle can change states, according to the following probabilities:

[102] Jargon: Later we will say that $\begin{pmatrix} 1 & 1 \end{pmatrix}$ is a "left eigenvector" of A with "eigenvalue" 1.



That is, if the particle is currently in state 1 then it has an 80% chance to stay in state 1 and a 20% chance to transition to state 2. If the particle is in state 2 then it has a 70% chance to stay and a 30% chance to stay. This is why the columns of A must sum to 1.

Now let's consider the first few powers of A:

$$\begin{aligned} A &= \begin{pmatrix} .8 & .3 \\ .2 & .7 \end{pmatrix}, \\ A^2 &= \begin{pmatrix} .7 & .45 \\ .3 & .55 \end{pmatrix}, \\ A^3 &= \begin{pmatrix} .65 & .525 \\ .35 & .475 \end{pmatrix}, \\ &\vdots \\ A^{10} &= \begin{pmatrix} 0.600390625 & 0.5994140625 \\ 0.399609375 & 0.4005859375 \end{pmatrix}. \end{aligned}$$

Do you see any pattern here? It seems likely that

$$A^n \to \begin{pmatrix} .6 & .6 \\ .4 & .4 \end{pmatrix} \text{ as } n \to \infty,$$

but the entries of the matrices look complicated. Nevertheless, at the end of this section we will obtained exact formulas for the entries of each power A^n.

We have shown that each column of each power A^n sums to 1. This fact has a probabilistic interpretation. Let p_k and q_k be the probabilities that the particle is in state 1 or 2, respectively, after k seconds, and let p_0, q_0 denote the initial probabilities. Then I claim that[103]

$$\boxed{\mathbf{p}_n = \begin{pmatrix} p_n \\ q_n \end{pmatrix} = A \begin{pmatrix} p_{n-1} \\ q_{n-1} \end{pmatrix} = A\mathbf{p}_{n-1}.}$$

[103] Alternatively, suppose that we have an ensemble of particles and let x_n, y_n denote the **expected number** of particles in each state. Then the same theory will hold.

To prove this we use the *law of total probability* (which I won't explain here). Given any two events S and T, we have the following identities:

$$\begin{aligned} P(S) &= P(T)P(S|T) + P(T')P(S|T'), \\ P(S') &= P(T)P(S'|T) + P(T')P(S'|T'). \end{aligned}$$

Let S_n be the event that "the particle is in state 1 after n seconds", so that

$$p_n = P(S_n) \text{ and } q_n = P(S'_n).$$

The transition matrix A tells us that

$$\begin{aligned} P(S_n|S_{n-1}) &= .8, \\ P(S'_n|S_{n-1}) &= .2, \\ P(S_n|S'_{n-1}) &= .3, \\ P(S'_n|S'_{n-1}) &= .7, \end{aligned}$$

which are independent of n. Hence we have

$$\begin{aligned} p_n &= P(S_n) \\ &= P(S_{n-1})P(S_n|S_{n-1}) + P(S'_{n-1})P(S_n|S'_{n-1}) \\ &= p_{n-1}(.8) + q_{n-1}(.3) \end{aligned}$$

and

$$\begin{aligned} q_n &= P(S'_n) \\ &= P(S_{n-1})P(S'_n|S_{n-1}) + P(S'_{n-1})P(S'_n|S'_{n-1}) \\ &= p_{n-1}(.2) + q_{n-1}(.7), \end{aligned}$$

as desired. But enough about probability.

Given the initial distribution $\mathbf{p}_0 = (p_0, q_0)$, the distribution after n seconds is given by

$$\begin{aligned} \mathbf{p}_n &= A\mathbf{p}_{n-1}, \\ &= AA\mathbf{p}_{n-2} \\ &\vdots \\ &= AA\cdots A\mathbf{p}_0 \\ &= A^n\mathbf{p}_0. \end{aligned}$$

Our goal is to find explicit formulas for p_n and q_n in terms of p_0 and q_0.

Now comes the key trick. We have the following mysterious identitites:

$$A\begin{pmatrix}3\\2\end{pmatrix} = \begin{pmatrix}3\\2\end{pmatrix} \quad \text{and} \quad A\begin{pmatrix}1\\-1\end{pmatrix} = \frac{1}{2}\begin{pmatrix}1\\-1\end{pmatrix}. \tag{$*$}$$

Jargon: We say that $(3,2)$ and $(1,-1)$ are "eigenvectors" of A with corresponding "eigenvalues" 1 and $1/2$. More generally, if $A\mathbf{x} = \lambda\mathbf{x}$ for some vector $\mathbf{x}$ and scalar λ, then the action of A^n on $\mathbf{x}$ is easy to compute:

$$\begin{aligned} A^n\mathbf{x} &= (A^{n-1}A)\mathbf{x} \\ &= A^{n-1}(A\mathbf{x}) \\ &= A^{n-1}(\lambda\mathbf{x}) \\ &= \lambda A^{n-1}\mathbf{x} \\ &\vdots \\ &= \lambda^n\mathbf{x}. \end{aligned}$$

Once we know $(*)$, the rest of the solution is straightforward. First we want to express our initial condition $\mathbf{p}_0$ as a linear combination of eigenvectors. In other words, we want to find a and b such that

$$\begin{aligned} \mathbf{p}_0 &= a\begin{pmatrix}3\\2\end{pmatrix} + b\begin{pmatrix}1\\-1\end{pmatrix} \\ &= \begin{pmatrix}3 & 1\\2 & -1\end{pmatrix}\begin{pmatrix}a\\b\end{pmatrix}. \end{aligned}$$

Since the two eigenvectors are not parallel, the matrix of eigenvectors is invertible, hence

$$\begin{aligned} \begin{pmatrix}a\\b\end{pmatrix} &= \begin{pmatrix}3 & 1\\2 & -1\end{pmatrix}^{-1}\begin{pmatrix}p_0\\q_0\end{pmatrix} \\ &= -\frac{1}{5}\begin{pmatrix}-1 & -1\\-2 & 3\end{pmatrix}\begin{pmatrix}p_0\\q_0\end{pmatrix} \\ &= -\frac{1}{5}\begin{pmatrix}-p_0-q_0\\-2p_0+3q_0\end{pmatrix} \\ &= \begin{pmatrix}1/5\\p_0-3/5\end{pmatrix}. && p_0+q_0=1 \end{aligned}$$

Then we obtain the solution:

$$\begin{aligned} \mathbf{p}_0 &= \frac{1}{5}\begin{pmatrix}3\\2\end{pmatrix} + \left(p_0-\frac{3}{5}\right)\begin{pmatrix}1\\-1\end{pmatrix} \\ A^n\mathbf{p}_0 &= \frac{1}{5}A^n\begin{pmatrix}3\\2\end{pmatrix} + \left(p_0-\frac{3}{5}\right)A^n\begin{pmatrix}1\\-1\end{pmatrix} \\ \mathbf{p}_n &= \frac{1}{5}\begin{pmatrix}3\\2\end{pmatrix} + \left(p_0-\frac{3}{5}\right)\left(\frac{1}{2}\right)^n\begin{pmatrix}1\\-1\end{pmatrix} \\ \begin{pmatrix}p_n\\q_n\end{pmatrix} &= \begin{pmatrix}3/5+(p_0-3/5)/2^n\\2/5-(p_0-3/5)/2^n\end{pmatrix}. \end{aligned}$$

As $n \to \infty$ we observe that $\mathbf{p}_n \to (3/5, 2/5)$, regardless of the initial probabilities p_0 and q_0. The fact that the initial condition is irrelevant is sometimes called the "ergodic property" (or the "mixing property").

But we can do more. Suppose that $A\mathbf{x}_1 = \lambda_1 \mathbf{x}_1$ and $A\mathbf{x}_2 = \lambda_2 \mathbf{x}_2$ for some eigenvectors $\mathbf{x}_1, \mathbf{x}_2$ and eigenvalues λ_1, λ_2. We can express both of these conditions simultaneously by forming the matrices

$$X = \left(\begin{array}{c|c} \mathbf{x}_1 & \mathbf{x}_2 \end{array} \right) \quad \text{and} \quad \Lambda = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}.$$

Then we have

$$\begin{aligned} AX &= A \left(\begin{array}{c|c} \mathbf{x}_1 & \mathbf{x}_2 \end{array} \right) \\ &= \left(\begin{array}{c|c} A\mathbf{x}_1 & A\mathbf{x}_2 \end{array} \right) \\ &= \left(\begin{array}{c|c} \lambda_1 \mathbf{x}_1 & \lambda_2 \mathbf{x}_2 \end{array} \right) \\ &= \left(\begin{array}{c|c} \mathbf{x}_1 & \mathbf{x}_2 \end{array} \right) \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \\ &= X\Lambda. \end{aligned}$$

This equation holds even when A is $n \times n$ and X is $n \times 2$. If A is 2×2 and if the vectors $\mathbf{x}_1, \mathbf{x}_2$ are not parallel, then the matrix X is square and invertible, hence

$$\begin{aligned} AX &= X\Lambda \\ A &= X\Lambda X^{-1}. \end{aligned}$$

In this case, we say that we have "diagonalized" the matrix A. In our case, we have

$$\begin{pmatrix} .8 & .3 \\ .2 & .7 \end{pmatrix} = \begin{pmatrix} 3 & 1 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1/2 \end{pmatrix} \begin{pmatrix} 3 & 1 \\ 2 & -1 \end{pmatrix}^{-1}.$$

The powers of A behave well with respect to this factorization. This follows from two key properties. First, the powers of a diagonal matrix are easy to compute:

$$\Lambda^n = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}^n = \begin{pmatrix} \lambda_1^n & 0 \\ 0 & \lambda_2^n \end{pmatrix}.$$

Second, there is a miraculous cancellation in the powers of $X\Lambda X^{-1}$:

$$\begin{aligned} A^n &= (X\Lambda X^{-1})^n \\ &= (X\Lambda X^{-1})(X\Lambda X^{-1}) \cdots (X\Lambda X^{-1}) \\ &= X\Lambda (X^{-1}X)\Lambda (X^{-1}X) \cdots (X^{-1}X)\Lambda X^{-1} \\ &= X\Lambda\Lambda \cdots \Lambda X^{-1} \\ &= X\Lambda^n X^{-1}. \end{aligned}$$

Putting these together gives us explicit formulas for the entries of A^n:

$$\begin{aligned}
\begin{pmatrix} .8 & .3 \\ .2 & .7 \end{pmatrix}^n &= \begin{pmatrix} 3 & 1 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1/2 \end{pmatrix}^n \begin{pmatrix} 3 & 1 \\ 2 & -1 \end{pmatrix}^{-1} \\
&= \begin{pmatrix} 3 & 1 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} 1^n & 0 \\ 0 & (1/2)^n \end{pmatrix} \begin{pmatrix} 3 & 1 \\ 2 & -1 \end{pmatrix}^{-1} \\
&= \begin{pmatrix} 3 & 1 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} 1^n & 0 \\ 0 & (1/2)^n \end{pmatrix} \begin{pmatrix} -\frac{1}{5} \end{pmatrix} \begin{pmatrix} -1 & -1 \\ -2 & 3 \end{pmatrix} \\
&= \begin{pmatrix} 3 & 1 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} 1^n & 0 \\ 0 & (1/2)^n \end{pmatrix} \begin{pmatrix} -\frac{1}{5} \end{pmatrix} \begin{pmatrix} -1 & -1 \\ -2 & 3 \end{pmatrix} \\
&= \frac{1}{5} \begin{pmatrix} 3 & 1 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & (1/2)^n \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 2 & -3 \end{pmatrix} \\
&= \frac{1}{5} \begin{pmatrix} 3 & 1 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 2/2^n & -3/2^n \end{pmatrix} \\
&= \frac{1}{5} \begin{pmatrix} 3+2/2^n & 3-3/2^n \\ 2-2/2^n & 2+3/2^n \end{pmatrix}.
\end{aligned}$$

These exact formulas would be very difficult to obtain without the trick of eigenvalues and eigenvectors. By letting n go to infinity, we confirm our experimental observation that

$$A^n \to \frac{1}{5} \begin{pmatrix} 3 & 3 \\ 2 & 2 \end{pmatrix} = \begin{pmatrix} .6 & .6 \\ .4 & .4 \end{pmatrix} \text{ as } n \to \infty.$$

Finally, let me mention an alternative expression for A^n that is often more useful. For any vectors $\mathbf{x}_1, \mathbf{x}_2, \mathbf{y}_1, \mathbf{y}_2$ and scalars λ_1, λ_2, one can check that

$$\left(\begin{array}{c|c} \mathbf{x}_1 & \mathbf{x}_2 \end{array} \right) \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \left(\begin{array}{c} \mathbf{y}_1^T \\ \hline \mathbf{y}_2^T \end{array} \right) = \lambda_1 \mathbf{x}_1 \mathbf{y}_1^T + \lambda_2 \mathbf{x}_2 \mathbf{y}_2^T.$$

Thus in our case we have

$$\begin{pmatrix} .8 & .3 \\ .2 & .7 \end{pmatrix}^n = \begin{pmatrix} 3 \\ 2 \end{pmatrix} \begin{pmatrix} 1/5 & 1/5 \end{pmatrix} + \left(\frac{1}{2} \right)^n \begin{pmatrix} 1 \\ -1 \end{pmatrix} \begin{pmatrix} 2/5 & -3/5 \end{pmatrix}.$$

This expression emphasizes the fact that A^n converges to a rank 1 matrix:

$$A^n \to \begin{pmatrix} 3 \\ 2 \end{pmatrix} \begin{pmatrix} 1/5 & 1/5 \end{pmatrix} \text{ as } n \to \infty.$$

Remember: This section is just for motivation. I will explain all of the ideas later.

6.2 The Characteristic Polynomial

Let A be a square matrix over $\mathbb{R}$ or $\mathbb{C}$. We say that $\lambda \in \mathbb{C}$ is an *eigenvalue* of A when there exists a nonzero vector $\mathbf{x}$ satisfying $A\mathbf{x} = \lambda\mathbf{x}$. Let me emphasize this:

$$\boxed{\lambda \text{ is an eigenvalue of } A \quad \iff \quad \text{there exists some } \mathbf{x} \neq \mathbf{0} \text{ satisfying } A\mathbf{x} = \lambda\mathbf{x}.}$$

If $A\mathbf{x} = \lambda\mathbf{x}$ then we say that $\mathbf{x}$ is a λ-*eigenvector* of A.

It is not immediately clear that eigenvalues exist. Our first result will show that any matrix has at least one eigenvalue. To do this we will rewrite the definition of eigenvalues in terms of determinants.[104] The following equivalences follow from results in the previous chapter:

$$\begin{aligned}
\lambda \text{ is an eigenvalue of } A &\iff A\mathbf{x} = \lambda\mathbf{x} \text{ for some } x \neq \mathbf{0} \\
&\iff (\lambda\mathbf{x} - A\mathbf{x}) = \mathbf{0} \text{ for some } \mathbf{x} \neq \mathbf{0} \\
&\iff \lambda I\mathbf{x} - A\mathbf{x} = \mathbf{0} \text{ for some } \mathbf{x} \neq \mathbf{0} \\
&\iff (\lambda I - A)\mathbf{x} = \mathbf{0} \text{ for some } \mathbf{x} \neq \mathbf{0} \\
&\iff \text{the nullspace } \mathcal{N}(\lambda I - A) \text{ contains a nonzero vector} \\
&\iff \dim \mathcal{N}(\lambda I - A) \geq 1 \\
&\iff \text{the matrix } \lambda I - A \text{ is not invertible} \\
&\iff \det(\lambda I - A) = 0.
\end{aligned}$$

This last equivalence is the most convenient way to study eigenvalues. In summary,

$$\boxed{\lambda \text{ is an eigenvalue of } A \iff \det(\lambda I - A) = 0.}$$

If λ is an eigenvalue of an $n \times n$ matrix A, we observe that the set of λ-eigenvectors is a subspace of $\mathbb{R}^n$. Indeed, the λ-eigenvectors of A are just the vectors in the nullspace $\mathcal{N}(\lambda I - A)$. We say that

$$\mathcal{N}(\lambda I - A) = \text{the } \lambda\text{-}\textit{eigenspace} \text{ of } A.$$

When λ is **not** an eigenvalue the matrix $\lambda I - A$ is invertible, so in this case the "λ-eigenspace" is trivial: $\mathcal{N}(\lambda I - A) = \{\mathbf{0}\}$. Before going further with the theory, we compute the eigenvalues of a general 2×2 matrix:

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

We have

$$\begin{aligned}
\det(\lambda I - A) &= \det\left(\lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \begin{pmatrix} a & b \\ c & d \end{pmatrix}\right) \\
&= \det \begin{pmatrix} \lambda - a & -b \\ -c & \lambda - d \end{pmatrix} \\
&= (\lambda - a)(\lambda - d) - (-b)(-c) \\
&= \lambda^2 - (a + d)\lambda + (ad - bc).
\end{aligned}$$

Hence the eigenvalues of A are

$$\lambda = \frac{a + d \pm \sqrt{(a + d)^2 - 4(ad - bc)}}{2}.$$

[104] There is a popular textbook called *Linear Algebra Done Right* in which the author goes to great lengths to avoid the use of determinants in the theory of eigenvalues. There is another well-known textbook called *Linear Algebra Done Wrong*, which I greatly prefer.

Let $\Delta = (a+d)^2 - 4(ad-bc)$ denote the discriminant of this quadratic polynomial. If $\Delta = 0$ then the matrix A has only one eigenvalue. Now suppose that A has real entries. If $\Delta > 0$ then A has two distinct real eigenvalues and if $\Delta < 0$ then A has two distinct complex eigenvalues.

After finding the eigenvalues, it is an easy matter to find all of the eigenvectors. Consider our example from the previous section:

$$A = \begin{pmatrix} .8 & .3 \\ .2 & .7 \end{pmatrix}.$$

The eigenvalues are the roots of the polynomial equation

$$\begin{aligned} \det(\lambda I - A) &= 0 \\ \lambda^2 - (.8+.7)\lambda + (.8)(.7) - (.2)(.3) &= 0 \\ \lambda^2 - 1.5\lambda + 0.5 &= 0, \end{aligned}$$

which are

$$\begin{aligned} \lambda &= \frac{1}{2}(1.5 \pm \sqrt{(1.5)^2 - 4(0.5)}) \\ &= \frac{1}{2}(1.5 \pm \sqrt{0.25}) \\ &= \frac{1}{2}(1.5 \pm 0.5) \\ &= 1 \text{ and } 1/2. \end{aligned}$$

To find the 1-eigenvectors, we use row reduction to compute the nullspace of $1I - A$. First we observe that the matrix $1I - A$ has dependent rows (hence also dependent columns):

$$1I - A = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \begin{pmatrix} .8 & .3 \\ .2 & .7 \end{pmatrix} = \begin{pmatrix} .2 & -.3 \\ -.2 & .3 \end{pmatrix}.$$

Indeed, this must be the case because 1 is an eigenvalue. Then we compute the RREF:

$$(1I - A)\mathbf{x} = \mathbf{0} \quad \rightsquigarrow \quad \begin{pmatrix} .2 & -.3 \\ -.2 & .3 \end{pmatrix}\mathbf{x} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad \overset{\text{RREF}}{\rightsquigarrow} \quad \begin{pmatrix} 1 & -3/2 \\ 0 & 0 \end{pmatrix}\mathbf{x} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

It follows that there is a line of 1-eigenvectors:[105]

$$\mathbf{x} = t\begin{pmatrix} 3/2 \\ 1 \end{pmatrix}.$$

Next we compute the $(1/2)$-eigenspace:

$$\left(\frac{1}{2}I - A\right)\mathbf{x} = \mathbf{0} \quad \rightsquigarrow \quad \begin{pmatrix} -.3 & -.3 \\ -.2 & -.2 \end{pmatrix}\mathbf{x} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad \overset{\text{RREF}}{\rightsquigarrow} \quad \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}\mathbf{x} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

[105] In the previous section I chose $\mathbf{x} = (3, 2)$ to avoid fractions.

Thus we have also have a line of (1/2)-eigenvectors:[106]

$$\mathbf{x} = t\begin{pmatrix} -1 \\ 1 \end{pmatrix}$$

The procedure is the same for larger matrices. Given the eigenvalues, we can find all of the eigenvectors by row reduction. The hard part is to find the eigenvalues.[107] In general, we define the *characteristic polynomial* of a square matrix A:

$$\boxed{\chi_A(\lambda) := \det(\lambda I - A).}$$

This is, indeed, a polynomial in λ. Furthermore, if A is $n \times n$ then $\chi_A(\lambda)$ is a polynomial of degree n. In general the coefficients are quite complicated, but two of the coefficients have special names. We have

$$\boxed{\chi_A(\lambda) = \lambda^n - \operatorname{tr}(A)\lambda^{n-1} + \cdots + (-1)^n \det(A),}$$

where the *trace* of a square matrix is defined as the sum of its diagonal entries:

$$\boxed{\operatorname{tr}(A) = \operatorname{tr}\begin{pmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{n1} & \cdots & a_{nn} \end{pmatrix} := a_{11} + a_{22} + \cdots + a_{nn}.}$$

We already know this formula for 2×2 matrices, and the general case is not hard to check.

If matrices A and B satisfy $B = XAX^{-1}$ for some invertible matrix X, then I claim that A and B have the same characteristic polynomial:

$$\boxed{\chi_{XAX^{-1}}(\lambda) = \chi_A(\lambda).}$$

To prove this, we note that

$$\begin{aligned} \chi_B(\lambda) &= \det(\lambda I - B) \\ &= \det(\lambda XX^{-1} - XAX^{-1}) \\ &= \det(X(\lambda I - A)X^{-1}) \\ &= \cancel{\det(X)}\det(\lambda I - A)\cancel{\det(X)^{-1}} \\ &= \det(\lambda I - A) \\ &= \chi_A(\lambda). \end{aligned}$$

[106] In the previous example I chose $\mathbf{x} = (1, -1)$ because I didn't want a negative sign in the first coordinate.

[107] Solving polynomials equations is a non-linear problem. There are no exact algorithms, but there are reasonably good approximation schemes. The state of the art for computing eigenvalues is the *QR algorithm*, which doesn't use the characteristic polynomial at all.

By comparing the coefficients of $\chi_A(\lambda)$ and $\chi_B(\lambda)$, it follows that[108]

$$\boxed{\operatorname{tr}(XAX^{-1}) = \operatorname{tr}(A) \quad \text{and} \quad \det(XAX^{-1}) = \det(A).}$$

The eigenvalues of a square matrix A are the roots of the characteristic polynomial. It follows from the Fundamental Theorem of Algebra that

$$\boxed{\text{Every square matrix has at least one eigenvalue.}}$$

Indeed, since the characteristic polynomial $\chi_A(\lambda)$ has degree n, the FTA says that there exist complex numbers $\lambda_1, \ldots, \lambda_n \in \mathbb{C}$ such that

$$\chi_A(\lambda) = (\lambda - \lambda_1)(\lambda - \lambda_2) \cdots (\lambda - \lambda_n).$$

We can expand this to get

$$\boxed{\chi_A(\lambda) = \lambda^n - (\lambda_1 + \cdots + \lambda_n)\lambda^{n-1} + \cdots + (-1)^n \lambda_1 \cdots \lambda_n.}$$

Then comparing the coefficients with our previous expansion for $\chi_A(\lambda)$ gives

$$\boxed{\operatorname{tr}(A) = \lambda_1 + \cdots + \lambda_n \quad \text{and} \quad \det(A) = \lambda_1 \cdots \lambda_n.}$$

That is, the trace of A equals the sum of the eigenvalues (with multiplicities) and the determinant of A equals the product of the eigenvalues (with multiplicities). This is often useful.

Remarks:

- I guess we could say that every $n \times n$ matrix has n eigenvalues, but they need not be distinct. For example, the identity matrix I_n has characteristic polynomial

$$\det(\lambda I_n - I_n) = \det \begin{pmatrix} \lambda - 1 & & \\ & \ddots & \\ & & \lambda - 1 \end{pmatrix} = (\lambda - 1)^n,$$

  hence 1 is the only eigenvalue. The corresponding eigenspace is all of $\mathbb{R}^n$. Indeed, every vector $\mathbf{x} \in \mathbb{R}^n$ is a 1-eigenvector of the identity matrix: $I_n \mathbf{x} = \mathbf{x} = 1\mathbf{x}$.

- A real matrix has at least one complex eigenvalue, but it need not have any real eigenvalues. For example, consider the matrix that rotates counterclockwise by 90°:

$$A = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

  The characteristic polynomial is

$$\det(\lambda I - A) = \begin{pmatrix} \lambda & 1 \\ -1 & \lambda \end{pmatrix} = \lambda^2 + 1,$$

  hence the eigenvalues are $\pm i$. The corresponding eigenspaces are

$$\mathcal{N}(iI - A) = t \begin{pmatrix} 1 \\ -i \end{pmatrix} \quad \text{and} \quad \mathcal{N}(-iI - A) = t \begin{pmatrix} 1 \\ i \end{pmatrix}.$$

  Hence A is a real matrix with no real eigenvalues and no real eigenvectors.

[108] Of course, we already knew this property of the determinant.

6.3 Diagonalization

We say that a square matrix A is *diagonalizable* when it has a basis of eigenvectors. So far we have seen only diagonalizable matrices. Here is the simplest example of a matrix that is **not diagonalizable**:

$$A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

The characteristic polynomial is

$$\det(\lambda I - A) = \det \begin{pmatrix} \lambda - 1 & -1 \\ 0 & \lambda - 1 \end{pmatrix} = (\lambda - 1)^2,$$

hence 1 is the only eigenvalue. But the 1-eigenspace is only one dimensional:

$$(1I - A)\mathbf{x} = \mathbf{0} \quad \rightsquigarrow \quad \begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix} \mathbf{x} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad \rightsquigarrow \quad \mathbf{x} = t \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

Non-diagonalizable matrices are quite a nuisance. Fortunately, they are rare. The next result shows that any $n \times n$ matrix with n distinct eigenvalues is diagonalizable.

Theorem (Distinct Eigenvalues Implies Diagonalizable). Let A be an $n \times n$ matrix. Suppose that the characteristic polynomial factors as

$$\chi_A(\lambda) = (\lambda - \lambda_1)(\lambda - \lambda_2) \cdots (\lambda - \lambda_n),$$

where the complex numbers $\lambda_1, \ldots, \lambda_n \in \mathbb{C}$ are distinct. Furthermore, let $\mathbf{x}_1, \ldots, \mathbf{x}_n \in \mathbb{C}^n$ be some nonzero vectors satisfying $A\mathbf{x}_i = \lambda_i \mathbf{x}_i$.[109] Then I claim that $\mathbf{x}_1, \ldots, \mathbf{x}_n$ is a basis for $\mathbb{C}^n$.

Warning: This theorem is not sharp. If the characteristic polynomial of a matrix has a repeated factor then the matrix may or may not be diagonalizable. For example, the following two matrices both have characteristic polynomial $(\lambda - 1)^2$:

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

The matrix on the left is not diagonalizable, but the matrix on the right **is** diagonalizable. Indeed, every vector is a 1-eigenvector for the identity matrix. I will give a sharp characterization of diagonalizability in the next section.

Proof. It is enough to show that the set $\mathbf{x}_1, \ldots, \mathbf{x}_n$ is linearly independent. Then the subspace of $\mathbb{C}^n$ spanned by $\mathbf{x}_1, \ldots, \mathbf{x}_n$ is n-dimensional, hence it must be the whole space.

First we observe that the vectors $\mathbf{x}_1, \ldots, \mathbf{x}_n$ are distinct. Indeed, suppose we had $\mathbf{x}_i = \mathbf{x}_j = \mathbf{x}$ for some $i \neq j$. This would imply that

$$A\mathbf{x} = A\mathbf{x}$$

[109] Such vectors exist because $\lambda_1, \ldots, \lambda_n$ are eigenvalues.

$$\lambda_i \mathbf{x} = \lambda_j \mathbf{x}$$
$$(\lambda_i - \lambda_j)\mathbf{x} = \mathbf{0}.$$

But by assumption we have $\lambda_i - \lambda_j \neq 0$ and $\mathbf{x} \neq \mathbf{0}$, which gives a contradiction. This is no big deal; it just says that a given vector can't be an eigenvector for two different eigenvalues.

We will prove by induction on k that the set of vectors $\mathbf{x}_1, \ldots, \mathbf{x}_k$ is independent for any $1 \leq k \leq n$, and it will follow that the set $\mathbf{x}_1, \ldots, \mathbf{x}_n$ is independent. The result is trivial for $k = 1$ because any set containing one vector is by convention called independent. It is not logically necessary, but let's also consider the case $k = 2$, to get a feel for the general argument. Suppose that we have

$$b_1 \mathbf{x}_1 + b_2 \mathbf{x}_2 = \mathbf{0}$$

for some scalars $b_1, b_2 \in \mathbb{C}$. Our goal is to show that $b_1 = 0$ and $b_2 = 0$. First multiply both sides on the left by A to obtain

$$A(b_1 \mathbf{x}_1 + b_2 \mathbf{x}_2) = A\mathbf{0}$$
$$b_1 A\mathbf{x}_1 + b_2 A\mathbf{x}_2 = \mathbf{0}$$
$$b_1 \lambda_1 \mathbf{x}_1 + b_2 \lambda_2 \mathbf{x}_2 = \mathbf{0}.$$

Subtract λ_2 times the first equation from this equation to obtain

$$(b_1 \lambda_1 \mathbf{x}_1 + b_2 \lambda_2 \mathbf{x}_2) - \lambda_2(b_1 \mathbf{x}_1 + b_2 \mathbf{x}_2) = \mathbf{0}$$
$$b_1(\lambda_1 - \lambda_2)\mathbf{x}_1 + b_2(\lambda_2 - \lambda_2)\mathbf{x}_2 = \mathbf{0}$$
$$b_1(\lambda_1 - \lambda_2)\mathbf{x}_1 = \mathbf{0}.$$

Since $\lambda_1 - \lambda_2 \neq 0$ and $\mathbf{x}_1 \neq \mathbf{0}$ this implies that $b_1 = 0$. But then substituting into the first equation gives

$$b_1 \mathbf{x}_1 + b_2 \mathbf{x}_2 = \mathbf{0}$$
$$0\mathbf{x}_1 + b_2 \mathbf{x}_2 = \mathbf{0}$$
$$b_2 \mathbf{x}_2 = \mathbf{0},$$

which implies that $b_2 = 0$ because $\mathbf{x}_2 \neq \mathbf{0}$.

Now we prove the general case. Fix some $k \geq 2$ and suppose that we have

$$b_1 \mathbf{x}_1 + b_2 \mathbf{x}_2 + \cdots + b_k \mathbf{x}_k = \mathbf{0} \tag{1}$$

for some scalars $b_1, \ldots, b_k \in \mathbb{C}$. Our goal is to show that $b_1 = b_2 = \cdots = b_k = 0$. To do this we multiply both sides on the left by A to obtain

$$A(b_1 \mathbf{x}_1 + b_2 \mathbf{x}_2 + \cdots + b_k \mathbf{x}_k) = A\mathbf{0}$$
$$b_1 A\mathbf{x}_1 + b_2 A\mathbf{x}_2 + \cdots + b_k A\mathbf{x}_k = \mathbf{0}$$
$$b_1 \lambda_1 \mathbf{x}_1 + b_2 \lambda_2 \mathbf{x}_2 + \cdots + b_k \lambda_k \mathbf{x}_k = \mathbf{0}. \tag{2}$$

Then we consider the equation (2) $-\lambda_k(1)$:

$$\begin{aligned}\left(\sum_{i=1}^{k} b_i\lambda_i\mathbf{x}_i\right) - \lambda_k\left(\sum_{i=1}^{k} b_i\mathbf{x}_i\right) &= \mathbf{0}\\ \sum_{i=1}^{k} b_i(\lambda_i-\lambda_k)\mathbf{x}_k &= \mathbf{0}\\ 0\mathbf{x}_k + \sum_{i=1}^{k-1} b_i(\lambda_i-\lambda_k)\mathbf{x}_i &= \mathbf{0}\\ \sum_{i=1}^{k-1} b_i(\lambda_i-\lambda_k)\mathbf{x}_i &= \mathbf{0}.\end{aligned}$$

By induction, the vectors $\mathbf{x}_1,\ldots,\mathbf{x}_{k-1}$ are independent, hence for any $1\le i\le k-1$ we must have $b_i(\lambda_i-\lambda_k)=0$. But by assumption we have $\lambda_i\neq\lambda_k$, and hence $b_i=0$, for any $1\le i\le k-1$. Finally, we substitute back into equation (1) to obtain $0\mathbf{x}_1+\cdots+0\mathbf{x}_{k-1}+b_k\mathbf{x}_k=\mathbf{0}$, which implies that $b_k=0$ because $\mathbf{x}_k\neq\mathbf{0}$. □

This result implies that "almost all" matrices are diagonalizable. To see this, we will use the concept of the *discriminant* of a polynomial. For example, consider the general 2×2 matrix

$$A=\begin{pmatrix}a & b\\ c & d\end{pmatrix}.$$

The characteristic polynomial is

$$\chi_A(\lambda)=\lambda^2-(a+d)\lambda+(ad-bc),$$

and hence the eigenvalues are

$$\lambda=\frac{a+d\pm\sqrt{(a+d)^2-4(ad-bc)}}{2}.$$

The quantity $\Delta(a,b,c,d)=(a+d)^2-4(ad-bc)$ is called the discriminant of the characteristic polynomial. If $\Delta\neq 0$ then we observe that A has two distinct eigenvalues, hence is diagonalizable. If we choose the entries a,b,c,d of the matrix A at random then it would be quite unlikely to have $\Delta(a,b,c,d)=0$. To be more precise, we can view the set of 2×2 matrices as a 4-dimensional vector space:

$$\mathbb{C}^{2\times 2}=\text{the vector space of } 2\times 2 \text{ matrices with complex entries.}$$

Inside this 4-dimensional vector space, the set of matrices satisfying $\Delta(a,b,c,d)=0$ forms a "3-dimensional subset".[110] By analogy, consider a 2-dimensional plane in $\mathbb{R}^3$. A randomly

[110]For a general polynomial $f(x_1,\ldots,x_n)$ in n variables, the set of points $\mathbf{x}\in\mathbb{C}^n$ satisfying $f(\mathbf{x})=0$ forms an $(n-1)$-dimensional subset. I don't want to be too precise about this.

chosen point in $\mathbb{R}^3$ will not lie on this plane. Similarly, a randomly chosen point in a 4-dimensional vector space will not lie in a given 3-dimensional shape.

This discussion generalizes to square matrices of any size. Given an $n \times n$ matrix A with entries a_{ij}, there is a certain polynomial $\Delta(A)$ in the n^2 variables a_{ij} such that $\Delta(A) = 0$ if and only if A has a repeated eigenvalue. Since the equation $\Delta(A) = 0$ defines an $(n^2 - 1)$-dimensional subset of the n^2-dimensional space of $n \times n$ matrices, a randomly chosen matrix will have distinct eigenvalues, and hence will be diagonalizable.

Why do we call it diagonalization? Let A be a diagonalizable $n \times n$ matrix and let $\mathbf{x}_1, \ldots, \mathbf{x}_n \in \mathbb{C}^n$ be a basis of eigenvectors with corresponding eigenvalues $A\mathbf{x}_i = \lambda_i \mathbf{x}_i$. (Here we do not assume that the eigenvalues are distinct.) We can write the n equations $A\mathbf{x}_i = \lambda_i \mathbf{x}_i$ simultaneously as a matrix equation:

$$\begin{pmatrix} A\mathbf{x}_1 \mid \cdots \mid A\mathbf{x}_n \end{pmatrix} = \begin{pmatrix} \lambda_1\mathbf{x}_1 \mid \cdots \mid \lambda_n\mathbf{x}_n \end{pmatrix}$$

$$A \begin{pmatrix} \mathbf{x}_1 \mid \cdots \mid \mathbf{x}_n \end{pmatrix} = \begin{pmatrix} \mathbf{x}_1 \mid \cdots \mid \mathbf{x}_n \end{pmatrix} \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{pmatrix}$$

$$AX = X\Lambda,$$

where Λ is a **diagonal** matrix containing the eigenvalues. Since the columns of X are independent by assumption, the matrix X is invertible and we can write

$$\boxed{A = X\Lambda X^{-1}} \quad \text{or} \quad \boxed{X^{-1}AX = \Lambda.}$$

Thus we have used the eigenvector matrix X to convert A into the diagonal matrix Λ. In other words, we have "diagonalized" A. We will see below that this is extremely useful.

6.4 Evaluating a Polynomial at a Matrix

Matrices can be added, multiplied by scalars, and raised to powers. This allows us to consider polynomials of matrices. More precisely, we can "evaluate" polynomials at matrices. Consider a polynomial in one variable, with complex coefficients:

$$f(x) = b_0 + b_1 x + \cdots + b_k x^k.$$

Then for any $n \times n$ matrix A we define the $n \times n$ matrix $f(A)$ by[111]

$$f(A) := b_0 I_n + b_1 A + b_2 A^2 + \cdots + b_k A^k.$$

This evaluation behaves well with respect to eigenvalues and eigenvectors. That is, for any vector $\mathbf{x}$ and scalar λ, we have

$$\boxed{A\mathbf{x} = \lambda\mathbf{x} \quad \Longrightarrow \quad f(A)\mathbf{x} = f(\lambda)\mathbf{x}.}$$

Indeed, if $A\mathbf{x} = \lambda\mathbf{x}$ then we can show by induction that $A^m\mathbf{x} = \lambda^m\mathbf{x}$ for any $m \geq 0$:

[111] Here we use the convention that $A^0 = I_n$.

- **Base Case.** $A^0\mathbf{x} = I_n\mathbf{x} = \mathbf{x} = \lambda^0\mathbf{x}$.
- **Induction Step.** If $A^{m-1}\mathbf{x} = \lambda^{m-1}\mathbf{x}$ then

$$A^m\mathbf{x} = A(A^{m-1}\mathbf{x}) = A^m(\lambda^{m-1}\mathbf{x}) = \lambda^{m-1}(A\mathbf{x}) = \lambda^{m-1}\lambda\mathbf{x} = \lambda^m\mathbf{x}.$$

Then for any polynomial $f(x) = b_0 + b_1x + \cdots + b_kx^k$ we have

$$\begin{aligned} f(A)\mathbf{x} &= (b_0I_n + b_1A + \cdots + b_kA^k)\mathbf{x} \\ &= b_0I_n\mathbf{x} + b_1A\mathbf{x} + \cdots + b_kA^k\mathbf{x} \\ &= b_0\mathbf{x} + b_1\lambda\mathbf{x} + \cdots + b_k\lambda^k\mathbf{x} \\ &= (b_0 + b_1\lambda + \cdots + b_k\lambda^k)\mathbf{x} \\ &= f(\lambda)\mathbf{x}. \end{aligned}$$

Now suppose that the matrix A is a "root" of the polynomial $f(x)$. That is, suppose that $f(A)$ is the zero matrix. Then every eigenvalue of A is also a root of $f(x)$:

$$\boxed{f(A) = O \quad \Longrightarrow \quad \text{every eigenvalue of } A \text{ satisfies } f(\lambda) = 0.}$$

Indeed, if $A\mathbf{x} = \lambda\mathbf{x}$ and $f(A) = O$ then we have

$$f(\lambda)\mathbf{x} = f(A)\mathbf{x} = O\mathbf{x} = \mathbf{0}.$$

And if $\mathbf{x} \neq \mathbf{0}$ then this implies $f(\lambda) = 0$. Here are some examples.

Projections. Any matrix satisfying $P^2 = P$ has eigenvalues in the set $\{0, 1\}$. Indeed, if $P^2 - P = O$ then any eigenvalue λ of P satisfies

$$\begin{aligned} \lambda^2 - \lambda &= 0 \\ \lambda(\lambda - 1) &= 0 \\ \lambda &= 0 \text{ or } 1. \end{aligned}$$

This doesn't mean that both eigenvalues must occur. For example, the zero matrix satisfies $O^2 = O$ and its only eigenvalue is 0, while the identity matrix satisfies $I^2 = I$ and its only eigenvalue is 1.

At the end of this section we will show that any matrix satisfying $P^2 = P$ is diagonalizable. Assuming this for now, we can prove that any $n \times n$ matrix satisfying $P^2 = P$ is a (possibly non-orthogonal) projection matrix. To do this, let $\mathbf{x}_1, \ldots, \mathbf{x}_n$ be a basis of eigenvectors. Since the only possible eigenvalues are 1 and 0, we can sort the eigenvectors so that $P\mathbf{x}_i = 1\mathbf{x}_i = \mathbf{x}_i$ for $0 \leq i \leq r$ and $P\mathbf{x}_i = 0\mathbf{x}_i = \mathbf{0}$ for $r < i \leq n$.[112] This gives the factorization

$$P = X \left(\begin{array}{c|c} I_r & O_{r,n-r} \\ \hline O_{n-r,r} & O_{n-r}, n-r \end{array} \right) X^{-1}.$$

[112] We allow the possibilities $r = 0$ (all eigenvalues are 0) and $r = n$ (all eigenvalues are 1).

Now let A be the $n \times r$ matrix consisting of the first r columns of X and let B be the $r \times n$ matrix consisting of the first r rows of X^{-1}. Then we have

$$P = \left(\begin{array}{c|c} A & * \end{array} \right) \left(\begin{array}{c|c} I_r & O_{r,n-r} \\ \hline O_{n-r,r} & O_{n-r,n-r} \end{array} \right) \left(\begin{array}{c} B \\ \hline * \end{array} \right) = \left(\begin{array}{c|c} A & * \end{array} \right) \left(\begin{array}{c} B \\ \hline O \end{array} \right) = AB + O = AB.$$


And we also have

$$\left(\begin{array}{c|c} I_r & O_{r,n-r} \\ \hline O_{n-r,r} & I_{n-r,n-r} \end{array} \right) = I_n = X^{-1}X = \left(\begin{array}{c} B \\ \hline * \end{array} \right) \left(\begin{array}{c|c} A & * \end{array} \right) = \left(\begin{array}{c|c} BA & * \\ \hline * & * \end{array} \right),$$

which implies that $BA = I_r$. In summary:

$$\boxed{P^2 = P \quad \Longrightarrow \quad P = AB \text{ for some } A, B \text{ satisfying } BA = I_r, \text{ where } r = \operatorname{rank}(P).}$$

This is the projection onto the column space $U = \mathcal{C}(A)$, in a direction parallel to the null space $V = \mathcal{N}(B)$. Picture:



The projection is orthogonal if and only if $V = U^\perp$. In this case we have $\mathcal{N}(B) = \mathcal{C}(A)^\perp = \mathcal{N}(A^T)$, which impliess that $\mathcal{R}(A) = \mathcal{N}(B)^T = \mathcal{N}(A^T)^\perp = \mathcal{R}(A^T)$. Since $\mathcal{R}(B) = \mathcal{R}(A^T)$ we can find an invertible $r \times r$ matrix S of row operations such that $B = SA^T$. But then $BA = I_r$ implies $SA^TA = I_r$ and hence $S = (A^TA)^{-1}$. Finally, we conclude that

$$P = AB = ASA^T = A(A^TA)^{-1}A^T,$$

which agrees with our previous formula for orthogonal projections.


Reflections. Any matrix satisfying $F^2 = I$ has eigenvalues in the set $\{1, -1\}$. Indeed, if $F^2 - I = O$ then any eigenvalue λ of F satisfies

$$\begin{aligned} \lambda^2 - 1 &= 0 \\ \lambda^2 &= 1 \\ \lambda &= 1 \text{ or } -1. \end{aligned}$$

Consider the unique matrix P satisfying $F = 2P - I$ and $P = (F + I)/2$. We observe that

$$P^2 = \frac{1}{4}(F^2 + 2F + I^2) = \frac{1}{2}(I + 2F + I) = \frac{1}{4}(2F + 2I) = \frac{1}{2}(F + I) = P,$$

so that P is a projection. Let U and V be the 1-eigenspace and 0-eigenspace of P as in the previous example, then U is the 1-eigenspace of F and V is the (-1)-eigenspace of F. Geometrically, F is the reflection across the subspace U in the direction of V. Picture:



In terms of matrices, if F is a matrix of rank r satisfying $F^2 = I$ then we can find two $r \times (n-r)$ matrices A, B satisfying $BA = I_r$, such that

$$F = 2P - I = 2AB - I.$$

This is the reflection across the column space $U = \mathcal{C}(A)$, parallel to the nullspace $V = \mathcal{N}(B)$.

Rotations. Any matrix satisfying $R^n = I$ has eigenvalues in the set $\{e^{2\pi ik/n} : k \in \mathbb{Z}\}$. Indeed, since $R^n - I = O$, any eigenvalue λ of R must satisfy $\lambda^n - 1 = 0$, and hence must be an nth

root of unity. Such matrices can be quite complicated. For a simple example, we consider the 2×2 rotation matrix:

$$R_\theta = \begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix}.$$

The characteristic polynomial is

$$\chi_{R_\theta}(\lambda) = \lambda^2 - 2\cos\theta\lambda + 1,$$

hence the eigenvalues are[113]

$$\begin{aligned} \lambda &= \frac{2\cos\theta \pm \sqrt{4\cos^2\theta - 4}}{2} \\ &= \frac{2\cos\theta \pm 2\sqrt{\cos^2\theta - 1}}{2} \\ &= \frac{2\cos\theta \pm 2\sqrt{-\sin^2\theta}}{2} \\ &= \frac{2\cos\theta \pm 2i\sin\theta}{2} \\ &= \cos\theta \pm i\sin\theta \\ &= e^{\pm i\theta}. \end{aligned}$$

The case $\theta = 0$ corresponds to the identity matrix, with eigenvalues $(1, 1)$ and the case $\theta = \pi$ corresponds to the negative identity matrix, with eigenvalues $(-1, -1)$. In all other cases, the eigenvalues (and hence also the eigenvectors) are not real. If $\theta = 2\pi/n$ then the eigenvalues $\lambda = e^{\pm 2\pi i/n}$ satisfy $\lambda^n = 1$. This agrees with the fact that

$$(R_{2\pi/n})^n = I.$$

Next, we give an alternative proof for the existence of eigenvalues, which does not use determinants.[114]

Theorem (Existence of Eigenvalues). Let A be any $n \times n$ matrix with real or complex entries and consider an arbitrary nonzero vector $\mathbf{v} \in \mathbb{C}^n$. Since $\mathbb{C}^n$ is n-dimensional, the following $n + 1$ vectors must be linearly dependent:[115]

$$\mathbf{v}, A\mathbf{v}, A^2\mathbf{x}, \ldots, A^n\mathbf{v}.$$

In other words, we can find scalars $b_0, b_1, \ldots, b_n$, not all zero, such that

$$b_0\mathbf{v} + b_1A\mathbf{v} + b_2A^2\mathbf{v} + \cdots + b_nA^n\mathbf{v} = \mathbf{0}.$$

In fact, one of the scalars $b_1, \ldots, b_n$ must be nonzero, otherwise we would have $b_0\mathbf{v} = \mathbf{0}$ and $b_0 \neq 0$, which contradicts the fact that $\mathbf{v} \neq \mathbf{0}$. We can rewrite the previous equation as

$$(b_0I + b_1A + b_2A^2 + \cdots + b_nA^n)\mathbf{v} = \mathbf{0},$$

[113] It's a bit reckless to take square roots in this way, but it gives the correct answer.
[114] This proof is the motivation for Axler's approach in *Linear Algebra Done Right*.
[115] Indeed, **any** collection of $n + 1$ vectors in $\mathbb{C}^n$ is linearly dependent.

$$f(A) = \mathbf{0},$$

for the polynomial $f(x) = b_0 + b_1 x + b_2 x^2 + \cdots b_n x^n$, which has degree between 1 and n because not all of $b_1, \ldots, b_n$ are zero. Let's say $\deg(f) = k$. By the Fundamental Theorem of Algebra we can factor $f(x)$ as

$$f(x) = (x - \alpha_1)(x - \alpha_2) \cdots (x - \alpha_k),$$

for some complex numbers $\alpha_1, \ldots, \alpha_k \in \mathbb{C}$, not necessarily distinct. Now the equation $f(A)\mathbf{v} = \mathbf{0}$ becomes[116]

$$(A - \alpha_1 I)(A - \alpha_2 I) \cdots (A - \alpha_k I)\mathbf{v} = \mathbf{0}.$$

To save notation, let's write $A_i = A - \alpha_i I$. Thus we have

$$A_1 A_2 \cdots A_k \mathbf{v} = \mathbf{0}.$$

Since $\mathbf{v} \neq \mathbf{0}$, this implies that the matrix $A_1 \cdots A_k$ is not invertible. And since a product of invertible matrices is invertible, this implies that at least one of the factors, say A_i, is not invertible. Finally, since $A_i = A - \alpha_i I$ is not invertible, we conclude that α_i is an eigenvalue of A. In particular, we have shown that A has an eigenvalue. □

Building on this idea, we can give a sharper result about diagonalization. The proof is tricky so you can feel free to skip it.

Theorem (Existence of Diagonalization). A square matrix A is diagonalizable if and only if we have $f(A) = O$ for some polynomial with no repeated roots.

Proof. First suppose that A has a basis of eigenvectors $\mathbf{x}_1, \ldots, \mathbf{x}_n$ with corresponding eigenvalues $A\mathbf{x}_i = \lambda_i \mathbf{x}_i$. Some of these eigenvalues might be repeated. Let $\mu_1, \ldots, \mu_k$ be the list of eigenvalues with repetition removed, and consider the polynomial

$$f(x) = (x - \mu_1) \cdots (x - \mu_k),$$

which has no repeated roots. We will show that $f(A) = O$. To do this, we observe that the matrices $(A - \mu_i I)$ and $(A - \mu_j I)$ commute for any i, j:

$$(A - \mu_i I)(A - \mu_j I) = A^2 - (\mu_i + \mu_j)A + \mu_i \mu_j I = (A - \mu_j I)(A - \mu_i - I).$$

We will use this to show that $f(A)\mathbf{v} = \mathbf{0}$ for any eigenvector $\mathbf{v}$. Then since there exists a basis of eigenvectors, it will follow from this that $f(A)\mathbf{v} = \mathbf{0}$ for **any** vector $\mathbf{v}$, and hence $f(A)$ is the zero matrix. So let $\mathbf{v}$ be an eigenvector with eigenvalue μ_i.[117] Then we have[118]

$$f(A)\mathbf{v} = \left(\prod_j (A - \mu_j I) \right) \mathbf{v}$$

[116] There is a subtle point hiding here. Given a polynomial $f(x)$ and square matrices A, B, it is **not** generally true that $f(AB) = f(A)f(B)$. However, if $AB = BA$ then we do have $f(AB) = f(A)f(B)$. Since the matrices $A - \alpha_1 I, \ldots, A - \alpha_k I$ commute with each other, we are okay in this case.

[117] Recall that every eigenvalue is in the list $\mu_1, \ldots, \mu_k$.

[118] We used commutativity to pull the factor $A - \mu_i I$ to the right.

$$
\begin{aligned}
&= \left(\prod_{j\neq i}(A-\mu_j)\right)(A-\mu_i)\mathbf{v} \\
&= \left(\prod_{j\neq i}(A-\mu_j)\right)(A\mathbf{v}-\mu_{\mathbf{v}}) \\
&= \left(\prod_{j\neq i}(A-\mu_j)\right)\mathbf{0} \\
&= \mathbf{0}.
\end{aligned}
$$

Thus we have shown that a diagonalizable matrix A satisfies an equation $f(A) = O$ for some polynomial $f(x)$ with no repeated roots.

Conversely, suppose that an $n \times n$ matrix A satisfies $f(A) = O$ for some polynomial $f(x)$ with no repeated roots. Suppose that $\deg(f) = k$ and write

$$f(x) = (x-\lambda_1)\cdots(x-\lambda_k)$$

for some distinct complex numbers $\lambda_1, \ldots, \lambda_k \in \mathbb{C}$. We want to show that A has a basis of eigenvectors. First we define the null spaces

$$E_{\lambda_i} = \mathcal{N}(A-\lambda_i I) = \{\mathbf{x} : A\mathbf{x} = \lambda_i \mathbf{x}\}.$$

Note that $E_{\lambda_i} \neq \{\mathbf{0}\}$ if and only if λ_i is an eigenvalue, in which case E_{λ_i} is the corresponding eigenspace. We don't really care if all of the numbers λ_i are eigenvalues. Indeed, some of them might not be. Our goal is merely to show that the spaces $E_{\lambda_1}, \ldots, E_{\lambda_k}$ are big enough to fill up all of $\mathbb{C}^n$. To be precise, we will show that

- $E_{\lambda_i} \cap E_{\lambda_j} = \{\mathbf{0}\}$ for all $i \neq j$,
- $\mathbb{C}^n = \{\mathbf{x}_1 + \cdots + \mathbf{x}_k : \mathbf{x}_i \in E_{\lambda_i} \text{ for all } i\}$.

Then by concatenating bases for $E_{\lambda_1}, \ldots, E_{\lambda_k}$ we will obtain a basis for $\mathbb{C}^n$ that consists of eigenvectors of A. For the first statement, suppose that $\mathbf{x} \in E_{\lambda_i} \cap E_{\lambda_j}$ so that $\lambda_i \mathbf{x} = A\mathbf{x} = \lambda_j \mathbf{x}$. If $\mathbf{x} \neq \mathbf{0}$ then this implies that $\lambda_i = \lambda_j$ and hence $i \neq j$, because the λ_i are distinct. The second statement is trickier. First we consider the partial fraction expansion of $1/f(x)$:

$$\frac{1}{f(x)} = \frac{1}{(x-\lambda_1)\cdots(x-\lambda_k)} = \sum_i \frac{\alpha_i}{x-\lambda_i},$$

for some scalars $\alpha_1, \ldots, \alpha_k \in \mathbb{C}$, not necessarily distinct.[119] Now consider the polynomials

$$p_i(x) = \frac{\alpha_i f(x)}{x-\lambda_i} = \alpha_i \prod_{j\neq i}(x-\lambda_j),$$

[119] I won't prove the existence of the partial fraction expansion. It depends on the theory of greatest common divisors in the ring of polynomials.

and note that

$$\begin{aligned} p_1(x) + \cdots p_k(x) &= \frac{\alpha_1 f(x)}{x - \lambda_1} + \cdots + \frac{\alpha_k f(x)}{x - \lambda_k} \\ &= f(x) \cdot \sum_i \frac{\alpha_i}{x - \lambda_i} \\ &= f(x) \cdot \frac{1}{f(x)} \\ &= 1. \end{aligned}$$

Finally, consider any vector $\mathbf{x} \in \mathbb{C}^n$ and write $\mathbf{x}_i := p_i(A)\mathbf{x}$. On the one hand, by evaluating the polynomial equation $(x - \lambda_i)p_i(A) = \alpha_i f(x)$ at A we have

$$(A - \lambda_i I)\mathbf{x}_i = (A - \lambda_i I)p_i(A)\mathbf{x} = \alpha_i f(A)\mathbf{x} = O\mathbf{x} = \mathbf{0},$$

and hence $\mathbf{x}_i \in E_{\lambda_i}$. On the other hand, by evaluating the polynomial equation $p_1(x) + \cdots + p_k(x) = 1$ at A we have

$$\begin{aligned} \mathbf{x}_1 + \cdots + \mathbf{x}_k &= p_1(A)\mathbf{x} + \cdots + p_k(A)\mathbf{x} \\ &= (p_1(A) + \cdots p_k(A))\mathbf{x} \\ &= I\mathbf{x} \\ &= \mathbf{x}, \end{aligned}$$

as desired. □

That was a tricky proof, but it's a useful theorem. In particular, it implies that any matrix satisfying $P^2 = P$, and hence $P^2 - P = O$, is diagonalizable because the polynomial $x^2 - x = x(x-1)$ has distinct roots. Furthermore, any matrix satisfying $R^n = I$, and hence $R^n - I = O$, is diagonalizable because the polynomial $x^n - 1$ has distinct roots:

$$x^n - 1 = (x - 1)(x - e^{2\pi i/n})(x - e^{4\pi i/n}) \cdots (x - e^{2\pi i(n-1)/n}).$$

Finally, we examine what goes wrong for a specific non-diagonalizable matrix. Consider the following small matrices with repeated eigenvalues:

$$A = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix}$$

Each of these has characteristic polynomial $(x - \lambda)^2$:

$$(x - \lambda)^2 = \det \begin{pmatrix} x - \lambda & 0 \\ 0 & x - \lambda \end{pmatrix} = \det \begin{pmatrix} x - \lambda & -1 \\ 0 & x - \lambda \end{pmatrix}.$$

In the next section we will show that every matrix satisfies its own characteristic polynomial, which we can easily verify for these two matrices:

$$(A - \lambda I)^2 = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}^2 = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad (B - \lambda I)^2 = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}^2 = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}.$$

The matrix A also satisfies the polynomial $f(x) = x - \lambda$, which has no repeated roots. This confirms that A is diagonalizable; in fact, it is diagonal. On the other hand, the matrix B is not diagonalizable. This is easy to check directly. Instead, we will prove it indirectly, by showing that B cannot satisfy any polynomial with distinct roots. The basic reason is that

$$(B - \lambda I)^2 = O \quad \text{but} \quad B - \lambda I \neq O.$$

Indeed, consider any polynomial $g(x) = (x - \lambda_1) \cdots (x - \lambda_k)$ with distinct roots $\lambda_1, \ldots, \lambda_k$. If $g(B) = O$ then one of the matrices $B - \lambda_j I$ must be non-invertible, so that λ_j is an eigenvalue and hence $\lambda_j = \lambda$. Since the λ_i are distinct, this implies that the λ_i with $i \neq j$ are **not** eigenvalues. Then since $g(B) = O$ equals $B - \lambda I$ times a product of invertible matrices $B - \lambda_i I$ for $i \neq j$, we conclude that $B - \lambda I = O$. Contradiction.

Remark: In the next chapter we will say more about non-diagonalizable matrices.

6.5 The Functional Calculus

Why are diagonalizable matrices good? As we mentioned in the first section, if we can diagonalize a matrix A then we can find explicit formulas for the entries of its powers A^k. More generally, diagonalizing a matrix allows us to compute any polynomial evaluation of the matrix $f(A)$. We can even compute convergent power series, such as

$$\exp(A) := I + A + \frac{1}{2}A^2 + \cdots + \frac{1}{k!}A^k + \cdots .$$

To begin, suppose that an $n \times n$ matrix A has a basis of eigenvectors $\mathbf{x}_1, \ldots, \mathbf{x}_n$, with corresponding eigenvalues $A\mathbf{x}_i = \lambda_i \mathbf{x}_i$. (The eigenvalues are not necessarily distinct.) Then, as we showed in the previous section, we can write

$$A = X\Lambda X^{-1} = \begin{pmatrix} \mathbf{x}_1 \,|\, \cdots \,|\, \mathbf{x}_n \end{pmatrix} \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{pmatrix} \begin{pmatrix} \mathbf{x}_1 \,|\, \cdots \,|\, \mathbf{x}_n \end{pmatrix}^{-1}.$$

This factorization is compatible with polynomial evaluation. That is, for any polynomial $f(x)$, I claim that

$$\boxed{f(A) = X \cdot f(\Lambda) \cdot X^{-1}.}$$

If A (hence also Λ) is invertible, then we can even allow negative powers of x in the polynomial. Such expressions are called *Laurent polynomials*:

$$f(x) = b_{-\ell}x^{-\ell} + b_{-\ell+1}x^{-\ell+1} + \cdots + b_{k-1}x^{k-1} + b_k x^k \quad \text{for some } k, \ell \geq 0.$$

Actually, we will prove the more general fact that $A = XBX^{-1}$ implies $f(A) = X \cdot f(B) \cdot X^{-1}$ for any polynomial $f(x)$, and for any Laurent polynomial $f(x)$ when A (hence also B) is invertible. The first step is to prove that

$$\boxed{(XBX^{-1})^k = XB^kX^{-1} \text{ for all } k \geq 0, \text{ and also for } k < 0 \text{ when } B \text{ is invertible.}}$$

For this we use induction. When $k = 0$ we have $XB^0X^{-1} = XI_nX^{-1} = XX^{-1} = I_n = (XBX^{-1})^0$. Then for $k \geq 1$ we have

$$\begin{aligned}(XBX^{-1})^k &= (XBX^{-1})(XBX^{-1})^{k-1} \\ &= (XBX^{-1})(XB^{k-1}X^{-1}) && \text{induction} \\ &= XB(X^{-1}X)B^{k-1}X^{-1} \\ &= XBB^{k-1}X^{-1} \\ &= XB^kX^{-1}.\end{aligned}$$

If B is invertible, then for all $k \geq 0$ we also have

$$(XBX^{-1})^{-k} = [(XBX^{-1})^{-1}]^k = (XB^{-1}X^{-1})^k = X(B^{-1})^kX^{-1} = XB^{-k}X^{-1}.$$

Finally, for any polynomial $f(x) = b_0 + b_1x + \cdots + b_kx^k$ we have

$$\begin{aligned}f(A) &= b_0I + b_1A + b_2A^2 + \cdots + b_kA^k \\ &= b_0I + b_1(XBX^{-1}) + b_2(XBX^{-1})^2 + \cdots + b_k(XBX^{-1})^k \\ &= b_0(XX^{-1}) + b_1(XBX^{-1}) + b_2(XB^2X^{-1}) + \cdots + b_k(XB^kX^{-1}) \\ &= X(b_0I + b_1B + b_2B^2 + \cdots + b_kB^k)X^{-1} \\ &= X \cdot f(B) \cdot X^{-1}.\end{aligned}$$

The proof for Laurent polynomials is the same.

So far, this is not very useful. It becomes useful because of the following basic observation.

Multiplication of Diagonal Matrices is Easy. The formula for a general matrix product AB is complicated. However, multiplication of diagonal matrices is easy:

$$\begin{pmatrix} a_1 & & \\ & \ddots & \\ & & a_n \end{pmatrix} \begin{pmatrix} b_1 & & \\ & \ddots & \\ & & b_n \end{pmatrix} = \begin{pmatrix} a_1b_1 & & \\ & \ddots & \\ & & a_nb_n \end{pmatrix}.$$

It follows that for any diagonal matrix Λ and any (Laurent) polynomial $f(x)$ we have

$$f(\Lambda) = f\begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{pmatrix} = \begin{pmatrix} f(\lambda_1) & & \\ & \ddots & \\ & & f(\lambda_n) \end{pmatrix}.$$

This is the reason why diagonalization is a powerful technique.

Here is a first application. Recall the following results from the previous two sections:

- If the characteristic polynomial $\chi_A(x)$ has distinct roots then A is diagonalizable.

- If $f(A) = O$ for some polynomial $f(x)$ with distinct roots then A is diagonalizable.

The next theorem ties these results together.

The Cayley-Hamilton Theorem. Let A be a square matrix with characteristic polynomial $\chi_A(x) = \det(xI - A)$. Then we have

$$\boxed{\chi_A(A) = O.}$$

This is a strange idea, so let's first examine the 2×2 case. Consider the matrix

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix},$$

with characteristic polynomial

$$\chi_A(\lambda) = \lambda^2 - (a+d)\lambda + (ad - bc).$$

Then one can check (as Cayley and Hamilton did) that

$$\begin{aligned} \chi_A(A) &= A^2 - (a+d)A + (ad-bc)I \\ &= \begin{pmatrix} a & b \\ c & d \end{pmatrix}^2 - (a+d)\begin{pmatrix} a & b \\ c & d \end{pmatrix} + (ad-bc)\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ &= \text{some calculations} \\ &= \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}. \end{aligned}$$

Why on earth should this be true? It is because of diagonalization.

Proof of Cayley-Hamilton. Suppose first that A is diagonalizable, with $A = X\Lambda X^{-1}$. For any eigenvalue λ of A, the characteristic polynomial satisfies $\chi_A(\lambda) = 0$ by definition. Hence

$$\chi_A(A) = X \cdot \chi_A(\Lambda) \cdot X^{-1} = X \begin{pmatrix} \chi_A(\lambda_1) & & \\ & \ddots & \\ & & \chi_A(\lambda_n) \end{pmatrix} X^{-1} = XOX^{-1} = O.$$

The result for non-diagonalizable matrices follows by continuity. That is, any non-diagonalizable matrix is a limit of diagonalizable matrices. And the entries of the matrix $\chi_A(A)$ are continuous functions of the entries of A. But each entry of $\chi_A(A)$ is zero for any diagonal matrix. Hence the entries of the limit are zero.[120] □

Remark: The Cayley-Hamilton is actually more general than this. It holds over any commutative ring. As written, our proof only works over the complex numbers.

[120] This is a typical way to deal with non-diagonalizable matrices, i.e., view them as limits of diagonalizable matrices in the space $\mathbb{C}^{n \times n}$ of square matrices.

Next we consider two examples of infinite power series.

The Geometric Series. Consider an $n \times n$ matrix A. Suppose that A is diagonalizable with

$$A = X\Lambda X^{-1} = X \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{pmatrix} X^{-1}.$$

Evaluating A at the polynomial $f(x) = 1 - x$ gives

$$I - A = X(I - \Lambda)X^{-1} = X \begin{pmatrix} 1-\lambda_1 & & \\ & \ddots & \\ & & 1-\lambda_n \end{pmatrix} X^{-1}.$$

If none of the eigenvalues is 1 then $I - \Lambda$ (hence also $I - A$) is invertible, and we obtain

$$(I - A)^{-1} = X(I - \Lambda)^{-1}X^{-1} = X \begin{pmatrix} 1/(1-\lambda_1) & & \\ & \ddots & \\ & & 1/(1-\lambda_n) \end{pmatrix} X^{-1}.$$

On the other hand, for all $k \geq 0$ we can evaluate the polynomial $f(x) = 1 + x + \cdots + x^k$ at A to obtain

$$\begin{aligned} I + A + \cdots + A^k &= X(I + \Lambda + \cdots + \Lambda^k)X^{-1} \\ &= X \begin{pmatrix} 1+\lambda_1+\cdots+\lambda_1^k & & \\ & \ddots & \\ & & 1+\lambda_n+\cdots+\lambda_n^k \end{pmatrix} X^{-1}. \end{aligned}$$

Finally, suppose that the eigenvalues satisfy $0 < |\lambda_i| < 1$ for all i. Then the usual geometric series for scalars implies that

$$I + \Lambda + \cdots + \Lambda^k \to (I - \Lambda)^{-1} \quad \text{as } k \to \infty.$$

The convergence is componentwise in each entry of the matrix. For a fixed invertible matrix X, the function $B \mapsto XBX^{-1}$ is continuous in the matrix entries, hence

$$X(I + \Lambda + \cdots + \Lambda^k)X^{-1} \to X(I - \Lambda)^{-1}X^{-1} \quad \text{as } k \to \infty.$$

In summary, for a diagonalizable matrix A with eigenvalues satisfying $0 < |\lambda| < 1$, we have

$$I + A + \cdots + A^k \to (I - A)^{-1} \text{ componentwise.}$$

And by continuity, the result also holds for non-diagonalizable matrices.[121] On a previous homework you proved a weaker version of this result, using more difficult techniques. Diagonalization makes things easier because it turns matrix arithmetic into scalar arithmetic.

[121] Basically, this is because the eigenvalues depend continuously on the matrix entries. I don't want to get specific about it.

The Matrix Exponential. Given a square matrix A, the functional calculus allows us to define $f(A)$ for any power series $f(x) = a_0 + a_1 x + a_2^2 + \cdots$, as long as this power series converges when evaluated at the eigenvalues of A. For example, consider the power series definition of the exponential function

$$\exp(x) = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \cdots .$$

It is a basic theorem of analysis that $\exp(x)$ converges for any complex number $x \in \mathbb{C}$. In order to define $\exp(A)$ we first suppose that A is diagonalizable:

$$A = X\Lambda X^{-1} = X \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{pmatrix} X^{-1}.$$

For any $k \geq 0$ we have

$$\sum_{i=0}^{k} \frac{1}{i!} A^i = X \left(\sum_{i=0}^{k} \frac{1}{i!} \Lambda^i \right) X^{-1} = X \begin{pmatrix} \sum_{i=0}^{k} \frac{1}{i!}\lambda_1^i & & \\ & \ddots & \\ & & \sum_{i=0}^{k} \frac{1}{i!}\lambda_n^i \end{pmatrix} X^{-1}.$$

Since the power series for $\exp(x)$ converges everywhere, we have

$$\sum_{i=1}^{k} \frac{1}{i!} \cdot \Lambda_i \to \begin{pmatrix} \exp(\lambda_1) & & \\ & \ddots & \\ & & \exp(\lambda_n) \end{pmatrix} \quad \text{as } k \to \infty.$$

Then since conjugation by the fixed matrix X is continuous, we conclude that

$$\sum_{i=1}^{k} \frac{1}{i!} \cdot A^i \to X \begin{pmatrix} \exp(\lambda_1) & & \\ & \ddots & \\ & & \exp(\lambda_n) \end{pmatrix} X^{-1} \quad \text{as } k \to \infty.$$

This establishes the existence of the matrix exponential for any diagonalizable matrix A:[122]

$$\boxed{\exp(A) = I + A + \frac{1}{2!}A^2 + \frac{1}{3!}A^3 + \cdots .}$$

Let me warn you that

$$\exp(A + B) \neq \exp(A)\exp(B) \text{ for general matrices } A, B.$$

[122] We can also prove existence for non-diagonalizable matrices using a continuity argument, though this proof doesn't tell us how to compute $\exp(A)$ in the non-diagonalizable case. The computation of $\exp(A)$ for non-diagonalizable A uses the Jordan canonical form.

The proof of $\exp(x+y) = \exp(x)\exp(y)$ relied on the fact that scalars commute. If $AB = BA$ then this same proof carries over, and we have

$$\boxed{\exp(A+B) = \exp(A)\exp(B) \text{ for matrices satisfying } AB = BA.}$$

Later we will see that the matrix exponential is the key to solving differential equations. In that context we will consider the series

$$\exp(At) = I + At + \frac{t^2}{2!}A^2 + \frac{t^3}{3!}A^3 + \cdots,$$

where t is a real variable representing time.

For now, we present two example computations. First consider the matrix

$$A = \frac{1}{6}\begin{pmatrix} 5 & 4 \\ 2 & -2 \end{pmatrix}.$$

The characteristic polynomial is

$$\begin{aligned} \det(xI - A) &= (x - 5/6)(x + 2/6) - (-4/6)(-2/6) \\ &= x^2 - (1/2)x - (1/2) \\ &= (x-1)(x+1/2), \end{aligned}$$

hence the eigenvalues are 1 and $-1/2$. Since this 2×2 matrix has 2 distinct eigenvalues, we know that it is diagonalizable. After some computation we find the eigenvectors:

$$A\begin{pmatrix} 4 \\ 1 \end{pmatrix} = 1\begin{pmatrix} 4 \\ 1 \end{pmatrix} \quad \text{and} \quad A\begin{pmatrix} 1 \\ -2 \end{pmatrix} = -\frac{1}{2}\begin{pmatrix} 1 \\ -2 \end{pmatrix}.$$

Hence we obtain the diagonalization:

$$A = \left(\begin{array}{c|c} 4 & 1 \\ 1 & -2 \end{array}\right)\begin{pmatrix} 1 & 0 \\ 0 & -1/2 \end{pmatrix}\left(\begin{array}{c|c} 4 & 1 \\ 1 & -2 \end{array}\right)^{-1} = \frac{1}{9}\begin{pmatrix} 4 & 1 \\ 1 & -2 \end{pmatrix}\begin{pmatrix} 1 & 0 \\ 0 & -1/2 \end{pmatrix}\begin{pmatrix} 2 & 1 \\ 1 & -4 \end{pmatrix}.$$

Finally, we obtain the exponential:

$$\exp(A) = \frac{1}{9}\begin{pmatrix} 4 & 1 \\ 1 & -2 \end{pmatrix}\begin{pmatrix} \exp(1) & 0 \\ 0 & \exp(-1/2) \end{pmatrix}\begin{pmatrix} 2 & 1 \\ 1 & -4 \end{pmatrix}.$$

The last example is more interesting. Consider the matrix that rotates by 90°:

$$R = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

For any real number θ, we will show that

$$\exp(R\theta) = \begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos t \end{pmatrix} = \cos\theta \cdot \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \sin\theta \cdot \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$

which is the matrix that rotates by θ. This is a matrix version of Euler's formula

$$e^{i\theta} = \cos\theta + i\sin\theta,$$

where the 90° rotation matrix R plays the role of the imaginary unit i. We begin by computing the eigenvalues. The characteristic polynomial is

$$\det(xI - R) = \det\begin{pmatrix} x & 1 \\ -1 & x \end{pmatrix} = x^2 + 1,$$

hence there are two distinct eigenvalues: i and $-i$. It is no surprise that these are complex conjugates, since the complex eigenvalues of real matrices come in conjugate pairs. (See the homework.) With a bit of work, one finds the eigenvectors

$$R\begin{pmatrix} 1 \\ -i \end{pmatrix} = i\begin{pmatrix} 1 \\ -i \end{pmatrix} \quad \text{and} \quad R\begin{pmatrix} 1 \\ i \end{pmatrix} = -i\begin{pmatrix} 1 \\ i \end{pmatrix},$$

and hence the exponential:

$$\exp(R\theta) = \begin{pmatrix} 1 & 1 \\ -i & i \end{pmatrix}\begin{pmatrix} \exp(i\theta) & 0 \\ 0 & \exp(-i\theta) \end{pmatrix}\begin{pmatrix} 1 & 1 \\ -i & i \end{pmatrix}^{-1}.$$

Then some simplification using Euler's formula $e^{i\theta} = \cos\theta + i\sin\theta$ gives the desired result.

But this makes the result look like a miracle. We can gain more insight by looking at the real and imaginary parts of the complex eigenvalues. Let $\mathbf{x} = (1, -i)$, so that $\mathbf{x} = (1, 0) - i(0, 1)$. Since $R\mathbf{x} = i\mathbf{x}$ we must also have $\exp(R\theta)\mathbf{x} = \exp(i\theta)\mathbf{x}$,[123] and hence

$$\begin{aligned}
\exp(R\theta)\begin{pmatrix} 1 \\ 0 \end{pmatrix} - i\exp(R\theta)\begin{pmatrix} 0 \\ 1 \end{pmatrix} &= \exp(R\theta)\begin{pmatrix} 1 \\ -i \end{pmatrix} \\
&= \exp(i\theta)\begin{pmatrix} 1 \\ -i \end{pmatrix} \\
&= (\cos\theta + i\sin\theta)\begin{pmatrix} 1 \\ -i \end{pmatrix} \qquad \text{Euler's formula} \\
&= \begin{pmatrix} \cos\theta + i\sin\theta \\ \sin\theta - i\cos\theta \end{pmatrix} \\
&= \begin{pmatrix} \cos\theta \\ \sin\theta \end{pmatrix} - i\begin{pmatrix} -\sin\theta \\ \cos\theta \end{pmatrix}.
\end{aligned}$$

Since the matrix $\exp(R\theta)$ has real entries, comparing real and imaginary parts gives

$$\exp(R\theta)\begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} \cos\theta \\ \sin\theta \end{pmatrix} \quad \text{and} \quad \exp(R\theta)\begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} -\sin\theta \\ \cos\theta \end{pmatrix},$$

which is the desired result.

Remark: In general, complex eigenvalues of real matrices lead to rotation. We will examine this in the next section.

[123] The proof that $A\mathbf{x} = \lambda\mathbf{x}$ implies $f(A)\mathbf{x} = f(\lambda)\mathbf{x}$ for polynomials $f(x)$ carries over to power series.

6.6 Complex Eigenvalues and Eigenvectors of Real Matrices

For any complex number $a + ib \in \mathbb{C}$ with $a, b \in \mathbb{R}$ we will denote its complex conjugate by

$$\overline{\alpha} = \overline{a + ib} = a - ib.$$

Recall that complex conjugation satisfies the following properties:

- $\overline{\alpha + \beta} = \overline{\alpha} + \overline{\beta}$,
- $\overline{\alpha\beta} = \overline{\alpha} \cdot \overline{\beta}$,
- $\overline{\alpha^k} = (\overline{\alpha})^k$,
- $\overline{\alpha} = \alpha$ if and only if $\alpha \in \mathbb{R}$.

Given a polynomial $f(x) = b_0 + b_1 x + \cdots + b_n x^n$ with **real coefficients** and a complex number $\alpha \in \mathbb{C}$, it follows from these properties that

$$\begin{aligned}
\overline{f(\alpha)} &= \overline{b_0 + b_1\alpha + \cdots + b_n\alpha^n} \\
&= \overline{b_0} + \overline{b_1} \cdot \overline{\alpha} + \cdots + \overline{b_n} \cdot \overline{\alpha^n} \\
&= b_0 + b_1\overline{\alpha} + \cdots + b_n(\overline{\alpha})^n \\
&= f(\overline{\alpha}).
\end{aligned}$$

In particular, we see that α is a root of $f(x)$ if and only if $\overline{\alpha}$ is a root of $f(x)$. Indeed, if $f(\alpha) = 0$ then

$$f(\overline{\alpha}) = \overline{f(\alpha)} = \overline{0} = 0,$$

and if $f(\overline{\alpha}) = 0$ then

$$f(\alpha) = \overline{\overline{f(\alpha)}} = \overline{f(\overline{\alpha})} = \overline{0} = 0.$$

It follows from this that

the non-real roots of a real polynomial come in conjugate pairs.

And, as an interesting consequence,

every real polynomial of odd degree has as least one real root.

We will apply these observations to eigenvalues of real matrices.

Complex Eigenvalues of a Real Matrix. Let A be an $n \times n$ matrix with **real entries**, so the characteristic polynomial $\chi_A(x) = \det(xI - A)$, has **real coefficients**. According to the previous result, the characteristic polynomial can be factored as

$$\chi_A(x) = (x - \lambda_1) \cdots (x - \lambda_{n-2m})(x - \alpha_1)(x - \overline{\alpha_1}) \cdots (x - \alpha_m)(x - \overline{\alpha_m}),$$

for some real numbers $\lambda_i \in \mathbb{R}$ and non-real complex numbers $\alpha_i \in \mathbb{C}$. If n is even, then the matrix A need not have any real eigenvalues. For example, the real matrix

$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$

has characteristic polynomial $(x-i)(x+i)$. On the other hand, if n is odd then the number $n-2m$ must be ≥ 1, so that A has at least one real eigenvalue.

Complex Eigenvectors. If a real matrix A has a real eigenvalue λ, then the corresponding eigenvectors are real.[124] Indeed, the space of λ-eigenvectors is the null space $\mathcal{N}(\lambda I - A)$, which can be computed by elimination over $\mathbb{R}$. What about complex eigenvalues? Suppose that a real $n \times n$ matrix A has a complex eigenvalue $\lambda \in \mathbb{C}$, and let $\mathbf{x} \in \mathbb{C}^n$ be a corresponding eigenvector:

$$A\mathbf{x} = \lambda\mathbf{x}.$$

If λ is not real then $\mathbf{x}$ cannot have real entries. Indeed, if $\mathbf{x} \in \mathbb{R}^n$ then since A has real entries we would have $\lambda\mathbf{x} = A\mathbf{x} \in \mathbb{R}^n$ which implies that $\lambda \in \mathbb{R}$. Let us suppose that $\lambda = a + ib$ with $a, b \in \mathbb{R}$ and $b \neq 0$. Then we can write

$$\mathbf{x} = \mathbf{u} + i\mathbf{v}$$

for unique real vectors $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ with $\mathbf{v} \neq \mathbf{0}$. By expanding the equation $A\mathbf{x} = \lambda\mathbf{x}$ we obtain

$$\begin{aligned} A\mathbf{u} + iA\mathbf{v} &= A(\mathbf{u} + i\mathbf{v}) \\ &= A\mathbf{x} \\ &= \lambda\mathbf{x} \\ &= (a+ib)(\mathbf{u} + i\mathbf{v}) \\ &= (a\mathbf{u} - b\mathbf{v}) + i(b\mathbf{u} + a\mathbf{v}). \end{aligned}$$

Since the vectors $A\mathbf{u}$, $A\mathbf{v}$, $a\mathbf{u} - \mathbf{b}\mathbf{v}$ and $b\mathbf{u} + a\mathbf{v}$ have real entries, it follows by comparing real and imaginary parts that

$$\begin{cases} A\mathbf{u} &= a\mathbf{u} &- b\mathbf{v}, \\ A\mathbf{v} &= b\mathbf{u} &+ a\mathbf{v}, \end{cases}$$

which can be expressed as a matrix equation:

$$A \begin{pmatrix} \mathbf{u} \mid \mathbf{v} \end{pmatrix} = \begin{pmatrix} \mathbf{u} \mid \mathbf{v} \end{pmatrix} \begin{pmatrix} a & b \\ -b & a \end{pmatrix}.$$

Next, I claim that the vectors $\mathbf{u}$ and $\mathbf{v}$ are linearly independent over $\mathbb{C}$.[125] To see this, we note that the conjugate vector $\overline{\mathbf{x}} = \mathbf{u} - i\mathbf{v}$ is an eigenvector of A corresponding to the conjugate eigenvalue $\overline{\lambda} = a - ib$. Indeed, since A has real entries, conjugating both sides of the equation $A\mathbf{x} = \lambda\mathbf{x}$ gives[126]

$$A\overline{\mathbf{x}} = \overline{A}\overline{\mathbf{x}} = \overline{A\mathbf{x}} = \overline{\lambda\mathbf{x}} = \overline{\lambda}\overline{\mathbf{x}}.$$

Since $\lambda \neq \overline{\lambda}$, the vectors $\mathbf{x}, \overline{\mathbf{x}} \in \mathbb{C}$ correspond to different eigenvalues, hence they are linearly independent over $\mathbb{C}$. But then since

$$\begin{pmatrix} \mathbf{x} \mid \overline{\mathbf{x}} \end{pmatrix} = \begin{pmatrix} \mathbf{u} \mid \mathbf{v} \end{pmatrix} \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix}, \quad \text{where } \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix} \text{ is invertible,}$$

[124] Technically, every λ-eigenvector is a scalar multiple of a real vector. You could take a real λ-eigenvector $\mathbf{x}$ and scale it to get a complex λ-eigenvector $i\mathbf{x}$, but why would you want to do that?

[125] In this section we will only discuss linear independence over $\mathbb{C}$, which implies linear independence over $\mathbb{R}$.

[126] The equation $\overline{A\mathbf{x}} = \overline{A}\overline{\mathbf{x}}$ needs to be checked. It follows from the standard properties.

we conclude that $\mathbf{u}$ and $\mathbf{v}$ are linearly independent. In particular, we have

$$A = \begin{pmatrix} \mathbf{u} \mid \mathbf{v} \end{pmatrix} \begin{pmatrix} a & b \\ -b & a \end{pmatrix} \begin{pmatrix} \mathbf{u} \mid \mathbf{v} \end{pmatrix}^{-1}.$$

Furthermore, for any $\mathbf{y} \in \mathbb{C}^n$ we have

$$\begin{pmatrix} \mathbf{y} \mid \mathbf{x} \mid \overline{\mathbf{x}} \end{pmatrix} = \begin{pmatrix} \mathbf{y} \mid \mathbf{u} \mid \mathbf{v} \end{pmatrix} \begin{pmatrix} 1 & & \\ & 1 & 1 \\ & i & -i \end{pmatrix},$$

which implies that the set $\mathbf{y}, \mathbf{x}, \overline{\mathbf{x}}$ is independent if and only if $\mathbf{y}, \mathbf{u}, \mathbf{v}$ is independent.

Real Diagonalizable Matrices. Finally, we discuss diagonalization of real matrices. Let A be a real $n \times n$ matrix. As we saw above, the characteristic polynomial can be factored as

$$\chi_A(x) = (x - \lambda_1) \cdots (x - \lambda_{n-2m})(x - \alpha_1)(x - \overline{\alpha_1}) \cdots (x - \alpha_m)(x - \overline{\alpha_m}),$$

where $\lambda_1, \ldots, \lambda_{n-2m}$ are real and $\alpha_1, \ldots, \alpha_m \in \mathbb{C}$ are non-real.

Suppose that A is diagonalizable over $\mathbb{C}$. This means that we can find nonzero vectors $\mathbf{y}_1, \ldots, \mathbf{y}_{n-2m} \in \mathbb{C}^n$ and $\mathbf{x}_1, \ldots, \mathbf{x}_m \in \mathbb{C}^n$ such that $A\mathbf{y}_i = \lambda_i \mathbf{y}_i$ and $A\mathbf{x}_i = \alpha_i \mathbf{x}_i$, hence also $A\overline{\mathbf{x}_i} = \overline{\alpha_i \mathbf{x}_i}$, and such that

$$\mathbf{y}_1, \ldots, \mathbf{y}_{n-2m}, \mathbf{x}_1, \overline{\mathbf{x}_1}, \ldots, \mathbf{x}_m, \overline{\mathbf{x}_m}$$

is a basis for $\mathbb{C}^n$. If X is the $n \times n$ (invertible) matrix with these column vectors, then we have

$$A = X \begin{pmatrix} \lambda_1 & & & & & & & & \\ & \ddots & & & & & & & \\ & & \lambda_{n-2m} & & & & & & \\ & & & \alpha_1 & & & & & \\ & & & & \overline{\alpha_1} & & & & \\ & & & & & \ddots & & & \\ & & & & & & \alpha_m & & \\ & & & & & & & \overline{\alpha_m} \end{pmatrix} X^{-1}.$$

Now we will eliminate the complex numbers from this factorization. Since the eigenvalues $\lambda_1, \ldots, \lambda_{n-2m}$ are real, we can choose the eignevectors $\mathbf{y}_1, \ldots, \mathbf{y}_{n-2m}$ to be real. Next we write $\mathbf{x}_i = \mathbf{u}_i + i\mathbf{v}_i$ for real vectors $\mathbf{u}_i, \mathbf{v}_i$ with $\mathbf{v}_i \neq \mathbf{0}$. From the previous remarks we see that

$$\mathbf{y}_1, \ldots, \mathbf{y}_{n-2m}, \mathbf{u}_1, \mathbf{v}_1, \ldots, \mathbf{u}_m, \mathbf{v}_m$$

is a basis for $\mathbb{C}^n$ consisting of real vectors. Furthermore, if Y is the (invertible) matrix with these columns, then we have

$$A = Y \begin{pmatrix} \lambda_1 & & & & & & & \\ & \ddots & & & & & & \\ & & \lambda_{n-2m} & & & & & \\ & & & a_1 & b_1 & & & \\ & & & -b_1 & a_1 & & & \\ & & & & & \ddots & & \\ & & & & & & a_m & b_m \\ & & & & & & -b_m & a_m \end{pmatrix} Y^{-1}.$$

This is not quite a "diagonalization", but it has the virtue using only real numbers.

6.7 Normal Operators

In this chapter we have studied the evaluation of polynomials (also power series and Laurent polynomials) at matrices. This discussion has left out one important operation; namely, the transpose and conjugate transpose. In this final section we consider the relationship between eigenvalues and (conjugate) transposition.

The main role is played by *normal matrices*. We say that a matrix A is normal when it commutes with its (conjugate) transpose:

$$A^*A = AA^*.$$

These matrices are extremely common in applications and include the following four families:

- Real symmetric matrices $A^T = A$.
- Complex Hermitian matrices $A^* = A$.
- Real orthogonal matrices $A^T = A^{-1}$.
- Complex unitary matrices $A^* = A^{-1}$.

Of course, these families could be dealt with separately. The reason to combine them under the concept of normal matrices is because of the following fundamental theorem, which we will prove in the next chapter.

The Spectral Theorem. Let A be a square matrix over $\mathbb{R}$ or $\mathbb{C}$. Then

$$A^*A = AA^* \quad \Longleftrightarrow \quad A \text{ has an orthonormal basis of eigenvectors.}$$

Actually, some people think it undignified to call this the Spectral Theorem. They say that the true Spectral Theorem applies to operators on infinite dimensional Hilbert spaces. Recall,

if V is a real or complex Hilbert space and if $A : V \to V$ is a bounded[127] linear operator then there exists a unique bounded linear operator $A^* : V \to V$ satisfying

$$\langle A\mathbf{x}, \mathbf{y}\rangle = \langle \mathbf{x}, A^*\mathbf{y}\rangle \quad \text{for all } \mathbf{x}, \mathbf{y} \in V.$$

As with many results in functional analysis, the proof is 80% algebra and 20% analysis, which is mostly plausible from geometric intuition.

Anyway, it is convenient to state and prove the results of this section in a language that applies also to Hilbert spaces. Our first theorem was proved by Cauchy in 1829, as part of his extension of the Principal Axes Theorem to higher dimensions. Cauchy's original proof was quite complicated, but today's proof is a one-liner.[128]

Cauchy's Reality Theorem. A real symmetric matrix has real eigenvalues.

Actually, we will prove the following more general statement, since it has the same proof.

Theorem. A self-adjoint operator on a complex inner product space has real eigenvalues.

Proof. Let V be a real or complex inner product space and let $A : V \to V$ be an operator satisfying $A^* = A$. If $A\mathbf{x} = \lambda\mathbf{x}$ for some scalar λ and nonzero vector $\mathbf{x} \neq \mathbf{0}$ then we have

$$\begin{aligned}
\lambda\|\mathbf{x}\|^2 &= \lambda\langle \mathbf{x}, \mathbf{x}\rangle \\
&= \langle \mathbf{x}, \lambda\mathbf{x}\rangle \\
&= \langle \mathbf{x}, A\mathbf{x}\rangle \\
&= \langle A^*\mathbf{x}, \mathbf{x}\rangle \\
&= \langle A\mathbf{x}, \mathbf{x}\rangle && A^* = A \\
&= \langle \lambda\mathbf{x}, \mathbf{x}\rangle \\
&= \overline{\lambda}\langle \mathbf{x}, \mathbf{x}\rangle \\
&= \overline{\lambda}\|\mathbf{x}\|^2.
\end{aligned}$$

Since $\|\mathbf{x}\| \neq 0$ this implies that $\overline{\lambda} = \lambda$, and hence λ is real. □

The next theorem has a similar proof.

Theorem. Unitary (and real orthogonal) operators have eigenvalues of length 1. That is, they have eigenvalues of the form $e^{i\theta}$.

Proof. Let V be a real or complex inner product space and let $A : V \to V$ be an operator satisfying $A^*A = I$. If $A\mathbf{x} = \lambda\mathbf{x}$ for some scalar λ and nonzero vector $\mathbf{x} \neq \mathbf{0}$ then we have

$$\|\mathbf{x}\|^2 = \langle \mathbf{x}, \mathbf{x}\rangle$$

[127] This means that A sends the unit ball to a bounded set. It is equivalent to A being continuous.

[128] "Dazzled by the brilliance of the new theory of determinants, mathematicians overlooked simple inner product considerations", Hawkins, *The Mathematics of Frobenius in Context*, page 98.

$$
\begin{aligned}
&= \langle \mathbf{x}, I\mathbf{x} \rangle \\
&= \langle \mathbf{x}, A^*A\mathbf{x} \rangle && A^*A = I \\
&= \langle A\mathbf{x}, A\mathbf{x} \rangle \\
&= \langle \lambda\mathbf{x}, \lambda\mathbf{x} \rangle \\
&= \lambda\langle \lambda\mathbf{x}, \mathbf{x} \rangle \\
&= \overline{\lambda}\lambda\langle \mathbf{x}, \mathbf{x} \rangle \\
&= |\lambda|^2 \|\mathbf{x}\|^2.
\end{aligned}
$$

Since $\|\mathbf{x}\| \neq 0$ this implies that $|\lambda| = 1$. □

Though it doesn't involve eigenvalues, we should probably include the following result.

Theorem. Unitary operators preserve lengths and angles.

Proof. This follows from the fact that unitary operators preserve the inner product. If $A^*A = I$ then for all vectors $\mathbf{x}, \mathbf{y}$ we have

$$\langle A\mathbf{x}, A\mathbf{y} \rangle = \langle \mathbf{x}, A^*A\mathbf{y} \rangle = \langle \mathbf{x}, I\mathbf{y} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle.$$

□

We have seen that the important families of normal matrices have quite restricted eigenvalues:

- Real symmetric matrices $A^T = A$ and complex Hermitian matrices $A^* = A$ matrices have real eigenvalues.
- Real orthogonal matrices $A^TA = I$ and complex unitary matrices $A^*A = I$ have eigenvalues of the form $e^{i\theta}$.

On the other hand, a general normal matrix can have any eigenvalues you want. Indeed, consider any complex numbers $\lambda_1, \ldots, \lambda_n \in \mathbb{C}$ and let Λ be the diagonal matrix with these numbers on the diagonal. Then for any unitary matrix $U^*U = I$, the matrix

$$A = U\Lambda U^*$$

is normal and has eigenvalues $\lambda_1, \ldots, \lambda_n$.

What about eigen**vectors**? In this case, the key property is shared by all normal operators. This result is a precursor to the Spectral Theorem.

Theorem (Normal with Distinct Eigenvalues ⇒ Orthogonal Eigenvectors). Let $A^*A = AA^*$ be a normal operator on an inner product space. Then

$$A\mathbf{x} = \lambda\mathbf{x} \text{ and } A\mathbf{y} = \mu\mathbf{y} \text{ with } \lambda \neq \mu \quad \Longrightarrow \quad \langle \mathbf{x}, \mathbf{y} \rangle = 0.$$

We will work up to the proof by a series of lemmas.

Lemma 1. If $A^*A = AA^*$ then $\langle A\mathbf{x}, A\mathbf{y}\rangle = \langle A^*\mathbf{x}, A^*\mathbf{y}\rangle$ for all $\mathbf{x}, \mathbf{y}$.

Proof. We have $\langle A\mathbf{x}, A\mathbf{y}\rangle = \langle \mathbf{x}, A^*A\mathbf{y}\rangle = \langle \mathbf{x}, AA^*\mathbf{y}\rangle = \langle A^*\mathbf{x}, A^*\mathbf{y}\rangle$. □

Lemma 2. If $A^*A = AA^*$ then we have $A\mathbf{x} = 0$ if and only if $A^*\mathbf{x} = \mathbf{0}$.

Proof. Putting $\mathbf{y} = \mathbf{x}$ in Lemma 1 gives $\|A\mathbf{x}\|^2 = \langle A\mathbf{x}, A\mathbf{x}\rangle = \langle A^*\mathbf{x}, A^*\mathbf{x}\rangle = \|A^*\mathbf{x}\|^2$. Hence

$$A\mathbf{x} = \mathbf{0} \quad \Longleftrightarrow \quad \|A\mathbf{x}\| = 0 \quad \Longleftrightarrow \quad \|A^*\mathbf{x}\| = 0 \quad \Longleftrightarrow \quad A^*\mathbf{x} = \mathbf{0}.$$

□

Lemma 3. Let $A^*A = AA^*$. Then for any vector $\mathbf{x}$ and scalar λ we have

$$A\mathbf{x} = \lambda\mathbf{x} \quad \Longleftrightarrow \quad A^*\mathbf{x} = \overline{\lambda}\mathbf{x}.$$

Proof. Consider the matrix $B = \lambda I - A$, with $B^* = \overline{\lambda}I - A^*$. Then B is normal:

$$\begin{aligned} B^*B &= (\overline{\lambda}I - A^*)(\lambda I - A) \\ &= \overline{\lambda}\lambda I - \overline{\lambda}A - \lambda A^* + A^*A \\ &= \overline{\lambda}\lambda I - \overline{\lambda}A - \lambda A^* + AA^* && A^*A = AA^* \\ &= (\lambda I - A)(\overline{\lambda}I - A^*) \\ &= BB^*. \end{aligned}$$

Hence applying Lemma 2 gives

$$A\mathbf{x} = \lambda\mathbf{x} \quad \Longleftrightarrow \quad B\mathbf{x} = \mathbf{0} \quad \Longleftrightarrow \quad B^*\mathbf{x} = \mathbf{0} \quad \Longleftrightarrow \quad A^*\mathbf{x} = \overline{\lambda}\mathbf{x}.$$

Proof of the Theorem. Let $A^*A = AA^*$ and suppose that $A\mathbf{x} = \lambda\mathbf{x}$ and $A\mathbf{y} = \mu\mathbf{y}$ with $\lambda \neq \mu$. Then from Lemma 3 we have $A^*\mathbf{x} = \overline{\lambda}\mathbf{x}$, hence

$$\begin{aligned} \lambda\langle \mathbf{x}, \mathbf{y}\rangle &= \langle \overline{\lambda}\mathbf{x}, \mathbf{y}\rangle \\ &= \langle A^*\mathbf{x}, \mathbf{y}\rangle && \text{Lemma 3} \\ &= \langle \mathbf{x}, A\mathbf{y}\rangle \\ &= \langle \mathbf{x}, \mu\mathbf{y}\rangle \\ &= \mu\langle \mathbf{x}, \mathbf{y}\rangle. \end{aligned}$$

Finally, since $(\lambda - \mu)\langle \mathbf{x}, \mathbf{y}\rangle = 0$ and $\lambda \neq \mu$ we have $\langle \mathbf{x}, \mathbf{y}\rangle = 0$. □

In particular, this shows that an $n \times n$ normal matrix $A^*A = AA^*$ with n distinct eigenvalues has an orthogonal basis of eigenvectors. The Spectral Theorem says that this is still true even when A has repeated eigenvalues. The hard part is to show that there are **enough** eigenvectors to fill up the whole space. See the next chapter.

NOTE TO SELF: Maybe I should discuss the "polarization" $A = (A + A^*)/2 + (A - A^*)/2$, i.e., the "real and imaginary parts" of A. This plays a role later in the proof of the Spectral Theorem.

7 Factorization Theorems

7.1 Gram-Schmidt and QR Factorization

Most of the theorems in this chapter deal with orthonormal bases. In this section we lay the groundwork by showing how any basis can be converted into an orthonormal basis. The procedure is quite general. First we consider an infinite dimensional inner product space V over $\mathbb{R}$ or $\mathbb{C}$. Afterwards we will consider finite dimensional spaces and matrices.

Given any linearly independent set $\mathbf{a}_1, \mathbf{a}_2, \ldots \in V$ the Gram-Schmidt procedure produces linearly independent vectors $\mathbf{b}_1, \mathbf{b}_2, \ldots \in V$ with the following properties:[129]

- $\langle \mathbf{b}_i, \mathbf{b}_j \rangle = 0$ for $i \neq j$,
- $\text{Span}\{\mathbf{a}_1, \ldots, \mathbf{a}_k\} = \text{Span}\{\mathbf{b}_1, \ldots, \mathbf{b}_k\}$.

The definition is recursive:

- First set $\mathbf{b}_1 := \mathbf{a}_1$.
- Then for any $k \geq 1$ set $\mathbf{b}_{k+1} := \mathbf{a}_{k+1} - \text{Proj}_k(\mathbf{a}_{k+1})$, where $\text{Proj}_k : V \to V$ is the orthogonal projection onto the subspace spanned by $\mathbf{b}_1, \ldots, \mathbf{b}_k$. To be precise, we set

$$\mathbf{b}_{k+1} := \mathbf{a}_{k+1} - \frac{\langle \mathbf{a}_{k+1}, \mathbf{b}_1 \rangle}{\langle \mathbf{b}_1, \mathbf{b}_1 \rangle} \mathbf{b}_1 - \cdots - \frac{\langle \mathbf{a}_{k+1}, \mathbf{b}_k \rangle}{\langle \mathbf{b}_k, \mathbf{b}_k \rangle} \mathbf{b}_k.$$

You will prove on the homework that this procedure has the desired properties. Afterwards, we can easily turn $\mathbf{b}_1, \mathbf{b}_2, \ldots \in V$ into an ortho**normal** set by dividing each $\mathbf{b}_k$ by its length.

Before applying this to matrices, we give one application to infinite dimensional function spaces. Consider the real Hilbert space $L^2[-1, 1]$ with inner product

$$\langle f(x), g(x) \rangle = \int_{-1}^{1} f(x) g(x) \, dx.$$

And consider the "obvious" basis $1, x, x^2, \ldots \in L^2[-1, 1]$.[130] Note that these functions are not orthogonal. For example,

$$\langle 1, x^2 \rangle = \int_{-1}^{1} x^2 \, dx = \frac{1}{3} x^3 \Big|_{-1}^{1} = \frac{1}{3}(1)^3 - \frac{1}{3}(-1)^3 = \frac{1}{3} + \frac{1}{3} = \frac{2}{3} \neq 0.$$

Applying the Gram-Schmidt procedure to the non-orthogonal basis $1, x, x^2, \ldots$ produces the orthogonal basis of *Legendre polynomials*: $P_0(x), P_1(x), P_2(x), \ldots$. These are used in physics in the study of spherically symmetric potentials. For example, they determine the "shapes"

[129] If $\mathbf{a}_1, \mathbf{a}_2, \ldots$ is a Hilbert space basis, with appropriate convergence properties, then the vectors $\mathbf{b}_1, \mathbf{b}_2, \ldots$ will also be a Hilbert space basis, though we won't prove this.

[130] It can be shown that this is, indeed, a Hilbert space basis.

of electron orbitals. To be precise, we first define the *associated Legendre function* for integers $\ell, m \in \mathbb{Z}$ with $0 \le m \le \ell$:

$$P_\ell^m(x) = (1-x^2)^{m/2} \cdot \frac{d^m}{dx^m} P_\ell(x).$$

Then the radial equation for the shape of the (ℓ, m)-orbital is[131]

$$\rho = (\text{constant}) \cdot |P_\ell^m(\cos\theta)|.$$

Now we turn to matrices. The matrix form of Gram-Schmidt is called QR factorization. Given an invertible $n \times n$ matrix A, we will produce a unitary matrix $Q^*Q = I$ and an upper-triangular matrix R such that $A = QR$. If A has real entries then Q and R will have real entries. In this case $Q^TQ = I$ is real orthogonal.

Let $\mathbf{a}_1, \ldots, \mathbf{a}_n$ be a basis for $\mathbb{C}^n$. Then the Gram-Schmidt basis $\mathbf{b}_1, \ldots, \mathbf{b}_n$ satisfies

$$\begin{aligned}
\mathbf{a}_1 &= \mathbf{b}_1, \\
\mathbf{a}_2 &= \mathbf{b}_2 + \frac{\langle \mathbf{a}_2, \mathbf{b}_1 \rangle}{\langle \mathbf{b}_1, \mathbf{b}_1 \rangle} \mathbf{b}_1, \\
&\vdots \\
\mathbf{a}_n &= \mathbf{b}_n + \frac{\langle \mathbf{a}_n, \mathbf{b}_{n-1} \rangle}{\langle \mathbf{b}_{n-1}, \mathbf{b}_{n-1} \rangle} \mathbf{b}_{n-1} + \cdots + \frac{\langle \mathbf{a}_n, \mathbf{b}_1 \rangle}{\langle \mathbf{b}_1, \mathbf{b}_1 \rangle} \mathbf{b}_1,
\end{aligned}$$

which can be expressed as a matrix equation:

$$A = BU$$

$$\begin{pmatrix} \mathbf{a}_1 \,\middle|\, \cdots \,\middle|\, \mathbf{a}_n \end{pmatrix} = \begin{pmatrix} \mathbf{b}_1 \,\middle|\, \cdots \,\middle|\, \mathbf{b}_n \end{pmatrix} \begin{pmatrix} 1 & \frac{\langle \mathbf{a}_2, \mathbf{b}_1 \rangle}{\langle \mathbf{b}_1, \mathbf{b}_1 \rangle} & \cdots & \cdots & \frac{\langle \mathbf{a}_n, \mathbf{b}_1 \rangle}{\langle \mathbf{b}_1, \mathbf{b}_1 \rangle} \\ & 1 & & & \vdots \\ & & \ddots & & \vdots \\ & & & 1 & \frac{\langle \mathbf{a}_n, \mathbf{b}_{n-1} \rangle}{\langle \mathbf{b}_{n-1}, \mathbf{b}_{n-1} \rangle} \\ & & & & 1 \end{pmatrix}$$

By construction, the columns of B are orthogonal. We can make them ortho**normal** by scaling the kth column $\mathbf{b}_k$ by $1/\|\mathbf{b}_k\|$. If S is the diagonal matrix with entries $1/\|\mathbf{b}_k\|$, then the matrix $Q = BS$ has orthonormal columns $\mathbf{q}_k = \mathbf{b}_k/\|\mathbf{b}_k\|$, hence $Q^*Q = I$. To convert $A = BU$ into $A = QR$ we define $R = S^{-1}U$ so that

$$\begin{aligned}
A &= BU \\
&= B(SS^{-1})U \\
&= (BS)(S^{-1}U) \\
&= QR.
\end{aligned}$$

[131] This example is just for fun. See Griffiths, *Introduction to Quantum Mechanics*, Equation 4.32.

It turns out that the matrix $R = S^{-1}U$ has a nice form. To see this, we first observe that

$$\langle \mathbf{a}_k, \mathbf{b}_k \rangle = \langle \mathbf{b}_k + \text{stuff orthogonal to } \mathbf{b}_k, \mathbf{b}_k \rangle = \langle \mathbf{b}_k, \mathbf{b}_k \rangle = \|\mathbf{b}_k\|^2,$$

which implies that

$$\langle \mathbf{a}_k, \mathbf{q}_k \rangle = \left\langle \mathbf{a}_k, \frac{\mathbf{b}_k}{\|\mathbf{b}_k\|} \right\rangle = \frac{1}{\|\mathbf{b}_k\|} \langle \mathbf{a}_k, \mathbf{b}_k \rangle = \frac{1}{\|\mathbf{b}_k\|} \|\mathbf{b}_k\|^2 = \|\mathbf{b}_i\|.$$

Furthermore, for any $1 \leq i < k$ we have

$$\|\mathbf{b}_i\| \cdot \frac{\langle \mathbf{a}_k, \mathbf{b}_i \rangle}{\langle \mathbf{b}_i, \mathbf{b}_i \rangle} = \|\mathbf{b}_i\| \cdot \frac{\langle \mathbf{a}_k, \mathbf{b}_i \rangle}{\|\mathbf{b}_i\|^2} = \frac{1}{\|\mathbf{b}_i\|} \cdot \langle \mathbf{a}_k, \mathbf{b}_i \rangle = \left\langle \mathbf{a}_k, \frac{\mathbf{b}_i}{\|\mathbf{b}_i\|} \right\rangle = \langle \mathbf{a}_k, \mathbf{q}_i \rangle.$$

Putting these together gives

$$\begin{aligned} R &= S^{-1}U \\ &= \begin{pmatrix} \|\mathbf{b}_1\| & & \\ & \ddots & \\ & & \|\mathbf{b}_n\| \end{pmatrix} \begin{pmatrix} 1 & \frac{\langle \mathbf{a}_2, \mathbf{b}_1 \rangle}{\langle \mathbf{b}_1, \mathbf{b}_1 \rangle} & \cdots & \cdots & \frac{\langle \mathbf{a}_n, \mathbf{b}_1 \rangle}{\langle \mathbf{b}_1, \mathbf{b}_1 \rangle} \\ & 1 & & & \vdots \\ & & \ddots & & \vdots \\ & & & 1 & \frac{\langle \mathbf{a}_n, \mathbf{b}_{n-1} \rangle}{\langle \mathbf{b}_{n-1}, \mathbf{b}_{n-1} \rangle} \\ & & & & 1 \end{pmatrix} \\ &= \begin{pmatrix} \langle \mathbf{a}_1, \mathbf{q}_1 \rangle & \cdots & \langle \mathbf{a}_n, \mathbf{q}_1 \rangle \\ & \ddots & \vdots \\ & & \langle \mathbf{a}_n, \mathbf{q}_n \rangle \end{pmatrix}. \end{aligned}$$

In summary, for any $n \times n$ invertible matrix A with columns $\mathbf{a}_i$, we can find an $n \times n$ unitary matrix $Q^*Q = I$ with columns $\mathbf{q}_i$, such that

$$\begin{pmatrix} \mathbf{a}_1 \mid \cdots \mid \mathbf{a}_n \end{pmatrix} = \begin{pmatrix} \mathbf{q}_1 \mid \cdots \mid \mathbf{q}_n \end{pmatrix} \begin{pmatrix} \langle \mathbf{a}_1, \mathbf{q}_1 \rangle & \cdots & \langle \mathbf{a}_n, \mathbf{q}_1 \rangle \\ & \ddots & \vdots \\ & & \langle \mathbf{a}_n, \mathbf{q}_n \rangle \end{pmatrix}$$

$$A = QR.$$

And if A is real then we can choose Q and R with real entries.

Due to rounding errors, the matrix Q computed from Gram-Schmidt is only approximately orthogonal. It is worth mentioning another method, due to Householder, that gives an exactly orthogonal matrix. This method also has an interesting theoretical consequence:

Any real orthogonal matrix $A^TA = I$ is a composition of reflections.

This method uses the *Householder reflection matrices*:

$$H_{\mathbf{v}} = I - 2\frac{\mathbf{v}\mathbf{v}^T}{\|\mathbf{v}\|^2} \quad \text{for } \mathbf{v} \in \mathbb{R}^n.$$

Recall that


$$P_{\mathbf{v}} = \mathbf{v}(\mathbf{v}\mathbf{v}^T)^{-1}\mathbf{v}^T = \frac{\mathbf{v}\mathbf{v}^T}{\mathbf{v}^T\mathbf{v}} = \frac{\mathbf{v}\mathbf{v}^T}{\|\mathbf{v}\|^2}$$

is the matrix that projects orthogonally onto the line spanned by $\mathbf{v}$, hence $I - P_{\mathbf{v}}$ is the matrix that projects onto the orthogonal hyperplane $\mathbf{v}^\perp = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{x}^T\mathbf{v} = 0\}$. From the remarks in the previous chapter, this implies that $2P_{\mathbf{v}} - I$ is the matrix that reflects across the line $\mathbf{v}$ and $2(I - P\mathbf{v}) - I = I - 2P_{\mathbf{v}} = H_{\mathbf{v}}$ is the matrix that reflects across the hyperplane $\mathbf{v}^\perp$. Since $P_{\mathbf{v}}^2 = P_{\mathbf{v}}$ and $P_{\mathbf{v}}^T = P_{\mathbf{v}}$, we find that

$$H_{\mathbf{v}}^{-1} = H_{\mathbf{v}} \quad \text{and } H_{\mathbf{v}}^T = H_{\mathbf{v}}.$$

In particular, $H_{\mathbf{v}}$ is an orthogonal matrix.

The key trick of the Householder algorithm is that $H_{\mathbf{v}}\mathbf{a} = \mathbf{r}$ for any $\mathbf{a}, \mathbf{r}$ satisfying $\|\mathbf{a}\| = \|\mathbf{r}\|$ and $\mathbf{a} - \mathbf{r} = \mathbf{v}$. Picture:



Here is the algorithm.

Householder QR. We are given an invertible matrix A with first column $\mathbf{a} \in \mathbb{R}^n$.

- Let $\mathbf{r} := (\|\mathbf{a}\|, 0, \ldots, 0)$, $\mathbf{v} := \mathbf{a} - \mathbf{r}$ and $H_1 := H_{\mathbf{v}}$, so that $H_1\mathbf{a} = \mathbf{r}$. Then we have

$$H_1 A = \left(\begin{array}{c|c|c|c} H_1\mathbf{a} & * & \cdots & * \end{array} \right) = \left(\begin{array}{c|c|c|c} \mathbf{r} & * & \cdots & * \end{array} \right) = \left(\begin{array}{c|ccc} \|\mathbf{a}\| & * & \cdots & * \\ \hline 0 & & & \\ \vdots & & A' & \\ 0 & & & \end{array} \right),$$

  for some matrix A' of size $(n-1) \times (n-1)$.

- Let $\mathbf{a}' \in \mathbb{R}^{n-1}$ be the first column of A', let $\mathbf{r}' = (\|\mathbf{a}'\|, 0, \ldots, 0) \in \mathbb{R}^{n-1}$ and let $\mathbf{v}' = \mathbf{a}' - \mathbf{r}'$, so that $H_{\mathbf{v}'}\mathbf{a}' = \mathbf{r}'$. Then the matrix

$$H_2 := \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & H_{\mathbf{v}'} & \\ 0 & & & \end{array}\right),$$

satisfies

$$\begin{aligned} H_2H_1A &= \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & H_{\mathbf{v}'} & \\ 0 & & & \end{array}\right)\left(\begin{array}{c|ccc} \|\mathbf{a}\| & * & \cdots & * \\ \hline 0 & & & \\ \vdots & & A' & \\ 0 & & & \end{array}\right) \\ &= \left(\begin{array}{c|ccc} \|\mathbf{a}\| & * & \cdots & * \\ \hline 0 & & & \\ \vdots & & H_2A' & \\ 0 & & & \end{array}\right) \\ &= \left(\begin{array}{c|c|ccc} \|\mathbf{a}\| & * & * & \cdots & * \\ \hline 0 & \|\mathbf{a}'\| & * & \cdots & * \\ \hline 0 & 0 & & & \\ \vdots & \vdots & & A'' & \\ 0 & 0 & & & \end{array}\right), \end{aligned}$$

for some matrix A'' of size $(n-2) \times (n-2)$. We observe that the matrix H_2 is itself a Householder reflection matrix. To see this, let $\mathbf{w} = \left(\begin{array}{c|c} 0 & (\mathbf{v}')^T \end{array}\right)^T$, so that $\|\mathbf{w}\| = \|\mathbf{v}'\|$. Then we have

$$\begin{aligned} H_{\mathbf{w}} &= I - 2\frac{\mathbf{w}\mathbf{w}^T}{\|\mathbf{w}\|} \\ &= I - \frac{2}{\|\mathbf{v}'\|}\left(\begin{array}{c} 0 \\ \hline \mathbf{v}' \end{array}\right)\left(\begin{array}{c|c} 0 & (\mathbf{v}')^T \end{array}\right) \\ &= \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & I_{n-1} & \\ 0 & & & \end{array}\right) - \left(\begin{array}{c|ccc} 0 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & -2\frac{\mathbf{v}'(\mathbf{v}')^T}{\|\mathbf{v}'\|} & \\ 0 & & & \end{array}\right) \end{aligned}$$

$$= \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & I_{n-1} - 2\frac{\mathbf{v}'(\mathbf{v}')^T}{\|\mathbf{v}'\|} & \\ 0 & & & \end{array}\right)$$

$$= \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & H_{\mathbf{v}'} & \\ 0 & & & \end{array}\right)$$

$$= H_2.$$

- Continuing in this way for $n-1$ steps gives an upper triangular matrix:

$$H_{n-1}\cdots H_2 H_1 A = \begin{pmatrix} \|\mathbf{a}\| & * & \cdots & \cdots & * \\ & \|\mathbf{a}'\| & & & \vdots \\ & & \ddots & & \vdots \\ & & & \|\mathbf{a}^{(n-1)}\| & * \\ & & & & b \end{pmatrix} = R,$$

  where each H_i is a Householder matrix $H_{\mathbf{v}_i}$ for some vector $\mathbf{v}_i \in \mathbb{R}$. Note that the diagonal entries of R are nonzero since we have assumed that A is invertible. The real number b can be positive or negative.

- Finally, since each Householder reflection is equal to its own inverse, we obtain

$$\begin{aligned} H_{n-1}\cdots H_2 H_1 A &= R \\ A &= H_1 H_2 \cdots H_{n-1} R \\ A &= QR. \end{aligned}$$

As a consequence, we will prove that every real orthogonal matrix is a composition of reflections. Suppose that $A^T A = I$ and consider the Householder factorization

$$H_{n-1}\cdots H_2 H_1 A = R.$$

Now each matrix on the left is orthogonal. Since a product of orthogonal matrices is orthogonal, we conclude that R is also orthogonal. In particular, the rows of R are orthonormal. Since R is also upper-triangular, this implies that R is diagonal:

$$R = \begin{pmatrix} \|\mathbf{a}\| & & & \\ & \|\mathbf{a}'\| & & \\ & & \ddots & \\ & & & \|\mathbf{a}^{(n-1)}\| \\ & & & & b \end{pmatrix}$$

Finally, since each row of R has length 1, we conclude that

$$R = \begin{pmatrix} 1 & & & & \\ & 1 & & & \\ & & \ddots & & \\ & & & 1 & \\ & & & & \pm 1 \end{pmatrix}$$

If the last entry is $+1$ then $R = I$ and we obtain

$$A = H_1 H_2 \cdots H_{n-1} I = H_1 H_2 \cdots H_{n-1},$$

which shows that A is a product of $n-1$ reflection matrices. If the last entry of R is -1, then $H_n := R$ equals the Householder matrix $H_{\mathbf{e}_n}$, where $\mathbf{e}_n = (0, \ldots, 0, 1)$. In this case we see that A is a product of n reflection matrices:

$$A = H_1 H_2 \cdots H_{n-1} R = H_1 H_2 \cdots H_{n-1} H_n.$$

Remark: There is some restriction on n. Since each reflection matrix has determinant -1, a product of n reflection matrices has determinant $(-1)^n$. Hence an orthogonal matrix A satisfying $\det(A) = +1$ can only be expressed as an even product of reflections and an orthogonal matrix satisfying $\det(A) = -1$ can only be expressed as an odd product of reflections.

7.2 Schur Triangularization

Given a square matrix A, we always want to find a simpler matrix B that is *similar* to A. That is, we want to find a simpler matrix B and an invertible matrix X such that $A = XBX^{-1}$. Then for any polynomial function $f(x)$ (more generally, for power series or Laurent polynomials) we can compute

$$f(A) = X \cdot f(B) \cdot X^{-1}.$$

The nicest possible situation is when B is diagonal and X is orthogonal or unitary: $X^{-1} = X^T$ or $X^{-1} = X^*$. This is the subject of the Spectral Theorem in the next section. But diagonalization is not always possible. There are three different theorems for dealing with non-diagonalizable matrices:

- Schur triangularization.
- Jordan normal form.
- Singular value decomposition.

We will deal with all three of these in this chapter. We begin with Schur triangularization.

We say that a matrix is upper-triangular if all entries below the main diagonal are zero:[132]

$$T = \begin{pmatrix} t_{11} & * & \cdots & * \\ & t_{22} & & \vdots \\ & & \ddots & * \\ & & & t_{nn} \end{pmatrix}$$

These matrices have some nice properties:

- The eigenvalues of T are the diagonal entries. Indeed, the characteristic polynomial is
$$\chi_T(x) = (x - t_{11})(x - t_{22}) \cdots (x - t_{nn}).$$
- Products and sums of upper-triangular matrices behave as products and sums for the diagonal entries. Thus for any polynomial $f(x)$ we have
$$f(T) = \begin{pmatrix} f(t_{11}) & * & \cdots & * \\ & f(t_{22}) & & \vdots \\ & & \ddots & * \\ & & & f(t_{nn}) \end{pmatrix}$$
Unfortunately, the entries above the diagonal are messy.
- If T is invertible then T^{-1} is also upper-triangular, and the previous formula also applies for Laurent polynomials $f(x)$.
- If the largest eigenvalue satisfies $|\lambda| < 1$ then one can show that $T^k \to O$ as $k \to \infty$, though the proof is a bit tricky.[133]

Here is our main theorem.

Theorem (Schur Triangularization). For **any** square matrix A over $\mathbb{R}$ or $\mathbb{C}$, there exists an upper-triangular matrix T and a unitary matrix $U^{-1} = U^*$ such that

$$A = UTU^{-1}$$
$$A = UTU^*$$
$$\left(\begin{array}{c|c|c|c} \mathbf{a}_1 & \cdots & \mathbf{a}_n \end{array} \right) = \left(\begin{array}{c|c|c|c} \mathbf{u}_1 & \ldots & \mathbf{u}_n \end{array} \right) \begin{pmatrix} t_{11} & * & \cdots & * \\ & t_{22} & & \vdots \\ & & \ddots & * \\ & & & t_{nn} \end{pmatrix} \left(\begin{array}{c} \mathbf{u}_1^* \\ \hline \vdots \\ \hline \mathbf{u}_n^* \end{array} \right).$$

Even if A is real, the matrices U and T will generally have complex entries. However, if A is a real matrix with real eigenvalues then we can choose U and T to be real.

[132] Similarly, a lower-triangular matrix has zeros above the main diagonal.
[133] This is easy for diagonalizable matrices.

Proof. We use induction on the size of A. First we note that the theorem is trivially true for 1×1 matrices, i.e., for scalars: $(a) = (1)(a)(1)$. Now let A have shape $n \times n$ for some $n \geq 2$. We have seen that every real matrix has a (possibly complex) eigenvalue. Let $t_{11} \in \mathbb{C}$ be an eigenvalue of A and let $\mathbf{u}_1$ be a corresponding eigenvector of length 1.[134] Now let U_1 be any unitary matrix with first column $\mathbf{u}_1$:

$$U_1 = \left(\begin{array}{c|c|c|c} \mathbf{u}_1 & \cdots & \mathbf{u}_n \end{array} \right).$$

To find such a matrix, we first complete $\mathbf{u}_1$ to a basis, $\mathbf{u}_1, \mathbf{x}_2, \ldots, \mathbf{x}_n$, then apply Gram-Schmidt to convert this into an orthonormal basis $\mathbf{u}_1, \mathbf{u}_2, \ldots, \mathbf{u}_n$.[135] Since these vectors satisfy $\mathbf{u}_1^* \mathbf{u}_1 = 1$ and $\mathbf{u}_i^* \mathbf{u}_1 = 0$ for $i \geq 2$, we observe that

$$\begin{aligned}
U_1^* A U_1 &= \left(\begin{array}{c} \mathbf{u}_1^* \\ \hline \vdots \\ \hline \mathbf{u}_n^* \end{array} \right) \left(\begin{array}{c|c|c|c} A\mathbf{u}_1 & A\mathbf{u}_2 & \cdots & A\mathbf{u}_n \end{array} \right) \\
&= \left(\begin{array}{c} \mathbf{u}_1^* \\ \hline \vdots \\ \hline \mathbf{u}_n^* \end{array} \right) \left(\begin{array}{c|c|c|c} t_{11}\mathbf{u}_1 & A\mathbf{u}_2 & \cdots & A\mathbf{u}_n \end{array} \right) \\
&= \left(\begin{array}{c|c|c|c} t_{11}\mathbf{u}_1^*\mathbf{u}_1 & * & \cdots & * \\ t_{11}\mathbf{u}_2^*\mathbf{u}_1 & & & \\ \vdots & & & \\ t_{11}\mathbf{u}_n^*\mathbf{u}_1 & & & \end{array} \right) \\
&= \left(\begin{array}{c|ccc} t_{11} & * & \cdots & * \\ \hline 0 & & & \\ \vdots & & A_2 & \\ 0 & & & \end{array} \right),
\end{aligned}$$

for some matrix A_2 of shape $(n-1) \times (n-1)$. By induction there exists an $(n-1) \times (n-1)$ unitary matrix U_2 such that $T_2 := U_2^* A_2 U_2$ is upper-triangular. Now define the matrix

$$U := U_1 \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & U_2 & \\ 0 & & & \end{array} \right).$$

[134] Just take any eigenvector and scale it. If t_{11} is real and if A has real entries then we can choose $\mathbf{u}_1$ to have real entries, in which case we can choose U_1 to have real entries.

[135] Any set of independent vectors can be completed to a basis using Steinitz exchange.

We observe that this matrix is unitary:

$$\begin{aligned}
U^*U &= \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & U_2^* & \\ 0 & & & \end{array}\right) U_1^*U_1 \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & U_2 & \\ 0 & & & \end{array}\right) \\
&= \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & U_2^* & \\ 0 & & & \end{array}\right) \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & U_2 & \\ 0 & & & \end{array}\right) \\
&= \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & U_2^*U_2 & \\ 0 & & & \end{array}\right) \\
&= \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & I_{n-1} & \\ 0 & & & \end{array}\right) \\
&= I_n.
\end{aligned}$$

And we observe that the matrix $T := U^*AU$ is upper triangular, as desired:

$$\begin{aligned}
T &= U^*AU \\
&= \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & U_2^* & \\ 0 & & & \end{array}\right) U_1^*AU_1 \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & U_2 & \\ 0 & & & \end{array}\right) \\
&= \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & U_2^* & \\ 0 & & & \end{array}\right) \left(\begin{array}{c|ccc} t_{11} & * & \cdots & * \\ \hline 0 & & & \\ \vdots & & A_2 & \\ 0 & & & \end{array}\right) \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & U_2 & \\ 0 & & & \end{array}\right) \\
&= \left(\begin{array}{c|ccc} t_{11} & * & \cdots & * \\ \hline 0 & & & \\ \vdots & & U_2^*A_2U_2 & \\ 0 & & & \end{array}\right)
\end{aligned}$$

$$= \left(\begin{array}{c|ccc} t_{11} & * & \cdots & * \\ \hline 0 & & & \\ \vdots & & T_2 & \\ 0 & & & \end{array}\right).$$

□

Before moving on, I will mention one application. If $A = XTX^{-1}$ for some (upper or lower) triangular matrix T, then the eigenvalues of A are the diagonal entries of T. One could imagine using this to **compute** the eigenvalues of A. Unfortunately, the proof of Schur triangularization assumes that we already know the eigenvalues of A.

Nevertheless, this is still the good idea, and it is behind the most powerful algorithm for computing eigenvalues. This algorithm uses the QR factorization (which does not assume knowledge of the eigenvalues) in a surprising way to recursively approximate the Schur decomposition, and hence the eigenvalues. It was discovered in the late 1950s by Francis and Kublanovskaya. I will present only the most basic version. The real world version uses extra tricks and optimizations.

The QR Algorithm for Computing Eigenvalues. Given a square matrix A, we recursively define unitary matrices $Q_1, Q_2, \ldots$ and upper-triangular matrices $R_1, R_2, \ldots$ as follows:

- Compute a QR factorization: $A = Q_1 R_1$.
- Next, compute a QR factorization of the matrix $R_1 Q_1$:[136]

$$R_1 Q_1 = Q_2 R_2.$$

- Continue to compute Q_{k+1} and R_{k+1} from the matrix $R_k Q_k$:

$$R_k Q_k = Q_{k+1} R_{k+1}.$$

Let's write $A_1 := A = Q_1 R_1$ and $A_k := Q_k R_k$. Since the Q in the QR factorization is unitary, we have $R_k = Q_k^* A_k$ and hence

$$A_{k+1} = R_k Q_k = Q_k^* A_k Q_k.$$

This implies that the sequence of matrices $A = A_1, A_2, \ldots$ all have the same eigenvalues. The theorem says the following.

Theorem. Suppose that A has eigenvalues with distinct absolute values:[137]

$$|\lambda_1| > |\lambda_2| > \cdots > |\lambda_n|.$$

[136] This is a strange idea, but it leads to great results.

[137] There are modified versions of the algorithm that work for all square matrices.

Then the matrix $A_k = R_k Q_k$ in the QR algorithm converges to an upper triangular matrix, whose diagonal entries are the eigenvalues of A.

It is difficult to find a proof of this written down.[138] The only full proof I can find is in Wilkinson, *The Algebraic Eigenvalue Problem* (1965), page 516. Here is a sketch.

Sketch of a Proof. Define $\tilde{Q}_k := Q_1 Q_2 \cdots Q_k$ and $\tilde{R}_k = R_1 R_2 \cdots R_k$. Since the sets of unitary matrices and upper triangular matrices are closed under multiplication,[139] we see that $\tilde{Q}_k$ is unitary and $\tilde{R}_k$ is upper triangular. I claim that $A_{k+1} = \tilde{Q}_k^* A \tilde{Q}_k$ and $A^k = \tilde{Q}_k \tilde{R}_k$. Indeed, we have

$$\begin{aligned} A_{k+1} &= Q_k^* A_k Q_k \\ &= Q_k^* Q_{k-1}^* A_{k-1} Q_{k-1} Q_k \\ &\vdots \\ &= Q_k^* \cdots Q_2^* Q_1^* A Q_1 Q_2 \cdots Q_k \\ &= (Q_1 \cdots Q_k)^* A (Q_1 \cdots Q_k) \\ &= \tilde{Q}_k^* A \tilde{Q}_k \end{aligned}$$

and

$$\begin{aligned} A^k &= (Q_1 R_1) \cdots (Q_1 R_1) \\ &= Q_1 (R_1 Q_1) \cdots (R_1 Q_1) R_1 \\ &= Q_1 (Q_2 R_2) \cdots (Q_2 R_2) R_1 \\ &= Q_1 Q_2 (R_2 Q_2) \cdots (R_2 Q_2) R_2 R_1 \\ &= Q_1 Q_2 (Q_3 R_3) \cdots (Q_3 R_3) R_2 R_1 \\ &\vdots \\ &= (Q_1 Q_2 \cdots Q_k)(R_k \cdots R_2 R_1) \\ &= \tilde{Q}_k \tilde{R}_k. \end{aligned}$$

From our assumption that A has distinct eigenvalues, we can diagonalize A as

$$A = X \Lambda X^{-1} = X \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{pmatrix} X^{-1}.$$

By multiplying X^{-1} on the left by elementary matrices we can write $X^{-1} = L\Gamma$ where Γ is upper triangular and L is lower triangular with 1s on the diagonal. The key to the whole proof is to observe that $\Lambda^k L \Lambda^{-k}$ is lower triangular and converges to the identity matrix as

[138] Pure math books tend not to discuss it and applied math books tend not to prove it.
[139] Jargon: These sets are groups.

$k \to \infty$. Indeed, if ℓ_{ij} is the ij entry of L (so that $\ell_{ii} = 1$ and $\ell_{ij} = 0$ when $i < j$) then the ij entry of $\Lambda^k L \Lambda^{-k}$ is

$$(\Lambda^k L \Lambda^{-k})_{ij} = \begin{cases} 0 & i < j, \\ 1 & i = j, \\ \ell_{ij}(\lambda_i/\lambda_j)^k & i > j. \end{cases}$$

Since we have assumed that $|\lambda_1| > \cdots > |\lambda_n|$, it follows that the entries below the diagonal go to zero as $k \to \infty$.

By combining these ingredients, Wilkinson shows that the sequences $\tilde{Q}_k$ and $\tilde{R}_k$ converge, and that the sequence Q_k converges to a diagonal matrix, hence $A_k = Q_k R_k$ converges to an upper triangular matrix. Let's say $\tilde{Q}_k \to U$ and $A_k \to T$, for unitary U and upper triangular T. Then in the limit we obtain the Schur triangularization:

$$A = \tilde{Q}_k A_{k+1} \tilde{Q}_k^* \to UTU^*.$$

□

Remark: The proof uses the fact QR factorization is unique up to multiplication with a unitary diagonal matrix D: $QR = (QD)(D^{-1}R)$. This follows from the fact that any unitary upper triangular matrix must be diagonal.

7.3 The Spectral Theorem

The Spectral Theorem might be viewed as the "fundamental theorem" of spectral theory. It characterizes the best kind of matrices; namely, those that possess an orthonormal basis of eigenvectors. Here is the statement.

The Spectral Theorem. Let A be a square matrix with real or complex entries. Then

$$\boxed{A \text{ has an orthonormal basis of eigenvectors} \iff A \text{ is normal, i.e., } A^*A = AA^*.}$$

Remarks: This theorem also applies to infinite dimensional Hilbert spaces, with suitable notions of continuity and convergence. The original version of the theorem applies to real symmetric matrices $A^T = A$, in which case we can choose the eigenvectors to be real.

You will find many different proofs in the literature. One proof uses the Schur Triangularization, which is convenient because we proved this in the previous section.

Proof using Schur Triangularization. One direction is easy. If A has an orthonormal basis of eigenvectors then we can find unitary $U^*U = I$ and diagonal Λ such that $A = U\Lambda U^*$. But then A is normal because diagonal matrices are normal:

$$\begin{aligned} A^*A &= (U\Lambda U^*)^*(U\Lambda U^*) \\ &= U\Lambda^* U^* U \Lambda U^* \\ &= U\Lambda^*\Lambda U^* \end{aligned}$$

$$
\begin{aligned}
&= U\Lambda\Lambda^*U^* && \Lambda^*\Lambda = \Lambda^*\Lambda \\
&= (U\Lambda U^*)(U\Lambda^*U^*) \\
&= AA^*.
\end{aligned}
$$

Conversely, suppose that $A^*A = AA^*$. By Schur's Theorem, any square matrix A can be written as $A = UTU^*$ where $U^*U = I$ is unitary and T is upper-triangular. Thus we have

$$T = U^*AU,$$

which implies that $T^*T = TT^*$ by the previous argument. Finally, one can check that any upper-triangular matrix T satisfying $T^*T = TT^*$ must be diagonal. This would make a good homework exercise. The proofs for self-adjoint and unitary matrices are easier. If $A^* = A$ then $T = U^*AU$ implies that $T^* = T$. This says that the above diagonal-entries of T are the conjugates of the below-diagonal entries of T, which are zero, and hence T is diagonal. If $A^*A = I$ then $T = U^*AU$ implies that $T^*T = I$, which says that the rows of T are orthogonal. By working from the bottom row to the top, this implies that the above-diagonal entries of T must be zero, hence T is diagonal. □

However, I think it is useful to give a proof from scratch. I will give the full proof for matrices, and I will sketch out the proof for operators on Hilbert spaces. For full details see John B. Conway, *A Course in Functional Analysis*, Chapter 2.

Proof of the Spectral Theorem, not using Schur Triangularization.

Step 1 for Matrices. If $A^*A = AA^*$ then A has an eigenvalue. Indeed, **any** square matrix has an eigenvalue by the Fundamental Theorem of Algebra.

Step 1 for Operators. If $A^*A = AA^*$ and if the set $\{A\mathbf{u} : \|\mathbf{u}\| = 1\}$ is compact (in which case we say that A is a *compact normal operator*) then A has an eigenvalue. I'll just sketch the proof. Since the set $\{A\mathbf{u} : \|\mathbf{u}\| = 1\}$ is compact,[140] the *operator norm* is finite:

$$\|A\| = \sup\{\|A\mathbf{u}\| : \|\mathbf{u}\| = 1\} < \infty.$$

Since $\|A\|$ exists we can find a sequence of unit vectors $\mathbf{v}_1, \mathbf{v}_2, \ldots$ such that $\langle A\mathbf{v}_n, \mathbf{v}_n\rangle \to \lambda$ as $n \to \infty$ for some $\lambda \in \mathbb{C}$ satisfying $|\lambda| = \|A\|$. Then since the set $\{A\mathbf{u} : \|\mathbf{u}\| = 1\}$ is compact we can find a subsequence $\mathbf{v}_{n_1}, \mathbf{v}_{n_2}, \ldots$ so that $A\mathbf{v}_{n_k} \to \mathbf{v}$ as $k \to \infty$ for some nonzero vector $\mathbf{v}$ satisfying $A\mathbf{v} = \lambda\mathbf{v}$, and hence λ is an eigenvalue.

Step 2 for Matrices. Let $A^*A = AA^*$, with shape $n \times n$. By Step 1 we can find an eigenvalue λ_1. Choose any unit length eigenvector $\mathbf{u}_1$ so that $A\mathbf{u}_1 = \lambda_1\mathbf{u}_1$, and complete this to an orthonormal basis $\mathbf{u}_1, \mathbf{u}_2, \ldots, \mathbf{u}_n$. In particular, we have $\langle \mathbf{u}_1, \mathbf{u}_j\rangle$ for all $j > 1$. Since A is normal, I claim that we also have $\langle \mathbf{u}_1, A\mathbf{u}_j\rangle = 0$ for all $j > 1$. Indeed, if A is normal then

[140] This set is the image of the unit ball under A. In finite dimensions this set is an ellipsoid.

we showed in the previous chapter that $A\mathbf{u}_1 = \lambda_1 \mathbf{u}_1$ implies $A^*\mathbf{u}_1 = \bar{\lambda_1}\mathbf{u}_1$. But then we must have

$$\langle \mathbf{u}_1, A\mathbf{u}_j \rangle = \langle A^*\mathbf{u}_1, \mathbf{u}_j \rangle = \langle \bar{\lambda_1}\mathbf{u}_1, \mathbf{u}_j \rangle = \lambda_1 \langle \mathbf{u}_1, \mathbf{u}_j \rangle = 0.$$

If U is the (unitary) matrix with columns $\mathbf{u}_1, \ldots, \mathbf{u}_n$ then we have

$$A = U \left(\begin{array}{c|ccc} \lambda_1 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & A' & \\ 0 & & & \end{array} \right) U^*,$$

for some matrix A' of size $(n-1) \times (n-1)$. The displayed matrix containing λ and A' equals U^*AU. This implies that it is normal, and, in particular, the matrix A' is normal. Now we apply induction exactly as in the proof of Schur Triangularization.

Step 2 for Operators. (This is just a sketch; the details fill pages.) Let $A : V \to V$ be a compact normal operator and set $A_1 := A$. By Step 1 there exists an eigenvalue $\lambda_1 \in \mathbb{C}$ such that $|\lambda_1| = \|A_1\|$. If A_1 is not the zero operator then we have $\|A_1\| \neq 0$ and hence $\lambda_1 \neq 0$. Consider the eigenspace $V_1 = ker(\lambda_1 I - A_1) \subseteq V$ and its orthogonal complement $V_1^\perp \subseteq V$. Since A_1 is a compact operator, one can show that the eigenspace V_1 is finite dimensional, say $\dim V_1 = n_1$. Pick an orthonormal basis $\mathbf{u}_{11}, \ldots, \mathbf{u}_{1n_1} \in V_1$. Then since A_1 is normal, the same proof as for matrices shows that A_1 sends $V_1^\perp$ to itself.[141] Let $A_2 : V_1^\perp \to V_1^\perp$ denote the restriction of A_1 to $V_1^\perp$. One can show that A_2 is compact and normal, with $\|A_2\| < \|A_1\|$. If $A_2 \neq 0$ then by Step 1 there exists a nonzero eigenvalue $\lambda_2 \in \mathbb{C}$ with $|\lambda_2| = \|A_2\|$ and with a finite dimensional eigenspace $V_2 = ker(\lambda_2 I - A_2)$. Say $\dim A_2 = n_2$ and pick an orthonormal basis $\mathbf{u}_{21}, \ldots, \mathbf{u}_{2n_2} \in V_2$. Continuing in this way we obtain a sequence of finite dimensional eigenspaces $V_1, V_2, \ldots$ corresponding to eigenvalues with $|\lambda_1| > |\lambda_2| > \cdots$. Finally, one can show that the concatenation of the bases $\mathbf{u}_{k1}, \mathbf{u}_{k2}, \ldots, \mathbf{u}_{kn_k} \in V_k$ is an orthonormal basis for V. The remaining issue is to show that every vector $\mathbf{v} \in V$ can be expressed as a convergent series of eigenvectors $\mathbf{v} = \mathbf{v}_1 + \mathbf{v}_2 + \cdots$, with $\mathbf{v}_k \in V_k$. □

In applications it is usually safe to ignore convergence issues and just the problem at an algebraic level. For example,

7.4 The Singular Value Decomposition

A form of "generalized diagonalization" that applies to rectangular matrices and non-diagonalizable square matrices.

For any $m \times n$ matrix A and $n \times m$ matrix B, the square matrices AB $(m \times m)$ and BA $(n \times n)$ have the same nonzero eigenvalues. If $m < n$ then the matrix BA has $n - m$ extra zero eigenvalues compared to AB.

For any $m \times n$ matrix A, the eigenvalues of A^TA are real and non-negative: $\lambda_1 \geq \cdots \geq \lambda_n \geq 0$. The singular values of A are the non-negative real square roots of the eigenvalues: $\sigma_i = \sqrt{\lambda_i}$. Equivalently, $\sigma_1^2, \ldots, \sigma_n^2$ are the eigenvalues of A^TA.

[141] This is the key point in the proof.

Properties: The largest singular value σ_1 is the operator norm $\|A\|$. The product of the singular values is $\sqrt{\det(A^TA)}$.

Let Σ be the $n \times n$ diagonal matrix of singular values, so $\Lambda = \Sigma^2 = \Sigma^T\Sigma = \Sigma\Sigma^T$ is the diagonal matrix of eigenvalues. From the spectral theorem there exists a unitary (orthogonal) matrix V such that $A^TA = V\Lambda V^T = V\Sigma\Sigma^TV^T = (V\Sigma)(V\Sigma)^T$. The columns $\mathbf{v}_i$ of V are the eigenvectors of A^TA.

Suppose that A^TA has rank r, which is also the rank of A and AA^T, so that there are r nonzero singular values $\sigma_1 \geq \cdots \geq \sigma_r$. From our first remark, A^TA and AA^T have the same non-zero eigenvalues. Define $\mathbf{u}_i = (A\mathbf{v}_i)/\sigma_i$ for $1 \leq i \leq r$. Then $\mathbf{u}_i$ are the eigenvectors of AA^T corresponding to the eigenvalues $\sigma_1^2 \geq \cdots \geq \sigma_r^2$. Complete the $\mathbf{u}_i$ to a basis of $\mathbb{R}^m$ arbitrarily and let U the $m \times m$ unitary (orthogonal) matrix with columns $\mathbf{u}_i$. Then we have $A = U\Sigma V^*$. (I guess we have to pad Σ with some zeros.) This is the singular value decomposition (SVD).

Geometry: A sends the unit ball in $\mathbb{R}^n$ to an ellipsoid in $\mathbb{R}^m$. The singular values of A are the radii of the ellipsoid.

Eckart-Young Theorem. Write $A = \sum_{i=1}^r \sigma_i \mathbf{u}_i \mathbf{v}_i^T$. Then $A_k = \sum_{i=1}^k \sigma_i \mathbf{u}_i \mathbf{v}_i^T$ is the best rank k approximation to A. That is, for any rank k matrix B we have $\|A - A_k\| \leq \|A - B\|$.

Application: Principal Component Analysis. (Total Least Squares.)

Maybe put all of this in a separate chapter.

7.5 Jordan Canonical Form

If $\chi_A(x) = \prod(x - \lambda_i)^{n_i}$ and $E_{\lambda_i} = \mathcal{N}((\lambda_i I - A)^{n_i}) = n_i$ is the generalized eigenspace then $\dim E_{\lambda_i} = n_i$.

The companion matrix.

8 Applications of Spectral Theory

8.1 The Principal Axes Theorem

The earliest example of the Spectral Theorem goes all the way back to the birth of analytic geometry. It was known to Descartes and Fermat in the early 1600s and was applied by Euler in the 1700s to the mechanics of rotating bodies.[142]

You may have seen this theorem in school: Any polynomial equation of the form

$$f(x, y) = a + bx + cy + dx^2 + exy + fy^2 = 0$$

can be brought into standard form by a translation and a rotation. The standard forms are

$$\begin{aligned} \text{parabola} &: y = ax^2 \text{ or } x = ay^2, \\ \text{ellipse} &: x^2/a^2 + y^2/b^2 = 1, \\ \text{hyperbola} &: \pm(x^2/a^2 - y^2/b^2) = 1. \end{aligned}$$

[142] *Highlights in the History of Spectral Theory*, Steen.

The Principal Axes Theorem generalizes this to higher dimensions.

Theorem (Principal Axes Theorem). Consider a general polynomial $f(x_1, \ldots, x_n)$ of degree 2 in n variables. This can be expressed as

$$f(\mathbf{x}) = b + \mathbf{b}^T\mathbf{x} + \mathbf{x}^T B\mathbf{x},$$

for some scalar b, vector $\mathbf{b}$ and **symmetric** matrix B. If B^{-1} exists, then we can find a change of variables $\mathbf{u} = Q\mathbf{x} + \mathbf{t}$, where $Q^TQ = I$ is an orthogonal matrix[143] and $\mathbf{t}$ is a (translation) vector, such that

$$f(u_1, \ldots, u_n) = a + \lambda_1 u_1^2 + \lambda_2 u^2 + \cdots + \lambda_n u_n^2.$$

What if B^{-1} doesn't exist?

Proof. Let $\mathbf{u} = Q\mathbf{x} + \mathbf{t}$ for some invertible matrix Q and vector $\mathbf{t}$. Then we have

$$\begin{aligned}
f(\mathbf{u}) &= f(Q\mathbf{x} + \mathbf{t}) \\
&= b + \mathbf{b}^T(Q\mathbf{x} + \mathbf{t}) + (Q\mathbf{x} + \mathbf{t})^T B(Q\mathbf{x} + \mathbf{t}) \\
&= b + \mathbf{b}^TQ\mathbf{x} + \mathbf{b}^T\mathbf{t} + (\mathbf{x}^TQ^T + \mathbf{t}^T)B(Q\mathbf{x} + \mathbf{t}) \\
&= b + \mathbf{b}^TQ\mathbf{x} + \mathbf{b}^T\mathbf{t} + \mathbf{x}^TQ^TBQ\mathbf{x} + \mathbf{x}^TQ^TB\mathbf{t} + \mathbf{t}^TBQ\mathbf{x} + \mathbf{t}^TB\mathbf{t} \\
&= b + \mathbf{b}^TQ\mathbf{x} + \mathbf{b}^T\mathbf{t} + \mathbf{x}^TQ^TBQ\mathbf{x} + 2\mathbf{t}^TBQ\mathbf{x} + \mathbf{t}^TB\mathbf{t} \qquad (*) \\
&= (b + \mathbf{b}^T\mathbf{t} + \mathbf{t}^TB\mathbf{t}) + (\mathbf{b}^TQ + 2\mathbf{t}BQ)\mathbf{x} + \mathbf{x}^TQ^TBQ\mathbf{x}.
\end{aligned}$$

Step $(*)$ uses the facts that $B^T = B$ and that $\mathbf{t}^TBQ\mathbf{x}$ is a scalar, hence

$$t^TBQ\mathbf{x} = (t^TBQ\mathbf{x})^T = \mathbf{x}^TQ^TB^T\mathbf{t} = \mathbf{x}^TQ^TB\mathbf{x}.$$

If B^{-1} exists, then we can eliminate the linear terms by taking

$$\begin{aligned}
\mathbf{b}^TQ + 2\mathbf{t}BQ &= \mathbf{0}^T \\
2\mathbf{t}BQ &= -\mathbf{b}^TQ \\
\mathbf{t}BQ &= -\frac{1}{2}\mathbf{b}^TQ \\
\mathbf{t} &= -\frac{1}{2}\mathbf{b}^TQQ^{-1}B^{-1} \\
\mathbf{t} &= -\frac{1}{2}\mathbf{b}^TB^{-1}.
\end{aligned}$$

Finally, since B is symmetric, the Spectral Theorem says that we can choose orthogonal $Q^TQ = I$ so that Q^TBQ is diagonal:

Choose $\mathbf{t}$ so that $\mathbf{b}^T + 2\mathbf{t}^TB = \mathbf{0}^T$. Assume B invertible.

[143] If the coefficients of f are complex then $Q^*Q = I$ is unitary, but we are usually interested in the real case.

8.2 Positive Definite Matrices

If $\langle \mathbf{x}, B\mathbf{x} \rangle \geq 0$ (enough to assume $\in \mathbb{R}$) for all $\mathbf{x} \in \mathbb{C}^n$ then $B^* = B$.

Proof: We need to show that $\langle B\mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{x}, B\mathbf{y} \rangle$ for all $\mathbf{x}, \mathbf{y}$. First note that $\langle B\mathbf{x}, \mathbf{x} \rangle = \langle \mathbf{x}, B\mathbf{x} \rangle^* = \langle \mathbf{x}, B\mathbf{x} \rangle$, so $\langle \mathbf{x}, (B - B^*)\mathbf{x} \rangle$ for all $\mathbf{x}$. We need to show that $\langle \mathbf{x}, T\mathbf{x} \rangle = 0$ for all $\mathbf{x}$ implies $T = O$. Indeed, we have

$$0 = \langle \mathbf{x} + \mathbf{y}, T(\mathbf{x} + \mathbf{y}) \rangle = \langle \mathbf{x}, T\mathbf{y} \rangle + \langle \mathbf{y}, T\mathbf{x} \rangle + 0 + 0$$

and

$$0 = \langle \mathbf{x} + i\mathbf{y}, T(\mathbf{x} + i\mathbf{y}) \rangle = i\langle \mathbf{x}, T\mathbf{y} \rangle - i\langle \mathbf{y}, T\mathbf{x} \rangle + 0 + 0.$$

Divide the second equation by i and add them to obtain $2\langle \mathbf{x}, T\mathbf{y} \rangle = 0$ and hence $\langle \mathbf{x}, T\mathbf{y} \rangle = 0$ for all $\mathbf{x}, \mathbf{y}$.

In principle, our proof of the Spectral Theorem gives an algorithm to factor a semi-definite matrix $B = A^T A$, but is probably not the most efficient method since it assumes that we already know the eigenvalues. The *Cholesky factorization* is a method to factor $B = A^T A$ that avoids having to compute eigenvalues.

8.3 Differential Equations

The matrix exponential encodes the solution to linear systems of differential equations. To begin, recall the power series definition of the exponential function:

$$\exp(x) := 1 + x + \frac{1}{2}x^2 + \cdots + \frac{1}{k!}x^k + \cdots .$$

It is a basic theorem of analysis that this series converges uniformly for any complex number $x \in \mathbb{C}$. It was invented by Euler because of the following special properties. For any complex numbers $x, y \in \mathbb{C}$ we have

$$\begin{aligned}
\exp(x)\exp(y) &= \left(\sum_{i \geq 0} \frac{1}{i!} \cdot x^i \right) \left(\sum_{j \geq 0} \frac{1}{j!} \cdot x^j \right) \\
&= \sum_{k \geq 0} \left(\sum_{i+j=k} \frac{1}{i!} \cdot x^i \cdot \frac{1}{j!} \cdot x^j \right) \\
&= \sum_{k \geq 0} \frac{1}{k!} \cdot \left(\sum_{i+j=k} \frac{k!}{i!j!} x^i x^j \right) \\
&= \sum_{k \geq 0} \frac{1}{k!} \cdot (x + y)^k \\
&= \exp(x + y).
\end{aligned}$$

This property suggests that $\exp(x) = e^x$ for some number e, which Euler calculated to be ≈ 2.71828. Furthermore, the power series $\exp(x)$ is equal to its own derivative:

$$\frac{d}{dx}\exp(x) = \frac{d}{dx}\left(1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \cdots \right)$$

$$
\begin{aligned}
&= 0 + 1 + \frac{1}{2!} \cdot 2x + \frac{1}{3!} \cdot 3x^2 + \frac{1}{4!} \cdot 4x^3 + \cdots \\
&= 0 + 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \cdots \\
&= \exp(x).
\end{aligned}
$$

Conversely, let $f : \mathbb{C} \to \mathbb{C}$ be any function satisfying $\frac{d}{dx} f(x) = f(x)$. Suppose that $f(x)$ has a convergent power series expansion near $x = 0$:

$$f(x) = a_0 + a_1 x + a_2 x^2 + \cdots .$$

The equation $\frac{d}{dx} f(x) = f(x)$ tells us that

$$a_0 + a_1 x + a_2 x^2 + \cdots = a_1 + 2a_1 x + 3a_2 x^2 + \cdots .$$

Then comparing coefficients tells us that $a_k = (k+1)a_{k-1}$ for all $k \geq 0$, which has the unique solution $a_k = a_0/k!$. Hence we must have $f(x) = a_0 \exp(x)$.

Now consider a vector of functions $\mathbf{x}(t) = (x_1(t), x_2(t), \ldots, x_n(t))$. We can think of this as a parametrized path in n-dimensional space: $\mathbf{x} : \mathbb{R} \to \mathbb{R}^n$.[144] A linear system of ordinary differential equations has the form

$$
\left\{ \begin{array}{rcccccc}
x_1'(t) & = & a_{11}x_1(t) & + & \cdots & + & a_{1n}x_n(t) \\
 & \vdots & & & & & \\
x_n'(t) & = & a_{n1}x_1(t) & + & \cdots & + & a_{nn}x_n(t)
\end{array} \right\} \quad \rightsquigarrow \quad \mathbf{x}'(t) = A\mathbf{x}(t),
$$

for some $n \times n$ matrix A of constants. We can think of $\mathbf{x}'(t)$ as the velocity vector of the path $\mathbf{x}(t)$, and we can think of A as specifying a vector field on $\mathbb{R}^n$, with value $A\mathbf{x}$ at the point $\mathbf{x}$. A solution to the equation $\mathbf{x}'(t) = A\mathbf{x}(t)$ is any path $\mathbf{x}(t)$ in $\mathbb{R}^n$ that flows along the vector field defined by A. For any initial point $\mathbf{x}(0) \in \mathbb{R}^n$

The companion matrix:
`https://math.stackexchange.com/questions/348498/jordan-basis-of-a-when-a-is-the-companion`

8.4 Graph Theory

Powers of the adjacency matrix. Acyclic directed graphs:

Paper: acyclic digraphs and eigenvalues of (0,1)-matrices

8.5 Markov Chains

Perron-Frobenius, Page Rank

[144] I guess we'll work with real numbers to make visualization easier.

8.6 Singular Value Decomposition

8.7 Total Least Squares

Given a matrix of n data points in any dimensional space:

$$X = \left(\begin{array}{c|c|c} & & \\ \mathbf{x}_1 & \cdots & \mathbf{x}_n \\ & & \end{array} \right)$$

For any vector $\mathbf{a}$ let $P_{\mathbf{a}} = \mathbf{a}\mathbf{a}^T/\|\mathbf{a}\|^2$ be projection onto the line $\mathbf{a}$ and $Q_{\mathbf{a}} = I - P_{\mathbf{a}}$ be projection onto the hyperplane $\mathbf{a}^\perp$. For any i we have

$$\begin{aligned} \mathbf{x}_i &= P_{\mathbf{a}}\mathbf{x}_i + Q_{\mathbf{a}}\mathbf{x}_i \\ \|\mathbf{x}_i\|^2 &= \|P_{\mathbf{a}}\mathbf{x}_i\|^2 + \|Q_{\mathbf{a}}\mathbf{x}_i\|^2 \\ \sum_i \|\mathbf{x}_i\|^2 &= \sum \|P_{\mathbf{a}}\mathbf{x}_i\|^2 + \sum \|Q_{\mathbf{a}}\mathbf{x}_i\|^2. \end{aligned}$$

Goal: Choose $\mathbf{a}$ to minimize $\sum \|Q_{\mathbf{a}}\mathbf{x}_i\|^2$. Since $\sum_i \|\mathbf{x}_i\|^2$ is fixed by the data, this is the same as maximizing $\sum \|P_{\mathbf{a}}\mathbf{x}_i\|^2$. But

$$\|P_{\mathbf{a}}\mathbf{x}_i\|^2 = \frac{1}{\|a\|^2}|\mathbf{a}^T\mathbf{x}_i|^2 = \frac{1}{\|\mathbf{a}\|^2}\mathbf{a}^T\mathbf{x}_i\overline{\mathbf{a}^T\mathbf{x}_i} = \frac{1}{\|\mathbf{a}\|^2}\mathbf{a}^T\mathbf{x}_i\mathbf{x}_i^T\mathbf{a},$$

hence

$$\sum \|P_{\mathbf{a}}\mathbf{x}_i\|^2 = \frac{1}{\|\mathbf{a}\|^2}\mathbf{a}^T XX^T\mathbf{a} = \frac{1}{\|\mathbf{a}\|^2}(X^T\mathbf{a})^T(X^T\mathbf{a}) = \frac{\|X^T\mathbf{a}\|^2}{\|\mathbf{a}\|^2}.$$

This is maximized by letting $\mathbf{a}$ be an eigenvector for the largest (real) eigenvalue of XX^T.

Proof. By S.T., XX^T can be unitarily diagonalized: $XX^T\mathbf{u}_i = \sigma_i^2\mathbf{u}_i$. Let

$$\mathbf{a} = c_1\mathbf{u}_1 + \cdots + c_n\mathbf{u}_n,$$

so that

$$\frac{1}{\|\mathbf{a}\|^2}\mathbf{a}^T XX^T\mathbf{a} = \frac{\sigma_1^2c_1^2 + \cdots + \sigma_n^2c_n^2}{c_1^2 + \cdots + c_n^2}.$$

Maximum when $c_1 = 1$ and $c_2 = \cdots = c_n = 0$. Maximum under constraint $c_1 = 0$ gives $c_2 = 1$ and $c_3 = \cdots = c_n = 0$, etc.