

Every process function lifts to a unitary process

Draft note. Not submitted. Not refereed. Canonical source of the paper; `lift.pdf` is built from this file.

Abstract

A process function is a classical map w from the outputs of several parties to their inputs such that every choice of local deterministic interventions has exactly one fixed point. Baumeler, Costa, Ralph, Wolf and Zych showed that every process function extends to a bijection by adding a source and a sink, and left open whether every such bijection quantizes to a valid unitary quantum process; Daher Ahmed and Kunjwal (2026) state this as a conjecture and prove it for a subclass. We give a short argument that it holds for every process function with finite alphabets: the permutation unitary $U|o,e\rangle = |w(o)+e, o\rangle$ composed with arbitrary local unitaries induces a unitary from source to sink. The key combinatorial fact, that two distinct output strings always disagree at some party whose input w assigns equally, is the pairwise-exclusivity property of Dourdent et al. (2026). The philosophical sections read the result as saying that fixed-point consistency is already a quantization condition. The result is a draft: the exclusivity lemma and the identity $M M^\dagger = 1$ (complex amplitudes, any finite number of parties, any finite alphabets, local ancillas) are machine-checked in Lean 4 with Mathlib (theorems `PFUL.induced_mul_conjTranspose`, `PFUL.induced_conjTranspose_mul` and `PFUL.induced_unitary`, the last stating membership in the unitary group; standard axioms only). The step from "unitary under local unitaries" to "valid process" (Stinespring, Appendix A) is a standard paper argument and is not formalised.

1. Result and short proof

Setting. Parties $k = 1..n$. Party k has a finite input alphabet I_k , identified with the cyclic group $Z_{|I_k|}$, and a finite output alphabet O_k . A process function is $w = (w_k): \prod O_k \rightarrow \prod I_k$ such that for every tuple of local functions $f_k : I_k \rightarrow O_k$ the map $o \rightarrow (f_k(w_k(o)))_k$ has exactly one fixed point. The source has output space $P = \prod I_k$, the sink has input space $F = \prod O_k$, and

$$U |o\rangle_0 |e\rangle_P = |w(o) + e\rangle_I |o\rangle_F \quad (\text{componentwise addition mod } |I_k|).$$

U is a permutation of basis states, hence unitary, for every map w (Baumeler et al. 2019, Theorem 2).

Lemma (exclusivity). If w is a process function and $s \neq s'$ in $\prod O_k$, there is a party k with $s_k \neq s'_k$ and $w_k(s) = w_k(s')$.

Proof. Otherwise, for every k , $w_k(s) = w_k(s')$ implies $s_k = s'_k$. Define $f_k(x) = s_k$ if $x = w_k(s)$, $f_k(x) = s'_k$ if $x = w_k(s')$ (and arbitrary otherwise); the two clauses agree when $w_k(s) = w_k(s')$. Then s and s' are two distinct fixed points of the same intervention, a contradiction. (This is the forward direction of Dourdent et al. 2026, Theorem 7.)

Theorem. Let V_k be any unitary from $I_k(x) X_k$ to $O_k(x) Y_k$, with local ancilla spaces X_k, Y_k ($|I_k||X_k| = |O_k||Y_k|$). Composing U with the V_k (feeding each I_k into V_k and each O_k back into U) gives the operator M from $P(x) X$ to $F(x) Y$ with entries

$$M[(s,y),(e,x)] = \text{prod}_k V_k[(s_k, y_k), (w_k(s) + e_k, x_k)].$$

M is unitary.

Proof. $(M M^\dagger)[(s,y),(s',y')] = \sum_{e,x} \text{prod}_k \sum_{\{e_k, x_k\}} V_k[(s_k, y_k), (w_k(s) + e_k, x_k)] \text{conj} V_k[(s'_k, y'_k), (w_k(s') + e_k, x_k)]$. If $w_k(s) = w_k(s')$, substituting $i = w_k(s) + e_k$ turns the k -th factor into the inner product of rows (s_k, y_k) and (s'_k, y'_k) of the unitary V_k , i.e. a Kronecker delta. If $s = s'$ every factor is of this kind, giving $\delta_{y,y'}$. If $s \neq s'$, the Lemma gives a party with $w_k(s) = w_k(s')$ and $s_k \neq s'_k$, whose factor is 0. So $M M^\dagger = 1$, and M is square, hence unitary.

In the language of Araújo, Feix, Navascués and Brukner (2017, Definition 1), the process $|U\rangle\rangle\langle\langle U|$ is therefore *pure*. Section 2 spells out why this implies validity for arbitrary local CPTP maps and instruments, including ancillas shared or entangled with outside systems. Fixing the source to $|0\rangle$ and tracing the sink recovers the diagonal process matrix of w .

Converse (numerical only). If w is not a process function, the exclusivity property fails (Dourdent et al. 2026, Theorem 7); we have not proved that unitarity then fails in general, but numerically (code/check.py) every non-process function in the 2- and 3-party binary cases fails unitarity for random local unitaries. The "open/closed" contrast is the point: U is a bijection for every w ; closing the loop is unitary exactly when w is consistent.

Checks. code/check.py enumerates all 12 binary 2-party and 744 binary 3-party process functions, verifies $\|MM^\dagger - 1\| < 1e-13$ for Haar-random local unitaries (with and without qubit ancillas) and for ternary 2-party examples, and verifies that non-process functions fail. Lean status is in lean/STATUS.md.

2. From unitarity to validity

Section 1 shows that the lift is *pure* in the sense of Araújo et al. (unitaries in, unitary out). Appendix A derives full process validity from this (any local CPTP maps, local and shared/entangled ancillas, instruments) using Stinespring dilation. Appendix B covers input and output alphabets of different sizes. Appendix C treats stochastic classical processes: mixtures of process functions are purifiable, but a non-deterministic extremal point of the Baumeler–Wolf polytope numerically is not.

3. Backward causation, before the formalism

The linguistic objection to backward causation is that a cause is, by definition, earlier than its effect. Dummett argued in 1954 and again in 1964 that this is a stipulation about the word, not a discovery about the world. Black's bilking argument is the better objection. If an earlier event is supposed to be the effect of a later one, an agent who sees the earlier event can intervene so as to prevent the later one. Either the agent cannot intervene, which wants an explanation, or the later event was not necessary for the earlier one. Lewis's reply splits the modal: one can, relative to the local facts and the laws, and one cannot, relative to the whole past. That dissolves a verbal paradox. It does not say which histories the laws admit.

General relativity makes the question physical. Gödel's 1949 solution has closed timelike curves. Morris, Thorne and Yurtsever showed that a traversable wormhole, if one could be held open, can be converted into a time machine by relative motion of the mouths. The Cauchy problem on the resulting spacetime is the grandfather paradox in differential equations. Friedman, Morris, Novikov, Thorne and collaborators proposed, in 1990, that

only self-consistent solutions occur. Echeverria, Klinkhammer and Thorne then integrated the hard-sphere billiard on a wormhole and found initial data with one consistent continuation, data with two, and data for which none was apparent. Carlini, Frolov, Mensky, Novikov and Solodukhin derived a version of self-consistency from stationarity of the action in a model, and found the same underdetermination. Novikov's principle removes inconsistent solutions. It does not choose among consistent ones. Earman's survey of the classical problem remains the right warning: a consistency condition that is imposed after the dynamics is not a dynamics.

4. Three quantum replies, and why none is the criterion

Deutsch's 1991 model asks for a fixed point of the partial trace around the curve. In finite dimension a fixed point exists, so no particular action is forbidden, and the grandfather paradox becomes a mixed state. The cost is nonlinearity. The model clones unknown states. Aaronson and Watrous showed that the computational power is PSPACE. Bennett, Leung, Smith and Smolin argued that the power is an artefact of demanding a fixed point for a distribution that an ordinary preparation would not produce. Tolkord and Verch showed that the Deutsch condition can be met to arbitrary precision in quantum field theory on a spacetime with no closed timelike curve at all. A Deutsch fixed point is not diagnostic of a loop.

Lloyd, Maccone and collaborators replace the fixed point by post-selection, the P-CTC model. Inconsistent branches are the ones that fail the post-selection. The grandfather paradox is resolved by the experiment not happening. Process functions sit inside this class: the event list appears conditional on the post-selection succeeding. That is a simulation. It is not an account of why the inconsistent branch was never a physical option. Post-selection is a filter on samples, of the same shape as a measure-zero axiom.

The process-matrix formalism of Oreshkov, Costa and Brukner is the third reply, and the one this note uses. A process is a map from local instruments to probabilities, required to give a probability for every choice of instruments, with no global causal order assumed. Classical deterministic processes in that class are process functions. Baumeler and Wolf described the polytope. Three parties suffice for a violation of a causal inequality. The Lugano function, found by Araújo and Feix and studied by Baumeler and Wolf,

$$w(a,b,c) = (\text{not } b \text{ and } c, \text{ not } c \text{ and } a, \text{ not } a \text{ and } b),$$

is the standard witness. Each party's input depends on the other two outputs. The dependence graph is a cycle. Every deterministic intervention has exactly one fixed point. Two parties are not enough: the twelve binary two-party process functions are all causally ordered. Tobar and Costa extended the characterisation to any number of parties.

5. What the fixed-point clause is doing

The clause is an attempt at the criterion the filters do not give. A map from outputs to inputs is admitted only when every local function, freely chosen, composes with it to give exactly one fixed point. No fixed point is the grandfather case: the intervention has nowhere to land. More than one is the bootstrap: the loop underdetermines its own contents. The freedom is the content of the no-new-physics principle in Baumeler, Costa, Ralph, Wolf and Zych. Any operation possible in an ordinary region remains possible in a region that does not itself contain the closed curve. Consistency is not allowed to forbid the operation. It is allowed only to constrain the global solution.

Dourdent, Leitherer, Boghiu, Simonov, Kunjwal and Acín have since said the same thing without quantifying over interventions. A list of events is the event list of a process function exactly when it is complete and pairwise exclusive. If any two of several questions can be answered jointly, so can all of them. That is a multipartite form of Specker's principle. It is a constraint on the process, not an extra axiom that paradoxical histories have measure zero. Cyclic causation is not the enemy. Cyclic signalling is.

6. Reversibility, and the quantization question

Reversibility is not free on the original alphabet. The Lugano function is two-to-one. Of the 256 binary two-party maps, twelve are process functions and none equals its input-output reverse. Baumeler, Costa, Ralph, Wolf and Zych, Theorem 2, restore reversibility by a source, a region with trivial input, and a sink, a region with trivial output. The extended process can be read backwards. The form used here is the permutation unitary

$$U |o, e\rangle = |w(o) + e, o\rangle.$$

The inverse reads the sink as the output string and the source as the difference between the input and $w(o)$. A bijection of a finite set is a unitary. A unitary on a fixed basis is not yet a quantum process. Parties must be able to insert instruments that are not diagonal in that basis. A Hadamard is the test case. If the induced map from source to sink ceases to be trace-preserving, the classical consistency clause has not lifted, and quantum theory needs an extra filter. If it remains trace-preserving for every instrument, the fixed-point clause was already the quantum consistency clause.

Daher Ahmed and Kunjwal conjectured, on 30 September 2026, that every process function purifies to a unitary process. They prove it under mutual exclusivity of control conditions, and under a second sufficient condition, unambiguity, and they note that the bare permutation unitary of the reversible extension may fail to be a valid process. The argument below is that it does not fail. Araújo, Feix, Navascués and Brukner had already written down, in 2017, a purification of the Lugano process by exactly this construction (the standard $|x, y\rangle \rightarrow |x, y + f(x)\rangle$ trick, credited there to Baumeler and Wolf); a routed-circuit reconstruction is due to Vanrietvelde, Ormrod, Kristjánsson and Barrett. Baumeler, Gilani and Rashid (2022) proved that the diagonal (dephased) process matrix of every process function is valid, and remarked, without a separate proof for the coherent operator, that the reversible extensions are therefore unitarily extensible. The contribution claimed here is a short general proof for the coherent permutation unitary, for every process function.

7. Time symmetry and retrocausality

Price has argued that a time-symmetric ontology for quantum theory should be retrocausal, on the assumption that the quantum state is real. Leifer and Pusey replaced that assumption with lambda-mediation: correlations between a preparation and a later measurement are mediated by the ontic state of the system. No retrocausality, operational time symmetry, and lambda-mediation then imply a timelike Bell factorization, which sequential measurements violate. Maudlin's reply is that the operational time-reverse of an experiment is not always itself an experiment of the same kind. The no-go is conditional.

The dilation in this note is adjacent to that argument, not a solution of it. A process function that violates a causal inequality already has each party's input depending on another party's output. The later setting is an argument of the earlier input. Lambda-mediation, in the strict past-screening sense, fails because of the cycle, not because a retrocausal hidden variable was added by hand. Read forwards, the source is an input and the sink is an output.

Read backwards, they exchange roles, and the inverse of the permutation unitary is the reversed process. The objection that the bare Lugano function is not an involution on its own three bits is an objection to refusing the source and the sink. Whether the dependence is the retrocausality Price asked for, or only an operational time-reverse, is a reading of the same operator. The mathematics does not choose the reading. It does remove one cheap reply: one cannot say that the cyclic classical model dies when instruments become unitary.

8. What this does not settle

Embeddability is untouched. A unitary process on a finite alphabet is not a solution of Einstein's equation, and it is not a local Hamiltonian flow on a wormhole. The billiard has the wrong multiplicity. Post-selection realises the event list conditional on the post-selection succeeding. The theorem says the list has a unitary dilation. It does not say the dilation is a spacetime.

Purifiability in the sense of open quantum problem 43, whether every extensively causal process is purifiable, is related but not identical. The theorem puts every process function, including those that violate causal inequalities, inside the unitarily extendible fragment. It does not classify processes that are not deterministic classical functions.

Freedom is not settled. Nothing here decides whether an agent whose output is an argument of an earlier input has a choice. The fixed-point clause says the choice, if made, completes uniquely. It does not say the choice was open. Whether this is an adequate reply to Black depends on whether consistency conditions may rule out interventions, or only rule out histories given the interventions. The note takes the first reading, and the formalism is built for it.

The combinatorial lemma is finite for any fixed alphabet and is the piece an attacker has to break. The linear algebra after it is row orthogonality. See lean/STATUS.md for exactly what is and is not machine-checked.

Appendix A. Unitarity under local unitaries implies validity

Definitions (AFNB 2017, Sec. 2). A process with source P , sink F and parties k is an operator $W \geq 0$ on $P(x) F(x) \prod_k (I_k(x) O_k)$. For local maps $A_k : L(I_k(x) X_k) \rightarrow L(O_k(x) Y_k)$, the induced map is the link product $G_A = W * (A_1(x) \dots (x) A_n)$, a map $L(P(x) X) \rightarrow L(F(x) Y)$ with $X = (x)_k X_k$ and $Y = (x)_k Y_k$. W is *valid* if G_A is CPTP whenever every A_k is CPTP, for all finite-dimensional local ancillas X_k, Y_k . W is *pure* (their Definition 1) if G_A is unitary whenever every A_k is unitary. Their Theorem 2 says W is pure iff $W = |U\rangle\rangle\langle\langle U|$ for a unitary U . For pure Choi vectors the link product composes operators: if $A_k(\rho) = V_k \rho V_k^\dagger$, then $G_A(\rho) = M \rho M^\dagger$, where M is the operator of Section 1.

Claim. $W = |U\rangle\rangle\langle\langle U|$ with U the permutation lift of a process function w is a valid process.

Step 1 (local unitaries). By the Theorem of Section 1, G_A is conjugation by a unitary M whenever every A_k is conjugation by a unitary V_k . This holds for any ancilla dimensions with $|I_k||X_k| = |O_k||Y_k|$. (Machine-checked, both sides, including the dimension count: `PFUL.induced_mul_conjTranspose`, `PFUL.induced_conjTranspose_mul`, `PFUL.induced_unitary`; see lean/STATUS.md.)

Step 2 (local CPTP maps, via Stinespring). Let $A_k : L(I_k X_k) \rightarrow L(O_k Y_k)$ be CPTP. Stinespring dilation gives an isometry $S_k : I_k X_k \rightarrow O_k Y_k E_k$ with $A_k(\rho) = \text{Tr}_{\{E_k\}}[S_k \rho S_k^\dagger]$. Pad the input with E'_k of dimension $|O_k||Y_k||E_k|$ and the output with E''_k of dimension $|I_k||X_k|$. The two sides then have equal dimension, and the isometry $\psi(x) |0\rangle \rightarrow (S_k \psi)(x) |0\rangle$, defined on the $|0\rangle$ -slice, extends to a unitary $V_k : I_k X_k E'_k \rightarrow O_k Y_k E_k E''_k$ by completing orthonormal bases. So $A_k(\rho) = \text{Tr}_{\{E_k E''_k\}}[V_k(\rho V_k^\dagger)]$.

$(x) |0\rangle\langle 0|_{\{E'_k\}} V_k^\dagger]$. The ancillas E' , E , E'' never enter U , and the link product is multilinear and commutes with appending fixed states and with partial traces on systems it does not contract. Hence

$$G_A(\rho) = \text{Tr}_{\{E, E''\}} [M(\rho(x) |0\rangle\langle 0|_{\{E'\}}) M^\dagger],$$

where M is the operator of Step 1 for the unitaries V_k , with enlarged local ancillas $X_k E'_k$ and $Y_k E_k E''_k$. Appending a state, a unitary conjugation and a partial trace are each CPTP, so G_A is CPTP.

Step 3 (shared and entangled ancillas). An ancilla R that is shared between parties, or entangled with any outside system, is covered by complete positivity. The joint state lives on $P \times R$; $G_A(x) \text{id}_R$ is CPTP because G_A is; outputs are normalised states, and every probability that a later measurement on $F \times Y \times R$ assigns is non-negative and sums to 1. Pre-shared entanglement between laboratories is the special case in which part of X_j and part of X_k start in an entangled state supplied through the source. It is also covered, because Step 2 already allows arbitrary input states on $P(x) \times X$.

Step 4 (instruments). Let $\{A_k^{a_k}\}$ be CP maps summing to a CPTP map A_k . The link product of positive operators (Choi operators) is positive, so each $G_a = W * (x)_k A_k^{a_k}$ is CP. Multilinearity gives $\sum_a G_a = G_A$, which is CPTP by Step 2. Then $p(a) = \text{Tr} G_a(\rho) \geq 0$ and $\sum_a p(a) = 1$ for every input ρ . This is the validity condition of Oreshkov, Costa and Brukner, applied to the process seen by the parties once the source is prepared and the sink discarded. Fixing the source in $|0\rangle$ and tracing the sink yields $W_w = \sum_o |o\rangle\langle o|_O (x) |w(o)\rangle\langle w(o)|_I$, the classical process matrix of w .

So purity gives validity with no further assumption. The only input specific to process functions is Step 1.

Appendix B. Unequal input and output alphabets

Nothing in Section 1 needs $|I_k| = |O_k|$.

- $U : O(x) \times P \rightarrow I(x) \times F$, with $P = \prod I_k$ and $F = \prod O_k$, maps $|o, e\rangle$ to $|w(o) + e, o\rangle$. Both sides have dimension $\prod |I_k| |O_k|$, and U is a permutation of basis states, so U is unitary for every w . Only I_k needs a group structure $(Z_{|I_k|})$; O_k is an arbitrary finite set.
- The exclusivity Lemma and its proof never compare I_k with O_k .
- A party with $|I_k| \neq |O_k|$ cannot apply a unitary $I_k \rightarrow O_k$. Its most general deterministic operation is a CPTP map, or a unitary with ancillas, $I_k \times X_k \rightarrow O_k \times Y_k$ with $|I_k| |X_k| = |O_k| |Y_k|$. The Theorem is stated in exactly that generality (and so is the Lean theorem, whose local types I_k, O_k, X_k, Y_k are independent). The Gram computation gives $M M^\dagger = 1$, and M is square because $\dim(P \times X) = \prod |I_k| |X_k| = \prod |O_k| |Y_k| = \dim(F \times Y)$. Appendix A then applies verbatim.

What holds. For any finite input and output alphabets, with any sizes, the permutation lift of a process function is a pure and valid process. **What is not covered.** Infinite or continuous alphabets (Baumeler et al. 2019 also treat continuous variables), and lifts built from a different bijection than U .

Appendix C. Stochastic classical processes

Question. Does every logically consistent classical process admit a unitary extension, in the sense of AFNB purifiability? The process may be stochastic, not just a deterministic process function.

What is known (verified sources).

- Baumeler and Wolf (NJP 18, 013036 (2016), <https://arxiv.org/abs/1507.01714>) show the classical processes form a polytope. With two binary parties it has 12 extremal points, all deterministic. With three binary parties it has 710,760 extremal points, of which only 744 are deterministic, i.e. process functions. The rest are "proper mixtures" of logically inconsistent deterministic processes. One explicit example is $E_{\text{ex1}} = (C + C)/2$, the uniform mixture of the circular identity channel C ($x_1 = a_3, x_2 = a_1, x_3 = a_2$) and its bitwise negation $\bar{C}$.
- Araújo, Feix, Navascués and Brukner (Quantum 1, 10 (2017), <https://arxiv.org/abs/1611.08535>) define purifiability (their Definition 3). Their Theorem 4 is an exact criterion, and Section 5 gives a necessary condition, $\dim V_W \geq \text{rank } W$.
- Baumeler, Gilani and Rashid (Quantum 6, 673 (2022), <https://arxiv.org/abs/2104.06234>) assert unitary extensibility for process functions only, via reversible embeddings, without a separate proof.
- Daher Ahmed and Kunjwal (<https://arxiv.org/abs/2610.00579>) conjecture that process functions are purifiable. Tobar and Costa (CQG 37, 205011 (2020), <https://arxiv.org/abs/2001.02511>) treat reversible deterministic dynamics.
- I found no source that claims to settle purifiability of the non-deterministic extremal points.

C.1 Proposition (proved). Every convex mixture of process functions is purifiable.

Proof. Let $W = \sum_j p_j W_{\{w_j\}}$. Take source $P' = C^{\wedge J} \otimes P$ and sink $F' = C^{\wedge J} \otimes F$, and define $U|o, e, j\rangle = |w_j(o) + e\rangle_I |o, j\rangle_{F'}$. With local unitaries inserted, the induced operator is the block sum $\oplus_j M_j$. Each M_j is unitary by the Theorem of Section 1, so the process is pure. Feed $\sum_j \sqrt{p_j} |j\rangle \otimes |0\rangle$ into P' . This is a fixed pure state, equivalently $|0\rangle$ followed by a unitary that is absorbed into U . Then trace out F' . Because the sink keeps a copy of j , the cross terms vanish and the result is $\sum_j p_j W_{\{w_j\}} = W$. ■

The control register must enter through the source and leave through the sink. If it is not handed to the sink, the coherence between branches survives, and the result is no longer the classical mixture. Hence the whole deterministic-extrema polytope of Baumeler and Wolf is purifiable. This covers every two-party classical process with binary inputs and outputs, since there all extremal points are deterministic.

C.2 Numerical result: E_{ex1} appears not to be purifiable. Scripts are in code/stochastic/.

- E_{ex1} is logically consistent and is a vertex of the three-party binary polytope (active-constraint rank 64 of 64). So it is not a mixture of process functions, and C.1 does not apply.
- Its process matrix has rank 16, and $\dim V_W = 27 \geq 16$. AFNB's necessary condition therefore does not rule it out.
- Applying their full Theorem 4: write the candidate vectors w_i in coordinates on V_W . The quadratic conditions then require a 16-dimensional subspace S of $C^{\wedge 27}$ on which 46 explicit sesquilinear forms vanish.
- For at least one form, the Hermitian part has inertia that allows totally isotropic subspaces of dimension at most $12 < 16$. So no such S exists, and E_{ex1} is not purifiable.
- Control: the same pipeline applied to Lugano, which is purifiable, gives a bound of $29 \geq 8$.
- Status: this is a floating-point computation with clean spectral gaps (about $1e-17$ versus at least $5e-3$). It is not an exact or symbolic certificate.

Answer (current status). No, if the numerics are right. Not every logically consistent classical process admits a unitary extension. Mixtures of process functions do. The non-deterministic extremal point E_{ex1} is numerically

not purifiable.

Open.

- An exact certificate for E_{ex1} , e.g. rational arithmetic on the 46 forms.
- Classifying which of the 709,936 non-deterministic extremal points are purifiable.
- Whether purifiability of a classical process implies that it lies in the deterministic-extrema polytope.

References

1. M. Dummett, Can an effect precede its cause?, *Aristotelian Society Supplementary Volume* 28, 27 (1954); Bringing about the past, *Philosophical Review* 73, 338 (1964).
2. M. Black, Why cannot an effect precede its cause?, *Analysis* 16, 49 (1956).
3. D. Lewis, The paradoxes of time travel, *American Philosophical Quarterly* 13, 145 (1976).
4. K. Gödel, An example of a new type of cosmological solutions of Einstein's field equations of gravitation, *Reviews of Modern Physics* 21, 447 (1949).
5. M. S. Morris, K. S. Thorne and U. Yurtsever, Wormholes, time machines, and the weak energy condition, *Physical Review Letters* 61, 1446 (1988).
6. J. Friedman, M. S. Morris, I. D. Novikov, F. Echeverria, G. Klinkhammer, K. S. Thorne and U. Yurtsever, Cauchy problem in spacetimes with closed timelike curves, *Physical Review D* 42, 1915 (1990).
7. F. Echeverria, G. Klinkhammer and K. S. Thorne, Billiard balls in wormhole spacetimes with closed timelike curves: classical theory, *Physical Review D* 44, 1077 (1991).
8. A. Carlini, V. P. Frolov, M. B. Mensky, I. D. Novikov and H. H. Solodukhin, Time machines: the Principle of Self-Consistency as a consequence of the Principle of Minimal Action, *International Journal of Modern Physics D* 4, 557 (1995).
9. J. Earman, *Bangs, Crunches, Whimpers, and Shrieks: Singularities and Acausalities in Relativistic Spacetimes*, Oxford University Press (1995).
10. D. Deutsch, Quantum mechanics near closed timelike lines, *Physical Review D* 44, 3197 (1991).
11. S. Aaronson and J. Watrous, Closed timelike curves make quantum and classical computing equivalent, *Proceedings of the Royal Society A* 465, 631 (2009).
12. C. H. Bennett, D. Leung, G. Smith and J. A. Smolin, Can closed timelike curves or nonlinear quantum mechanics improve quantum state discrimination or help solve hard problems?, *Physical Review Letters* 103, 170502 (2009).
13. S. Lloyd et al., Closed timelike curves via postselection: theory and experimental test of consistency, *Physical Review Letters* 106, 040403 (2011).
14. O. Oreshkov, F. Costa and Č. Brukner, Quantum correlations with no causal order, *Nature Communications* 3, 1092 (2012).
15. Ä. Baumeler and S. Wolf, The space of logically consistent classical processes without causal order, *New Journal of Physics* 18, 013036 (2016).
16. Ä. Baumeler, F. Costa, T. C. Ralph, S. Wolf and M. Zych, Reversible time travel with freedom of choice, *Classical and Quantum Gravity* 36, 224002 (2019), arXiv:1703.00779.
17. M. Araújo, A. Feix, M. Navascués and Č. Brukner, A purification postulate for quantum mechanics with indefinite causal order, *Quantum* 1, 10 (2017).
18. A. Vanrietvelde, N. Ormrod, H. Kristjánsson and J. Barrett, Consistent circuits for indefinite causal order, arXiv:2206.10042 (2022).

19. H. Price, Does time-symmetry imply retrocausality? How the quantum world says yes, *Studies in History and Philosophy of Modern Physics* 43, 75 (2012).
20. M. S. Leifer and M. F. Pusey, Is a time symmetric interpretation of quantum theory possible without retrocausality?, *Proceedings of the Royal Society A* 473, 20160607 (2017).
21. T. Maudlin, A tale of two tensors, or, how not to argue for retrocausality, preprint (2017).
22. E. Specker, Die Logik nicht gleichzeitig entscheidbarer Aussagen, *Dialectica* 14, 239 (1960).
23. H. Dourdent, A. Leitherer, E.-C. Boghiu, K. Simonov, R. Kunjwal and A. Acín, What makes a causal loop consistent?, arXiv:2609.39735 (2026).
24. N. Daher Ahmed and R. Kunjwal, Characterizing unitaries via quasi-process functions, arXiv:2610.00579 (2026).
25. J. Faye, Backward causation, *Stanford Encyclopedia of Philosophy*, substantive revision 28 October 2025.
26. L. M. Tolksdorf and R. Verch, Quantum physics, fields and closed timelike curves, *Communications in Mathematical Physics* 357, 319 (2018).
27. Ä. Baumeler, A. S. Gilani and J. Rashid, Unlimited non-causal correlations and their relation to non-locality, *Quantum* 6, 673 (2022).
28. G. Tobar and F. Costa, Reversible dynamics with closed time-like curves and freedom of choice, *Classical and Quantum Gravity* 37, 205011 (2020), arXiv:2001.02511.
29. H. Dourdent, K. Simonov, A. Leitherer, E.-C. Boghiu, R. Kunjwal, S. Halder, R. Augusiak and A. Acín, Paradox-free classical non-causality and unambiguous non-locality without entanglement are equivalent, arXiv:2512.23599 (2025).
30. Ä. Baumeler, Causal loops: logically consistent correlations, time travel, and computation, PhD thesis, Università della Svizzera italiana (2017).