

Complex Analysis

Dr. James Carter

carter42@iu.edu

Outline I

1 Introduction

- Course Objectives

2 Domains

3 Plane Curves

- Length
- Jordan Curves and Domains

4 Calculus in the Plane

- Various Derivatives
- Integration
- Green's Theorem and Identities

5 Harmonic Functions

- Properties and Principles
- Mean Value Properties
- Maximum Principles

6 Complex Numbers and Functions

- Complex Numbers

Outline II

- Analytic Functions
- Complex Differentiation
- The Cauchy-Riemann Equations
- The Exponential and Related Functions
- Harmonic Conjugates

7 Integration

- Complex Line Integrals
- Integration Theorems

8 Complex Series

- Singularities
- Laurent Series
- Laurent Expansion

9 Calculus of Residues

- Computation of Residues
- Argument Principle

Outline III

- Argument Principle
- Real Integrals

Introduction

Objectives Overview I

① Complex numbers

Exercises

- Mostly basic; but when it isn't...

Exercises

- Mostly basic; but when it isn't...
- PLTS: proof left to student

- Semi-qualitative approach

- Semi-qualitative approach
- Details later

- Semi-qualitative approach
- Details later
- Often the case that specifics are ignored/irrelevant

- Semi-qualitative approach
- Details later
- Often the case that specifics are ignored/irrelevant
- Polar coordinates frequently ‘used’; not explicitly (in this book anyway)

- Often ‘evaluating’ integrals

- Often ‘evaluating’ integrals
- however, usually considered as complete objects (this represents the area of a circle, so I know the answer)

Domains

Objectives

- Geography of the plane

Objectives

- Geography of the plane
- Real valued functions of two variables

Objectives

- Geography of the plane
- Real valued functions of two variables
 - Derivatives (partial, gradient, directional)

Objectives

- Geography of the plane
- Real valued functions of two variables
 - Derivatives (partial, gradient, directional)
 - Integrals (line, double)

Algebra in The Plane

- ① The real plane is denoted as \mathbb{R}^2

Algebra in The Plane

- ① The real plane is denoted as \mathbb{R}^2
- ② The set of all ordered pairs $\{(x, y) : x, y \in \mathbb{R}\}$

Algebra in The Plane

- ① The real plane is denoted as \mathbb{R}^2
- ② The set of all ordered pairs $\{(x, y) : x, y \in \mathbb{R}\}$
- ③ Addition and subtraction are defined component-wise

Algebra in The Plane

- ① The real plane is denoted as \mathbb{R}^2
- ② The set of all ordered pairs $\{(x, y) : x, y \in \mathbb{R}\}$
- ③ Addition and subtraction are defined component-wise
- ④ Scalar multiplication is identical to vectors

Geometry in The Plane

Definition

If $z = (x, y) \in \mathbb{R}^2$, the **norm (absolute value, modulus)** of z is

$$|z| = \sqrt{x^2 + y^2}$$

Geometry in The Plane

Definition

If $z = (x, y) \in \mathbb{R}^2$, the **norm (absolute value, modulus)** of z is

$$|z| = \sqrt{x^2 + y^2}$$

Definition

Let $z = (x, y)$ and $z_0 = (x_0, y_0)$. Then the **distance** between z, z_0 is

$$|z - z_0| = \sqrt{(x - x_0)^2 + (y - y_0)^2}$$

Domains in the Plane

Definition

Let $r > 0$ and z_0 be a fixed point in \mathbb{R}^2 . The **disc of radius r with center z_0** is

$$D(z_0; r) = \{z \in \mathbb{R}^2 : |z - z_0| < r\}$$

Domains in the Plane

Definition

Let $r > 0$ and z_0 be a fixed point in \mathbb{R}^2 . The **disc of radius r with center z_0** is

$$D(z_0; r) = \{z \in \mathbb{R}^2 : |z - z_0| < r\}$$

Definition

A point z is an **interior point** of a set $\Omega \subset \mathbb{R}^2$ if and only if there is a disc with center z contained entirely within Ω .

Examples

Domain Examples:

- Let $\Omega = \mathbb{R}^2$

Examples

Domain Examples:

- Let $\Omega = \mathbb{R}^2$
- Let $\Omega = \{(x, y) : 0 \leq x \leq 1, 0 < y \leq 1\}$

Closed Discs

Definition

Let $r > 0$ and z_0 be a fixed point in \mathbb{R}^2 . The set

$$\bar{D}(z_0; r) = \{z \in \mathbb{R}^2 : |z - z_0| \leq r\}$$

is the **closed disc** of radius r with center z_0 .

Closed Discs

Definition

Let $r > 0$ and z_0 be a fixed point in \mathbb{R}^2 . The set

$$\bar{D}(z_0; r) = \{z \in \mathbb{R}^2 : |z - z_0| \leq r\}$$

is the **closed disc** of radius r with center z_0 .

- Note that closed discs have points which are interior points and points which are not interior points.

Closed Discs

Definition

Let $r > 0$ and z_0 be a fixed point in \mathbb{R}^2 . The set

$$\bar{D}(z_0; r) = \{z \in \mathbb{R}^2 : |z - z_0| \leq r\}$$

is the **closed disc** of radius r with center z_0 .

- Note that closed discs have points which are interior points and points which are not interior points.
- Furthermore, a disc which is “partially closed” (includes some points on the circle but not all) is NOT a closed disc (or a closed set).

Open and Closed Subsets

Definition

A subset $\Omega \subset \mathbb{R}^2$ is **open** if and only if every point of Ω is an interior point.

Open and Closed Subsets

Definition

A subset $\Omega \subset \mathbb{R}^2$ is **open** if and only if every point of Ω is an interior point.

Definition

A subset $S \subset \mathbb{R}^2$ is **closed** if and only if the complement set $\mathbb{R}^2 \setminus S$ is open.

Connected and Disconnected Subsets

Definition

An open subset $\Omega \subset \mathbb{R}^2$ is **disconnected** if and only if it can be written as $\Omega = \Omega_1 \cup \Omega_2$ where Ω_1, Ω_2 are non-empty and disjoint.

Connected and Disconnected Subsets

Definition

An open subset $\Omega \subset \mathbb{R}^2$ is **disconnected** if and only if it can be written as $\Omega = \Omega_1 \cup \Omega_2$ where Ω_1, Ω_2 are non-empty and disjoint.

Definition

An open set $\Omega \subset \mathbb{R}^2$ is **connected** if and only if it is not disconnected.

Examples

Examples:

- $D(z_0; r)$

Examples

Examples:

- $D(z_0; r)$
- $D(z_0; r) \setminus \{z_0\}$

Examples

Examples:

- $D(z_0; r)$
- $D(z_0; r) \setminus \{z_0\}$
- $A = \{z : a < |z| < b\}$ for $a < b \in \mathbb{R}$

A Flawed Proof the Unit Disc Is Disconnected

Find the flaw in the following attempt to prove that $D(z_0; r)$ is disconnected:

- Let $U = \{z \in D(z_0; r) : x < 0\}$

Notes:

A Flawed Proof the Unit Disc Is Disconnected

Find the flaw in the following attempt to prove that $D(z_0; r)$ is disconnected:

- Let $U = \{z \in D(z_0; r) : x < 0\}$
- Let $V = \{z \in D(z_0; r) : x > 0\}$

Notes:

A Flawed Proof the Unit Disc Is Disconnected

Find the flaw in the following attempt to prove that $D(z_0; r)$ is disconnected:

- Let $U = \{z \in D(z_0; r) : x < 0\}$
- Let $V = \{z \in D(z_0; r) : x > 0\}$

Notes:

- $U = D(z_0; r) \cap \{z : x < 0\}$

A Flawed Proof the Unit Disc Is Disconnected

Find the flaw in the following attempt to prove that $D(z_0; r)$ is disconnected:

- Let $U = \{z \in D(z_0; r) : x < 0\}$
- Let $V = \{z \in D(z_0; r) : x > 0\}$

Notes:

- $U = D(z_0; r) \cap \{z : x < 0\}$
- $V = D(z_0; r) \cap \{z : x > 0\}$

A Flawed Proof the Unit Disc Is Disconnected

Find the flaw in the following attempt to prove that $D(z_0; r)$ is disconnected:

- Let $U = \{z \in D(z_0; r) : x < 0\}$
- Let $V = \{z \in D(z_0; r) : x > 0\}$

Notes:

- $U = D(z_0; r) \cap \{z : x < 0\}$
- $V = D(z_0; r) \cap \{z : x > 0\}$
- Hence both U, V are each the finite intersection of open sets, hence are themselves open.

Domain

Definition

A subset $\Omega \subset \mathbb{R}^2$ is a **domain** if and only if it is open and connected.

Domain

Definition

*A subset $\Omega \subset \mathbb{R}^2$ is a **domain** if and only if it is open and connected.*

Loosely speaking, open sets possibly containing holes but only 1 component.

Domain

Definition

*A subset $\Omega \subset \mathbb{R}^2$ is a **domain** if and only if it is open and connected.*

Loosely speaking, open sets possibly containing holes but only 1 component. Ask: "Would the paint bucket tool in MS Paint work?"

The Boundary and Boundary Points

Definition

A set $S \subset \mathbb{R}^2$ is **bounded** if and only if there exists $r > 0$ such that $S \subset D(0; r)$.

Note that a boundary point of S is also a boundary point of $\mathbb{R}^2 \setminus S$.

The Boundary and Boundary Points

Definition

A set $S \subset \mathbb{R}^2$ is **bounded** if and only if there exists $r > 0$ such that $S \subset D(0; r)$.

Definition

A point $z \in S \subset \mathbb{R}^2$ is a **boundary point** if and only if every disc $D(z; r)$ contains points in S and points not in S .

Note that a boundary point of S is also a boundary point of $\mathbb{R}^2 \setminus S$.

The Boundary and Boundary Points

Definition

A set $S \subset \mathbb{R}^2$ is **bounded** if and only if there exists $r > 0$ such that $S \subset D(0; r)$.

Definition

A point $z \in S \subset \mathbb{R}^2$ is a **boundary point** if and only if every disc $D(z; r)$ contains points in S and points not in S .

Note that a boundary point of S is also a boundary point of $\mathbb{R}^2 \setminus S$.

Definition

The **boundary** of a set $S \subset \mathbb{R}^2$ is the set of all boundary points of S ; denoted by ∂S .

Examples

Examples:

- $D(0; r); \partial D(0; r) = C(0; r) = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}$

Examples

Examples:

- $D(0; r); \partial D(0; r) = C(0; r) = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}$
- $H = \{(x, y) \in \mathbb{R}^2 : y > 0\}; \partial H = \{(x, y) \in \mathbb{R}^2 : y = 0\}$

Examples

Examples:

- $D(0; r); \partial D(0; r) = C(0; r) = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}$
- $H = \{(x, y) \in \mathbb{R}^2 : y > 0\}; \partial H = \{(x, y) \in \mathbb{R}^2 : y = 0\}$
- $\mathbb{R}^2; \partial \mathbb{R}^2 = \emptyset$

Plane Curves

Conventions and Definitions I

- $0 = (0, 0)$

Conventions and Definitions II

Circle

Uniqueness of Parameterizations

- Warning: parameterizations are not unique.

Uniqueness of Parameterizations

- Warning: parameterizations are not unique.
- Consider $\alpha : [0, 2\pi] \rightarrow C(0; r)$ with $\alpha(t) = (r \cos(t), r \sin(t))$ and $\beta : [0, 2\pi] \rightarrow C(0; r)$ with $\beta(t) = (r \cos(3t), r \sin(3t))$.

Uniqueness of Parameterizations

- Warning: parameterizations are not unique.
- Consider $\alpha : [0, 2\pi] \rightarrow C(0; r)$ with $\alpha(t) = (r \cos(t), r \sin(t))$ and $\beta : [0, 2\pi] \rightarrow C(0; r)$ with $\beta(t) = (r \cos(3t), r \sin(3t))$.
- Also $\gamma : [0, 1] \rightarrow C(0; r)$ with $\gamma(t) = (r \cos(2\pi t), r \sin(2\pi t))$.

Parameterization of Curves

Definition

Let $\Gamma \subset \mathbb{R}^2$ and $[a, b] \subset \mathbb{R}$ with $a < b$. Any function $\alpha = (\alpha_1, \alpha_2) : [a, b] \rightarrow \Gamma$ is a **parameterization** of Γ if and only if

Parameterization of Curves

Definition

Let $\Gamma \subset \mathbb{R}^2$ and $[a, b] \subset \mathbb{R}$ with $a < b$. Any function $\alpha = (\alpha_1, \alpha_2) : [a, b] \rightarrow \Gamma$ is a **parameterization** of Γ if and only if

- ① α is continuous; that is α_i are all continuous

Parameterization of Curves

Definition

Let $\Gamma \subset \mathbb{R}^2$ and $[a, b] \subset \mathbb{R}$ with $a < b$. Any function

$\alpha = (\alpha_1, \alpha_2) : [a, b] \rightarrow \Gamma$ is a **parameterization** of Γ if and only if

- ① α is continuous; that is α_i are all continuous
- ② α is surjective; that is for every $z \in \Gamma$ there exists $t \in [a, b]$ such that $\alpha(t) = z$

Parameterization of Curves

Definition

Let $\Gamma \subset \mathbb{R}^2$ and $[a, b] \subset \mathbb{R}$ with $a < b$. Any function

$\alpha = (\alpha_1, \alpha_2) : [a, b] \rightarrow \Gamma$ is a **parameterization** of Γ if and only if

- ① α is continuous; that is α_i are all continuous
- ② α is surjective; that is for every $z \in \Gamma$ there exists $t \in [a, b]$ such that $\alpha(t) = z$

Definition

α is called **smooth** or **continuously differentiable** if in addition

Parameterization of Curves

Definition

Let $\Gamma \subset \mathbb{R}^2$ and $[a, b] \subset \mathbb{R}$ with $a < b$. Any function

$\alpha = (\alpha_1, \alpha_2) : [a, b] \rightarrow \Gamma$ is a **parameterization** of Γ if and only if

- ① α is continuous; that is α_i are all continuous
- ② α is surjective; that is for every $z \in \Gamma$ there exists $t \in [a, b]$ such that $\alpha(t) = z$

Definition

α is called **smooth** or **continuously differentiable** if in addition

- ① α'_i all exist and are continuous for every $t \in [a, b]$

Parameterization of Curves

Definition

Let $\Gamma \subset \mathbb{R}^2$ and $[a, b] \subset \mathbb{R}$ with $a < b$. Any function

$\alpha = (\alpha_1, \alpha_2) : [a, b] \rightarrow \Gamma$ is a **parameterization** of Γ if and only if

- ① α is continuous; that is α_i are all continuous
- ② α is surjective; that is for every $z \in \Gamma$ there exists $t \in [a, b]$ such that $\alpha(t) = z$

Definition

α is called **smooth** or **continuously differentiable** if in addition

- ① α'_i all exist and are continuous for every $t \in [a, b]$
- ② for each $t \in [a, b]$, the velocity vector $\alpha'(t) = \langle \alpha'_1(t), \alpha'_2(t) \rangle \neq \langle 0, 0 \rangle$

Parameterization of Curves

Definition

Let $\Gamma \subset \mathbb{R}^2$ and $[a, b] \subset \mathbb{R}$ with $a < b$. Any function

$\alpha = (\alpha_1, \alpha_2) : [a, b] \rightarrow \Gamma$ is a **parameterization** of Γ if and only if

- ① α is continuous; that is α_i are all continuous
- ② α is surjective; that is for every $z \in \Gamma$ there exists $t \in [a, b]$ such that $\alpha(t) = z$

Definition

α is called **smooth** or **continuously differentiable** if in addition

- ① α'_i all exist and are continuous for every $t \in [a, b]$
- ② for each $t \in [a, b]$, the velocity vector $\alpha'(t) = \langle \alpha'_1(t), \alpha'_2(t) \rangle \neq \langle 0, 0 \rangle$
- ③ if $\alpha(a) = \alpha(b)$, then $\alpha'(a) = \alpha'(b)$

Definitions

Definition

A parameterization α is **simple** if and only if $\alpha|_{(a,b)}$ is injective.

Definitions

Definition

A parameterization α is **simple** if and only if $\alpha|_{(a,b)}$ is injective.

Definition

A parameterization α is **closed** if $\alpha(a) = \alpha(b)$.

Examples

- The line from z_0 to z_1 is parameterized by $\gamma(t) = (1 - t)z_0 + tz_1$ where $t \in [0, 1]$

Examples

- The line from z_0 to z_1 is parameterized by $\gamma(t) = (1 - t)z_0 + tz_1$ where $t \in [0, 1]$
- The circle $C(0; 1)$ is often parameterized by $\alpha(t) = (r \cos(t), r \sin(t))$ where $t \in [0, 2\pi]$ or $\beta(t) = (r \cos(2\pi t), r \sin(2\pi t))$ for $t \in [0, 1]$

Examples

- The line from z_0 to z_1 is parameterized by $\gamma(t) = (1 - t)z_0 + tz_1$ where $t \in [0, 1]$
- The circle $C(0; 1)$ is often parameterized by $\alpha(t) = (r \cos(t), r \sin(t))$ where $t \in [0, 2\pi]$ or $\beta(t) = (r \cos(2\pi t), r \sin(2\pi t))$ for $t \in [0, 1]$
- The parameterization $\eta(t) = (t, t^3)$ for $t \in [-1, 1]$ is a simple, not closed parameterization between $(-1, -1)$ and $(1, 1)$

Definition

A parameterization $\alpha : [a, b] \rightarrow \Gamma$ is a **piecewise-smooth parameterization** if and only if there exists a finite set of values $a = a_0 < a_1 < a_2 < \cdots < a_{n-1} < a_n = b$ such that the function α restricted to the intervals $[a_{i-1}, a_i]$ gives a smooth parameterization of $\Gamma_i = \{\alpha(t) : t \in [a_{i-1}, a_i]\}$.

Piecewise-smooth Parameterizations II

Examples:

Smooth Curves

- Exercise in symbols

Smooth Curves

- Exercise in symbols
- If α is a smooth parameterization of Γ , then the pair (Γ, α) is called a **piecewise-smooth curve**.

Smooth Curves

- Exercise in symbols
- If α is a smooth parameterization of Γ , then the pair (Γ, α) is called a **piecewise-smooth curve**.
- It is customary to omit mention of α and speak of “the (smooth) curve Γ ”

Smooth Curves

- Exercise in symbols
- If α is a smooth parameterization of Γ , then the pair (Γ, α) is called a **piecewise-smooth curve**.
- It is customary to omit mention of α and speak of “the (smooth) curve Γ ”
- The situation is delicate however; see the Peano curve

Motivation

- Consider the line from z_0 to z_1 parameterized by
 $\alpha(t) = tz_1 + (1 - t)z_0 = z_0 + t(z_1 - z_0)$.

Motivation

- Consider the line from z_0 to z_1 parameterized by
 $\alpha(t) = tz_1 + (1 - t)z_0 = z_0 + t(z_1 - z_0)$.
- Then $\alpha'(t) = z_1 - z_0$ (independent of t)

Motivation

- Consider the line from z_0 to z_1 parameterized by
 $\alpha(t) = tz_1 + (1 - t)z_0 = z_0 + t(z_1 - z_0)$.
- Then $\alpha'(t) = z_1 - z_0$ (independent of t)
- So

$$\int_0^1 \alpha'(t) dt = (z_1 - z_0)(1 - 0)$$

Definition

Definition

If Γ is a smooth curve given by $\alpha : [a, b] \rightarrow \Gamma$, then the **length** of α is

$$\text{length } (\alpha) = \int_a^b |\alpha'(t)| dt$$

Definition

Definition

If Γ is a smooth curve given by $\alpha : [a, b] \rightarrow \Gamma$, then the **length** of α is

$$\text{length } (\alpha) = \int_a^b |\alpha'(t)| dt$$

Since α is smooth, the integrand is piecewise continuous and bounded, hence the integral exists.

Examples

- $C(0; r)$ with $\alpha(t) = (r \cos(t), r \sin(t))$ for $t \in [0, 2\pi]$

Examples

- $C(0; r)$ with $\alpha(t) = (r \cos(t), r \sin(t))$ for $t \in [0, 2\pi]$
- $C(0; r)$ with $\beta(t) = (r \cos(t), r \sin(t))$ for $t \in [0, 6\pi]$

Parameter Selection for Parameterization

- Using an arbitrary parameter for curves is nonsensical

Parameter Selection for Parameterization

- Using an arbitrary parameter for curves is nonsensical
- Prefer to use an intrinsic property of the curve itself

Parameter Selection for Parameterization

- Using an arbitrary parameter for curves is nonsensical
- Prefer to use an intrinsic property of the curve itself
- The desired property is arc-length

Parameter Selection for Parameterization

- Using an arbitrary parameter for curves is nonsensical
- Prefer to use an intrinsic property of the curve itself
- The desired property is arc-length
- Goal: adapt length integral to be new parameter

Adaptation of Length

- Denote the new parameterization of a given curve (Γ, α) by $\sigma(s)$

Adaptation of Length

- Denote the new parameterization of a given curve (Γ, α) by $\sigma(s)$
- $\sigma(s)$ will have the property that $\sigma(s_1) - \sigma(s_0) = s_1 - s_0$

Adaptation of Length

- Denote the new parameterization of a given curve (Γ, α) by $\sigma(s)$
- $\sigma(s)$ will have the property that $\sigma(s_1) - \sigma(s_0) = s_1 - s_0$
- $\sigma(s)$ will be “equivalent” to α

Equivalence of Intervals

Definition

Let $[a_1, b_1]$ and $[a_2, b_2]$ be intervals on the s, t axes respectively. A function $E : [a_1, b_1] \rightarrow [a_2, b_2]$ is an **equivalence** if and only if it is:

Equivalence of Intervals

Definition

Let $[a_1, b_1]$ and $[a_2, b_2]$ be intervals on the s, t axes respectively. A function $E : [a_1, b_1] \rightarrow [a_2, b_2]$ is an **equivalence** if and only if it is:

- ① bijective

Equivalence of Intervals

Definition

Let $[a_1, b_1]$ and $[a_2, b_2]$ be intervals on the s, t axes respectively. A function $E : [a_1, b_1] \rightarrow [a_2, b_2]$ is an **equivalence** if and only if it is:

- ① bijective
- ② continuous

Equivalence of Intervals

Definition

Let $[a_1, b_1]$ and $[a_2, b_2]$ be intervals on the s, t axes respectively. A function $E : [a_1, b_1] \rightarrow [a_2, b_2]$ is an **equivalence** if and only if it is:

- ① bijective
- ② continuous
- ③ increasing

Equivalence of Intervals

Definition

Let $[a_1, b_1]$ and $[a_2, b_2]$ be intervals on the s, t axes respectively. A function $E : [a_1, b_1] \rightarrow [a_2, b_2]$ is an **equivalence** if and only if it is:

- ① *bijection*
- ② *continuous*
- ③ *increasing*
- ④ *piecewise differentiable with everywhere positive first derivative*

Equivalence of Intervals

Definition

Let $[a_1, b_1]$ and $[a_2, b_2]$ be intervals on the s, t axes respectively. A function $E : [a_1, b_1] \rightarrow [a_2, b_2]$ is an **equivalence** if and only if it is:

- ① bijective
- ② continuous
- ③ increasing
- ④ piecewise differentiable with everywhere positive first derivative

This is a formalization of the idea that $E(a_1) = a_2$, $E(b_1) = b_2$, and as s increases from a_1 to b_1 , it is also the case that $t = E(s)$ increases from a_2 to b_2 .

Theorem

Theorem

Let $\alpha : [a, b] \rightarrow \Gamma$ be a piecewise-smooth parameterization with length $L > 0$. Then there exists a unique piecewise-smooth parameterization $\sigma : [0, L] \rightarrow \Gamma$ equivalent to α which satisfies:

Theorem

Theorem

Let $\alpha : [a, b] \rightarrow \Gamma$ be a piecewise-smooth parameterization with length $L > 0$. Then there exists a unique piecewise-smooth parameterization $\sigma : [0, L] \rightarrow \Gamma$ equivalent to α which satisfies:

- ① the tangent vector $\sigma'(s)$ has unit length; $|\sigma'(s)| = 1$

Theorem

Theorem

Let $\alpha : [a, b] \rightarrow \Gamma$ be a piecewise-smooth parameterization with length $L > 0$. Then there exists a unique piecewise-smooth parameterization $\sigma : [0, L] \rightarrow \Gamma$ equivalent to α which satisfies:

- ① the tangent vector $\sigma'(s)$ has unit length; $|\sigma'(s)| = 1$
- ② the distance traveled along Γ from $\sigma(0)$ to $\sigma(s)$ is s

Proof

Proof.



Proof

Proof.

- Note that if σ satisfies property (1), then the distance from $\sigma(0)$ to $\sigma(s)$ is $\int_0^s |\sigma'(t)|dt = s$. It is sufficient to prove the result for smooth parameterizations; hence assume α is smooth. Define $F(\tau) = \int_a^\tau |\alpha'(t)|dt$ for $\tau \in [a, b]$. By Calculus, $F'(\tau) = |\alpha'(\tau)| > 0$ and continuous.



Proof

Proof.

- Note that if σ satisfies property (1), then the distance from $\sigma(0)$ to $\sigma(s)$ is $\int_0^s |\sigma'(t)|dt = s$. It is sufficient to prove the result for smooth parameterizations; hence assume α is smooth. Define $F(\tau) = \int_a^\tau |\alpha'(t)|dt$ for $\tau \in [a, b]$. By Calculus, $F'(\tau) = |\alpha'(\tau)| > 0$ and continuous.
- Define E as the inverse of F . By the Chain Rule, $E'(s) > 0$ and E is an equivalence from $[0, L]$ to $[a, b]$.



Proof

Proof.

- Note that if σ satisfies property (1), then the distance from $\sigma(0)$ to $\sigma(s)$ is $\int_0^s |\sigma'(t)|dt = s$. It is sufficient to prove the result for smooth parameterizations; hence assume α is smooth. Define $F(\tau) = \int_a^\tau |\alpha'(t)|dt$ for $\tau \in [a, b]$. By Calculus, $F'(\tau) = |\alpha'(\tau)| > 0$ and continuous.
- Define E as the inverse of F . By the Chain Rule, $E'(s) > 0$ and E is an equivalence from $[0, L]$ to $[a, b]$.
- Define $\sigma(s) = \alpha(E(s))$ and verify $|\sigma'(s)| = 1$ by the Chain Rule.



Definitions

Definition

A **Jordan curve** is a simple, closed, piecewise smooth curve.

Definitions

Definition

A **Jordan curve** is a simple, closed, piecewise smooth curve.

Definition

A **Jordan domain** is a bounded domain Ω whose boundary $\partial\Omega$ is a Jordan curve.

Jordan Curve Theorem

Theorem (Jordan Curve Theorem)

If Γ is a Jordan curve, then its complement $\mathbb{R}^2 \setminus \Gamma$ consists of two disjoint domains, one bounded (the “inside”) and one unbounded (the “outside”); both have Γ as boundary. If a point inside Γ is joined by a path to a point outside Γ , then the path must cross Γ .

Jordan Curve Theorem

Theorem (Jordan Curve Theorem)

If Γ is a Jordan curve, then its complement $\mathbb{R}^2 \setminus \Gamma$ consists of two disjoint domains, one bounded (the “inside”) and one unbounded (the “outside”); both have Γ as boundary. If a point inside Γ is joined by a path to a point outside Γ , then the path must cross Γ .

Proof.

Omitted. And difficult for generic Jordan curves.



Jordan Curve Theorem

Theorem (Jordan Curve Theorem)

If Γ is a Jordan curve, then its complement $\mathbb{R}^2 \setminus \Gamma$ consists of two disjoint domains, one bounded (the “inside”) and one unbounded (the “outside”); both have Γ as boundary. If a point inside Γ is joined by a path to a point outside Γ , then the path must cross Γ .

Proof.

Omitted. And difficult for generic Jordan curves. □

A Jordan domain is *positively oriented* if the given parameterization has the interior to the left.

Holes in Domains

Jordan domains can have holes

Holes in Domains

Jordan domains can have holes

Holes in Domains

Jordan domains can have holes

Definition

A Jordan domain Ω is **k-connected** if its boundary consists of k Jordan curves.

Holes in Domains

Jordan domains can have holes

Definition

A Jordan domain Ω is **k-connected** if its boundary consists of k Jordan curves.

Note that the complement of a k -connected Jordan domain is therefore k disjoint connected components.

Normal Vectors I

Definition

Let Ω be a Jordan domain with boundary $\partial\Omega$ and Γ one of the Jordan curves comprising $\partial\Omega$ with parameterization $\sigma : [0, L] \rightarrow \Gamma$

is called the **outward normal vector**.

Normal Vectors II

Example:

Calculus in the Plane

Overview

- The study of plane curves is the study of functions $f : \mathbb{R} \rightarrow \mathbb{R}^2$

Overview

- The study of plane curves is the study of functions $f : \mathbb{R} \rightarrow \mathbb{R}^2$
- The study of plane calculus is the study of functions $g : \mathbb{R}^2 \rightarrow \mathbb{R}$

Overview

- The study of plane curves is the study of functions $f : \mathbb{R} \rightarrow \mathbb{R}^2$
- The study of plane calculus is the study of functions $g : \mathbb{R}^2 \rightarrow \mathbb{R}$
- In particular, how these functions change in the vicinity of a point $z \in \mathbb{R}^2$ - differentiation

Overview

- The study of plane curves is the study of functions $f : \mathbb{R} \rightarrow \mathbb{R}^2$
- The study of plane calculus is the study of functions $g : \mathbb{R}^2 \rightarrow \mathbb{R}$
- In particular, how these functions change in the vicinity of a point $z \in \mathbb{R}^2$ - differentiation
- The Chain Rule will be used offensively often

Objective

- The goal is to determine the values of $q(\zeta)$ for $\zeta \in \Omega$ given:

Objective

- The goal is to determine the values of $q(\zeta)$ for $\zeta \in \Omega$ given:
- $q(\zeta)$ exists and is twice differentiable on $\bar{\Omega}$

Objective

- The goal is to determine the values of $q(\zeta)$ for $\zeta \in \Omega$ given:
- $q(\zeta)$ exists and is twice differentiable on $\bar{\Omega}$
- $\Delta q(z)$ is known for all $z \in \Omega$

Objective

- The goal is to determine the values of $q(\zeta)$ for $\zeta \in \Omega$ given:
- $q(\zeta)$ exists and is twice differentiable on $\bar{\Omega}$
- $\Delta q(z)$ is known for all $z \in \Omega$
- $q(z)$ is known for all $z \in \Omega$

Objective

- The goal is to determine the values of $q(\zeta)$ for $\zeta \in \Omega$ given:
- $q(\zeta)$ exists and is twice differentiable on $\bar{\Omega}$
- $\Delta q(z)$ is known for all $z \in \Omega$
- $q(z)$ is known for all $z \in \Omega$
- $\nabla q(z)$ is known for all $z \in \partial\Omega$

Objective

- The goal is to determine the values of $q(\zeta)$ for $\zeta \in \Omega$ given:
- $q(\zeta)$ exists and is twice differentiable on $\bar{\Omega}$
- $\Delta q(z)$ is known for all $z \in \Omega$
- $q(z)$ is known for all $z \in \Omega$
- $\nabla q(z)$ is known for all $z \in \partial\Omega$

This will be “accomplished” using Green III

Notation

- Ω - domain (open, connected)

Notation

- Ω - domain (open, connected)
- function u assigns value $u(z)$ to the point $z \in \Omega$

Notation

- Ω - domain (open, connected)
- function u assigns value $u(z)$ to the point $z \in \Omega$
- Since $z = (x, y)$, $u(z) = u(x, y)$

Notation

- Ω - domain (open, connected)
- function u assigns value $u(z)$ to the point $z \in \Omega$
- Since $z = (x, y)$, $u(z) = u(x, y)$
- $u : \Omega \rightarrow \mathbb{R}$

Notation

- Ω - domain (open, connected)
- function u assigns value $u(z)$ to the point $z \in \Omega$
- Since $z = (x, y)$, $u(z) = u(x, y)$
- $u : \Omega \rightarrow \mathbb{R}$
- For $S \subset \mathbb{R}$, $u^{-1}(S) = \{z \in \mathbb{R}^2 : u(z) \in S\}$

Continuity

Definition

Let $U \subset \mathbb{R}^2$. A function $u : \Omega \rightarrow \mathbb{R}$ is **continuous** if and only if for every open interval $S \subset \mathbb{R}$, the set $u^{-1}(S)$ is open in \mathbb{R}^2 .

Differentiability

- Assume familiarity with partial derivatives for multivariable functions:

Differentiability

- Assume familiarity with partial derivatives for multivariable functions:
- If $u(z) = u(x, y)$, then $u_x(x, y)$ and $u_y(x, y)$

Differentiability

- Assume familiarity with partial derivatives for multivariable functions:
- If $u(z) = u(x, y)$, then $u_x(x, y)$ and $u_y(x, y)$
- Notation: $\mathfrak{C}^k(\Omega)$

Differentiability

- Assume familiarity with partial derivatives for multivariable functions:
- If $u(z) = u(x, y)$, then $u_x(x, y)$ and $u_y(x, y)$
- Notation: $\mathfrak{C}^k(\Omega)$
- In particular, $u \in \mathfrak{C}^1(\Omega)$ implies $u_x, u_y \in \mathfrak{C}(\Omega)$

Differentiability

- Assume familiarity with partial derivatives for multivariable functions:
- If $u(z) = u(x, y)$, then $u_x(x, y)$ and $u_y(x, y)$
- Notation: $\mathfrak{C}^k(\Omega)$
- In particular, $u \in \mathfrak{C}^1(\Omega)$ implies $u_x, u_y \in \mathfrak{C}(\Omega)$
- That is, $\mathfrak{C}^1(\Omega)$ is the set of all continuously differentiable functions on Ω

Theorem I

Theorem (Linear Approximation)

Let $u \in \mathfrak{C}^1(\Omega)$ and $z_0 = (x_0, y_0) \in \Omega$. For each point $z = (x, y) \in \Omega$, we may write

$$u(z) = u(z_0) + u_x(z_0)(x - x_0) + u_y(z_0)(y - y_0) + \epsilon_1(x - x_0) + \epsilon_2(y - y_0)$$

where $\epsilon_i \rightarrow 0$ as $z \rightarrow z_0 \in \Omega$.

Notation:

$$u(z) = L(z; z_0) + E(z; z_0)$$

Note: $E(z; z_0)$ tends to zero rapidly; specifically

$$\lim_{z \rightarrow z_0} \frac{|E(z; z_0)|}{|z - z_0|} = 0$$

Theorem II

Theorem

Let $u \in \mathfrak{C}^1(\Omega)$, $z_0 \in \Omega$. Then u is continuous at z_0 . That is

$$\mathfrak{C}^1(\Omega) \subset \mathfrak{C}(\Omega)$$

Theorem (Chain Rule)

Let $u \in \mathfrak{C}^1(\Omega)$ and $\alpha(t) = (\alpha_1(t), \alpha_2(t))$ parameterize $\Gamma \subset \Omega$. Then

$$\frac{d}{dt} u(\alpha(t)) = u_x(\alpha(t))\alpha'_1(t) + u_y(\alpha(t))\alpha'_2(t)$$

Directional Derivatives I

Definition

Let $z = (x, y)$ be a point in the domain Ω , V a unit vector with tail at z , and u a function defined on Ω . Then the **derivative of u at z in the direction V** is denoted $u'(z; V)$ and given by

$$u'(z; V) = \frac{d}{ds} u(z + sV) \Big|_{s=0}$$

Alternatively

$$u'(z; V) = \lim_{s \rightarrow 0} \frac{u(z + sV) - u(z)}{s} \quad (1)$$

Directional Derivatives II

Theorem

Let $u \in \mathfrak{C}^1(\Omega)$ and $z \in \Omega$. If $V = \langle k_1, k_2 \rangle$ is a unit vector, then

$$u'(z; V) = u_x(z)k_1 + u_y(z)k_2$$

Gradient Vector

Definition

Let Ω be a domain and $u \in \mathfrak{C}^1(\Omega)$. Then the **gradient vector** of u at the point z is

$$\nabla u(z) = \langle u_x(z), u_y(z) \rangle$$

Fact:

$$u'(z_0; V) = \nabla u(z_0) \cdot V$$

Notation: $\nabla = \left\langle \frac{\partial}{\partial x}, \frac{\partial}{\partial y} \right\rangle$

Outward Normal Derivative

Motivation: For a Jordan domain Ω and function $u \in \mathfrak{C}^1(\Omega^+)$, what is the rate of change of z as z crosses $\partial\Omega$ in the normal direction $N(z_0)$?

Outward Normal Derivative

Motivation: For a Jordan domain Ω and function $u \in \mathfrak{C}^1(\Omega^+)$, what is the rate of change of z as z crosses $\partial\Omega$ in the normal direction $N(z_0)$?

Answer: $u'(z_0; N(z_0))$, denoted as $\frac{\partial u}{\partial n}(z_0)$.

Outward Normal Derivative

Motivation: For a Jordan domain Ω and function $u \in \mathcal{C}^1(\Omega^+)$, what is the rate of change of z as z crosses $\partial\Omega$ in the normal direction $N(z_0)$?

Therefore

$$\frac{\partial u}{\partial n}(z_0) = u'(z_0; N(z_0)) = \nabla u(z_0) \cdot N(z_0)$$

Outward Normal Derivative

Motivation: For a Jordan domain Ω and function $u \in \mathcal{C}^1(\Omega^+)$, what is the rate of change of z as z crosses $\partial\Omega$ in the normal direction $N(z_0)$?

Therefore

$$\frac{\partial u}{\partial n}(z_0) = u'(z_0; N(z_0)) = \nabla u(z_0) \cdot N(z_0)$$

Treat $(\partial u / \partial n)(z)$ as a function of z defined on the curves that make up $\partial\Omega$.

The Laplacian I

- If u_{xy} and u_{yx} exist and are continuous, then they are equal.

Note:

$$\mathfrak{C}(\Omega) \supset \mathfrak{C}^1(\Omega) \supset \mathfrak{C}^2(\Omega) \cdots \supset \mathfrak{C}^\infty(\Omega)$$

where $\mathfrak{C}^\infty(\Omega)$ is the set of functions with continuous partial derivatives of all order.

The Laplacian II

Definition

Let $u \in \mathcal{C}^2(\Omega)$. Then the **Laplacian** of u is the second derivative, that is $u_{xx} + u_{yy}$ and denoted

$$\Delta u(z) = \Delta u(x, y) = u_{xx}(x, y) + u_{yy}(x, y)$$

Definition

A function $u \in \mathcal{C}^2(\Omega)$ is **harmonic** if and only if $\Delta u = 0$.

Line Integrals

- $p = p(x, y), q = q(x, y)$ continuous, real valued

Line Integrals

- $p = p(x, y), q = q(x, y)$ continuous, real valued
- $\alpha : [a, b] \rightarrow \Gamma$ a parameterization; $\alpha(t) = (\alpha_1(t), \alpha_2(t))$

Line Integrals

- $p = p(x, y), q = q(x, y)$ continuous, real valued
- $\alpha : [a, b] \rightarrow \Gamma$ a parameterization; $\alpha(t) = (\alpha_1(t), \alpha_2(t))$

$$\int_{\Gamma} [p(x, y) dx + q(x, y) dy]$$

Line Integrals

- $p = p(x, y), q = q(x, y)$ continuous, real valued
- $\alpha : [a, b] \rightarrow \Gamma$ a parameterization; $\alpha(t) = (\alpha_1(t), \alpha_2(t))$

$$\int_{\Gamma} [p(x, y) dx + q(x, y) dy]$$

$$I_{\alpha} = \int_{t=a}^{t=b} [p(\alpha(t))\alpha'_1(t)dt + q(\alpha(t))\alpha'_2(t) dt]$$

Line Integrals

- $p = p(x, y), q = q(x, y)$ continuous, real valued
- $\alpha : [a, b] \rightarrow \Gamma$ a parameterization; $\alpha(t) = (\alpha_1(t), \alpha_2(t))$

$$\int_{\Gamma} [p(x, y) dx + q(x, y) dy]$$

$$I_{\alpha} = \int_{t=a}^{t=b} [p(\alpha(t))\alpha'_1(t)dt + q(\alpha(t))\alpha'_2(t) dt]$$

- by calculus and $(x, y) \in \Gamma \Rightarrow x = \alpha_1(t), y = \alpha_2(t)$

Definition

Theorem

Let α, β be equivalent parameterizations of Γ . Then

$$I_\alpha = I_\beta$$

Definition

Theorem

Let α, β be equivalent parameterizations of Γ . Then

$$I_\alpha = I_\beta$$

Proof.

Apply the definition of equivalence and the Chain Rule. □

Example

- Let $p(x, y) = x^2y$, $q(x, y) = 2$ and $\Gamma = \{z : |z| = 1, y \geq 0\}$

Example

- Let $p(x, y) = x^2y$, $q(x, y) = 2$ and $\Gamma = \{z : |z| = 1, y \geq 0\}$
- Compute

$$\int_{\Gamma} [p \, dx + q \, dy]$$

Example

- Let $p(x, y) = x^2y$, $q(x, y) = 2$ and $\Gamma = \{z : |z| = 1, y \geq 0\}$
- Compute

$$\int_{\Gamma} [p \, dx + q \, dy]$$

- $\alpha(t) = (\cos(2\theta), \sin(2\theta))$, $\theta \in [0, \pi/2]$

Example

- Let $p(x, y) = x^2y$, $q(x, y) = 2$ and $\Gamma = \{z : |z| = 1, y \geq 0\}$
- Compute

$$\int_{\Gamma} [p \, dx + q \, dy]$$

- $\alpha(t) = (\cos(2\theta), \sin(2\theta))$, $\theta \in [0, \pi/2]$
- $\beta(t) = (-t, \sqrt{1 - t^2})$, $t \in [-1, 1]$

Example

- Let $p(x, y) = x^2y$, $q(x, y) = 2$ and $\Gamma = \{z : |z| = 1, y \geq 0\}$
- Compute

$$\int_{\Gamma} [p \, dx + q \, dy]$$

- $\alpha(t) = (\cos(2\theta), \sin(2\theta))$, $\theta \in [0, \pi/2]$
- $\beta(t) = (-t, \sqrt{1 - t^2})$, $t \in [-1, 1]$
- Find the equivalence between α, β

Exact Differentials

- $p \, dx + q \, dy$ - differential, first order differential, one-form

Exact Differentials

- $p \, dx + q \, dy$ - differential, first order differential, one-form
- Given $u \in \mathfrak{C}^1(\Omega)$, $du = u_x \, dx + u_y \, dy$

Exact Differentials

- $p \, dx + q \, dy$ - differential, first order differential, one-form
- Given $u \in \mathcal{C}^1(\Omega)$, $du = u_x \, dx + u_y \, dy$
- Given $p \, dx + q \, dy$, does there exist u such that $u_x = p$ and $u_y = q$?

Exact Differentials

- $p \, dx + q \, dy$ - differential, first order differential, one-form
- Given $u \in \mathcal{C}^1(\Omega)$, $du = u_x \, dx + u_y \, dy$
- Given $p \, dx + q \, dy$, does there exist u such that $u_x = p$ and $u_y = q$?
- Integrals of such differentials are “nice”

Exact Differential Diagram

$$u(x, y) = C$$
$$p(x, y) dx + q(x, y) dy = 0$$
$$\frac{\partial}{\partial x} \swarrow \qquad \qquad \qquad \searrow \frac{\partial}{\partial y}$$
$$p(x, y) dx \qquad \qquad + \qquad \qquad q(x, y) dy = 0$$
$$\frac{\partial}{\partial y} \downarrow \qquad \qquad \qquad \downarrow \frac{\partial}{\partial x}$$
$$p_y \qquad \qquad \qquad q_x$$

Figure: Exact Equation Model

Path Independence Theorem

Theorem

Let Γ be a piecewise-smooth curve from z_0 to z_1 lying inside a domain Ω .
Let $u \in \mathfrak{C}^1(\Omega)$, then

$$\int_{\Gamma} du = u(z_1) - u(z_0)$$

Path Independence Theorem

Theorem

Let Γ be a piecewise-smooth curve from z_0 to z_1 lying inside a domain Ω .
Let $u \in \mathfrak{C}^1(\Omega)$, then

$$\int_{\Gamma} du = u(z_1) - u(z_0)$$

Proof.

Denote the parameterization by $\alpha : [a, b] \rightarrow \Gamma$. Apply the definition of du and the Chain Rule. □

Double Integrals

- Suppose $\Omega \subset \mathbb{R}^2$ is a bounded domain and $u = u(x, y) \in \mathfrak{C}(\bar{\Omega})$.

Double Integrals

- Suppose $\Omega \subset \mathbb{R}^2$ is a bounded domain and $u = u(x, y) \in \mathfrak{C}(\bar{\Omega})$.
- If $u(x, y) \geq 0$ for all $(x, y) \in \bar{\Omega}$, then

$$\iint_{\Omega} u \, dA$$

represents the volume of the right cylinder whose base is $\bar{\Omega}$ and whose height above (x, y) is $u(x, y)$.

Double Integrals

- Suppose $\Omega \subset \mathbb{R}^2$ is a bounded domain and $u = u(x, y) \in \mathfrak{C}(\bar{\Omega})$.
- If $u(x, y) \geq 0$ for all $(x, y) \in \bar{\Omega}$, then

$$\iint_{\Omega} u \, dA$$

represents the volume of the right cylinder whose base is $\bar{\Omega}$ and whose height above (x, y) is $u(x, y)$.

- Alternate notation:

$$\iint_{\Omega} u \, dx \, dy$$

Computation of Double Integrals

- Let Ω be the triangular region with vertices $(0, 0)$, $(1, 0)$, $(1, 1)$ and $u(x, y) = xy^2$

Computation of Double Integrals

- Let Ω be the triangular region with vertices $(0, 0)$, $(1, 0)$, $(1, 1)$ and $u(x, y) = xy^2$
- Method: iterated integrals

Computation of Double Integrals

- Let Ω be the triangular region with vertices $(0, 0)$, $(1, 0)$, $(1, 1)$ and $u(x, y) = xy^2$
- Method: iterated integrals
- Not always successful - Ω or u may not be “nice enough” for this to work

Green's Theorem

Theorem (Green's Theorem)

Let Ω be a k -connected Jordan domain and suppose $p(x, y), q(x, y)$ are functions in $\mathfrak{C}^1(\Omega^+)$ where Ω^+ is a domain containing both Ω and $\partial\Omega$.

Then

$$\int_{\partial\Omega} [p \, dx + q \, dy] = \iint_{\Omega} (q_x - p_y) \, dxdy \quad (2)$$

Green's Theorem

Theorem (Green's Theorem)

Let Ω be a k -connected Jordan domain and suppose $p(x, y), q(x, y)$ are functions in $\mathfrak{C}^1(\Omega^+)$ where Ω^+ is a domain containing both Ω and $\partial\Omega$.

Then

$$\int_{\partial\Omega} [p \, dx + q \, dy] = \iint_{\Omega} (q_x - p_y) \, dxdy \quad (2)$$

General proof is hard due to potentially “ugly” boundaries of arbitrary Jordan domains.

Green's Theorem

Theorem (Green's Theorem)

Let Ω be a k -connected Jordan domain and suppose $p(x, y), q(x, y)$ are functions in $C^1(\Omega^+)$ where Ω^+ is a domain containing both Ω and $\partial\Omega$.

Then

$$\int_{\partial\Omega} [p \, dx + q \, dy] = \iint_{\Omega} (q_x - p_y) \, dxdy \quad (2)$$

General proof is hard due to potentially “ugly” boundaries of arbitrary Jordan domains. Proof below will be for the case Ω is a rectangular region.

Green's Theorem

Theorem (Green's Theorem)

Let Ω be a k -connected Jordan domain and suppose $p(x, y), q(x, y)$ are functions in $C^1(\Omega^+)$ where Ω^+ is a domain containing both Ω and $\partial\Omega$.

Then

$$\int_{\partial\Omega} [p \, dx + q \, dy] = \iint_{\Omega} (q_x - p_y) \, dxdy \quad (2)$$

General proof is hard due to potentially “ugly” boundaries of arbitrary Jordan domains. Proof below will be for the case Ω is a rectangular region. Relates the integral over the boundary to the integral over the interior.

Proof of Green's Theorem

Proof.

Let $\Omega = \{(x, y) : x_0 < x < x_1, y_0 < y < y_1\}$ and denote $\partial\Omega = \Gamma_i$ for $i = 1, 2, 3, 4$ with positive orientation. Then compute $-\iint_{\Omega} p_y dx dy$ iteratively to get $\int_{\partial\Omega} p(x, y) dx$ and observing that $dx = 0$ on vertical edges. Now compute $\iint_{\Omega} q_x dx dy$ iteratively to get $\int_{\partial\Omega} q dy$ using a similar observation about dy on $\partial\Omega$. Integral properties give the desired result from here. □

Example

- Let Ω be the triangular region with vertices $(1, 0), (2, 0), (2, 1)$ with positive orientation.

Example

- Let Ω be the triangular region with vertices $(1, 0), (2, 0), (2, 1)$ with positive orientation.
- Compute

$$\int_{\partial\Omega} [ydx - xdy]$$

Example

- Let Ω be the triangular region with vertices $(1, 0), (2, 0), (2, 1)$ with positive orientation.
- Compute

$$\int_{\partial\Omega} [ydx - xdy]$$

- iteratively

Example

- Let Ω be the triangular region with vertices $(1, 0), (2, 0), (2, 1)$ with positive orientation.
- Compute

$$\int_{\partial\Omega} [ydx - xdy]$$

- iteratively
- using Green's theorem

Prerequisites and Review

- Assume that Ω is a Jordan domain, $p, q \in \mathfrak{C}^2(\Omega^+)$

Prerequisites and Review

- Assume that Ω is a Jordan domain, $p, q \in \mathfrak{C}^2(\Omega^+)$
- Recall: $\nabla p(x, y) = \langle p_x, p_y \rangle$ and $\Delta p = p_{xx} + p_{yy}$

Prerequisites and Review

- Assume that Ω is a Jordan domain, $p, q \in \mathfrak{C}^2(\Omega^+)$
- Recall: $\nabla p(x, y) = \langle p_x, p_y \rangle$ and $\Delta p = p_{xx} + p_{yy}$
- Therefore $\Delta = \nabla \cdot \nabla$

Prerequisites and Review

- Assume that Ω is a Jordan domain, $p, q \in \mathfrak{C}^2(\Omega^+)$
- Recall: $\nabla p(x, y) = \langle p_x, p_y \rangle$ and $\Delta p = p_{xx} + p_{yy}$
- Therefore $\Delta = \nabla \cdot \nabla$
- Consider

$$\int_{\partial\Omega} \frac{\partial q}{\partial n} ds$$

Prerequisites and Review

- Assume that Ω is a Jordan domain, $p, q \in \mathfrak{C}^2(\Omega^+)$
- Recall: $\nabla p(x, y) = \langle p_x, p_y \rangle$ and $\Delta p = p_{xx} + p_{yy}$
- Therefore $\Delta = \nabla \cdot \nabla$
- Consider

$$\int_{\partial\Omega} \frac{\partial q}{\partial n} ds$$

- Recall that Ω will be parameterized by arc-length; then $N = \langle \sigma'_2(s), -\sigma'_1(s) \rangle$ so that $\frac{\partial q}{\partial n} ds = q_x dy - q_y dx$

Prerequisites and Review

- Assume that Ω is a Jordan domain, $p, q \in \mathfrak{C}^2(\Omega^+)$
- Recall: $\nabla p(x, y) = \langle p_x, p_y \rangle$ and $\Delta p = p_{xx} + p_{yy}$
- Therefore $\Delta = \nabla \cdot \nabla$
- Consider

$$\int_{\partial\Omega} \frac{\partial q}{\partial n} ds$$

- Recall that Ω will be parameterized by arc-length; then $N = \langle \sigma'_2(s), -\sigma'_1(s) \rangle$ so that $\frac{\partial q}{\partial n} ds = q_x dy - q_y dx$
- Thus $\frac{\partial q}{\partial n} ds$ is a “hybrid notation” for a line integral

Green's Identities I

Theorem (Green I)

Let Ω be a k connected Jordan domain and $p, q \in \mathfrak{C}^2(\Omega^+)$. Then

$$\iint_{\Omega} \nabla p \cdot \nabla q \, dx dy = \int_{\partial\Omega} p \frac{\partial q}{\partial n} \, ds - \iint_{\Omega} p \Delta q \, dx dy \quad (3)$$

Interpretation: two dimensional integration by parts

Green's Identities II

Theorem (Green II)

Let Ω be a k connected Jordan domain and $p, q \in \mathfrak{C}^2(\Omega^+)$. Then

$$\int_{\partial\Omega} \left(p \frac{\partial q}{\partial n} - q \frac{\partial p}{\partial n} \right) ds = \iint_{\Omega} (p \Delta q - q \Delta p) dx dy \quad (4)$$

Proof.

Note that the left side of 4 is symmetric in p, q , therefore exchanging roles produces the same value. Performing the swap on the right side and subtracting gives the identity. □

Green's Identities III

Theorem (Green III)

Let Ω be a k connected Jordan domain with $\zeta \in \Omega$ and $q \in \mathfrak{C}^2(\Omega^+)$. Let $z = (x, y)$, $r = |z - \zeta|$, then

$$q(\zeta) = \frac{1}{2\pi} \iint_{\Omega} \ln r \Delta q(z) dx dy - \frac{1}{2\pi} \int_{\partial\Omega} \left(\ln r \frac{\partial q}{\partial n}(z) - q(z) \frac{\partial \ln r}{\partial n} \right) ds \quad (5)$$

Application

Theorem (Inside-Outside Theorem)

Let Ω be a k connected Jordan domain and $q \in \mathfrak{C}^2(\Omega^+)$. Then

$$\int_{\partial\Omega} \frac{\partial q}{\partial n} ds = \iint_{\Omega} \Delta q dx dy \quad (6)$$

Interpretation: $\int_{\partial\Omega} \frac{\partial q}{\partial n} ds$ is “net change” of q across $\partial\Omega$ (outside) is equal to “total change” of q over Ω $\iint_{\Omega} \Delta q dx dy$ (inside).

Application

Theorem (Inside-Outside Theorem)

Let Ω be a k connected Jordan domain and $q \in \mathfrak{C}^2(\Omega^+)$. Then

$$\int_{\partial\Omega} \frac{\partial q}{\partial n} ds = \iint_{\Omega} \Delta q dx dy \quad (6)$$

Proof.

Take $p = 1$ in Green I. □

Interpretation: $\int_{\partial\Omega} \frac{\partial q}{\partial n} ds$ is “net change” of q across $\partial\Omega$ (outside) is equal to “total change” of q over Ω $\iint_{\Omega} \Delta q dx dy$ (inside).

Natural Log Lemma

Lemma

Fix $\zeta \in \mathbb{R}^2$ and define $r = |z - \zeta|$ for any $z \in \mathbb{R}^2$. Then

Natural Log Lemma

Lemma

Fix $\zeta \in \mathbb{R}^2$ and define $r = |z - \zeta|$ for any $z \in \mathbb{R}^2$. Then

- ① $\ln r$ is constant on circles $C(\zeta; r)$ centered at ζ

Natural Log Lemma

Lemma

Fix $\zeta \in \mathbb{R}^2$ and define $r = |z - \zeta|$ for any $z \in \mathbb{R}^2$. Then

- ① $\ln r$ is constant on circles $C(\zeta; r)$ centered at ζ
- ② $\ln r$ is harmonic as a function of z in the punctured plane $\mathbb{R}^2 \setminus \{\zeta\}$; that is $\Delta \ln r = 0$

Natural Log Lemma

Lemma

Fix $\zeta \in \mathbb{R}^2$ and define $r = |z - \zeta|$ for any $z \in \mathbb{R}^2$. Then

- ① $\ln r$ is constant on circles $C(\zeta; r)$ centered at ζ
- ② $\ln r$ is harmonic as a function of z in the punctured plane $\mathbb{R}^2 \setminus \{\zeta\}$; that is $\Delta \ln r = 0$

Proof.



Natural Log Lemma

Lemma

Fix $\zeta \in \mathbb{R}^2$ and define $r = |z - \zeta|$ for any $z \in \mathbb{R}^2$. Then

- ① $\ln r$ is constant on circles $C(\zeta; r)$ centered at ζ
- ② $\ln r$ is harmonic as a function of z in the punctured plane $\mathbb{R}^2 \setminus \{\zeta\}$; that is $\Delta \ln r = 0$

Proof.

- ① Observe that r is constant on $C(\zeta; r)$



Natural Log Lemma

Lemma

Fix $\zeta \in \mathbb{R}^2$ and define $r = |z - \zeta|$ for any $z \in \mathbb{R}^2$. Then

- ① $\ln r$ is constant on circles $C(\zeta; r)$ centered at ζ
- ② $\ln r$ is harmonic as a function of z in the punctured plane $\mathbb{R}^2 \setminus \{\zeta\}$; that is $\Delta \ln r = 0$

Proof.

- ① Observe that r is constant on $C(\zeta; r)$
- ② Let $z = (x, y)$, $\zeta = (\xi, \eta)$ and compute r using Euclidean distance. Then use properties of natural log and partial derivatives.



Green III Proof I

Recall:

Theorem (Green III)

Let Ω be a k connected Jordan domain with $\zeta \in \Omega$ and $q \in C^2(\Omega^+)$. Let $z = (x, y)$, $r = |z - \zeta|$, then

$$q(\zeta) = \frac{1}{2\pi} \iint_{\Omega} \ln r \Delta q(z) dx dy - \frac{1}{2\pi} \int_{\partial\Omega} \left(\ln r \frac{\partial q}{\partial n}(z) - q(z) \frac{\partial \ln r}{\partial n} \right) ds \quad (7)$$

Proof strategy: Apply Green II with $p = \ln r$ on $\Omega \setminus \bar{D}(\zeta; \epsilon)$ and let $\epsilon \rightarrow 0$.

Green III Proof II

Proof Continued

Proof.



Proof Continued

Proof.

- ① Consider $\int_0^{2\pi} \left(\ln \epsilon \frac{\partial q}{\partial n} + q \frac{1}{\epsilon} \right) \epsilon d\theta$



Proof Continued

Proof.

- ① Consider $\int_0^{2\pi} \left(\ln \epsilon \frac{\partial q}{\partial n} + q \frac{1}{\epsilon} \right) \epsilon d\theta$
- ② Observe that there exists $\theta_\epsilon \in [0, 2\pi]$ such that $\int_0^{2\pi} q d\theta = \int_0^{2\pi} q(\epsilon, \theta) d\theta = 2\pi q(\epsilon, \theta_\epsilon)$ by the Mean Value Theorem



Proof Continued

Proof.

- ① Consider $\int_0^{2\pi} \left(\ln \epsilon \frac{\partial q}{\partial n} + q \frac{1}{\epsilon} \right) \epsilon d\theta$
- ② Observe that there exists $\theta_\epsilon \in [0, 2\pi]$ such that $\int_0^{2\pi} q d\theta = \int_0^{2\pi} q(\epsilon, \theta) d\theta = 2\pi q(\epsilon, \theta_\epsilon)$ by the Mean Value Theorem
- ③ Since q is continuous, $2\pi q(\epsilon, \theta_\epsilon) \rightarrow 2\pi q(\zeta)$ as $\epsilon \rightarrow 0$



Proof Continued

Proof.

- ① Consider $\int_0^{2\pi} \left(\ln \epsilon \frac{\partial q}{\partial n} + q \frac{1}{\epsilon} \right) \epsilon d\theta$
- ② Observe that there exists $\theta_\epsilon \in [0, 2\pi]$ such that $\int_0^{2\pi} q d\theta = \int_0^{2\pi} q(\epsilon, \theta) d\theta = 2\pi q(\epsilon, \theta_\epsilon)$ by the Mean Value Theorem
- ③ Since q is continuous, $2\pi q(\epsilon, \theta_\epsilon) \rightarrow 2\pi q(\zeta)$ as $\epsilon \rightarrow 0$
- ④ Consider $\epsilon \ln \epsilon \int_0^{2\pi} \frac{\partial q}{\partial n} d\theta$.



Proof Continued

Proof.

- ① Consider $\int_0^{2\pi} \left(\ln \epsilon \frac{\partial q}{\partial n} + q \frac{1}{\epsilon} \right) \epsilon d\theta$
- ② Observe that there exists $\theta_\epsilon \in [0, 2\pi]$ such that $\int_0^{2\pi} q d\theta = \int_0^{2\pi} q(\epsilon, \theta) d\theta = 2\pi q(\epsilon, \theta_\epsilon)$ by the Mean Value Theorem
- ③ Since q is continuous, $2\pi q(\epsilon, \theta_\epsilon) \rightarrow 2\pi q(\zeta)$ as $\epsilon \rightarrow 0$
- ④ Consider $\epsilon \ln \epsilon \int_0^{2\pi} \frac{\partial q}{\partial n} d\theta$.
- ⑤ Claim: $\epsilon \ln \epsilon \rightarrow 0$ as $\epsilon \rightarrow 0$ by L'Hôpital's Rule



Proof Continued Again

Proof.



Interpretation: theoretical use only!

Proof Continued Again

Proof.

- ① Now 8 reduces to

$$2\pi q(\zeta) = \lim_{\epsilon \rightarrow 0} \iint_{\Omega \setminus \bar{D}(\zeta; \epsilon)} \ln r \Delta q \, dx dy - \int_{\partial\Omega} \left(\ln r \frac{\partial q}{\partial n} - q \frac{\partial \ln r}{\partial n} \right) ds \quad (9)$$



Interpretation: theoretical use only!

Proof Continued Again

Proof.

- ① Now 8 reduces to

$$2\pi q(\zeta) = \lim_{\epsilon \rightarrow 0} \iint_{\Omega \setminus \bar{D}(\zeta; \epsilon)} \ln r \Delta q \, dx dy - \int_{\partial\Omega} \left(\ln r \frac{\partial q}{\partial n} - q \frac{\partial \ln r}{\partial n} \right) ds \quad (9)$$

- ② The limit must exist and equal

$$\iint_{\Omega} \ln r \Delta q \, dx dy \quad (10)$$



Interpretation: theoretical use only!

Future Useage

Note that $\frac{\partial \ln r}{\partial n} = -\frac{\partial \ln r}{\partial r}$ on the boundary of a disc (that is, on circles).
This fact will be treated as “common knowledge” from this point onwards.

Harmonic Functions

Objectives

- Properties

Objectives

- Properties
- Principles

Examples

- Constant functions

Verify $\Delta u = 0$ as exercise. Also construct/guess/determine two harmonic functions such that the product is not harmonic.

Examples

- Constant functions
- $u(x, y) = x, v(x, y) = y$

Verify $\Delta u = 0$ as exercise. Also construct/guess/determine two harmonic functions such that the product is not harmonic.

Examples

- Constant functions
- $u(x, y) = x, v(x, y) = y$
- $u(x, y) = x^2 - y^2$

Verify $\Delta u = 0$ as exercise. Also construct/guess/determine two harmonic functions such that the product is not harmonic.

Examples

- Constant functions
- $u(x, y) = x, v(x, y) = y$
- $u(x, y) = x^2 - y^2$
- $u(x, y) = e^x \sin(y), v(x, y) = e^x \cos(y)$

Verify $\Delta u = 0$ as exercise. Also construct/guess/determine two harmonic functions such that the product is not harmonic.

Examples

- Constant functions
- $u(x, y) = x, v(x, y) = y$
- $u(x, y) = x^2 - y^2$
- $u(x, y) = e^x \sin(y), v(x, y) = e^x \cos(y)$
- $u(x, y) = v(x, y) + w(x, y)$ where v, w are harmonic functions

Verify $\Delta u = 0$ as exercise. Also construct/guess/determine two harmonic functions such that the product is not harmonic.

Fundamental Theorem

Theorem

Let Ω be a k connected Jordan domain and u a harmonic function in $C^2(\Omega^+)$. Then

Fundamental Theorem

Theorem

Let Ω be a k connected Jordan domain and u a harmonic function in $C^2(\Omega^+)$. Then

① $\int_{\partial\Omega} \frac{\partial u}{\partial n}(z) ds = 0$

Fundamental Theorem

Theorem

Let Ω be a k connected Jordan domain and u a harmonic function in $C^2(\Omega^+)$. Then

- ① $\int_{\partial\Omega} \frac{\partial u}{\partial n}(z) ds = 0$
- ② for $\zeta \in \Omega$, $z \in \partial\Omega$,

$$u(\zeta) = -\frac{1}{2\pi} \int_{\partial\Omega} \left(\ln r \frac{\partial u}{\partial n}(z) - u(z) \frac{\partial \ln r}{\partial n} \right) ds \quad (11)$$

where $r = |z - \zeta|$.

Fundamental Theorem

Theorem

Let Ω be a k connected Jordan domain and u a harmonic function in $C^2(\Omega^+)$. Then

- ① $\int_{\partial\Omega} \frac{\partial u}{\partial n}(z) ds = 0$
- ② for $\zeta \in \Omega$, $z \in \partial\Omega$,

$$u(\zeta) = -\frac{1}{2\pi} \int_{\partial\Omega} \left(\ln r \frac{\partial u}{\partial n}(z) - u(z) \frac{\partial \ln r}{\partial n} \right) ds \quad (11)$$

where $r = |z - \zeta|$.

Proof.

Note that $\Delta u = 0$ and apply Theorem 13 (part 1) then use Green III (part 2). □

Temperature

- Harmonic functions are those that exactly describe steady-state temperatures in a domain.

Temperature

- Harmonic functions are those that exactly describe steady-state temperatures in a domain.
- If a Jordan domain is made of a heat conducting material with either sources or sinks at each point of the boundary, then the temperature at an interior point is constant along specific curves -isotherms

Temperature

- Harmonic functions are those that exactly describe steady-state temperatures in a domain.
- If a Jordan domain is made of a heat conducting material with either sources or sinks at each point of the boundary, then the temperature at an interior point is constant along specific curves -isotherms
- Note that heat flows throughout the domain (not along isotherms) from hot spots to cold spots

Temperature

- Harmonic functions are those that exactly describe steady-state temperatures in a domain.
- If a Jordan domain is made of a heat conducting material with either sources or sinks at each point of the boundary, then the temperature at an interior point is constant along specific curves -isotherms
- Note that heat flows throughout the domain (not along isotherms) from hot spots to cold spots
- Assume that the temperature does not vary with time (only location)

Temperature

- Harmonic functions are those that exactly describe steady-state temperatures in a domain.
- If a Jordan domain is made of a heat conducting material with either sources or sinks at each point of the boundary, then the temperature at an interior point is constant along specific curves -isotherms
- Note that heat flows throughout the domain (not along isotherms) from hot spots to cold spots
- Assume that the temperature does not vary with time (only location)
- Goal: if u is the temperature, then $\Delta u = 0$

Mathematical Perspective

- For any small portion of the domain, the heat flow in must equal the heat flow out (why?)

Mathematical Perspective

- For any small portion of the domain, the heat flow in must equal the heat flow out (why?)
- Suppose Γ is a small, closed loop inside the domain.

Mathematical Perspective

- For any small portion of the domain, the heat flow in must equal the heat flow out (why?)
- Suppose Γ is a small, closed loop inside the domain.
- The net heat flow across Γ must be zero: $\int_{\Gamma} \frac{\partial u}{\partial n} ds = 0$

Mathematical Perspective

- For any small portion of the domain, the heat flow in must equal the heat flow out (why?)
- Suppose Γ is a small, closed loop inside the domain.
- The net heat flow across Γ must be zero: $\int_{\Gamma} \frac{\partial u}{\partial n} ds = 0$
- This is based on the assumption $\Delta u = 0$

Characterization of Harmonic Functions

Theorem

Let Ω be a plane domain and $u \in \mathfrak{C}^2(\Omega)$. Then u is harmonic in Ω if and only if for every Jordan curve Γ inside Ω whose interior lies inside Ω ,

$$\int_{\Gamma} \frac{\partial u}{\partial n} \, ds = 0 \tag{12}$$

Characterization of Harmonic Functions

Theorem

Let Ω be a plane domain and $u \in \mathcal{C}^2(\Omega)$. Then u is harmonic in Ω if and only if for every Jordan curve Γ inside Ω whose interior lies inside Ω ,

$$\int_{\Gamma} \frac{\partial u}{\partial n} ds = 0 \tag{12}$$

Hence a steady state temperature inside a Jordan domain is harmonic.
And conversely!

Bump Principle

Lemma (Bump Principle)

Let Ω be a domain and $q \in \mathfrak{C}(\Omega)$ with $q(z) \geq 0$ for all $z \in \Omega$. Then

$$\iint_{\Omega} q \, dx dy = 0 \Leftrightarrow q \equiv 0 \quad (13)$$

Boundary Mean-Value Theorem

Theorem (Boundary Mean-Value Theorem)

Let u be harmonic in a domain Ω , $\zeta \in \Omega$ and suppose $\bar{D}(\zeta; R) \subset \Omega$. Then $u(\zeta)$ is determined by the values $u(z)$ for $z \in \partial D$:

$$u(\zeta) = \frac{1}{2\pi R} \int_{\partial D} u(z) dz = \frac{1}{2\pi} \int_0^{2\pi} u(R, \theta) d\theta \quad (14)$$

Note: $u(R, \theta)$ is defined using polar coordinates centered at ζ .

Solid Mean-Value Theorem

Theorem (Solid Mean-Value Theorem)

Let u be harmonic in a domain Ω , $\zeta \in \Omega$ and suppose $\bar{D}(\zeta; R) \subset \Omega$. Then

$$u(\zeta) = \frac{1}{2\pi R^2} \iint_D u \, dx \, dy \quad (15)$$

Solid Mean-Value Theorem

Theorem (Solid Mean-Value Theorem)

Let u be harmonic in a domain Ω , $\zeta \in \Omega$ and suppose $\bar{D}(\zeta; R) \subset \Omega$. Then

$$u(\zeta) = \frac{1}{2\pi R^2} \iint_D u \, dx \, dy \quad (15)$$

Proof.

Apply Theorem 17 to a disc with $0 < r \leq R$; multiply both sides by $r \, dr$ and integrate from $r = 0$ to $r = R$. Change coordinates to rectangular and evaluate. □

Characterization of Harmonic Functions

Harmonic functions are characterized by their boundary values.

Definition

Let Ω be a domain. Then $u \in \mathcal{C}^2(\Omega)$ has the **circumferential mean value property** in Ω if and only if for each $D = D(\zeta; R)$ with $\bar{D} \subset \Omega$

$$u(\zeta) = \frac{1}{2\pi R} \int_{\partial D} u \, ds \quad (16)$$

Characterization Theorem

Theorem

Let Ω be a domain and $u \in \mathfrak{C}^2(\Omega)$. Then u is harmonic if and only if it has the circumferential mean value property in Ω .

Characterization Theorem

Theorem

Let Ω be a domain and $u \in \mathfrak{C}^2(\Omega)$. Then u is harmonic if and only if it has the circumferential mean value property in Ω .

Proof.

Differentiate defining equation (in polar coordinates centered at ζ) and multiply by r ; simplify. □

Strong Maximum Principle

It is not possible for a non-constant harmonic function to attain its maximum value at an interior point of Ω ; hence the maximum value is only attainable on $\partial\Omega$.

Lemma (Strong Maximum Principle)

A non-constant harmonic function on a domain Ω does not assume its maximum or minimum on Ω .

Strong Maximum Principle

It is not possible for a non-constant harmonic function to attain its maximum value at an interior point of Ω ; hence the maximum value is only attainable on $\partial\Omega$.

Lemma (Strong Maximum Principle)

A non-constant harmonic function on a domain Ω does not assume its maximum or minimum on Ω .

For example, $u(x, y) = 1 - x^2 - y^2$ has maximum value on \mathbb{R}^2 at $(0, 0)$ by definition.

Strong Maximum Proof

Proof.

Assume u attains its maximum, c , at $\zeta \in \Omega$. If u is not constant, then the set $A = \{w : u(w) < c\}$ is non-empty and open by continuity. Furthermore $B = \{t : u(t) = c\}$ is not open or else $\Omega = A \cup B$ and Ω is disconnected. Hence there exists some $\zeta_1 \in \Omega$ and $r > 0$ such that $\bar{D}(\zeta_1; r) \subset \Omega$ with $u(\zeta_1) = c$ and at least one $z \in \partial D$ with $u(z) < c$. Using 2 on the arc near z gives a set of values z such that $u(z) < c$ for all z on this arc. But then

$$u(\zeta_1) = \frac{1}{2\pi} \int_0^{2\pi} u(r, \theta) d\theta < \frac{1}{2\pi} \int_0^{2\pi} u(\zeta_1) d\theta = u(\zeta_1)$$

which contradicts the expected value for $u(\zeta_1)$ from Theorem 17 as u is harmonic. □

Applying this argument to the function $w = -u$ proves the minimum value of a harmonic function must also occur on the boundary.

Weak Maximum Principle

Lemma (Weak Maximum Principle)

Let Ω be a bounded domain with $u \in \mathfrak{C}(\bar{\Omega})$ and u harmonic on Ω . Then either u is constant on $\bar{\Omega}$ or u assumes its maximum and minimum values on $\partial\Omega$ only.

Weak Maximum Principle

Lemma (Weak Maximum Principle)

Let Ω be a bounded domain with $u \in \mathfrak{C}(\bar{\Omega})$ and u harmonic on Ω . Then either u is constant on $\bar{\Omega}$ or u assumes its maximum and minimum values on $\partial\Omega$ only.

Proof.

Observe that $\bar{\Omega}$ is compact, hence $\{u(z) : z \in \bar{\Omega}\}$ is a closed interval of finite length. Note that for any $z \in \Omega$, $u(z)$ is not the maximum value by the strong maximum principle applied to a disc $D(z; r) \subset \Omega$. □

Application of Maximum Principles

Theorem

Suppose Ω is a bounded domain and u, v are continuous functions on $\bar{\Omega}$ with $u(z) = v(z)$ for all $z \in \partial\Omega$. Then $u(\zeta) = v(\zeta)$ for all $\zeta \in \bar{\Omega}$.

Application of Maximum Principles

Theorem

Suppose Ω is a bounded domain and u, v are continuous functions on $\bar{\Omega}$ with $u(z) = v(z)$ for all $z \in \partial\Omega$. Then $u(\zeta) = v(\zeta)$ for all $\zeta \in \bar{\Omega}$.

Proof.

Note that $u - v$ is harmonic in Ω and vanishes on $\partial\Omega$, hence has maximum value 0. □

Definition

Definition

*A harmonic function with domain \mathbb{R}^2 is called **entire**.*

Definition

*A harmonic function with domain \mathbb{R}^2 is called **entire**.*

Note that no consideration is given to unbounded domains which are not \mathbb{R}^2 aside from the Strong Maximum Principle.

Harnack's Inequality

Lemma (Harnack's Inequality)

Let $D = D(z_0; R)$ be an open disc and u be harmonic on D such that $u(z) \geq 0$ for all $z \in D$. Then for all $z \in D$

$$0 \leq u(z) \leq \left(\frac{R}{R - |z - z_0|} \right)^2 u(z_0) \quad (17)$$

Interpretation: the growth rate of harmonic functions is limited by the distance to the boundary of a disc.

Proof of Harnack's Inequality

Proof.

Apply Theorem 18 to a disc $D'(z; R - |z - z_0|)$

$$0 \leq u(z) = \frac{1}{\pi(R - |z - z_0|)^2} \iint_{D'} u \, dx \, dy$$

but since $D' \subset D$

$$\iint_{D'} u \, dx \, dy \leq \iint_D u \, dx \, dy = \pi R^2 u(z_0)$$



Note that the growth rate bound is independent of derivatives of u - an unusual situation wholly dependent on the harmonic nature of u .

Liouville's Theorem

Theorem (Liouville's Theorem)

If u is entire, harmonic and bounded either above or below, then u is constant.

Liouville's Theorem

Theorem (Liouville's Theorem)

If u is entire, harmonic and bounded either above or below, then u is constant.

- Alternative phrasing: non-constant harmonic functions have unbounded images.

Liouville's Theorem

Theorem (Liouville's Theorem)

If u is entire, harmonic and bounded either above or below, then u is constant.

- Alternative phrasing: non-constant harmonic functions have unbounded images.
- Using heat: a non-constant steady state temperature achieves every temperature somewhere.

Liouville's Theorem Proof

Proof.

Suppose u is bounded below: $u(z) \geq \alpha$ for some α . Then $v = u - \alpha \geq 0$ is harmonic and constant if and only if u is constant. Therefore WLOG $\alpha = 0$. Choose any two points $z, \zeta \in \mathbb{R}^2$. For any $R > |z - \zeta|$,

$$0 \leq u(z) \leq \left(\frac{R}{R - |z - \zeta|} \right)^2 u(\zeta)$$

As $R \rightarrow \infty$, $0 \leq u(z) \leq u(\zeta)$. Interchanging z, ζ gives the reverse inequality hence $u(z) = u(\zeta)$. If u is bounded above, apply the argument to the harmonic function $w = -u$. □

Complex Numbers and Functions

Objectives

- Construct C

Objectives

- Construct C
- Complex functions

Objectives

- Construct C
- Complex functions
 - Derivatives

Objectives

- Construct C
- Complex functions
 - Derivatives
 - Analytic

Objectives

- Construct C
- Complex functions
 - Derivatives
 - Analytic
 - Transcendental

Construction

- Endow \mathbb{R}^2 with a sensible multiplication

Construction

- Endow \mathbb{R}^2 with a sensible multiplication
- Let $\mathbb{C} = \{z = x + yi : x, y \in \mathbb{R}\}$

Construction

- Endow \mathbb{R}^2 with a sensible multiplication
- Let $\mathbb{C} = \{z = x + yi : x, y \in \mathbb{R}\}$
- $x + yi = \xi + \eta i$ if and only if both $x = \xi, y = \eta$

Construction

- Endow \mathbb{R}^2 with a sensible multiplication
- Let $\mathbb{C} = \{z = x + yi : x, y \in \mathbb{R}\}$
- $x + yi = \xi + \eta i$ if and only if both $x = \xi$, $y = \eta$
- $(x + yi) \pm (\xi + \eta i) = (x \pm \xi) + (y \pm \eta)i$

Construction

- Endow \mathbb{R}^2 with a sensible multiplication
- Let $\mathbb{C} = \{z = x + yi : x, y \in \mathbb{R}\}$
- $x + yi = \xi + \eta i$ if and only if both $x = \xi$, $y = \eta$
- $(x + yi) \pm (\xi + \eta i) = (x \pm \xi) + (y \pm \eta)i$
- Define $i^2 = -1$; $1i = i$, $a(bi) = (ab)i$, $ai = ia$

Construction

- Endow \mathbb{R}^2 with a sensible multiplication
- Let $\mathbb{C} = \{z = x + yi : x, y \in \mathbb{R}\}$
- $x + yi = \xi + \eta i$ if and only if both $x = \xi$, $y = \eta$
- $(x + yi) \pm (\xi + \eta i) = (x \pm \xi) + (y \pm \eta)i$
- Define $i^2 = -1$; $1i = i$, $a(bi) = (ab)i$, $ai = ia$
- $(x + yi) \cdot (\xi + \eta i) = (x\xi - y\eta) + (x\eta + y\xi)i$

Division

Definition

If $z = x + yi$, then $\bar{z} = x - yi$ is the **conjugate** of z .

Division

Definition

If $z = x + yi$, then $\bar{z} = x - yi$ is the **conjugate** of z .

- First, $\frac{\zeta}{z} = \frac{1}{z} \cdot \zeta$ for $z \neq 0$

Division

Definition

If $z = x + yi$, then $\bar{z} = x - yi$ is the **conjugate** of z .

- First, $\frac{\zeta}{z} = \frac{1}{z} \cdot \zeta$ for $z \neq 0$
- Multiply $1/z$ by $1 = \frac{\bar{z}}{\bar{z}}$; simplify

Division

Definition

If $z = x + yi$, then $\bar{z} = x - yi$ is the **conjugate** of z .

- First, $\frac{\zeta}{z} = \frac{1}{z} \cdot \zeta$ for $z \neq 0$
- Multiply $1/z$ by $1 = \frac{\bar{z}}{\bar{z}}$; simplify
- Example(s)

Division

Definition

If $z = x + yi$, then $\bar{z} = x - yi$ is the **conjugate** of z .

- First, $\frac{\zeta}{z} = \frac{1}{z} \cdot \zeta$ for $z \neq 0$
- Multiply $1/z$ by $1 = \frac{\bar{z}}{\bar{z}}$; simplify
- Example(s)

Compute the following:

Division

Definition

If $z = x + yi$, then $\bar{z} = x - yi$ is the **conjugate** of z .

- First, $\frac{\zeta}{z} = \frac{1}{z} \cdot \zeta$ for $z \neq 0$
- Multiply $1/z$ by $1 = \frac{\bar{z}}{\bar{z}}$; simplify
- Example(s)

Compute the following:

- $z\bar{z} = |z|^2$

Division

Definition

If $z = x + yi$, then $\bar{z} = x - yi$ is the **conjugate** of z .

- First, $\frac{\zeta}{z} = \frac{1}{z} \cdot \zeta$ for $z \neq 0$
- Multiply $1/z$ by $1 = \frac{\bar{z}}{\bar{z}}$; simplify
- Example(s)

Compute the following:

- $z\bar{z} = |z|^2$
- $z\bar{\zeta} - \bar{z}\bar{\zeta}$

The Complex Plane

- Identify $(x, y) \in \mathbb{R}^2$ with $z = x + yi \in \mathbb{C}$

The Complex Plane

- Identify $(x, y) \in \mathbb{R}^2$ with $z = x + yi \in \mathbb{C}$
- Hence point multiplication (not a dot product) is sensible

Definition

Definition

Let Ω be a subset of \mathbb{C} . A **complex function** $f : \Omega \rightarrow \mathbb{C}$ is a rule that assigns to each $z \in \Omega$ a complex number $f(z)$.

Examples:

Definition

Definition

Let Ω be a subset of \mathbb{C} . A **complex function** $f : \Omega \rightarrow \mathbb{C}$ is a rule that assigns to each $z \in \Omega$ a complex number $f(z)$.

Examples:

- $f(z) = z^2 + (1 - i)z - (2 + 3i)$

Definition

Definition

Let Ω be a subset of \mathbb{C} . A **complex function** $f : \Omega \rightarrow \mathbb{C}$ is a rule that assigns to each $z \in \Omega$ a complex number $f(z)$.

Examples:

- $f(z) = z^2 + (1 - i)z - (2 + 3i)$
- $g(z) = \bar{z}$

Definition

Definition

Let Ω be a subset of \mathbb{C} . A **complex function** $f : \Omega \rightarrow \mathbb{C}$ is a rule that assigns to each $z \in \Omega$ a complex number $f(z)$.

Examples:

- $f(z) = z^2 + (1 - i)z - (2 + 3i)$
- $g(z) = \bar{z}$
- $h(z) = i - z^3$

Real and Imaginary Parts

- if $w = f(z)$ is a complex number, then $w = u + iv$

Real and Imaginary Parts

- if $w = f(z)$ is a complex number, then $w = u + iv$
- but w varies with z ; hence $u = u(z)$ and $v = v(z)$ also

Real and Imaginary Parts

- if $w = f(z)$ is a complex number, then $w = u + iv$
- but w varies with z ; hence $u = u(z)$ and $v = v(z)$ also
- therefore $f(z) = u(z) + iv(z)$ is the sum of the real and imaginary functions

Complex Limits

Definition

Given $z_0 \in \mathbb{C}$, the symbols

$$\lim_{z \rightarrow z_0} f(z) = w$$

mean that for all $\epsilon > 0$, there exists $\delta > 0$ such that $z \in D(z_0; \delta)$ implies $f(z) \in D(w; \epsilon)$.

Complex Limits

Definition

Given $z_0 \in \mathbb{C}$, the symbols

$$\lim_{z \rightarrow z_0} f(z) = w$$

mean that for all $\epsilon > 0$, there exists $\delta > 0$ such that $z \in D(z_0; \delta)$ implies $f(z) \in D(w; \epsilon)$.

- This is the two dimensional statement of the $\epsilon - \delta$ definition of limit from introductory Calculus.

Complex Limits

Definition

Given $z_0 \in \mathbb{C}$, the symbols

$$\lim_{z \rightarrow z_0} f(z) = w$$

mean that for all $\epsilon > 0$, there exists $\delta > 0$ such that $z \in D(z_0; \delta)$ implies $f(z) \in D(w; \epsilon)$.

- This is the two dimensional statement of the $\epsilon - \delta$ definition of limit from introductory Calculus.
- Complex limits are compatible with arithmetic operations

Continuity

Definition

*The function $f : \Omega \rightarrow \mathbb{C}$ is **continuous** at z_0 if and only if*

Continuity

Definition

The function $f : \Omega \rightarrow \mathbb{C}$ is **continuous** at z_0 if and only if

- ① $z_0 \in \Omega$ hence $f(z_0)$ is defined

Continuity

Definition

The function $f : \Omega \rightarrow \mathbb{C}$ is **continuous** at z_0 if and only if

- ① $z_0 \in \Omega$ hence $f(z_0)$ is defined
- ② $\lim_{z \rightarrow z_0} f(z) = f(z_0)$

Continuity

Definition

The function $f : \Omega \rightarrow \mathbb{C}$ is **continuous** at z_0 if and only if

- ① $z_0 \in \Omega$ hence $f(z_0)$ is defined
- ② $\lim_{z \rightarrow z_0} f(z) = f(z_0)$

Algebraic combinations of continuous functions are also continuous.

Continuity

Definition

The function $f : \Omega \rightarrow \mathbb{C}$ is **continuous** at z_0 if and only if

- ① $z_0 \in \Omega$ hence $f(z_0)$ is defined
- ② $\lim_{z \rightarrow z_0} f(z) = f(z_0)$

Algebraic combinations of continuous functions are also continuous.

Note that $f(z) = u(z) + iv(z)$ is continuous at z_0 if and only if both u, v are continuous at z_0 .

Definition

Definition

Let $f : \Omega \rightarrow \mathbb{C}$ and $z_0 \in \Omega$. Then $f'(z_0)$ is the value

$$f'(z_0) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} \quad (18)$$

provided the limit exists.

Definition

Definition

Let $f : \Omega \rightarrow \mathbb{C}$ and $z_0 \in \Omega$. Then $f'(z_0)$ is the value

$$f'(z_0) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} \quad (18)$$

provided the limit exists.

Also denoted $\frac{df}{dz}(z_0)$

Definition

Definition

Let $f : \Omega \rightarrow \mathbb{C}$ and $z_0 \in \Omega$. Then $f'(z_0)$ is the value

$$f'(z_0) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} \quad (18)$$

provided the limit exists.

Also denoted $\frac{df}{dz}(z_0)$

Note that this complex limit is path independent.

Definition

Definition

Let $f : \Omega \rightarrow \mathbb{C}$ and $z_0 \in \Omega$. Then $f'(z_0)$ is the value

$$f'(z_0) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} \quad (18)$$

provided the limit exists.

Also denoted $\frac{df}{dz}(z_0)$

Note that this complex limit is path independent.

Additionally, the value $f'(z_0)$ is not to be interpreted as “slope”.

Derivative Rules

Theorem (Complex Derivative Rules)

Derivative Rules

Theorem (Complex Derivative Rules)

① $[f(z) \pm g(z)]' = f'(z) \pm g'(z)$

Derivative Rules

Theorem (Complex Derivative Rules)

- ① $[f(z) \pm g(z)]' = f'(z) \pm g'(z)$
- ② $[f(z)g(z)]' = f'(z)g(z) + f(z)g'(z)$

Derivative Rules

Theorem (Complex Derivative Rules)

- ① $[f(z) \pm g(z)]' = f'(z) \pm g'(z)$
- ② $[f(z)g(z)]' = f'(z)g(z) + f(z)g'(z)$
- ③ $\left[\frac{f(z)}{g(z)} \right]' = \frac{g(z)f'(z) - f(z)g'(z)}{(g(z))^2}$

Chain Rule

Composition is defined in the usual way.

Chain Rule

Composition is defined in the usual way.

Theorem (Chain Rule)

Suppose that $f'(z)$ and $g'(\zeta)$ exist where $g(\zeta) = z$. Then

$$(f \circ g)'(\zeta) = f'(g(\zeta))g'(\zeta) = f'(z)g'(\zeta) \quad (19)$$

Chain Rule

Composition is defined in the usual way.

Theorem (Chain Rule)

Suppose that $f'(z)$ and $g'(\zeta)$ exist where $g(\zeta) = z$. Then

$$(f \circ g)'(\zeta) = f'(g(\zeta))g'(\zeta) = f'(z)g'(\zeta) \quad (19)$$

Proof.

Simple case ($g(\zeta) = a\zeta + \beta$): multiply the difference quotient of the composite function by $1 = \frac{z-z_0}{z-z_0}$.



Chain Rule

Composition is defined in the usual way.

Theorem (Chain Rule)

Suppose that $f'(z)$ and $g'(\zeta)$ exist where $g(\zeta) = z$. Then

$$(f \circ g)'(\zeta) = f'(g(\zeta))g'(\zeta) = f'(z)g'(\zeta) \quad (19)$$

Proof.

Simple case ($g(\zeta) = a\zeta + \beta$): multiply the difference quotient of the composite function by $1 = \frac{z-z_0}{z-z_0}$.



The exponent formula works: $f(z) = z^n$ has $f'(z) = nz^{n-1}$ for n a natural number. PLTS.

Analytic Functions

Definition

*Let $f : \Omega \rightarrow \mathbb{C}$ be a function on Ω . Then f is **analytic** at $z_0 \in \Omega$ if and only if*

Analytic Functions

Definition

Let $f : \Omega \rightarrow \mathbb{C}$ be a function on Ω . Then f is **analytic** at $z_0 \in \Omega$ if and only if

- ① $f'(z)$ exists for all z in an open neighborhood of z_0

Analytic Functions

Definition

Let $f : \Omega \rightarrow \mathbb{C}$ be a function on Ω . Then f is **analytic** at $z_0 \in \Omega$ if and only if

- ① $f'(z)$ exists for all z in an open neighborhood of z_0
- ② f' is continuous in an open neighborhood of z_0

Analytic Functions

Definition

Let $f : \Omega \rightarrow \mathbb{C}$ be a function on Ω . Then f is **analytic** at $z_0 \in \Omega$ if and only if

- ① $f'(z)$ exists for all z in an open neighborhood of z_0
- ② f' is continuous in an open neighborhood of z_0

That is, f is continuously complex differentiable in an open neighborhood of z_0 .

Analytic Functions

Definition

Let $f : \Omega \rightarrow \mathbb{C}$ be a function on Ω . Then f is **analytic** at $z_0 \in \Omega$ if and only if

- ① $f'(z)$ exists for all z in an open neighborhood of z_0
- ② f' is continuous in an open neighborhood of z_0

That is, f is continuously complex differentiable in an open neighborhood of z_0 .

Extend to “analytic on Ω ” in obvious way.

Goursat's Theorem

Theorem (Goursat's Theorem)

Examples

- constant functions

Examples

- constant functions
- polynomial functions

Examples

- constant functions
- polynomial functions
- rational functions

Examples

- constant functions
- polynomial functions
- rational functions
- counter-example: $f(z) = \bar{z}$ is nowhere analytic. PLTS.

Motivation

Goal: determine analyticity of $f(z) = u(z) + iv(z)$ based on partial derivatives of u, v .

Motivation

Goal: determine analyticity of $f(z) = u(z) + iv(z)$ based on partial derivatives of u, v .

Lemma (Cauchy-Riemann Equations)

If f is analytic at $z \in \Omega$, then

$$f'(z) = u_x(z) + iv_x(z) = v_y(z) - iu_y(z) \quad (20)$$

so that

$$u_x = v_y \quad u_y = -v_x$$

Conversely, if $u, v \in \mathfrak{C}^1(\Omega)$ and satisfy these equations for all $z \in \Omega$, then $f = u + iv$ is analytic in Ω .

Cauchy-Riemann Equations Proof Outline

Proof.

Let z approach z_0 vertically, horizontally. Observe $f'(z_0)$ is unique. Apply Theorem 3 and the Cauchy-Riemann equations. Using the triangle inequality, $\epsilon_i \rightarrow 0$ as $z \rightarrow z_0$ hence f' exists; moreover is continuous. □

CR Equations in Polar Coordinates

Lemma

For $f(z) = u(r, \theta) + iv(r, \theta)$,

$$u_r = \frac{1}{r} v_\theta \quad u_\theta = -rv_r \quad (21)$$

CR Equations in Polar Coordinates

Lemma

For $f(z) = u(r, \theta) + iv(r, \theta)$,

$$u_r = \frac{1}{r} v_\theta \quad u_\theta = -rv_r \quad (21)$$

Proof.

Apply the Chain Rule.



Analytic and Harmonic Functions

Lemma

If $f(z) = u(z) + iv(z)$ is analytic with $u_x = v_y$ and $u_y = -v_x$, then u, v are harmonic functions.

Analytic and Harmonic Functions

Lemma

If $f(z) = u(z) + iv(z)$ is analytic with $u_x = v_y$ and $u_y = -v_x$, then u, v are harmonic functions.

That is, the real and imaginary parts of analytic functions are themselves harmonic functions. What??!!

The Exponential Function

Definition

For $z = x + iy \in \mathbb{C}$, define $\exp : \mathbb{C} \rightarrow \mathbb{C}$ by

$$\exp z = e^x(\cos y + i \sin y)$$

The Exponential Function

Definition

For $z = x + iy \in \mathbb{C}$, define $\exp : \mathbb{C} \rightarrow \mathbb{C}$ by

$$\exp z = e^x(\cos y + i \sin y)$$

Verify analyticity

The Exponential Function

Definition

For $z = x + iy \in \mathbb{C}$, define $\exp : \mathbb{C} \rightarrow \mathbb{C}$ by

$$\exp z = e^x(\cos y + i \sin y)$$

Verify analyticity

Use calculation to determine $[\exp z]'$

The Exponential Function

Definition

For $z = x + iy \in \mathbb{C}$, define $\exp : \mathbb{C} \rightarrow \mathbb{C}$ by

$$\exp z = e^x(\cos y + i \sin y)$$

Verify analyticity

Use calculation to determine $[\exp z]'$

Compute $\exp(z + 2\pi i)$.

Exponential Properties

- While $\exp(z)\exp(w) = \exp(z + w)$, the proof relies on trigonometric identities now.

Exponential Properties

- While $\exp(z)\exp(w) = \exp(z+w)$, the proof relies on trigonometric identities now.
- $e^{iy} = \cos(y) + i\sin(y)$

Exponential Properties

- While $\exp(z)\exp(w) = \exp(z+w)$, the proof relies on trigonometric identities now.
- $e^{iy} = \cos(y) + i\sin(y)$
- $e^z e^w = e^{z+w}$

Exponential Properties

- While $\exp(z)\exp(w) = \exp(z+w)$, the proof relies on trigonometric identities now.
- $e^{iy} = \cos(y) + i\sin(y)$
- $e^z e^w = e^{z+w}$
- $[e^z]' = e^z$

Exponential Properties

- While $\exp(z)\exp(w) = \exp(z+w)$, the proof relies on trigonometric identities now.
- $e^{iy} = \cos(y) + i\sin(y)$
- $e^z e^w = e^{z+w}$
- $[e^z]' = e^z$
- Polar form: changing $(x, y) \rightarrow (r, \theta)$ gives

$$z = x + iy = r(\cos \theta + i\sin \theta) = r \operatorname{cis} \theta = re^{i\theta}$$

Exponential Properties

- While $\exp(z)\exp(w) = \exp(z+w)$, the proof relies on trigonometric identities now.
- $e^{iy} = \cos(y) + i\sin(y)$
- $e^z e^w = e^{z+w}$
- $[e^z]' = e^z$
- Polar form: changing $(x, y) \rightarrow (r, \theta)$ gives

$$z = x + iy = r(\cos \theta + i\sin \theta) = r \operatorname{cis} \theta = re^{i\theta}$$

- Polar multiplication:

$$z_1 z_2 = r_1 r_2 e^{\theta_1 + \theta_2}$$

Uniqueness of Argument

- $\arg z$ is defined so that the range is $[0, 2\pi]$

Uniqueness of Argument

- $\arg z$ is defined so that the range is $[0, 2\pi]$
- Justification:

$$e^{i(\theta+2k\pi)} = e^{i\theta} e^{i2k\pi} = e^{i\theta} 1$$

Complex Roots

- If $z = re^{i\theta}$, then $z^n = r^n e^{in\theta}$

Complex Roots

- If $z = re^{i\theta}$, then $z^n = r^n e^{in\theta}$
- Goal: Solve $w^n = z$ for given $z = re^{i\theta} \in \mathbb{C}$

Complex Roots

- If $z = re^{i\theta}$, then $z^n = r^n e^{in\theta}$
- Goal: Solve $w^n = z$ for given $z = re^{i\theta} \in \mathbb{C}$
- Define: $\zeta_k = r^{1/n} e^{i(\theta+2k\pi)/n}$ for $n = 0, 1, \dots, n - 1$

Complex Roots

- If $z = re^{i\theta}$, then $z^n = r^n e^{in\theta}$
- Goal: Solve $w^n = z$ for given $z = re^{i\theta} \in \mathbb{C}$
- Define: $\zeta_k = r^{1/n} e^{i(\theta+2k\pi)/n}$ for $n = 0, 1, \dots, n - 1$
- Claim: $\zeta_k^n = z$; compute this

Complex Roots

- If $z = re^{i\theta}$, then $z^n = r^n e^{in\theta}$
- Goal: Solve $w^n = z$ for given $z = re^{i\theta} \in \mathbb{C}$
- Define: $\zeta_k = r^{1/n} e^{i(\theta+2k\pi)/n}$ for $n = 0, 1, \dots, n - 1$
- Claim: $\zeta_k^n = z$; compute this
- Claim: $\{\zeta_k\}$ are all distinct

Uniqueness of Roots

Proof.

Suppose $\zeta_k = \zeta_m$ for $0 \leq k < m \leq n - 1$. Then

$$\zeta_k \zeta_m^{-1} = 1 = (r^{1/n} e^{i(\theta+2\pi k)/n})(r^{-1/n} e^{-i(\theta+2\pi m)/n}) = e^{(2i(k-m)\pi)/n}$$

But $e^{2\pi i t} = 1$ if and only if t is an integer. Since $(k - m)/n$ is not an integer, contradiction. Furthermore, if $w^n = z$, then $w \in \{\zeta_k\}$: consider the factorization of $w^n - z = 0$ using $\{\zeta_k\}$. □

Roots of Unity

Definition

Define $\omega_k = e^{2\pi ik/n}$ so that ω_k is a solution to $z^n = 1$ for $k = 0, 1, \dots, n - 1$. Then ω_k is an **nth root of unity**.

Roots of Unity

Definition

Define $\omega_k = e^{2\pi ik/n}$ so that ω_k is a solution to $z^n = 1$ for $k = 0, 1, \dots, n - 1$. Then ω_k is an **nth root of unity**.

Lemma

If $\zeta^n = z$, then all solutions are given by

$$\zeta\omega_0, \zeta\omega_1, \dots, \zeta\omega_{n-1}$$

The Logarithm

- Recall that e^z is periodic, hence not injective

The Logarithm

- Recall that e^z is periodic, hence not injective
- Therefore a complete inverse function does not exist:

The Logarithm

- Recall that e^z is periodic, hence not injective
- Therefore a complete inverse function does not exist:
- It is reasonable to assign $\log 1 = 0$ and $\log 1 = 2\pi i$

Theorem

Let $S = \{z : -\pi < \operatorname{Im} z \leq \pi\}$. Then $\exp : S \rightarrow \mathbb{C} \setminus \{0\}$ in a bijective fashion. In particular, the line $y = \pi$ is mapped onto the negative real axis.

Theorem

Let $S = \{z : -\pi < \operatorname{Im} z \leq \pi\}$. Then $\exp : S \rightarrow \mathbb{C} \setminus \{0\}$ in a bijective fashion. In particular, the line $y = \pi$ is mapped onto the negative real axis.

This is the same idea used to force \sqrt{x} to be a function: restrict the domain.

Proof

Proof.

Suppose $w \neq 0$. Then $w = re^{i\theta}$ for some r, θ with $\theta \in (-\pi, \pi]$. Define $x = \ln r, y = \theta$ and let $z = x + iy$. Then $z \in S$ by construction and $e^z = w$ hence \exp is surjective.

Suppose $e^{z_1} = e^{z_2}$. Then $1 = e^{z_1 - z_2}$ but then $z_1 - z_2 = 2\pi ik$ for some integer k . Since $z_1, z_2 \in S$, $|z_1 - z_2| < 2\pi$ thus $k = 0$.



Definition

Definition

For $z = re^{i\theta}$ with $r > 0$

$$\log z = \ln|z| + i \arg z = \ln r + i\theta$$

Definition

Definition

For $z = re^{i\theta}$ with $r > 0$

$$\log z = \ln|z| + i \arg z = \ln r + i\theta$$

This function is not single valued since $\arg z$ is not single valued.

Definition

Definition

For $z = re^{i\theta}$ with $r > 0$

$$\log z = \ln|z| + i \arg z = \ln r + i\theta$$

This function is not single valued since $\arg z$ is not single valued.

Denote $S_\pi = \{z : -\pi < \operatorname{Im} z \leq \pi\}$. Then $\log : \mathbb{C} \setminus \{0\} \rightarrow S_\pi$ is the **principal branch** of the logarithm function.

Definition

Definition

For $z = re^{i\theta}$ with $r > 0$

$$\log z = \ln|z| + i \arg z = \ln r + i\theta$$

This function is not single valued since $\arg z$ is not single valued.

Denote $S_\pi = \{z : -\pi < \operatorname{Im} z \leq \pi\}$. Then $\log : \mathbb{C} \setminus \{0\} \rightarrow S_\pi$ is the **principal branch** of the logarithm function.

In general the range of \log can be any strip of width 2π . The choice of branch is up to the user but must be declared.

Properties of Logarithm

- On any strip of width 2π , \exp is the inverse of \log and vice versa.

Properties of Logarithm

- On any strip of width 2π , \exp is the inverse of \log and vice versa.
- $\log z$ is continuous on $\{z = re^{i\theta} : r > 0, -\pi < \theta < \pi\}$.

Properties of Logarithm

- On any strip of width 2π , \exp is the inverse of \log and vice versa.
- $\log z$ is continuous on $\{z = re^{i\theta} : r > 0, -\pi < \theta < \pi\}$.
- Every branch of $\log z$ is analytic: verify using Lemma 7.

Properties of Logarithm

- On any strip of width 2π , \exp is the inverse of \log and vice versa.
- $\log z$ is continuous on $\{z = re^{i\theta} : r > 0, -\pi < \theta < \pi\}$.
- Every branch of $\log z$ is analytic: verify using Lemma 7.
- $[\log z]' = 1/z$: verify using Chain Rule and inverse function relationship with $\exp z$.

Properties of Logarithm

- On any strip of width 2π , \exp is the inverse of \log and vice versa.
- $\log z$ is continuous on $\{z = re^{i\theta} : r > 0, -\pi < \theta < \pi\}$.
- Every branch of $\log z$ is analytic: verify using Lemma 7.
- $[\log z]' = 1/z$: verify using Chain Rule and inverse function relationship with $\exp z$.
- $\log z_1 z_2 = \log z_1 + \log z_2 + 2\pi i k$

Complex Trigonometry

- Recall $e^{i\theta} = \cos \theta + i \sin \theta$ so $e^{-i\theta} = \cos \theta - i \sin \theta$

Complex Trigonometry

- Recall $e^{i\theta} = \cos \theta + i \sin \theta$ so $e^{-i\theta} = \cos \theta - i \sin \theta$
- Solve this system for $\cos \theta, \sin \theta$

Complex Trigonometry

- Recall $e^{i\theta} = \cos \theta + i \sin \theta$ so $e^{-i\theta} = \cos \theta - i \sin \theta$
- Solve this system for $\cos \theta, \sin \theta$
- Replace $\theta \rightarrow z$ to define $\cos z, \sin z$

Complex Trigonometry

- Recall $e^{i\theta} = \cos \theta + i \sin \theta$ so $e^{-i\theta} = \cos \theta - i \sin \theta$
- Solve this system for $\cos \theta, \sin \theta$
- Replace $\theta \rightarrow z$ to define $\cos z, \sin z$
- Analytic automatically

Complex Trigonometry

- Recall $e^{i\theta} = \cos \theta + i \sin \theta$ so $e^{-i\theta} = \cos \theta - i \sin \theta$
- Solve this system for $\cos \theta, \sin \theta$
- Replace $\theta \rightarrow z$ to define $\cos z, \sin z$
- Analytic automatically
- Derivatives follow giving expected results

Complex Trigonometry

- Recall $e^{i\theta} = \cos \theta + i \sin \theta$ so $e^{-i\theta} = \cos \theta - i \sin \theta$
- Solve this system for $\cos \theta, \sin \theta$
- Replace $\theta \rightarrow z$ to define $\cos z, \sin z$
- Analytic automatically
- Derivatives follow giving expected results
- Remaining functions defined in usual way

General Complex Exponents

What is i^i ?

General Complex Exponents

What is i^i ?

- Define $z^w = \exp(w \log z)$

General Complex Exponents

What is i^i ?

- Define $z^w = \exp(w \log z)$
- Note: this is a multi-valued expression so does not represent a single number

General Complex Exponents

What is i^i ?

- Define $z^w = \exp(w \log z)$
- Note: this is a multi-valued expression so does not represent a single number
- Fact: $[z^w]' = wz^{w-1}$: PLTS

Analytic Function Construction

- So far: polynomials, Cauchy-Riemann equations

Analytic Function Construction

- So far: polynomials, Cauchy-Riemann equations
- Coming later: power series

Analytic Function Construction

- So far: polynomials, Cauchy-Riemann equations
- Coming later: power series
- Now showing: harmonic conjugates

Analytic Function Construction

- So far: polynomials, Cauchy-Riemann equations
- Coming later: power series
- Now showing: harmonic conjugates

Definition

Suppose that u is a real valued harmonic function defined on Ω . Then v is a **harmonic conjugate** if

$$f(z) = u(z) + iv(z)$$

is analytic on Ω .

Construction Process

Lemma

If Ω is simply connected and $z, z_0 \in \Omega$ then

$$\int_{z_0}^z -u_y \, dx + u_x \, dy$$

is path independent.

Construction Process

Lemma

If Ω is simply connected and $z, z_0 \in \Omega$ then

$$\int_{z_0}^z -u_y \, dx + u_x \, dy$$

is path independent.

- Suppose u is a real valued harmonic function defined on Ω

Construction Process

Lemma

If Ω is simply connected and $z, z_0 \in \Omega$ then

$$\int_{z_0}^z -u_y \, dx + u_x \, dy$$

is path independent.

- Suppose u is a real valued harmonic function defined on Ω
- Note: $dv = v_x \, dx + v_y \, dy$ and $v(z) - v(z_0) = \int_{z_0}^z dv$ is path independent

Construction Process

Lemma

If Ω is simply connected and $z, z_0 \in \Omega$ then

$$\int_{z_0}^z -u_y \, dx + u_x \, dy$$

is path independent.

- Suppose u is a real valued harmonic function defined on Ω
- Note: $dv = v_x \, dx + v_y \, dy$ and $v(z) - v(z_0) = \int_{z_0}^z dv$ is path independent
- Applying the previous lemma,

$$v(z) - v(z_0) = \int_{z_0}^z -u_y \, dx + u_x \, dy$$

where $v(z_0)$ is arbitrary (real).

Example

- $u(x, y) = x^2 - y^2$ is harmonic in \mathbb{C} (verify)

Example

- $u(x, y) = x^2 - y^2$ is harmonic in \mathbb{C} (verify)
- Note: $dv = 2y \, dx + 2x \, dy$

Example

- $u(x, y) = x^2 - y^2$ is harmonic in \mathbb{C} (verify)
- Note: $dv = 2y \, dx + 2x \, dy$
- By Figure 1, $\psi(x, y) = 2xy + c$ satisfies $d\psi = dv$

Example

- $u(x, y) = x^2 - y^2$ is harmonic in \mathbb{C} (verify)
- Note: $dv = 2y \, dx + 2x \, dy$
- By Figure 1, $\psi(x, y) = 2xy + c$ satisfies $d\psi = dv$
- Therefore, $v(x, y) = \psi(x, y)$ for $c \in \mathbb{R}$ is a harmonic conjugate of $u(x, y)$

Example

- $u(x, y) = x^2 - y^2$ is harmonic in \mathbb{C} (verify)
- Note: $dv = 2y \, dx + 2x \, dy$
- By Figure 1, $\psi(x, y) = 2xy + c$ satisfies $d\psi = dv$
- Therefore, $v(x, y) = \psi(x, y)$ for $c \in \mathbb{R}$ is a harmonic conjugate of $u(x, y)$
- Hence $f(z) = u(x, y) + iv(x, y) = z^2 + ic$ is analytic

Summary Theorem

Theorem

Given a real harmonic function $u(x, y)$ on a simply connected domain Ω , there exist infinitely many harmonic conjugates $v(x, y)$ each differing by an additive constant. Hence the given u is the real part of infinitely many analytic functions

$$f(z) = u(z) + iv(z)$$

on Ω each differing by an additive constant.

Summary Theorem

Theorem

Given a real harmonic function $u(x, y)$ on a simply connected domain Ω , there exist infinitely many harmonic conjugates $v(x, y)$ each differing by an additive constant. Hence the given u is the real part of infinitely many analytic functions

$$f(z) = u(z) + iv(z)$$

on Ω each differing by an additive constant.

Note that this implies $\ln z$ does not have a harmonic conjugate on $\mathbb{C} \setminus \{0\}$ yet is harmonic on this domain.

Integration

Objectives

- Complex line integrals

Objectives

- Complex line integrals
- Theorems about integration

Objectives

- Complex line integrals
- Theorems about integration
- Complex versions of previous theorems about harmonic functions

Basics

- Ω is a plane domain

Basics

- Ω is a plane domain
- Γ is a piecewise-smooth curve in Ω

Basics

- Ω is a plane domain
- Γ is a piecewise-smooth curve in Ω
- $f(z)$ is (usually) analytic

Basics

- Ω is a plane domain
- Γ is a piecewise-smooth curve in Ω
- $f(z)$ is (usually) analytic
- Question: What does

$$\int_{\Gamma} f(z) \, dz$$

mean and how to compute it?

Basics

- Ω is a plane domain
- Γ is a piecewise-smooth curve in Ω
- $f(z)$ is (usually) analytic
- Question: What does

$$\int_{\Gamma} f(z) \, dz$$

mean and how to compute it?

- Strategy: Separate into real and imaginary parts; use previous knowledge

Strategy In Action

- $f(z) = u(z) + iv(z)$ and $dz = dx + idy$

Strategy In Action

- $f(z) = u(z) + iv(z)$ and $dz = dx + idy$

$$\int_{\Gamma} f(z) dz = \int_{\Gamma} (u(z) dx - v(z) dy) + i \int_{\Gamma} (v(z) dx + u(z) dy)$$

Strategy In Action

- $f(z) = u(z) + iv(z)$ and $dz = dx + idy$

$$\int_{\Gamma} f(z) dz = \int_{\Gamma} (u(z) dx - v(z) dy) + i \int_{\Gamma} (v(z) dx + u(z) dy)$$

- Compute using parameterization (as usual)

Example

(get from another book)

Fundamental Theorem of Calculus

Theorem

Let $g(x)$ be a continuous real-valued function $a \leq x \leq b$. Define

$$G(x) = \int_a^x g(t) dt$$

Then

Fundamental Theorem of Calculus

Theorem

Let $g(x)$ be a continuous real-valued function $a \leq x \leq b$. Define

$$G(x) = \int_a^x g(t) dt$$

Then

- ① $G(x)$ is continuously differentiable on (a, b)

Fundamental Theorem of Calculus

Theorem

Let $g(x)$ be a continuous real-valued function $a \leq x \leq b$. Define

$$G(x) = \int_a^x g(t) dt$$

Then

- ① $G(x)$ is continuously differentiable on (a, b)
- ② $G'(x) = g(x)$

Fundamental Theorem of Calculus

Theorem

Let $g(x)$ be a continuous real-valued function $a \leq x \leq b$. Define

$$G(x) = \int_a^x g(t) dt$$

Then

- ① $G(x)$ is continuously differentiable on (a, b)
- ② $G'(x) = g(x)$
- ③ if also $F'(x) = g(x)$, then $F(x) - G(x)$ is constant and

$$\int_a^b g(t) dt = G(b) - G(a)$$

Theorem

Let $f(z)$ be continuous in a domain Ω and suppose that $F(z)$ is analytic with $F'(z) = f(z)$ for all $z \in \Omega$. If Γ is a curve from z_0 to z_1 in Ω , then

$$\int_{\Gamma} f(z) dz = \int_{\Gamma} F'(z) dz = F(z_1) - F(z_0) \quad (22)$$

Notes

- Writing $dF = F'(z)$, this is the complex version of Theorem 8

Notes

- Writing $dF = F'(z)$, this is the complex version of Theorem 8
- Need to know $f(z) = F'(z)$ for some analytic $F(z)$ beforehand

Notes

- Writing $dF = F'(z)$, this is the complex version of Theorem 8
- Need to know $f(z) = F'(z)$ for some analytic $F(z)$ beforehand
- The Cauchy Integral Theorem 30 will void the previous point

Crucial Example

- Compute $\int_{\Gamma} \frac{dz}{z}$ where Γ is the unit circle parameterized in the counter clockwise direction.

Crucial Example

- Compute $\int_{\Gamma} \frac{dz}{z}$ where Γ is the unit circle parameterized in the counter clockwise direction.
- Warning: this is not $\log z$ due to branch concerns

Crucial Example

- Compute $\int_{\Gamma} \frac{dz}{z}$ where Γ is the unit circle parameterized in the counter clockwise direction.
- Warning: this is not $\log z$ due to branch concerns
- Ignoring this warning would give a value of $\log z_1 - \log z_0 = 0$, a clear contradiction

Crucial Example

- Compute $\int_{\Gamma} \frac{dz}{z}$ where Γ is the unit circle parameterized in the counter clockwise direction.
- Warning: this is not $\log z$ due to branch concerns
- Ignoring this warning would give a value of $\log z_1 - \log z_0 = 0$, a clear contradiction
- Parameterize: $z = e^{i\theta}$ for $\theta \in [0, 2\pi]$ so that $dz = ie^{i\theta}d\theta$

Crucial Example

- Compute $\int_{\Gamma} \frac{dz}{z}$ where Γ is the unit circle parameterized in the counter clockwise direction.
- Warning: this is not $\log z$ due to branch concerns
- Ignoring this warning would give a value of $\log z_1 - \log z_0 = 0$, a clear contradiction
- Parameterize: $z = e^{i\theta}$ for $\theta \in [0, 2\pi]$ so that $dz = ie^{i\theta}d\theta$
- Manually compute (no theorems):

$$\int_{\Gamma} \frac{dz}{z} = \int_0^{2\pi} e^{-i\theta} (ie^{i\theta}) d\theta = 2\pi i$$

Theorem (ML Inequality)

Let $f(z)$ be continuous and defined on Γ . Suppose that $|f(z)| \leq M$ for all $z \in \Gamma$ and $L = \text{length } \Gamma$. Then

$$\left| \int_{\Gamma} f(z) dz \right| \leq \int_{\Gamma} |f(z)| |dz| = \int_{\Gamma} |f(z)| ds \leq ML \quad (23)$$

Integral Estimates

Theorem (ML Inequality)

Let $f(z)$ be continuous and defined on Γ . Suppose that $|f(z)| \leq M$ for all $z \in \Gamma$ and $L = \text{length } \Gamma$. Then

$$\left| \int_{\Gamma} f(z) dz \right| \leq \int_{\Gamma} |f(z)| |dz| = \int_{\Gamma} |f(z)| ds \leq ML \quad (23)$$

Common application: show some integral is zero.

Proof

Proof.

Denote $I = \int_{\Gamma} f(z) dz$. Then there exists ω such that $I = |I|e^{i\omega}$. Thus

$$|I| = \int_{\Gamma} e^{-i\omega} f(z) dz$$

can be parameterized by $z(t)$. The integrand can be written as $U(t) + iV(t)$ and all functions, $|I|$, U , V , are real. □

Cauchy Integral Theorem

Theorem

Let $f(z)$ be analytic in a domain Ω and let Γ be a closed Jordan curve inside Ω whose interior is contained in Ω so that $f(z)$ is analytic on and inside Γ . Then

$$\int_{\Gamma} f(z) \, dz = 0 \tag{24}$$

Proof

Proof.

Apply Green's Theorem 2 and the Cauchy-Riemann equations.

Examples

- $\int_{\Gamma} \cos(\sin z) dz$

For $\Gamma = C(0; 1)$.

Examples

- $\int_{\Gamma} \cos(\sin z) dz$
- $\int_{\Gamma} e^{z^2} dz$

For $\Gamma = C(0; 1)$.

Strong Cauchy Integral Theorem

Theorem

Let $f(z)$ be analytic in a domain Ω and let Γ be a closed curve in Ω which can be shrunk to a point within Ω . Then

$$\int_{\Gamma} f(z) \, dz = 0 \tag{25}$$

Corollary

Let $f(z)$ be analytic in Ω and Γ_1, Γ_2 be piecewise smooth curves from z_0 to z_1 all lying inside Ω such that all points between Γ_1, Γ_2 also lie in Ω . Then

$$\int_{\Gamma_1} f(z) dz = \int_{\Gamma_2} f(z) dz \quad (26)$$

Corollary

Let $f(z)$ be analytic in Ω and Γ_1, Γ_2 be piecewise smooth curves from z_0 to z_1 all lying inside Ω such that all points between Γ_1, Γ_2 also lie in Ω . Then

$$\int_{\Gamma_1} f(z) dz = \int_{\Gamma_2} f(z) dz \quad (26)$$

Proof.

Note that $\Gamma = \Gamma_1 + (-\Gamma_2)$ is a closed loop. □

Corollary

If $f(z)$ is analytic in the simply connected domain Ω and z, z_0 are points of Ω , then

$$F(z) = \int_{z_0}^z f(\zeta) d\zeta \quad (27)$$

is a value independent of the path from z_0 to z in Ω hence defines a continuous function $F = F(z)$ in Ω .

Corollary

If $f(z)$ is analytic in the simply connected domain Ω and z, z_0 are points of Ω , then

$$F(z) = \int_{z_0}^z f(\zeta) d\zeta \quad (27)$$

is a value independent of the path from z_0 to z in Ω hence defines a continuous function $F = F(z)$ in Ω .

- It is true that $F(z)$ is analytic and $F'(z) = f(z)$.

Corollary

If $f(z)$ is analytic in the simply connected domain Ω and z, z_0 are points of Ω , then

$$F(z) = \int_{z_0}^z f(\zeta) d\zeta \quad (27)$$

is a value independent of the path from z_0 to z in Ω hence defines a continuous function $F = F(z)$ in Ω .

- It is true that $F(z)$ is analytic and $F'(z) = f(z)$.
- A different choice of z_0 results in a different F ; how much?

Complex Series

Objectives

- Singularities

Objectives

- Singularities
- Laurent series

Questions

- Given $f(x) = x \sin(1/x)$, what value at 0 guarantees continuity of f ?

Questions

- Given $f(x) = x \sin(1/x)$, what value at 0 guarantees continuity of f ?
- Give $g(z)$ analytic on $\Omega \setminus \{z_0\}$, what value at z_0 guarantees analyticity on $\Omega \cup \{z_0\}$?

Questions

- Given $f(x) = x \sin(1/x)$, what value at 0 guarantees continuity of f ?
- Give $g(z)$ analytic on $\Omega \setminus \{z_0\}$, what value at z_0 guarantees analyticity on $\Omega \cup \{z_0\}$?
- Is this even possible??

Assumptions, Singularity, Notation

f is analytic on $\Omega' = \Omega \setminus \{z_0\}$

Assumptions, Singularity, Notation

f is analytic on $\Omega' = \Omega \setminus \{z_0\}$

Definition

If f is analytic on Ω' and z_0 is an interior point of Ω , then z_0 is an isolated singularity.

Types of Singularities

Definition

If f is analytic on Ω' and there exists w_0 such that $\lim_{z \rightarrow z_0} f(z) = w_0$, then z_0 is a **removable singularity**.

Types of Singularities

Definition

If f is analytic on Ω' and there exists w_0 such that $\lim_{z \rightarrow z_0} f(z) = w_0$, then z_0 is a **removable singularity**.

Definition

If f is analytic on Ω' and $\lim_{z \rightarrow z_0} |f(z)| = \infty$, then z_0 is a **pole**.

Types of Singularities

Definition

If f is analytic on Ω' and there exists w_0 such that $\lim_{z \rightarrow z_0} f(z) = w_0$, then z_0 is a **removable singularity**.

Definition

If f is analytic on Ω' and $\lim_{z \rightarrow z_0} |f(z)| = \infty$, then z_0 is a **pole**.

Definition

If f is analytic on Ω' and z_0 is a singular point which is not removable and not a pole, then z_0 is an **essential singularity**.

Examples

- If f is analytic on Ω , then $\lim_{z \rightarrow z_0} f(z) = w_0$ for all z_0 ; puncture and replace appropriately

Examples

- If f is analytic on Ω , then $\lim_{z \rightarrow z_0} f(z) = w_0$ for all z_0 ; puncture and replace appropriately
- Let $g(z) = \sin(z)/z$ on $\mathbb{C} \setminus \{0\}$; define $g(0) = 1$.

Examples

- If f is analytic on Ω , then $\lim_{z \rightarrow z_0} f(z) = w_0$ for all z_0 ; puncture and replace appropriately
- Let $g(z) = \sin(z)/z$ on $\mathbb{C} \setminus \{0\}$; define $g(0) = 1$.
- $1/z, 1/z^2, z - 1$ all have poles

Examples

- If f is analytic on Ω , then $\lim_{z \rightarrow z_0} f(z) = w_0$ for all z_0 ; puncture and replace appropriately
- Let $g(z) = \sin(z)/z$ on $\mathbb{C} \setminus \{0\}$; define $g(0) = 1$.
- $1/z, 1/z^2, z - 1$ all have poles
- $h(z) = e^{1/z}$ on $\mathbb{C} \setminus \{0\}$

Examples

- If f is analytic on Ω , then $\lim_{z \rightarrow z_0} f(z) = w_0$ for all z_0 ; puncture and replace appropriately
- Let $g(z) = \sin(z)/z$ on $\mathbb{C} \setminus \{0\}$; define $g(0) = 1$.
- $1/z, 1/z^2, z - 1$ all have poles
- $h(z) = e^{1/z}$ on $\mathbb{C} \setminus \{0\}$
- Compute $\lim_{z \rightarrow 0^+} h(z)$ and $\lim_{z \rightarrow 0^-} h(z)$

Theorem

Theorem

Let $f(z)$ be analytic on Ω' where z_0 is an interior point of Ω which is a removable singularity. Then there exists w_0 such that defining $f(z_0) = w_0$ guarantees f is analytic at z_0 , hence on all of Ω .

Theorem

Theorem

Let $f(z)$ be analytic on Ω' where z_0 is an interior point of Ω which is a removable singularity. Then there exists w_0 such that defining $f(z_0) = w_0$ guarantees f is analytic at z_0 , hence on all of Ω .

Note that there is no mention of integrals.

Theorem

Theorem

Let $f(z)$ be analytic on Ω' where z_0 is an interior point of Ω which is a removable singularity. Then there exists w_0 such that defining $f(z_0) = w_0$ guarantees f is analytic at z_0 , hence on all of Ω .

Note that there is no mention of integrals. Warning: the proof is long.
Also z, z_0, ρ are constants.

Proof

Let \hat{f} be the extension of f . Clearly \hat{f} is continuous on Ω . Choose $C(z_0; R) \subset \Omega'$ and define

$$F(z) = \frac{1}{2\pi i} \int_{C(z_0; R)} \frac{f(\zeta)}{\zeta - z} d\zeta$$

for $|z - z_0| < R$.

Claim: F is analytic at z_0 . This is an integral of Cauchy type, hence differentiation under the integral sign is permitted thus $F'(z)$ exists.

Claim: $F = f$ on $0 < |z - z_0| < R$. Suppose z_1 satisfies $0 < |z_1 - z_0| < R$ and consider

$$F(z_1) - f(z_1) = \frac{1}{2\pi i} \int_{C(z_0; R)} \frac{f(\zeta)}{\zeta - z_1} d\zeta - \frac{1}{2\pi i} \int_{C(z_1; r_1)} \frac{f(\zeta)}{\zeta - z_1} d\zeta$$

Proof Continued

Now compute

$$\frac{1}{2\pi i} \int_{C(z_0; \rho)} \frac{f(\zeta)}{\zeta - z_1} d\zeta$$

$\frac{f(\zeta)}{\zeta - z_1}$ is analytic on $D(z_0; R) \setminus (\bar{D}(z_1; r_1) \cup \bar{D}(z_0; \rho))$ and using the oriented boundary circles:

$$\int_{C(z_0; R)} - \int_{C(z_1; r_1)} - \int_{C(z_0; \rho)} = 0$$

Therefore $\frac{1}{2\pi i} \int_{C(z_0; \rho)} \frac{f(\zeta)}{\zeta - z_1} d\zeta = F(z_1) - f(z_1)$

Claim: $\left| \frac{f(\zeta)}{\zeta - z_1} \right| < M$ for some M . For $|\zeta - z_0| = \rho$, $|f(\zeta)|$ is bounded since $\lim_{\rho \rightarrow 0} f(\zeta) = w_0$ is finite. Furthermore, $|\zeta - z_1| \geq |z_1 - z_0| - \rho$ because ζ is 'far away' from z_1 (See graphic). Applying Theorem 29 to $|F(z_1) - f(z_1)|$ gives $|F(z_1) - f(z_1)| < M\rho$ for all sufficiently small ρ .

Proof Illustrations




Figure: Desmos

Key Proof Ideas

- Extending a function

Key Proof Ideas

- Extending a function
- Use of true circles (not arbitrary curves)

Singularities by Series

- Note that f is **not** analytic at z_0 by assumption and definition, hence does not have a Taylor series at z_0 .

Singularities by Series

- Note that f is **not** analytic at z_0 by assumption and definition, hence does not have a Taylor series at z_0 .
- Thus any series must have terms $b_k(z - z_0)^{-k}$

Singularities by Series

- Note that f is **not** analytic at z_0 by assumption and definition, hence does not have a Taylor series at z_0 .
- Thus any series must have terms $b_k(z - z_0)^{-k}$
- What is the radius of convergence (if this concept is even reasonable anymore)??

Convergence

- Consider $\sum b_k(z - z_0)^{-k} = \sum b_k \zeta^k$ for $\zeta = (z - z_0)^{-1}$

Convergence

- Consider $\sum b_k(z - z_0)^{-k} = \sum b_k \zeta^k$ for $\zeta = (z - z_0)^{-1}$
- This converges for some $|\zeta| < R$

Convergence

- Consider $\sum b_k(z - z_0)^{-k} = \sum b_k \zeta^k$ for $\zeta = (z - z_0)^{-1}$
- This converges for some $|\zeta| < R$
- Hence the original series converges for $|z - z_0| > 1/R$

Convergence

- Consider $\sum b_k(z - z_0)^{-k} = \sum b_k \zeta^k$ for $\zeta = (z - z_0)^{-1}$
- This converges for some $|\zeta| < R$
- Hence the original series converges for $|z - z_0| > 1/R$
- This is convergence *outside* some disc!

Notation



$$\sum_{n=-\infty}^{\infty} c_n(z - z_0)^n$$

Notation



$$\sum_{n=-\infty}^{\infty} c_n(z - z_0)^n$$



$$\sum_{n=0}^{\infty} a_n(z - z_0)^n + \sum_{n=1}^{\infty} \frac{b_n}{(z - z_0)^n}$$

The Laurent Expansion

- Assume f is analytic on $R_1 < |z - z_0| < R_2$

The Laurent Expansion

- Assume f is analytic on $R_1 < |z - z_0| < R_2$
- If $R_1 = 0$, this represents a *punctured* disc

The Laurent Expansion

- Assume f is analytic on $R_1 < |z - z_0| < R_2$
- If $R_1 = 0$, this represents a *punctured* disc

Theorem

Suppose f is analytic in the annulus $R_1 < |z - z_0| < R_2$. Then $f(z)$ is represented by a Laurent series

$$f(z) = \sum_{n=-\infty}^{\infty} c_n(z - z_0)^n$$

which converges to f throughout the annulus. Moreover

$$c_n = \frac{1}{2\pi i} \int_C \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta$$

where C is any circle centered at z_0 contained in the annulus.

Comments

- If f is analytic for $|z - z_0| < R_2$, then $c_{-n} = 0$ for $n = 1, 2, \dots$ and f has a Taylor series. Why?

Comments

- If f is analytic for $|z - z_0| < R_2$, then $c_{-n} = 0$ for $n = 1, 2, \dots$ and f has a Taylor series. Why?
- All c_n are independent of choice of C . Why?

Comments

- If f is analytic for $|z - z_0| < R_2$, then $c_{-n} = 0$ for $n = 1, 2, \dots$ and f has a Taylor series. Why?
- All c_n are independent of choice of C . Why?
- c_n not usually computed using this formula.

- The interior of the circles C_i are not completely contained within the annulus - on purpose.

- The interior of the circles C_i are not completely contained within the annulus - on purpose.
- It may be useful to denote the quantity $\frac{\zeta_1 - z_0}{z - z_0}$ as r

Proof Notes

- The interior of the circles C_i are not completely contained within the annulus - on purpose.
- It may be useful to denote the quantity $\frac{\zeta_1 - z_0}{z - z_0}$ as r
- Multiply by $-\frac{f(\zeta_1)}{2\pi i}$ and integrate

Examples

- $1/z$ is the Laurent series of $1/z$ at 0

Examples

- $1/z$ is the Laurent series of $1/z$ at 0
- $1/z$ is analytic at $z = 1$, hence $c_{-n} = 0$

Examples

- $1/z$ is the Laurent series of $1/z$ at 0
- $1/z$ is analytic at $z = 1$, hence $c_{-n} = 0$
- $e^{1/z}$ at $z = 0$

Examples

- $1/z$ is the Laurent series of $1/z$ at 0
- $1/z$ is analytic at $z = 1$, hence $c_{-n} = 0$
- $e^{1/z}$ at $z = 0$
- $1/(z - 1)$ at $z = 0$ in powers of $1/z$; try evaluating at 0

Examples

- $1/z$ is the Laurent series of $1/z$ at 0
- $1/z$ is analytic at $z = 1$, hence $c_{-n} = 0$
- $e^{1/z}$ at $z = 0$
- $1/(z - 1)$ at $z = 0$ in powers of $1/z$; try evaluating at 0
- $1/(z^2 + 1)$ at $z = i$

Lemma

Suppose f is analytic and has a pole at z_0 . Then $g(z) = 1/f(z)$ is analytic at z_0 which is an isolated zero. Conversely, if z_0 is an isolated zero of an analytic function f , then $g(z) = 1/f(z)$ has a pole at z_0 .

Lemma

Suppose f is analytic and has a pole at z_0 . Then $g(z) = 1/f(z)$ is analytic at z_0 which an isolated zero. Conversely, if z_0 is an isolated zero of an analytic function f , then $g(z) = 1/f(z)$ has a pole at z_0 .

Proof.

Consider $|f|$ and definitions. □

Order

- Use the same hypotheses as Lemma; $g(z)$ has isolated zero

Order

- Use the same hypotheses as Lemma; $g(z)$ has isolated zero
- $g(z) = (z - z_0)^m g_1(z)$ where $g_1(z_0) \neq 0$ is analytic in $V_\delta(z_0)$

Order

- Use the same hypotheses as Lemma; $g(z)$ has isolated zero
- $g(z) = (z - z_0)^m g_1(z)$ where $g_1(z_0) \neq 0$ is analytic in $V_\delta(z_0)$
- Then

$$f(z) = \frac{f_1(z)}{(z - z_0)^m}$$

has a **pole of order m at z_0**

Order

- Use the same hypotheses as Lemma; $g(z)$ has isolated zero
- $g(z) = (z - z_0)^m g_1(z)$ where $g_1(z_0) \neq 0$ is analytic in $V_\delta(z_0)$
- Then

$$f(z) = \frac{f_1(z)}{(z - z_0)^m}$$

has a **pole of order m at z_0**

- If $m = 1$, z_0 is a *simple pole*

Terminology

- Suppose z_0 is an isolated singularity of the analytic function f

Terminology

- Suppose z_0 is an isolated singularity of the analytic function f
- Consider $0 < |z - z_0| < R_2$

Terminology

- Suppose z_0 is an isolated singularity of the analytic function f
- Consider $0 < |z - z_0| < R_2$
- Write

$$f(z) = \sum_{n=-\infty}^{\infty} c_n(z - z_0)^n$$

Terminology

- Suppose z_0 is an isolated singularity of the analytic function f
- Consider $0 < |z - z_0| < R_2$
- Write

$$f(z) = \sum_{n=-\infty}^{\infty} c_n(z - z_0)^n$$

- The **polar part** is

$$\sum_{n=-1}^{\infty} c_n(z - z_0)^n = \sum_{n=1}^{\infty} c_{-n}(z - z_0)^{-n}$$

Classification of Singularities

Theorem

Suppose z_0 is an isolated singularity of the analytic function f which has polar part $\sum_{n=1}^{\infty} c_{-n}(z - z_0)^{-n} = \sum_{n=1}^{\infty} \frac{c_{-n}}{(z - z_0)^n}$. Then

Classification of Singularities

Theorem

Suppose z_0 is an isolated singularity of the analytic function f which has polar part $\sum_{n=1}^{\infty} c_{-n}(z - z_0)^{-n} = \sum_{n=1}^{\infty} \frac{c_{-n}}{(z - z_0)^n}$. Then

- ① z_0 is a removable singularity if and only if $c_{-n} = 0$ for all $n \geq 1$

Classification of Singularities

Theorem

Suppose z_0 is an isolated singularity of the analytic function f which has polar part $\sum_{n=1}^{\infty} c_{-n}(z - z_0)^{-n} = \sum_{n=1}^{\infty} \frac{c_{-n}}{(z - z_0)^n}$. Then

- ① z_0 is a removable singularity if and only if $c_{-n} = 0$ for all $n \geq 1$
- ② z_0 is a pole of order h if and only if the polar part is a sum $\sum_{n=1}^h c_{-n}(z - z_0)^{-n}$ where $c_{-h} \neq 0$

Classification of Singularities

Theorem

Suppose z_0 is an isolated singularity of the analytic function f which has polar part $\sum_{n=1}^{\infty} c_{-n}(z - z_0)^{-n} = \sum_{n=1}^{\infty} \frac{c_{-n}}{(z - z_0)^n}$. Then

- ① z_0 is a removable singularity if and only if $c_{-n} = 0$ for all $n \geq 1$
- ② z_0 is a pole of order h if and only if the polar part is a sum $\sum_{n=1}^h c_{-n}(z - z_0)^{-n}$ where $c_{-h} \neq 0$
- ③ z_0 is an essential singularity if and only if the polar part has infinitely many nonzero terms.

Examples Revisited

- $\sin(z)/z$ has a Taylor series

Examples Revisited

- $\sin(z)/z$ has a Taylor series
- $1/z$ has a (simple) pole

Examples Revisited

- $\sin(z)/z$ has a Taylor series
- $1/z$ has a (simple) pole
- $e^{1/z}$ has infinitely many nonzero terms in the polar part

Compactification

- Consider the set $\bar{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$; the Riemann sphere

Compactification

- Consider the set $\bar{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$; the Riemann sphere
- If f has a pole at z_0 , then $\lim_{z \rightarrow z_0} f(z) = \infty$

Compactification

- Consider the set $\bar{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$; the Riemann sphere
- If f has a pole at z_0 , then $\lim_{z \rightarrow z_0} f(z) = \infty$
- Expand the domain of f to $\bar{\mathbb{C}}$ and define $f(z_0) = \infty$

Compactification

- Consider the set $\bar{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$; the Riemann sphere
- If f has a pole at z_0 , then $\lim_{z \rightarrow z_0} f(z) = \infty$
- Expand the domain of f to $\bar{\mathbb{C}}$ and define $f(z_0) = \infty$
- Then f is continuous at z_0 !

Essential Singularities

Recall: z_0 an isolated singularity of analytic f is removable if and only if $|f(z)|$ is bounded in $V_\delta(z_0) \setminus \{z_0\}$.

Essential Singularities

Recall: z_0 an isolated singularity of analytic f is removable if and only if $|f(z)|$ is bounded in $V_\delta(z_0) \setminus \{z_0\}$.

Theorem (Cassorati-Weierstrass)

In every open neighborhood of an isolated essential singularity z_0 , the function f assumes values $f(z)$ arbitrarily close to every complex number w . That is, if D' is any punctured disc centered at z_0 and $D(w; \rho)$ is any disc about any point w , then there exists $z \in D'$ such that $f(z) \in D(w; \rho)$.

Application

Recall the surjective nature of \exp on horizontal strips of height 2π :

Theorem 25.

Consider $f(z) = e^{1/z}$. Let $w \in \mathbb{C} \setminus \{0\}$. Claim: for all $\delta > 0$, there exists $z_0 \in V = V_\delta(0) \setminus \{0\}$ such that $f(z_0) = w$. Note that $z \mapsto 1/z$ maps V to the set $Y = V_{1/\delta}(0) \setminus \{0\}^C$. Thus there exists $X \subset Y$ with “height” 2π . Since $\exp : X \rightarrow \mathbb{C} \setminus \{0\}$ is surjective, there exists $\zeta \in X$ with $e^\zeta = w$ so let $\zeta = 1/z$.

Calculus of Residues

Objectives

- Residue Theorem

Objectives

- Residue Theorem
- Argument Principle

Definition

Recall that $\int_{\Gamma} f(z) dz = 0$ if f is analytic on and inside Γ (simple, closed)

Definition

Recall that $\int_{\Gamma} f(z) dz = 0$ if f is analytic on and inside Γ (simple, closed)

Definition

Suppose f has an isolated singularity at z_0 . Then

$f(z) = \sum_{n=-\infty}^{\infty} c_n(z - z_0)^n$ is analytic on $0 < |z - z_0| < R_2$. The value c_{-1} is the **residue** of f at z_0 ; $c_{-1} = \text{Res}(f; z_0)$.

Definition

Recall that $\int_{\Gamma} f(z) dz = 0$ if f is analytic on and inside Γ (simple, closed)

Definition

Suppose f has an isolated singularity at z_0 . Then

$f(z) = \sum_{n=-\infty}^{\infty} c_n(z - z_0)^n$ is analytic on $0 < |z - z_0| < R_2$. The value c_{-1} is the **residue** of f at z_0 ; $c_{-1} = \text{Res}(f; z_0)$.

Now recall $c_n = \frac{1}{2\pi i} \int_C \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta$

Definition

Recall that $\int_{\Gamma} f(z) dz = 0$ if f is analytic on and inside Γ (simple, closed)

Definition

Suppose f has an isolated singularity at z_0 . Then

$f(z) = \sum_{n=-\infty}^{\infty} c_n(z - z_0)^n$ is analytic on $0 < |z - z_0| < R_2$. The value c_{-1} is the **residue** of f at z_0 ; $c_{-1} = \text{Res}(f; z_0)$.

Now recall $c_n = \frac{1}{2\pi i} \int_C \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta$

Choosing $n = -1$ and C to be a circle around z_0 containing no other singularities of f :

$$c_{-1} = \text{Res}(f; z_0) = \int_C f(\zeta) d\zeta$$

Theorem

The natural extension to a finite number of singularities is:

Theorem

The natural extension to a finite number of singularities is:

Theorem (Residue Theorem)

Suppose $f(z)$ is analytic in a domain Ω except for isolated singularities at z_1, \dots, z_m in Ω . Let Γ be a positively oriented simple closed curve in Ω which has all z_i in its interior. Then

$$\int_{\Gamma} f(z) dz = 2\pi i \sum_{i=1}^m \text{Res}(f; z_i)$$

Examples

- $\int_{\Gamma} dz/z = 2\pi i(1)$

Examples

- $\int_{\Gamma} dz/z = 2\pi i(1)$
- $\int_{\Gamma} (1/z + 1/(1-z)) dz$ where $\{0, 1\} \in \Gamma^{\circ}$:
 $\text{Res}(1/z; (0)) = 1, \text{Res}(1/(1-z); 1) = -1$

Examples

- $\int_{\Gamma} dz/z = 2\pi i(1)$
- $\int_{\Gamma} (1/z + 1/(1-z)) dz$ where $\{0, 1\} \in \Gamma^{\circ}$:
 $\text{Res}(1/z; (0)) = 1, \text{Res}(1/(1-z); 1) = -1$

Observation: Cauchy's Integral Formula is a special case of the Residue Theorem.

Example 1

Consider $\int_{|z|=2} \frac{e^z}{z^2(z-1)} dz$

Example 1

Consider $\int_{|z|=2} \frac{e^z}{z^2(z-1)} dz$




Figure: Desmos

- Recall $\frac{d}{dz} [f(z)]|_{z=\zeta} = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(z)}{(z-\zeta)^2} dz$

- Recall $\frac{d}{dz} [f(z)]|_{z=\zeta} = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(z)}{(z-\zeta)^2} dz$

- Hence

$$\int_{\Gamma_0} \frac{1}{z^2} \frac{e^z}{z-1} dz = \int_{\Gamma_0} \frac{e^z/(z-1)}{z^2} dz$$

- Recall $\frac{d}{dz} [f(z)] \Big|_{z=\zeta} = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(z)}{(z-\zeta)^2} dz$

- Hence

$$\int_{\Gamma_0} \frac{1}{z^2} \frac{e^z}{z-1} dz = \int_{\Gamma_0} \frac{e^z/(z-1)}{z^2} dz$$

- Let $\zeta = 0$, $f(z) = e^z/(z-1)$ and compute the value $-4\pi i$

- Recall $\frac{d}{dz} [f(z)]|_{z=\zeta} = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(z)}{(z-\zeta)^2} dz$

- Hence

$$\int_{\Gamma_0} \frac{1}{z^2} \frac{e^z}{z-1} dz = \int_{\Gamma_0} \frac{e^z/(z-1)}{z^2} dz$$

- Let $\zeta = 0$, $f(z) = e^z/(z-1)$ and compute the value $-4\pi i$

- Similarly

$$\int_{\Gamma_1} \frac{1}{z^2} \frac{e^z}{z-1} dz = \int_{\Gamma_1} \frac{e^z/z^2}{z-1} dz$$

- Recall $\frac{d}{dz} [f(z)]|_{z=\zeta} = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(z)}{(z-\zeta)^2} dz$

- Hence

$$\int_{\Gamma_0} \frac{1}{z^2} \frac{e^z}{z-1} dz = \int_{\Gamma_0} \frac{e^z/(z-1)}{z^2} dz$$

- Let $\zeta = 0, f(z) = e^z/(z-1)$ and compute the value $-4\pi i$
- Similarly

$$\int_{\Gamma_1} \frac{1}{z^2} \frac{e^z}{z-1} dz = \int_{\Gamma_1} \frac{e^z/z^2}{z-1} dz$$

- Let $\zeta = 1, f(z) = e^z/z^2$ and compute the value $2\pi ie$

- Recall $\frac{d}{dz} [f(z)]|_{z=\zeta} = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(z)}{(z-\zeta)^2} dz$

- Hence

$$\int_{\Gamma_0} \frac{1}{z^2} \frac{e^z}{z-1} dz = \int_{\Gamma_0} \frac{e^z/(z-1)}{z^2} dz$$

- Let $\zeta = 0, f(z) = e^z/(z-1)$ and compute the value $-4\pi i$
- Similarly

$$\int_{\Gamma_1} \frac{1}{z^2} \frac{e^z}{z-1} dz = \int_{\Gamma_1} \frac{e^z/z^2}{z-1} dz$$

- Let $\zeta = 1, f(z) = e^z/z^2$ and compute the value $2\pi ie$
- Hence the final value: $2\pi i(e-2)$

Example 2

- $\int_{|z|=1} \cot(z) dz = \int_{|z|=1} \frac{\cos z}{\sin z} dz$

Example 2

- $\int_{|z|=1} \cot(z) dz = \int_{|z|=1} \frac{\cos z}{\sin z} dz$
- Recall that $\sin z = zg(z)$ where $g(0) = 1$ by Taylor series

Example 2

- $\int_{|z|=1} \cot(z) dz = \int_{|z|=1} \frac{\cos z}{\sin z} dz$
- Recall that $\sin z = zg(z)$ where $g(0) = 1$ by Taylor series
- $\int_{|z|=1} \frac{\cos z}{\sin z} dz = \int_{|z|=1} \frac{\cos z/g(z)}{z} dz = 2\pi i \left. \frac{\cos z}{z} \right|_{z=0} = 2\pi i$

Summary

- If the denominator includes $(z - z_0)^k$, Cauchy Integral Formula probably works

Summary

- If the denominator includes $(z - z_0)^k$, Cauchy Integral Formula probably works
- If not, use residues

Techniques

- Laurent Series

Techniques

- Laurent Series
- Simple poles

Examples A

- $f(z) = e^{1/z} = 1 + 1/z + 1/z^2 + \dots$

Examples A

- $f(z) = e^{1/z} = 1 + 1/z + 1/z^2 + \dots$
- $g(z) = \sin(1/z^2) = 1/z^2 - 1/(3!z^6) + 1/(5!z^{10}) - \dots$

Examples B

Consider

$$f(z) = \frac{(z - i)^3(z + 1)}{z(z - 2)^4}$$

Examples B

Consider

$$f(z) = \frac{(z - i)^3(z + 1)}{z(z - 2)^4}$$

- Note that $z_0 \in \{0, 2\}$

Examples B

Consider

$$f(z) = \frac{(z-i)^3(z+1)}{z(z-2)^4}$$

- Note that $z_0 \in \{0, 2\}$
- Observe $zf(z)$ has no pole at 0

Examples B

Consider

$$f(z) = \frac{(z-i)^3(z+1)}{z(z-2)^4}$$

- Note that $z_0 \in \{0, 2\}$
- Observe $zf(z)$ has no pole at 0
- $\text{Res}(f; 0) = \lim_{z \rightarrow 0} zf(z) = zf(z)|_{z=0} = i/16$

Examples B

Consider

$$f(z) = \frac{(z-i)^3(z+1)}{z(z-2)^4}$$

- Note that $z_0 \in \{0, 2\}$
- Observe $zf(z)$ has no pole at 0
- $\text{Res}(f; 0) = \lim_{z \rightarrow 0} zf(z) = zf(z)|_{z=0} = i/16$
- Therefore $\int_{\Gamma} f(z) dz = -\pi/8$ provided $2 \notin \Gamma^o$

Double Pole Extension

- Suppose f has a pole of order 2 at z_0

Double Pole Extension

- Suppose f has a pole of order 2 at z_0
- Then

$$f(z) = \frac{c_{-2}}{(z - z_0)^2} + \frac{c_{-1}}{z - z_0} + c_0 + c_1(z - z_0) + \dots$$

Double Pole Extension

- Suppose f has a pole of order 2 at z_0
- Then

$$f(z) = \frac{c_{-2}}{(z - z_0)^2} + \frac{c_{-1}}{z - z_0} + c_0 + c_1(z - z_0) + \dots$$

- Observe that $(z - z_0)^2 f(z) = c_{-2} + c_{-1}(z - z_0) + c_0(z - z_0)^2 + \dots$

Double Pole Extension

- Suppose f has a pole of order 2 at z_0
- Then

$$f(z) = \frac{c_{-2}}{(z - z_0)^2} + \frac{c_{-1}}{z - z_0} + c_0 + c_1(z - z_0) + \dots$$

- Observe that $(z - z_0)^2 f(z) = c_{-2} + c_{-1}(z - z_0) + c_0(z - z_0)^2 + \dots$
- Therefore

$$c_{-1} = \lim_{z \rightarrow z_0} [(z - z_0)^2 f(z)]' = \text{Res}(f; z_0)$$

Double Pole Extension

- Suppose f has a pole of order 2 at z_0
- Then

$$f(z) = \frac{c_{-2}}{(z - z_0)^2} + \frac{c_{-1}}{z - z_0} + c_0 + c_1(z - z_0) + \dots$$

- Observe that $(z - z_0)^2 f(z) = c_{-2} + c_{-1}(z - z_0) + c_0(z - z_0)^2 + \dots$
- Therefore

$$c_{-1} = \lim_{z \rightarrow z_0} [(z - z_0)^2 f(z)]' = \text{Res}(f; z_0)$$

Can you find the pattern for a pole of order m ?

Example

Consider $f(z) = \frac{z+1}{z(z-i)^2}$ at $z = i, 0$

Example

Consider $f(z) = \frac{z+1}{z(z-i)^2}$ at $z = i, 0$

- $(z - i)^2 f(z) = 1 + 1/z$

Example

Consider $f(z) = \frac{z+1}{z(z-i)^2}$ at $z = i, 0$

- $(z - i)^2 f(z) = 1 + 1/z$
- $\frac{d}{dz} [1 + 1/z] = -1/z^2$

Example

Consider $f(z) = \frac{z+1}{z(z-i)^2}$ at $z = i, 0$

- $(z - i)^2 f(z) = 1 + 1/z$
- $\frac{d}{dz} [1 + 1/z] = -1/z^2$
- $-1/z^2|_{z=i} = 1 = \text{Res}(f; i)$

Example

Consider $f(z) = \frac{z+1}{z(z-i)^2}$ at $z = i, 0$

- $(z - i)^2 f(z) = 1 + 1/z$
- $\frac{d}{dz} [1 + 1/z] = -1/z^2$
- $-1/z^2|_{z=i} = 1 = \text{Res}(f; i)$
- $zf(z) = (z + 1)/(z - i)^2$

Example

Consider $f(z) = \frac{z+1}{z(z-i)^2}$ at $z = i, 0$

- $(z - i)^2 f(z) = 1 + 1/z$
- $\frac{d}{dz} [1 + 1/z] = -1/z^2$
- $-1/z^2|_{z=i} = 1 = \text{Res}(f; i)$
- $zf(z) = (z + 1)/(z - i)^2$
- $\lim_{z \rightarrow 0} (z + 1)/(z - i)^2 = -1 = \text{Res}(f; 0)$

Example Summary

Hence there are four possibilities for the value $\int_{\Gamma} f(z) dz$:

$$\int_{\Gamma} f(z) dz = \begin{cases} 0 & \text{no pole is in } \Gamma^\circ \\ 2\pi i & i \in \Gamma^\circ \text{ and } 0 \notin \Gamma^\circ \\ -2\pi i & 0 \in \Gamma^\circ \text{ and } i \notin \Gamma^\circ \\ 0 & 0, i \in \Gamma^\circ \end{cases}$$

Assumptions

- Ω - plane domain

Assumptions

- Ω - plane domain
- Γ - piecewise smooth, positively oriented, simple, closed, in Ω ; $\Gamma^\circ \subseteq \Omega$

Assumptions

- Ω - plane domain
- Γ - piecewise smooth, positively oriented, simple, closed, in Ω ; $\Gamma^\circ \subseteq \Omega$
- f - mostly analytic (finite number of poles) on Ω and NO poles on Γ

Assumptions

- Ω - plane domain
- Γ - piecewise smooth, positively oriented, simple, closed, in Ω ; $\Gamma^\circ \subseteq \Omega$
- f - mostly analytic (finite number of poles) on Ω and NO poles on Γ
- f has no zeros on Γ

Three-ish Numbers

① $\frac{1}{2\pi i} \int_{\Gamma} \frac{f'(z)}{f(z)} dz$

Three-ish Numbers

- ① $\frac{1}{2\pi i} \int_{\Gamma} \frac{f'(z)}{f(z)} dz$
- ② (number of zeros - the number of poles) of f inside Γ

Three-ish Numbers

- ① $\frac{1}{2\pi i} \int_{\Gamma} \frac{f'(z)}{f(z)} dz$
- ② (number of zeros - the number of poles) of f inside Γ

Note that the second must be an integer and the order matters.

Notation and Note

- $N_0(\Gamma)$ - the number of zeros of f inside Γ

Notation and Note

- $N_0(\Gamma)$ - the number of zeros of f inside Γ
- $N_\infty(\Gamma)$ - the number of poles of f inside Γ

Notation and Note

- $N_0(\Gamma)$ - the number of zeros of f inside Γ
- $N_\infty(\Gamma)$ - the number of poles of f inside Γ

Recall that counting is hard! Using the order (of poles and zeros) is the proper way to count these items.

Example

Consider $f(z) = \frac{(z-1/3)^2}{z^5(z-3)^4}$.

Example

Consider $f(z) = \frac{(z-1/3)^2}{z^5(z-3)^4}$.

- A zero of order 2 at $1/3$

Example

Consider $f(z) = \frac{(z-1/3)^2}{z^5(z-3)^4}$.

- A zero of order 2 at $1/3$
- A pole of order 5 at 0

Example

Consider $f(z) = \frac{(z-1/3)^2}{z^5(z-3)^4}$.

- A zero of order 2 at $1/3$
- A pole of order 5 at 0
- A pole of order 4 at 3

Example

Consider $f(z) = \frac{(z-1/3)^2}{z^5(z-3)^4}$.

- A zero of order 2 at $1/3$
- A pole of order 5 at 0
- A pole of order 4 at 3
- For $\Gamma = |z| = 1$, $N_0(\Gamma) = 2$, $N_\infty(\Gamma) = 5$

Example

Consider $f(z) = \frac{(z-1/3)^2}{z^5(z-3)^4}$.

- A zero of order 2 at $1/3$
- A pole of order 5 at 0
- A pole of order 4 at 3
- For $\Gamma = |z| = 1$, $N_0(\Gamma) = 2$, $N_\infty(\Gamma) = 5$
- For $\Gamma = |z| = 4$, $N_0(\Gamma) = 2$, $N_\infty(\Gamma) = 9$

Lemma

Let Γ be a piecewise smooth positively oriented simple closed curve and f an analytic function on and inside Γ except for a finite number of poles with no poles or zeros on Γ . Then

$$N_0(\Gamma) - N_\infty(\Gamma) = \frac{1}{2\pi i} \int_{\Gamma} \frac{f'(z)}{f(z)} dz$$

Lemma

Let Γ be a piecewise smooth positively oriented simple closed curve and f an analytic function on and inside Γ except for a finite number of poles with no poles or zeros on Γ . Then

$$N_0(\Gamma) - N_\infty(\Gamma) = \frac{1}{2\pi i} \int_{\Gamma} \frac{f'(z)}{f(z)} dz$$

Proof.



Example

Consider $f(z) = z^3$; $\Gamma = C(0; 1)$

Example

Consider $f(z) = z^3$; $\Gamma = C(0; 1)$

- $f'(z)/f(z) = 3/z$

Example

Consider $f(z) = z^3$; $\Gamma = C(0; 1)$

- $f'(z)/f(z) = 3/z$
- $\frac{1}{2\pi i} \int_{\Gamma} 3/z \, dz = 3$

Example

Consider $f(z) = z^3$; $\Gamma = C(0; 1)$

- $f'(z)/f(z) = 3/z$
- $\frac{1}{2\pi i} \int_{\Gamma} 3/z \, dz = 3$
- f has a zero of order 3 and no poles

Winding Numbers

- Let $f(t) = (\cos t, \sin t)$, $t \in [0, 6\pi]$

Winding Numbers

- Let $f(t) = (\cos t, \sin t)$, $t \in [0, 6\pi]$
- Suppose a camera at the origin is facing $(1, 0)$ and a rabbit runs the path traced by f

Winding Numbers

- Let $f(t) = (\cos t, \sin t)$, $t \in [0, 6\pi]$
- Suppose a camera at the origin is facing $(1, 0)$ and a rabbit runs the path traced by f
- Hence the location of the rabbit at time t is $(t, f(t))$

Winding Numbers

- Let $f(t) = (\cos t, \sin t)$, $t \in [0, 6\pi]$
- Suppose a camera at the origin is facing $(1, 0)$ and a rabbit runs the path traced by f
- Hence the location of the rabbit at time t is $(t, f(t))$
- If the camera takes a picture when the rabbit passes $(1, 0)$, how many photos are taken?

Winding Numbers

- Let $f(t) = (\cos t, \sin t)$, $t \in [0, 6\pi]$
- Suppose a camera at the origin is facing $(1, 0)$ and a rabbit runs the path traced by f
- Hence the location of the rabbit at time t is $(t, f(t))$
- If the camera takes a picture when the rabbit passes $(1, 0)$, how many photos are taken?
- What if $t \in [0, 5\pi]$?

Winding Numbers

- Let $f(t) = (\cos t, \sin t)$, $t \in [0, 6\pi]$
- Suppose a camera at the origin is facing $(1, 0)$ and a rabbit runs the path traced by f
- Hence the location of the rabbit at time t is $(t, f(t))$
- If the camera takes a picture when the rabbit passes $(1, 0)$, how many photos are taken?
- What if $t \in [0, 5\pi]$?
- What if $t \in [0, 7\pi]$?

Formalization

Definition

*Given a closed curve Γ and $z_0 \notin \Gamma$, define $n(\Gamma; z_0)$ to be the **winding number of Γ with respect to z_0***

Formalization

Definition

*Given a closed curve Γ and $z_0 \notin \Gamma$, define $n(\Gamma; z_0)$ to be the **winding number of Γ with respect to z_0***

Intuitively this quantity answers the question “How many times does Γ wind/wrap around z_0 ? ”

Lemma

Let Γ be a piecewise smooth closed curve that does not pass through z_0 . then

$$\frac{1}{2\pi i} \int_{\Gamma} \frac{f'(z)}{f(z)} dz$$

is an integer.

Lemma

Let Γ be a piecewise smooth closed curve that does not pass through z_0 . then

$$\frac{1}{2\pi i} \int_{\Gamma} \frac{f'(z)}{f(z)} dz$$

is an integer.

Note that the proof uses several “accessory functions”.

Proof I

Define

$$\phi(\tau) \int_a^\tau \frac{\gamma'(t)}{\gamma(t) - z_0} dt$$

and compute $\phi(b)$. Note that ϕ is continuous and

$$\phi'(\tau) = \frac{\gamma'(\tau)}{\gamma(\tau) - z_0}$$

wherever $\gamma'(t)$ is continuous. Define

$$\Phi(t) = (\gamma(t) - z_0)e^{-\phi(t)}$$

Observe that $\Phi(t)$ is continuous on $[a, b]$ except possibly for finitely many points. Also $\Phi'(t) = 0$ where Φ' exists. Therefore Φ is constant so that

$$\Phi(t) = \Phi(a) = (\gamma(a) - z_0)e^{-\phi(a)} = \gamma(a) - z_0$$

Proof II

Therefore

$$e^{\phi(t)} = \frac{\gamma(t) - z_0}{\gamma(a) - z_0}$$

Since Γ is closed it follows that $e^{\phi(b)} = 1$ and the result follows.

The Winding Number

- Define $n(\Gamma; z_0) = \frac{1}{2\pi i} \int_{\Gamma} \frac{dz}{z - z_0}$; *index*

The Winding Number

- Define $n(\Gamma; z_0) = \frac{1}{2\pi i} \int_{\Gamma} \frac{dz}{z - z_0}$; *index*
- Observe that $n(\Gamma; z) : \mathbb{C} \setminus \Gamma \rightarrow \mathbb{Z}$ is a function of z

The Winding Number

- Define $n(\Gamma; z_0) = \frac{1}{2\pi i} \int_{\Gamma} \frac{dz}{z - z_0}$; *index*
- Observe that $n(\Gamma; z) : \mathbb{C} \setminus \Gamma \rightarrow \mathbb{Z}$ is a function of z
- Note that $\mathbb{C} \setminus \Gamma$ is open and consists of 1 unbounded “piece” and a finite number of bounded “pieces”

The Winding Number

- Define $n(\Gamma; z_0) = \frac{1}{2\pi i} \int_{\Gamma} \frac{dz}{z - z_0}$; *index*
- Observe that $n(\Gamma; z) : \mathbb{C} \setminus \Gamma \rightarrow \mathbb{Z}$ is a function of z
- Note that $\mathbb{C} \setminus \Gamma$ is open and consists of 1 unbounded “piece” and a finite number of bounded “pieces”

Lemma

Let Γ be a closed curve and z_0, z_1 points in the same component of $\mathbb{C} \setminus \Gamma$. Then $n(\Gamma; z_0) = n(\Gamma; z_1)$; moreover if z_0 is outside Γ , then $n(\Gamma; z_0) = 0$.

Argument Principle Preliminaries

- Let Γ be a simple closed curve and f analytic, non-constant with $f'(z) \neq 0$ for all $z \in \Gamma$

Argument Principle Preliminaries

- Let Γ be a simple closed curve and f analytic, non-constant with $f'(z) \neq 0$ for all $z \in \Gamma$
- $f(\Gamma) = \{f(z) : z \in \Gamma\}$

Argument Principle Preliminaries

- Let Γ be a simple closed curve and f analytic, non-constant with $f'(z) \neq 0$ for all $z \in \Gamma$
- $f(\Gamma) = \{f(z) : z \in \Gamma\}$
- If Γ is parameterized by $\gamma : [a, b] \rightarrow \Gamma$, then $f(\Gamma) = f(\gamma(t))$

Argument Principle Preliminaries

- Let Γ be a simple closed curve and f analytic, non-constant with $f'(z) \neq 0$ for all $z \in \Gamma$
- $f(\Gamma) = \{f(z) : z \in \Gamma\}$
- If Γ is parameterized by $\gamma : [a, b] \rightarrow \Gamma$, then $f(\Gamma) = f(\gamma(t))$
- Therefore $f(\Gamma)$ is a curve

Question: If $w_0 \notin f(\Gamma)$, what is $n(f(\Gamma); w_0)$?

Question: If $w_0 \notin f(\Gamma)$, what is $n(f(\Gamma); w_0)$? Answer:

$$\frac{1}{2\pi i} \int_{f(\Gamma)} \frac{dw}{w - w_0} = \frac{1}{2\pi i} \int_{\Gamma} \frac{f'(z)}{f(z) - w_0} dz$$

Question: If $w_0 \notin f(\Gamma)$, what is $n(f(\Gamma); w_0)$? Answer:

$$\frac{1}{2\pi i} \int_{f(\Gamma)} \frac{dw}{w - w_0} = \frac{1}{2\pi i} \int_{\Gamma} \frac{f'(z)}{f(z) - w_0} dz$$

Note that $w = f(z)$ so that $dw = f'(z)dz$ by Change of Variables. Hence the answer is the same as the difference between the zeros and poles of $f(z) - w_0$ inside Γ .

Complete Statement

Theorem

Let Γ be a piecewise smooth positively oriented simple closed curve in the z -plane and f be analytic and non-constant on a domain containing Γ and its interior, except perhaps for a finite number of poles strictly inside Γ . Let w_0 be a point of the w -plane not on the curve $f(\Gamma)$. Then

$$N_{w_0}(\Gamma) - N_\infty(\Gamma) = \frac{1}{2\pi i} \int_{\Gamma} \frac{f'(z)}{f(z) - w_0} dz = n(f(\Gamma); w_0)$$

Examples

- z^k

Examples

- z^k
- $1/z$

Real Integrals

- Consider $I = \int_a^b f(x) dx$

Real Integrals

- Consider $I = \int_a^b f(x) dx$
- Process:

Real Integrals

- Consider $I = \int_a^b f(x) dx$
- Process:
- Relate I to a well-chosen complex line integral

Real Integrals

- Consider $I = \int_a^b f(x) dx$
- Process:
- Relate I to a well-chosen complex line integral
- Evaluate using residues

Real Integrals

- Consider $I = \int_a^b f(x) dx$
- Process:
- Relate I to a well-chosen complex line integral
- Evaluate using residues
- Extract answer

Real Integrals

- Consider $I = \int_a^b f(x) dx$
- Process:
- Relate I to a well-chosen complex line integral
- Evaluate using residues
- Extract answer

First define $\int_{\mathbb{R}} f(x) dx = \int_{-\infty}^{\infty} f(x) dx = \lim_{R \rightarrow \infty} \int_{-R}^R f(x) dx$; the **principal value**

Example: Type A

Lemma

Let $f(z)$ be analytic in a domain containing the closed half-plane $y \geq 0$ except for a finite number of poles in the open half-plane $y > 0$. Suppose further that $\lim_{|z| \rightarrow \infty} zf(z) = 0$ uniformly for $z \in \{w \in \mathbb{C} : \Im(w) \geq 0\}$. Then

$I = \int_{-\infty}^{\infty} f(x) dx$ is $2\pi i$ times the sum of residues of $f(z)$ at poles in the upper half-plane.

Example: Type A

Lemma

Let $f(z)$ be analytic in a domain containing the closed half-plane $y \geq 0$ except for a finite number of poles in the open half-plane $y > 0$. Suppose further that $\lim_{|z| \rightarrow \infty} zf(z) = 0$ uniformly for $z \in \{w \in \mathbb{C} : \Im(w) \geq 0\}$. Then

$I = \int_{-\infty}^{\infty} f(x) dx$ is $2\pi i$ times the sum of residues of $f(z)$ at poles in the upper half-plane.

- Evaluate $\int_{\mathbb{R}} \frac{dx}{1+x^2}$

Example: Type A

Lemma

Let $f(z)$ be analytic in a domain containing the closed half-plane $y \geq 0$ except for a finite number of poles in the open half-plane $y > 0$. Suppose further that $\lim_{|z| \rightarrow \infty} zf(z) = 0$ uniformly for $z \in \{w \in \mathbb{C} : \Im(w) \geq 0\}$. Then

$I = \int_{-\infty}^{\infty} f(x) dx$ is $2\pi i$ times the sum of residues of $f(z)$ at poles in the upper half-plane.

- Evaluate $\int_{\mathbb{R}} \frac{dx}{1+x^2}$
- Choose $f(z) = 1/(1+z^2)$; verify $zf(z) \rightarrow 0$

Example: Type A

Lemma

Let $f(z)$ be analytic in a domain containing the closed half-plane $y \geq 0$ except for a finite number of poles in the open half-plane $y > 0$. Suppose further that $\lim_{|z| \rightarrow \infty} zf(z) = 0$ uniformly for $z \in \{w \in \mathbb{C} : \Im(w) \geq 0\}$. Then

$I = \int_{-\infty}^{\infty} f(x) dx$ is $2\pi i$ times the sum of residues of $f(z)$ at poles in the upper half-plane.

- Evaluate $\int_{\mathbb{R}} \frac{dx}{1+x^2}$
- Choose $f(z) = 1/(1+z^2)$; verify $zf(z) \rightarrow 0$
- Note $f(z)$ has poles $i, -i$ and $\text{Res}(f; i) = -i/2$

Example: Type A

Lemma

Let $f(z)$ be analytic in a domain containing the closed half-plane $y \geq 0$ except for a finite number of poles in the open half-plane $y > 0$. Suppose further that $\lim_{|z| \rightarrow \infty} zf(z) = 0$ uniformly for $z \in \{w \in \mathbb{C} : \Im(w) \geq 0\}$. Then

$I = \int_{-\infty}^{\infty} f(x) dx$ is $2\pi i$ times the sum of residues of $f(z)$ at poles in the upper half-plane.

- Evaluate $\int_{\mathbb{R}} \frac{dx}{1+x^2}$
- Choose $f(z) = 1/(1+z^2)$; verify $zf(z) \rightarrow 0$
- Note $f(z)$ has poles $i, -i$ and $\text{Res}(f; i) = -i/2$
- Hence the integral has value $2\pi i(-i/2) = \pi$

Jordan's Lemma

Jordan's Lemma

Lemma (Jordan's Lemma)

Let $g(z)$ be analytic in a domain containing the closed upper half-plane except perhaps a finite number of poles in the open upper half-plane. Suppose further that

Then

$$\lim_{R \rightarrow \infty} \int_{C^+(0;R)} g(z) e^{ikz} dz = 0$$

Jordan's Lemma

Lemma (Jordan's Lemma)

Let $g(z)$ be analytic in a domain containing the closed upper half-plane except perhaps a finite number of poles in the open upper half-plane. Suppose further that

- (i) $g(z) \rightarrow 0$ uniformly as $|z| \rightarrow \infty$ in the closed upper half-plane

Then

$$\lim_{R \rightarrow \infty} \int_{C^+(0;R)} g(z) e^{ikz} dz = 0$$

Jordan's Lemma

Lemma (Jordan's Lemma)

Let $g(z)$ be analytic in a domain containing the closed upper half-plane except perhaps a finite number of poles in the open upper half-plane.

Suppose further that

- (i) $g(z) \rightarrow 0$ uniformly as $|z| \rightarrow \infty$ in the closed upper half-plane
- (ii) $k > 0$

Then

$$\lim_{R \rightarrow \infty} \int_{C^+(0;R)} g(z) e^{ikz} dz = 0$$

Jordan's Lemma

Lemma (Jordan's Lemma)

Let $g(z)$ be analytic in a domain containing the closed upper half-plane except perhaps a finite number of poles in the open upper half-plane. Suppose further that

- (i) $g(z) \rightarrow 0$ uniformly as $|z| \rightarrow \infty$ in the closed upper half-plane
- (ii) $k > 0$

Then

$$\lim_{R \rightarrow \infty} \int_{C^+(0;R)} g(z) e^{ikz} dz = 0$$

Corollary

$\int_{\mathbb{R}} g(x) e^{ikx} dx$ is $2\pi i$ times the sum of residues of $g(z) e^{ikz}$ in the upper half-plane.

Example: Type B

- Evaluate $\int_{\mathbb{R}} \frac{\cos(x)}{x^2+2x+4} dx$

Example: Type B

- Evaluate $\int_{\mathbb{R}} \frac{\cos(x)}{x^2+2x+4} dx$
- Choose $f(z) = \frac{e^{iz}}{z^2+2z+4}$

Example: Type B

- Evaluate $\int_{\mathbb{R}} \frac{\cos(x)}{x^2+2x+4} dx$
- Choose $f(z) = \frac{e^{iz}}{z^2+2z+4}$
- Compute $\text{Res}(f(z); -1 + i\sqrt{3}) = \frac{e^{-i-\sqrt{3}}}{2i\sqrt{3}}$

Example: Type B

- Evaluate $\int_{\mathbb{R}} \frac{\cos(x)}{x^2+2x+4} dx$
- Choose $f(z) = \frac{e^{iz}}{z^2+2z+4}$
- Compute $\text{Res}(f(z); -1 + i\sqrt{3}) = \frac{e^{-i-\sqrt{3}}}{2i\sqrt{3}}$
- Hence the integral evaluates as $\frac{\pi}{\sqrt{3}} e^{-i-\sqrt{3}}$

Example: Type B

- Evaluate $\int_{\mathbb{R}} \frac{\cos(x)}{x^2+2x+4} dx$
- Choose $f(z) = \frac{e^{iz}}{z^2+2z+4}$
- Compute $\text{Res}(f(z); -1 + i\sqrt{3}) = \frac{e^{-i-\sqrt{3}}}{2i\sqrt{3}}$
- Hence the integral evaluates as $\frac{\pi}{\sqrt{3}} e^{-i-\sqrt{3}}$
- Which has real part

$$\frac{\pi}{\sqrt{3}} e^{-\sqrt{3}} \cos(1)$$

Example: Type C

- Consider integrals of the form $\int_0^{2\pi} \Phi(\cos(\theta), \sin(\theta)) d\theta$ where Φ is a rational function with real coefficients.

Example: Type C

- Consider integrals of the form $\int_0^{2\pi} \Phi(\cos(\theta), \sin(\theta)) d\theta$ where Φ is a rational function with real coefficients.
- Change variables using Euler's identity to compute

$$\int_C \phi(z) dz$$

where $C = \{|z| = 1\}$

Example: Type C

- Consider integrals of the form $\int_0^{2\pi} \Phi(\cos(\theta), \sin(\theta)) d\theta$ where Φ is a rational function with real coefficients.
- Change variables using Euler's identity to compute

$$\int_C \phi(z) dz$$

where $C = \{|z| = 1\}$

- Observe that ϕ must be a rational function

Example: Type C

- Consider integrals of the form $\int_0^{2\pi} \Phi(\cos(\theta), \sin(\theta)) d\theta$ where Φ is a rational function with real coefficients.
- Change variables using Euler's identity to compute

$$\int_C \phi(z) dz$$

where $C = \{|z| = 1\}$

- Observe that ϕ must be a rational function
- If ϕ has a pole between $0, 2\pi$, this transformed integral is not possible.

Example

- Show $\int_0^{2\pi} \frac{\cos(3\theta)}{5-4\cos(\theta)} dz = \frac{\pi}{12}$

Example

- Show $\int_0^{2\pi} \frac{\cos(3\theta)}{5-4\cos(\theta)} dz = \frac{\pi}{12}$
- Note that substituting for $\cos(3\theta)$ results in a significantly more tedious integral