




🔐 PLANT HARDINESS ZONE 6 DATASET

COVER CROP DESCRIPTION

Upright small grain. Widely used in the Northeast. Best small grain for drought, heat, salty or alkaline soils. Not suitable for wet or acid soils. Winter barley typically needs vernalization (overwintering) to produce stalk/head/seed. May stay short and fail to flower if planted in spring. Quick growth and high biomass if fertility is good. Good weed suppressor, N scavenger, forage producer. Retains feed quality after heading. Good nurse crop for legumes. Use care when including as a cover crop in a small grain rotation (host for same diseases and pests). Bloom timing and height match crimson clover, rapeseed when used in a mix. More winter-hardy than oat, less than wheat and rye. Better fall growth and tillers more than wheat and rye but more prone to frost damage. More winter-hardy than oat, less than wheat and rye. Better fall growth and tillers more than wheat and rye but more prone to frost damage.









Barley, Winter - Salon [2020]



Barley, Winter - Salon [2020]

Growing Window Short Penetrates Plow Pan Nitrogen Scavenging Reduces Surface Compaction Lasting Residue Improve Soil Organic Matter Prevent Fall Soil Erosion Increase Soil Aggregation Prevent Spring Soil Erosion Good Grazing Forage Harvest Value Pollinator Food





Barley, Winter


Hordeum vulgare







♠ PLANT HARDINESS ZONE 6 DATASET

Growth Traits	
Duration	Annual
Zone Use	Common
Shape And Orientation	Erect
Dry Matter (Lbs/A/Yr)	2000 - 5000
Soil Texture	Medium
Soil pH	6 - 8.5
Soil Moisture Use	Medium
Hessian Fly Free Date?	No
Ease Of Establishment	
Establishes Quickly	
Early Spring Growth	
Flowering Trigger	Vernalization
Root Depth	Medium







Barley, Winter

Hordeum vulgare





PLANT HARDINESS ZONE 6 DATASET



Extended Comments

Basic Agronomics: More commonly grown south of the Mason-Dixon line and in SE PA.

Forage and Grazing: Failure to overwinter reliably in USDA hardiness zones 5 and colder means biomass production may be low, which potentially limits grazing/forage harvest value.

Insects: Terminating barler early or late showed no impact on arthropod (pest or beneficial) numbers in a subsequent soybean crop

References & Resources

Use of Cover Crops and Green Manures to Attract Beneficial Insects, University of Connecticut Integrated Pest Management Program

Fall Cover Crops, University of Delaware Cooperative Extension

Cover Crops and Green Manure Crops - Benefits, Selection, and Use, Rutgers Cooperative Research and Extension Pasture Production of Selected Forage Species, University of New Hampshire Cooperative Extension

Plant Cover Crops, University of Maryland Extension