

Discussion 3: Week 2 - Stats 100 A

Andrew Lizarraga

Department of Statistics and Data Science

July 2, 2024

Homework 1 Survey

- ▶ A majority of you indicated that Problem 2 and 3 are the most difficult.

Homework 1 Survey

- ▶ A majority of you indicated that Problem 2 and 3 are the most difficult.
- ▶ Remember if you're ever unclear about a topic, please feel free to address your questions to Prof. Wu during lectures.

Problem 2 (1)

► $X, Y \sim^{ind} U[0, 1]$

Problem 2 (1)

- ▶ $X, Y \sim^{ind} U[0, 1]$
- ▶ Equivalently (X, Y) is a random point in $[0, 1]^2$.

Problem 2 (1)

- ▶ $X, Y \sim^{ind} U[0, 1]$
- ▶ Equivalently (X, Y) is a random point in $[0, 1]^2$.

Determine $P(X^2 + Y^2 \leq 1)$ by measuring an area.

Problem 2 (1)

Problem 2 (2)

Suppose we repeat the experiment n times. For large n how, how often $X^2 + Y^2 \leq 1$? Suppose it happens m times, can you calculate π based on m and n ?

Problem 2 (2)

Problem 2 (3)

Calculate $P(X \geq \frac{1}{2})$ and $P(X \geq \frac{1}{2} \mid X + Y \geq 1)$

Problem 2 (3)

Problem 2 (4)

Let A be the event that $X \in [.2, .6]$ and B be the event that $Y \in [.3, .5]$. Show that $P(A \cap B) = P(A)P(B)$

Problem 2 (4)

Problem 3 (Buffon's Needle)

- ▶ The length of the needle is 1 unit.

Problem 3 (Buffon's Needle)

- ▶ The length of the needle is 1 unit.
- ▶ The length between parallel the parallel lines is also 1 unit.

Problem 3 (Buffon's Needle)

- ▶ The length of the needle is 1 unit.
- ▶ The length between parallel the parallel lines is also 1 unit.

Estimate π using the Buffon Needle experiment

Problem 3 (Buffon's Needle)

Problem 3 (Buffon's Needle)

Problem 3 (Lazzarini's Lie)

Thank You / Questions

Contact:

You may contact me at: **andrewlizarraga@g.ucla.edu**.