

# **PRIOR AUTHORIZATION POLICY**

Policy: Multiple Sclerosis (Injectable – Beta Interferon) – Avonex Prior

Authorization Policy

• Avonex® (interferon beta-1a intramuscular injection – Biogen)

**REVIEW DATE:** 07/23/2025

#### INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna COMPANIES AND/OR LINES OF BUSINESS ONLY PROVIDE UTILIZATION REVIEW SERVICES TO CLIENTS AND DO NOT MAKE COVERAGE DETERMINATIONS. REFERENCES TO STANDARD BENEFIT PLAN LANGUAGE AND COVERAGE DETERMINATIONS DO NOT APPLY TO THOSE CLIENTS. COVERAGE POLICIES ARE INTENDED TO PROVIDE GUIDANCE IN INTERPRETING CERTAIN STANDARD BENEFIT PLANS ADMINISTERED BY CIGNA COMPANIES. PLEASE NOTE, THE TERMS OF A CUSTOMER'S PARTICULAR BENEFIT PLAN DOCUMENT [GROUP SERVICE AGREEMENT, EVIDENCE OF COVERAGE, CERTIFICATE OF COVERAGE, SUMMARY PLAN DESCRIPTION (SPD) OR SIMILAR PLAN DOCUMENT] MAY DIFFER SIGNIFICANTLY FROM THE STANDARD BENEFIT PLANS UPON WHICH THESE COVERAGE POLICIES ARE BASED. FOR EXAMPLE, A CUSTOMER'S BENEFIT PLAN DOCUMENT MAY CONTAIN A SPECIFIC EXCLUSION RELATED TO A TOPIC ADDRESSED IN A COVERAGE POLICY. IN THE EVENT OF A CONFLICT, A CUSTOMER'S BENEFIT PLAN DOCUMENT ALWAYS SUPERSEDES THE INFORMATION IN THE COVERAGE POLICIES, IN THE ABSENCE OF A CONTROLLING FEDERAL OR STATE COVERAGE MANDATE, BENEFITS ARE ULTIMATELY DETERMINED BY THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT. COVERAGE DETERMINATIONS IN EACH SPECIFIC INSTANCE REQUIRE CONSIDERATION OF 1) THE TERMS OF THE APPLICABLE BENEFIT PLAN DOCUMENT IN EFFECT ON THE DATE OF SERVICE; 2) ANY APPLICABLE LAWS/REGULATIONS; 3) ANY RELEVANT COLLATERAL SOURCE MATERIALS INCLUDING COVERAGE POLICIES AND; 4) THE SPECIFIC FACTS OF THE PARTICULAR SITUATION. EACH COVERAGE REQUEST SHOULD BE REVIEWED ON ITS OWN MERITS. MEDICAL DIRECTORS ARE EXPECTED TO EXERCISE CLINICAL JUDGMENT WHERE APPROPRIATE AND HAVE DISCRETION IN MAKING INDIVIDUAL COVERAGE DETERMINATIONS. WHERE COVERAGE FOR CARE OR SERVICES DOES NOT DEPEND ON SPECIFIC CIRCUMSTANCES, REIMBURSEMENT WILL ONLY BE PROVIDED IF A REQUESTED SERVICE(S) IS SUBMITTED IN ACCORDANCE WITH THE RELEVANT CRITERIA OUTLINED IN THE APPLICABLE COVERAGE POLICY, INCLUDING COVERED DIAGNOSIS AND/OR PROCEDURE CODE(S). REIMBURSEMENT IS NOT ALLOWED FOR SERVICES WHEN BILLED FOR CONDITIONS OR DIAGNOSES THAT ARE NOT COVERED UNDER THIS COVERAGE POLICY (SEE "CODING INFORMATION" BELOW). WHEN BILLING, PROVIDERS MUST USE THE MOST APPROPRIATE CODES AS OF THE EFFECTIVE DATE OF THE SUBMISSION. CLAIMS SUBMITTED FOR SERVICES THAT ARE NOT ACCOMPANIED BY COVERED CODE(S) UNDER THE APPLICABLE COVERAGE POLICY WILL BE DENIED AS NOT COVERED. COVERAGE POLICIES RELATE EXCLUSIVELY TO THE ADMINISTRATION OF HEALTH BENEFIT PLANS. COVERAGE POLICIES ARE NOT RECOMMENDATIONS FOR TREATMENT AND SHOULD NEVER BE USED AS TREATMENT GUIDELINES. IN CERTAIN MARKETS, DELEGATED VENDOR GUIDELINES MAY BE USED TO SUPPORT MEDICAL NECESSITY AND OTHER COVERAGE DETERMINATIONS.

# CIGNA NATIONAL FORMULARY COVERAGE:

#### **OVERVIEW**

Avonex is indicated for the treatment of patients with relapsing forms of **multiple sclerosis** (MS) to include clinically isolated syndrome, relapsing remitting disease, and active secondary progressive disease in adults.<sup>1</sup>

### **Disease Overview**

MS is a chronic, inflammatory, demyelinating, autoimmune disease of the central nervous system (CNS) that impacts almost 1,000,000 people in the US.<sup>2-4</sup> The condition is marked by inflammation and demyelination, as well as degenerative alterations. Patients usually experience relapses and remissions in their neurological symptoms. For most patients, the onset of MS symptoms occurs when patients are 20 to 40 years of age; however, children can get MS and new onset disease can occur in older adults. The MS disease course is heterogeneous but has some patterns.

Approximately 85% to 90% of patients have a relapsing pattern at onset. However, this transitions over time in patients who are untreated to a worsening with very few or no relapses or magnetic resonance imaging (MRI) activity (secondary progressive MS). Around 10% to 15% of patients have a steady progression of symptoms over time (primary progressive MS), marked by some clinical manifestations or by MRI activity. Primary progressive MS is generally diagnosed in patients on the upper level of the typical age range (e.g., almost 40 years of age) and the distribution is equivalent among the two genders. Advances in the understanding of the MS disease process, as well as in MRI technology, spurned updated disease course descriptions in 2013,<sup>5</sup> as well as in 2017.<sup>6</sup> The revised disease courses are clinically isolated syndrome, relapsing remitting MS, primary progressive MS, and secondary progressive MS.<sup>2-6</sup> Clinically isolated syndrome is now more recognized among the course descriptions of MS. It is the first clinical presentation of MS that displays characteristics of inflammatory demyelination that may possibly be MS but has yet to fulfill diagnostic criteria. It is notable that the other MS designations can be further characterized considering whether patients have active disease (or not active), as well as if disease is worsening or stable. Disability in MS is commonly graded on the deterioration of mobility per the Expanded Disability Status Scale (EDSS) an ordinal scale that ranges from 0 to 10, with higher scores indicating greater disability.

## **Guidelines**

In September 2019, a consensus paper was updated by the MS Coalition that discusses the use of disease-modifying therapies in MS.<sup>2</sup> Many options from various disease classes, involving different mechanisms of action and modes of administration, have shown benefits in patients with MS.

#### **POLICY STATEMENT**

Prior Authorization is recommended for prescription benefit coverage of Avonex. All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with Avonex as well as the monitoring required for adverse events and long-term efficacy, approval requires Avonex to be prescribed by or in consultation with a physician who specializes in the condition being treated.

 Avonex® (interferon beta-1a intramuscular injection – Biogen)

is(are) covered as medically necessary when the following criteria is(are) met for FDA-approved indication(s) or other uses with supportive evidence (if applicable):

## **FDA-Approved Indication**

- **1. Multiple Sclerosis.** Approve for the duration noted below if the patient meets ONE of the following (A <u>or</u> B):
  - **A)** Initial Therapy. Approve for 1 year if the patient meets BOTH of the following (i <u>and</u> ii):
    - i. Patient has a relapsing form of multiple sclerosis; AND <a href="Note">Note</a>: Examples of relapsing forms of multiple sclerosis include clinically isolated syndrome, relapsing remitting disease, and active secondary progressive disease.
    - **ii.** Medication is prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis; OR
  - **B)** Patient is Currently Receiving Avonex for ≥ 1 Year. Approve for 1 year if the patient meets ALL of the following (i, ii, and iii):
    - i. Patient has a relapsing form of multiple sclerosis; AND <a href="Note">Note</a>: Examples of relapsing forms of multiple sclerosis include clinically isolated syndrome, relapsing remitting disease, and active secondary progressive disease.
    - ii. Patient meets ONE of the following (a or b):
      - a) Patient experienced a beneficial clinical response when assessed by at least one objective measure; OR
        Note: Examples include stabilization or reduced worsening in disease

activity as evaluated by magnetic resonance imaging (MRI) [absence or a decrease in gadolinium enhancing lesions, decrease in the number of new or enlarging T2 lesions]; stabilization or reduced worsening on the Expanded Disability State Scale (EDSS) score; achievement in criteria for No Evidence of Disease Activity-3 (NEDA-3) or NEDA-4; improvement on the fatigue symptom and impact questionnaire-relapsing multiple sclerosis (FSIQ-RMS) scale; reduction or absence of relapses; improvement or maintenance on the six-minute walk test or

- 12-Item MS Walking Scale; improvement on the Multiple Sclerosis Functional Composite (MSFC) score; and/or attenuation of brain volume loss
- b) Patient experienced stabilization, slowed progression, or improvement in at least one symptom such as motor function, fatigue, vision, bowel/bladder function, spasticity, walking/gait, or pain/numbness/tingling sensation; AND
- **iii.** Medication is prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis.

## **CONDITIONS NOT COVERED**

 Avonex® (interferon beta-1a intramuscular injection – Biogen)

is(are) considered not medically necessary for ANY other use(s) including the following (this list may not be all inclusive; criteria will be updated as new published data are available):

- Concurrent Use with Other Disease-Modifying Agents Used for Multiple Sclerosis. These agents are not indicated for use in combination (See <u>Appendix</u> for examples). Additional data are required to determine if use of diseasemodifying multiple sclerosis agents in combination is safe and provides added efficacy.
- **2. Non-Relapsing Forms of Multiple Sclerosis.** The efficacy of Avonex has not been established in patients with non-relapsing forms of multiple sclerosis. Note: An example of a non-relapsing form of multiple sclerosis is primary progressive multiple sclerosis.

#### REFERENCES

- 1. Avonex® intramuscular injection [prescribing information]. Cambridge, MA: Biogen; July 2023.
- 2. A Consensus Paper by the Multiple Sclerosis Coalition. The use of disease-modifying therapies in multiple sclerosis. September 2019.
- 3. McGinley MP, Goldschmidt C, Rae-Grant AD. Diagnosis and treatment of multiple sclerosis. A review. *JAMA*. 2021;325(8):765-779.
- 4. No authors listed. Drugs for multiple sclerosis. Med Lett Drugs Ther. 2021;63(1620):42-48.
- 5. Lublin FD, Reingold SC, Cohen JA, et al. Defining the clinical course of multiple sclerosis: the 2013 revisions. *Neurology*. 2014;83:278-286.
- 6. Thompson AJ, Banwell BL, Barkhof F, et al. Diagnosis of multiple sclerosis: 2017 revisions of the McDonald criteria. *Lancet Neurol*. 2018;17(2):162-173.

## History

Type of	Summary of Changes	Review
Revision		Date

Update	08/10/2023: No criteria changes. To the Appendix, Briumvi and	NA
	Tascenso ODT were added. Also, it was noted that Aubagio is	
	available as a generic.	
Annual Revision	No criteria changes.	11/08/2023
Annual Revision	To the Appendix, Ocrevus Zunovo was added.	10/09/2024
Early Annual	To the Appendix, Extavia was deleted. The Policy name was changed	07/23/2025
Revision	to add "Injectable - Beta Interferon".	

## **APPENDIX**

Medication	Mode of Administration	
Aubagio® (teriflunomide tablets, generic)	Oral	
Avonex® (interferon beta-1a intramuscular injection)	Injection (self-administered)	
Bafiertam® (monomethyl fumarate delayed-release capsules)	Oral	
Betaseron® (interferon beta-1b subcutaneous injection)	Injection (self-administered)	
Briumvi® (ublituximab-xiiy intravenous infusion)	Intravenous infusion	
Copaxone® (glatiramer acetate subcutaneous injection, generic)	Injection (self-administered)	
Gilenya® (fingolimod capsules, generic)	Oral	
Glatopa® (glatiramer acetate subcutaneous injection)	Injection (self-administered)	
Kesimpta® (ofatumumab subcutaneous injection)	Injection (self-administered)	
Lemtrada® (alemtuzumab intravenous infusion)	Intravenous infusion	
Mavenclad® (cladribine tablets)	Oral	
Mayzent® (siponimod tablets)	Oral	
Ocrevus® (ocrelizumab intravenous infusion)	Intravenous infusion	
Ocrevus Zunovo <sup>™</sup> (ocrelizumab and hyaluronidase-ocsq subcutaneous injection)	Subcutaneous Injection (not self- administered)	
Plegridy® (peginterferon beta-1a subcutaneous or intramuscular injection)	Injection (self-administered)	
Ponvory® (ponesimod tablets)	Oral	
Rebif® (interferon beta-1a subcutaneous injection)	Injection (self-administered)	
Tascenso ODT® (fingolimod orally disintegrating tablets)	Oral	
Tecfidera® (dimethyl fumarate delayed-release capsules, generic)	Oral	
Tyruko® (natalizumab-sztn intravenous infusion)	Intravenous infusion	
Tysabri® (natalizumab intravenous infusion)	Intravenous infusion	
Vumerity® (diroximel fumarate delayed-release capsules)	Oral	
Zeposia® (ozanimod capsules)	Oral	

<sup>&</sup>quot;Cigna Companies" refers to operating subsidiaries of The Cigna Group. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Evernorth Behavioral Health, Inc., Cigna Health Management, Inc., and HMO or service company subsidiaries of The Cigna Group.© 2025 The Cigna Group.