Australian Government
Australian Pesticides and Veterinary Medicines Authority

PERMIT TO ALLOW SUPPLY AND MINOR USE OF AN UNREGISTERED AGVET CHEMICAL

PRODUCT FOR CONTROL OF BROADLEAF WEEDS IN SEED PRODUCTION OF CARROT CROPS

PERMIT NUMBER – PER88504

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a Supplier (as indicated) to possess the product for the purposes of supply and to supply the product to a person who can use the product under permit. This permit also allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows the Permit Holder, the Supplier (if not one and the same) and any person stipulated below to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 6 APRIL 2021 TO 30 APRIL 2026

Permit Holder and Supplier:
SOUTH PACIFIC SEEDS PTY LTD
48 – 50 Willandra Avenue
GRIFFITH NSW 2680

Additional Suppliers:
AGPRO PROPAZINE 500
AGPRO NZ LTD
10 Polaris Place
AUCKLAND 2163

GESAMIL 500FW HERBICIDE
ORION CROP PROTECTION LIMITED
14-22 Triton Drive
AUCKLAND 2163

Persons who can use the product under this permit:
Employees of South Pacific Seeds or growers or spray contractors who are trained and experienced in the use of agricultural chemicals and who under the supervision of the permit holder

CONDITIONS OF USE

Products to be used:
AGPRO PROPAZINE 500
AN UNREGISTERED PRODUCT
Containing 494 g/L PROPAZINE as the only active constituent.

GESAMIL 500FW HERBICIDE
AN UNREGISTERED PRODUCT
Containing 500 g/L PROPAZINE as the only active constituent.

Directions for Use:

Crop	Pest / Disease	Rate
Carrot crops for seed production **ONLY.**	Broadleaf weeds, particularly Nightshade, Fat Hen and Wireweed.	2.2 L/ha

Critical Use Comments:
- Apply ONCE only per season, in spring.
- Apply by boomspray
- Avoid spray overlap
- Apply prior to emergence of weeds, prior to the crop developing a full canopy

Withholding Period:
Harvest: Do not harvest carrots for human consumption.
Grazing: Do not graze or use treated carrot roots, foliage or seed [or treated weeds] for stock-feed.

Resistant Weeds Warning GROUP C HERBICIDE

Agpro Propazine is a member of the Triazines group of herbicides. Propazine has the inhibitors of photosynthesis at photosystem II mode of action. For weed resistance management Propazine is a Group C Herbicide.

Some naturally occurring weed Group-C herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Propazine or other Group-C herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, the supplier accepts no liability for any losses that may result from the failure of Propazine to control resistant weeds.

Re-entry:
Do not re-enter treated areas until spray is dry

Protection of Crops, native and other non-target plants
DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants, crops, cropping lands or pastures.

Protection of Wildlife, Fish, Crustacea and Environment
This product is very highly toxic to algae and aquatic macrophytes.
DO NOT apply under meteorological conditions, or from spraying equipment which could be expected to cause spray to drift onto susceptible plants/crops, cropping lands, pastures, wetlands, water-bodies or water-courses.
DO NOT apply this product within 60m of natural or impounded lakes or dams.

DO NOT contaminate streams, rivers or waterways with the chemical or used container.
DO NOT use in channels or drains where roots of desirable plants or shrubs may be extended.
Wash sprayer thoroughly with clean water after use.

Storage and Disposal
KEEP OUT OF REACH OF CHILDREN**.**
Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight**.**
Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product

Safety directions:
May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. Wash hands after use.

First Aid
If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766

Jurisdiction:
NSW, SA, TAS only.

Additional Conditions:
This permit allows for the use of a product in a manner specified on the permit. Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit. Unless otherwise stated, the use of the product must be in accordance with the product label at **Attachment 1 or 2**.

Fate of produce from treated plants:
Treated weeds or produce from plants treated with this product must not be supplied to or otherwise made available for human or animal consumption. The produce plus other plant material must therefore be disposed of appropriately

Supply
The supplier must supply the product in a container that complies with the requirements of section 18 of the Agricultural and Veterinary Chemicals Code Regulations. Attached to this container must be a label which is identical in content and format to the label in **Attachment 1 or 2. The attachment labels must also CLEARLY state that " THIS PRODUCT IS NOT REGISTERED AND MUST ONLY BE USED UNDER PERMIT 88504."**

Issued by the Australian Pesticides and Veterinary Medicines Authority

Attachment 1

AGPRO DIRECT LABEL
AGPRO PROPAZINE 500

Issue Date: 11/04/2018

WARNING

Keep out of reach of children

ECOTOXIC

AGPRO PROPAZINE 500

Contains 494g/L propazine in the form of a suspension concentrate

For post-emergent control of selected weeds in carrots, pasnips and Radiata pine and Douglas fir seed beds.

Read the directions for use before using this product.

Registered pursuant to the ACVM Act 1997 No: P004608
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to HSNO Act 1996 No HSR000387
See www.epa.govt.nz for approval conditions

Date of manufacture:
Batch no:
Net contents: 5L, 20L

Registered and Distributed by:
AGPRO NZ Ltd
10 Polaris Place
East Tamaki
P O Box 58-963
Botany
Auckland 2163
Ph 09 273 3456
Fax 09 273 3457
www.agpro.co.nz

Contact AGPRO Direct on freephone 0508 536 536 or visit our website www.agpro.co.nz

Attachment 1

AGPRO DIRECT

LABEL
AGPRO PROPAZINE 500

Issue Date: 11/04/2018

WARNING
May be harmful if swallowed, inhaled or in contact with the skin. Do not get into eyes or on skin. In case of contact, flush skin or eyes with plenty of water for 15 minutes; for eyes get medical attention. Do not eat, drink or smoke while using. May cause mild skin irritation. May cause serious eye irritation. Suspected of causing cancer via exposure route. Very toxic to aquatic life with long lasting effects. Avoid contamination of any water supply with chemical or empty container. Very toxic to the soil environment.
Remove protective clothing and wash hands and face thoroughly before meals and after work.
Wash protective clothing after work.

PRECAUTIONS.
Store in the original unopened container in a cool dry place, avoiding sunlight and away from foodstuffs and drink containers.
SPILLAGE: Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely (such as in a suitable landfill).
CONTAINER DISPOSAL: Triple rinse container and add rinsate to spray tank. Burn in an appropriate incinerator, if circumstances such as wind direction permit. Otherwise crush or puncture and bury in a suitable landfill, or if appropriate, recycle.

FIRST AID
If swallowed do NOT induce vomiting.
For advice contact National Poisons Centre (0800 POISONS 764-766) or a doctor immediately.
If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water.
If splashed in eyes, wash out immediately with water.

GENERAL INSTRUCTIONS
AGPRO Propazine 500 is a selective herbicide for post emergence control of weeds in carrots, pasnips, Radiata pine and Douglas fir seed beds. This product is absorbed by the roots, although some foliar absorption does take place. The product works by inhibiting photosynthesis.

MIXING
Invert container several times before opening. Stir well. Add the full quantity of product to the partly filled spray tank while agitating. Fill tank and agitate to ensure thorough mixing and continue agitation while spraying. Agitate vigorously from the bottom if allowed to stand.

APPLICATION
AGPRO Propazine may be applied through all conventional spray equipment. Do not use strainers finer than 300 micrometers.

COMPATIBILITY
This product is compatible with most commonly used agricultural insecticides, fungicides and herbicides, however a test run is advisable.

SPRAY DRIFT WARNING
Beware: Apply this product carefully. Spray drift may cause serious damage to other desirable plants.
Ensure compliance with relevant regional air management plans.
Aerial application: To minimise the risk of damage from spray drift, especially where susceptible vegetation is within 500 metres, use low drift-nozzles which minimise the production of small droplets (< 100microns).
Preferable to spray in a light constant breeze that is away from sensitive crops.

Contact AGPRO Direct on freephone 0508 536 536 or visit our website **www.agpro.co.nz**

Attachment 1

AGPRO DIRECT

LABEL
AGPRO PROPAZINE 500

Issue Date: 11/04/2018

SHELF LIFE

When stored appropriately, this product should show no significant degradation for two years from the date of manufacture. Contact us for further information about the use of product that is older than this.

DIRECTIONS FOR USE

Carrots & Parsnips: Apply at 2-4L/ha in 500L of water. Use the lower rate on sandy or silty soils and the higher rate on heavy clay soils or soils heavy in organic matter. Must be applied within 2-3 days of sowing and prior to emergence of the crop or weeds.

Radiata Pine seedbeds: Apply 2L/ha in 500L of water. Apply within 2-3 days after sowing or 1 month after total emergence of the seedlings, ensuring that the seedbed has been freshly weeded.

Douglas fir seedbeds: Apply 2L/ha in 500L of water. Apply at least 6 weeks after total emergence of the seedlings, ensuring that the seedbed has been freshly weeded.

RESISTANCE MANAGEMENT

Some annual broadleaf weeds may be resistant. Use an alternative herbicide to remove them.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the New Zealand (Maximum Residue Limits of Agricultural Compounds) Food Standards.

CONDITIONS OF SALE

The use of AGPRO PROPAZINE 500 flowable herbicide being beyond the control of the manufacturer, no warranty expressed or implied, beyond that covered under the Fair Trading Act is given by AGPRO NZ Limited regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and AGPRO NZ Limited accepts no responsibility for any consequences whatsoever resulting from the use of this product.

Contact AGPRO Direct on freephone 0508 536 536 or visit our website **www.agpro.co.nz**

Attachment 2



ORION CROP PROTECTION

Attachment 2

PLEASE READ THE LABEL COMPLETELY BEFORE USE

HAZARD CLASSIFICATIONS: 6.1D, 6.3B, 6.4A, 6.7B, 9.1A, 9.2A
WARNING: TOXIC, ECOTOXIC
May be harmful if swallowed, inhaled or absorbed through the skin. May cause eye and mild skin irritation. Limited evidence of carcinogenic effects in animal feeding studies. Very toxic to aquatic organisms and to some plant species.

PRECAUTIONS
APPROVED HANDLER: This product must be under the control of an approved handler during use.
STORAGE: Keep out of reach of children. Store in the original, tightly closed container, in a cool, dry and well ventilated area away from feed or foodstuffs. Avoid temperatures below 5°C and above 35°C. Storage of certain quantities of this product trigger specific storage requirements (Refer to HazNote). When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of product that is older than this.
PERSONAL PROTECTION: Avoid contact with eyes and skin. Avoid inhalation of spray mist. When mixing or applying, wear overalls, boots, gloves and eye protection. Clean protective clothing daily after work. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work.
USE: Apply with well maintained and calibrated spray equipment.
DISPOSAL: Dispose of this product only by using in accordance with this label, or at a suitable landfill. Do not burn. Triple rinse container and add rinsate to the spray tank. Dispose of container to a suitable landfill or agricultural recycling scheme. Do not burn. Do not use container for any other purpose.
ENVIRONMENTAL: Use this product carefully. Do not contaminate aquatic environments with product, spray drift or empty container. Spray drift or equipment contamination may cause serious damage to desirable plants. Do not allow spray drift outside the target area to occur.

First Aid
If swallowed do not induce vomiting. If splashed in eyes, wash out immediately with running water for several minutes. If skin or hair contact occurs, remove contaminated clothing and wash with soap and running water. If inhaled, immediately move to fresh air and rest. Maintain half upright position if breathing is difficult. Call the National Poisons Centre **0800 POISON (0800 764 766)** (24 hrs) or a doctor immediately for advice.

Emergency Response
In a transport emergency call **111, Police or Fire Brigade**. For specialist advice in an emergency only, call **0800 CHEMCALL (0800 243 6225)** (24 hrs).

Spillage
Exclude non-essential people from the area. Wear protective clothing and prevent material from entering waterways. Absorb spill with inert material such as soil, sand, etc, and place in a waste container. Wash area with water and detergent, and absorb with further inert material. Dispose of safely to a suitable landfill.

Safety Data Sheet
Additional information is provided in the safety data sheet. Freephone 0800 ORIONCP (0800 674 6627) or visit www.orioncp.co.nz.

GENERAL INFORMATION
Gesamil 500FW is a selective herbicide for the pre-emergence control of many annual broadleaf and grass weeds. When applied according to the directions Gesamil 500FW will give prolonged control of weeds without tainting or harming the crop. Gesamil 500FW is effective on germinating weed seedlings and prevents weed competition with the establishing crop. Gesamil 500FW is not effective on deep -rooted perennial weeds such a Californian thistle, dock and couch.

Soil Conditions: For best results with Gesamil 500FW prepare a fine firm seedbed free from clods and undecayed vegetation.
Soil Moisture: Adequate soil moisture is necessary if Gesamil 500FW is to be effective. If dry conditions are being experienced then moisture should be applied prior to application of Gesamil 500FW. Light rain or light irrigation immediately after application will also improve results.

DIRECTIONS FOR USE
RATES AND TIMING OF APPLICATION
Carrots and Parsnips: For clay soils high in organic matter use 4 litres Gesamil 500FW per hectare. For sandy or silty soils use 2 litres Gesamil 500FW per hectare. Do not apply on very light soils with very little organic matter.
Pinus Species: (except small seeded species such as *P. contorta, P. mugo, P. sylvestris and P. uncinata*): Apply 2 litres Gesamil 500FW per hectare either as a post-sowing application (within 2 - 3 days of sowing) or to the freshly weeded seedbed one month after total emergence of the pines.
Douglas Fir: Apply 2 litres Gesamil 500FW per hectare as a post-emergence crop spray when crop seedlings are at least 6 weeks old and the seedbed is freshly weeded.
Time of Application: Gesamil 500FW must be applied within 2 -3 days of sowing and prior to the emergence of weeds or seedling crop. Gesamil 500FW can be applied to established carrots and parsnips provided that weeds are not present or to freshly weeded conifers one month after emergence. In carrots and parsnips if weeds have germinated use Gesagard® 500 FW.
Volume of Application: Apply in 500 litres of water per hectare.

APPLICATION METHODS
Gesamil 500FW can be applied with conventional ground spraying equipment. Care should be taken to avoid excessive overlapping. Gesamil 500FW is non-corrosive and almost non-abrasive. Do not use strainers finer than 300 micrometres. Wash equipment thoroughly after use.

MIXING
Gesamil 500FW mixes readily with water. To ensure a uniform suspension pour the correct amount into the partly filled tank and whilst stirring add the remainder of the water. Agitate thoroughly before and during spraying and before spraying after a stoppage.

SOIL RESIDUES
Subsequent crops may be safely planted after 4 - 5 months. Cultivation by ploughing to a depth of 15 cm is recommended to remove any residue of Gesamil 500FW from the soil surface.

WARRANTY
This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors beyond the manufacturer's control. The manufacturer declines all liability for damage resulting from improper storage or use of the product.

Registered pursuant to the ACVM Act 1997, No. P7219
See www.nzfsa.govt/acvm/ for registration conditions.
Approved pursuant to the HSNO Act 1996, No. HSR000399
See www.ermanz.govt.nz for approval conditions.

® GESAMIL and GESAGARD are Registered Trade Marks of a Syngenta Group Company.

Registrant: Orion Crop Protection Limited,
Triton Technical Park, Building B, 14 – 22 Triton Drive,
North Harbour, Auckland.