Australian Government
Australian Pesticides and Veterinary Medicines Authority

PERMIT TO ALLOW SUPPLY AND MINOR USE OF AN UNREGISTERED AGVET CHEMICAL PRODUCT FOR CONTROL OF ANNUAL BROADLEAVED WEEDS AND VARIOUS GRASSES IN BABY LEAF SPINACH AND BABY LEAF BEET

PERMIT NUMBER – PER92508

This permit is issued to the Permit Holder in response to an application granted by the APVMA under section 112 of the Agvet Codes of the jurisdictions set out below. This permit allows a Supplier (as indicated) to possess the product for the purposes of supply and to supply the product to a person who can use the product under permit. This permit also allows a person, as stipulated below, to use the product in the manner specified in this permit in the designated jurisdictions. This permit also allows the Permit Holder, the Supplier (if not one and the same) and any person stipulated below to claim that the product can be used in the manner specified in this permit.

THIS PERMIT IS IN FORCE FROM 15 JUNE 2022 TO 30 JUNE 2026

Permit Holder:
E.E. MUIR & SONS PTY LIMITED
18 Raymond Rd
LAVERTON NORTH VIC 3026

Supplier:
UPL AUSTRALIA PTY LTD
Level 3, 70 Hindmarsh Square
ADELAIDE SA 5000

Persons who can use the product under this permit:
Persons generally.

CONDITIONS OF USE

Products to be used:
CHLORONION SC HERBICIDE
AN UNREGISTERED PRODUCT
Containing: 430 g/L CHLORIDAZON as the only active constituent.

RESTRAINT:
DO NOT apply to sandy soils with low organic matter as crop damage may result.

Directions for Use:

Crop	Pest	Rate
Baby leaf spinach Baby leaf beet	Bitter cress, blackberry night-shade, black bindweed, capeweed, chamomile, charlock, chickweed, common sowthistle, cotula, deadnettle, docks, dwarf nettle, fat hen, field poppy, groundsel, marshmallow, mustard, pale persicaria, penny cress, persicaria, pig weed, potato-weed, scarlet pimpernel, shepherd's purse, sowthistle, speedwell, spurry, stagger weed, wild radish, winter grass	6.8 L/ha

Critical Use Comments:

- Apply after sowing of direct seeded crops, but before crop emergence.
- Apply by ground-based boomspray only, in a spray volume of 300-600 L/ha.
- If soil is dry, rain or irrigation is required within 6 to 7 days for residual activity.
- DO NOT apply more than 1 application per crop.
- Use in accordance with the permit instructions. For further information, refer to the product label at **Attachment 1**. Where the product label contradicts this permit, the permit must be followed.

SPRAY DRIFT RESTRAINTS
Specific definitions for terms used in this section of the permit can be found at apvma.gov.au/spraydrift

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The advisory buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are surface temperature inversion conditions present at the application site during the time of application. These conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

- spray droplets not smaller than a MEDIUM spray droplet size category
- minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for boom sprayers') are observed.

Buffer zones for boom sprayers

Droplet size	Application rate	Boom height above the target canopy	Bystander areas
MEDIUM	6.8 L/ha	0.5 m or lower	Not required
		1.0 m or lower	25 metres
COARSE		0.5 m or lower	Not required
		1.0 m or lower	10 metres

DO NOT apply by a vertical sprayer.
DO NOT apply by aircraft.

Withholding Period:
Harvest: Not required when used as directed.
Grazing: Do not graze or cut treated crops for stock feed.

Precautions:
Re-entry statement:
DO NOT enter treated areas until spray has dried. If prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

Protections:
Protection of wildlife, fish, crustaceans and environment
Do not contaminate streams, rivers or watercourses with the chemical or used containers.

Protection of crops, native and other non-target plants
In the case of crop failure, when the land must be replanted to another crop within 4 to 8 weeks after treatment, the following tolerant crops may be planted - potatoes, onion, maize, sorghum, safflower and sunflower. The following susceptible crops should not be planted until 4 to 8 weeks have elapsed following treatment with chloridazon - cucumbers, melons, tomatoes, snap beans, lima beans, lettuce, cabbages, grapes and carrots.

Storage and Disposal:
Keep out of reach of children. Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Safety Directions:
May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and preparing spray wear cotton overalls

buttoned to the neck and wrist (or equivalent clothing) and elbow length chemical resistant gloves. When using the prepared spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). In addition, if applying by open cab wear elbow length chemical resistant gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

First Aid Instructions:
If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766.

Jurisdiction:
NSW, TAS & VIC only.

Additional Conditions:
This permit allows for the use of a product in a manner specified on the permit. Persons who wish to prepare for use and/or use products for the purposes specified in this permit must read, or have read to them, the details and conditions of this permit. The product proposed to be used under this permit will carry a label as per **Attachment 1**.

The holder of the permit must notify the APVMA of new information, including relevant information in accordance with section 161 of the Schedule to the Agricultural and Veterinary Chemicals Code Act 1994, in accordance with the obligation imposed by that section.

Supply
The supplier must supply the product in a container that complies with the requirements of section 18 of the Agricultural and Veterinary Chemicals Code Regulations. Attached to this container must be a label which is identical in content and format to the label at **Attachment 1**. A copy of this permit must be supplied with the product.

Record keeping
Maintain records of all treatments performed under this permit. Details must include the date and locations where treatments occurred, the total amount of product used, and the names and addresses of the persons undertaking the use. These details must be maintained for a minimum period of two years from the date of expiry of this permit, and must be made available to the APVMA upon request.

Issued by the Australian Pesticides and Veterinary Medicines Authority
Note: 06/10/2023 – Permit changed to minor use permit. Permit updated to add NSW. S161 reporting condition added. Permit expiry extended to 31/10/2025. Permit issued as Version 2.
Note: 25/09/2025 – Permit expiry extended to 30/06/2026. Permit issued as Version 3.

Attachment 1

CHLORONION 20L 182W X 208H MM

CHLORONION™ SC
HERBICIDE

Selective pre- and post-emergent herbicide for fodder beet, silver beet, red beet, sugar beet and mangolds, and post emergent weed control in onions and chives.

Active Ingredient:
Chloronion SC contains 430 g/litre chloridazon in the form of a suspension concentrate.

GROUP	C1	HERBICIDE

Registered pursuant to the ACVM Act 1997, No. P009469.
See www.foodsafety.govt.nz for registration conditions
Approved pursuant to the HSNO Act 1996, Approval Code HSR101123

AGRECOVERY
Rural Recycling Programme

NET CONTENTS:
20
LITRES

Registrant:
UPL New Zealand Limited
PO Box 51584, Pakuranga 2140
Auckland, New Zealand
Phone: 0800 100 325
www.upl-ltd.com/nz
UPL/H/042-20/0221

UPL
OpenAg™

WARNING. KEEP OUT OF REACH OF CHILDREN
READ THE LABEL CAREFULLY AND SHAKE WELL BEFORE USING

CHLORONION 20L 182W X 208H MM

CHLORONION SC

This product MUST NOT be used for any purpose or in any manner contrary to the controls and conditions of this registration.
Any person purchasing, handling or disposing of this product must be suitably qualified.

HSNO Classifications: 6.5B, 9.1A, 9.2A
WARNING
May cause an allergic skin reaction.
Very toxic to aquatic organisms.
Very toxic in the soil environment.

PRECAUTIONS
Store in original container, tightly closed, away from foodstuffs and under lock and key.
Avoid contact with seeds, fertilisers and other pesticides.
When handling the concentrate wear full protective clothing such as overalls, gloves and face shield. Avoid contact with skin, eyes and inhalation of spray mist.
Wash splashes of concentrate from skin or eyes immediately.
When spraying, wear overalls and gloves.
DO NOT eat, drink or smoke while using.
Wash hands and face before eating, drinking or smoking and after work.
Wash protective clothing after use.
Apply with well-maintained and calibrated spray equipment.
DO NOT carry on a passenger service vehicle.
Application, storage and record keeping should be as described in the Management of Agrichemicals NZS8409.
Avoid contamination of any water supply with chemical or empty container.

CONTAINER DISPOSAL: Ensure the container is empty. Triple rinse empty container and add rinsate to the spray tank. Recycle empty container through Agrecovery (0800 247 326, www.agrecovery.co.nz). Otherwise crush and bury in a suitable landfill. DO NOT reuse this container for any other purpose.

PRODUCT DISPOSAL: Dispose of this product only by using according to the label, or at an approved landfill or other approved facility.

FIRST AID: If swallowed do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. DO NOT scrub the skin. If splashed in eyes, wash out immediately with water. If inhaled move the victim to fresh air immediately. Begin artificial respiration if breathing has stopped.

SPILLAGE
Wear suitable protective clothing including face shield, impervious gloves and boots. Prevent the product or spilled material from entering water bodies. Absorb liquid spills with inert material such as earth or sand and place in waste containers. Wash area with water and absorb with further inert material. Dispose of safely.

GENERAL INFORMATION
CHLORONION SC is a selective pre - and post-emergence herbicide providing a good residual action against a wide range of annual broadleaf weeds and some annual grasses with a contact action on a wide range of weeds up to the 2 leaf stage. Adequate soil moisture is required for good residual activity. The post-emergent effect can be enhanced with the addition of Peptoil Crop Oil Concentrate. **CHLORONION SC** remains effective in the soil for several weeks.

RESISTANCE MANAGEMENT
CHLORONION is a pyridazinone (Group C1) herbicide. Excessive use of any herbicide may lead to the development of resistant weeds. Follow label directions and use in programmes with herbicides from a different action group. Consult http://resistance.nzpps.org/ for alternate modes of action.

SPRAY DRIFT: Beware - apply this product carefully. Spray drift may cause serious damage to desirable plants. Ensure compliance with any relevant regional air management plans.

Peptoil® is a registered trademark of Drexel Company, USA
Chloronion™ Trade Mark of UPL New Zealand Limited.

24 HOUR EMERGENCY RESPONSE
0800 CHEM CALL
0800 243 622

BATCH NO:
DOM:

SUSCEPTIBLE WEEDS		MODERATELY SUSCEPTIBLE	RESISTANT
Annual poa Black nightshade Charlock Chick weed Docks Fathen Galinsoga Groundsel	Penny cress Rayless chamomile Redroot Speedwells Water purslane Wild carrot Wild radish Willow weed	Barnyard grass Cleavers Curled Dock Fumitory Spurrey Summer grass	Couch Thistles All weeds past the 2 leaf stage

DIRECTIONS FOR USE
Apply CHLORONION sc in 300-600 litres of water per hectare. Use the higher rate on heavy clay soils or soils rich in organic matter.

FODDER BEET, RED BEET, SILVER BEET, MANGOLDS:
PRE-EMERGENCE: Apply 7 to 10.5 litres per hectare as soon as possible afterseeding and before crop emergence, to a fine, moist soil surface. If weeds are present, mix with a suitable contact herbicide. If soil is dry, rain or irrigation is required within 6 to 7 days for residual activity.

POST-EMERGENCE: Apply 7 - 9 litres per hectare to a weed free soil after the first true leaf is visible on the beet. Do not spray in the 8 days before thinning. The addition of Peptoil Crop Oil Concentrate at 1 litre per hectare will enhance the knockdown on weed up to 2 true leaf.

ONIONS AND CHIVES:

POST-EMERGENCE: Apply the **CHLORONION SC** and the Agri-IPC at equal rates for best results. Apply 1 to 2 litres **CHLORONION SC** with 1 to 2 litres of Agri-IPC (400 g/litre formulation) and 1 litre Peptoil Crop Oil Concentrate per hectare at the flag stage. Repeat the application 2 to 3 weeks later. Weeds should not have passed the 4 true leaf stage. For best contact action, no rain should fall for 6 hours after application. Rain is required, however, within 7 days for good residual activity.

CAUTION: Crop damage may occur on light soils.

MIXING: Always shake the container before use. Pour the required amount directly into the partly filled spray tank with agitation running. Complete filling of the spray tank.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Limits of Agricultural Compounds

STORAGE
Store in a cool dry place avoiding temperatures above 35°C. When stored appropriately, this product should show no significant degradation for 2 years from the date of manufacture. See Safety Data Sheet for additional storage information.

SAFETY DATA SHEET
If additional hazard or safety information is required, refer to the Safety Data Sheet. For a copy visit our website at www.upl-ltd.com/nz

EXCLUSION OF LIABILITY: PLEASE READ
To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, UPL New Zealand Limited as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

UPL/H/042-20/0221

9 421022 425755