POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Tervigo®

Nematicide

syngenta®

ACTIVE CONSTITUENT: 20 g/L ABAMECTIN

GROUP	6	INSECTICIDE

For the control of Root Knot Nematode in Fruiting Vegetables, Cucurbits, Potatoes and Sweet Potatoes as per the Directions for Use

Syngenta Australia Pty Ltd
Level 1, 2 Lyonpark Road, Macquarie Park NSW 2113

In a transport emergency dial 000, Police or Fire Brigade
For specialist advice in an emergency only, call 1800 033 111 (24 hours)

APVMA Approval No.: 80620/130037

SC	Formulation type **Suspension Concentrate**

®

DIRECTIONS FOR USE

Restraints

DO NOT apply by aircraft.
DO NOT apply using hand held equipment (including low & high pressure hand wand / gun).
DO NOT apply using spraying equipment carried on the back of the user.
DO NOT apply to waterlogged soils.
DO NOT apply if heavy rains or storms that are likely to cause runoff are forecast within 3 days.
DO NOT irrigate to the point of runoff for at least 3 days after application.

Spray Drift Restraints:

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift

DO NOT allow bystanders to come into contact with the spray cloud.
DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.
DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.
DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:
- boom height is no higher than 0.5 m;
- spray droplets not smaller than a COARSE spray droplet size category

Application rate	Boom height above the target canopy	Mandatory downwind buffer zones				
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
60 mL/100 m row (banded over the row after planting - potatoes only)	0.5 m or lower	0 metres	145 metres	60 metres	0 metres	25 metres

Crop	**Pest**	**Rate**	**Critical Comments**
Fruiting vegetables, other than cucurbits (including Tomato, Capsicum, Chilli, Eggplant)	Root-knot nematode (*Meloidogyne* spp.)	**Planting hole drench** 400 mL/ 1000 plants **Drip/trickle irrigation** 120 mL/ 100 m of row Followed by	Apply TERVIGO® as a planting hole drench via a waterwheel at transplanting or via a trickle irrigation system immediately following transplanting. Product placement is important and trickle irrigation should be positioned around the plant roots. TERVIGO® application through drip/trickle irrigation should occur in the second quarter of the irrigation cycle after the soil has been partially wetted up.

Crop	Pest	Rate	Critical Comments
Cucurbits (including Cucumber, Zucchini, Rockmelon, Melon, Watermelon, Pumpkin, Asian melon)	Root-knot nematode (*Meloidogyne* spp.)	60 to 120 mL/ 100 m of row	Follow initial transplant treatment with up to 4 in crop applications via the drip/trickle irrigation system of 60 to 120 mL/100 metres of row at 14 day intervals. Use more applications and the high rate for longer growing cycles or under moderate to high root-knot nematode pressure where nematode counts, or paddock history indicate that the population is above economic threshold and yield loss is expected. DO NOT apply more than five (5) applications per crop. Refer to **Mixing and Application** under **General Instructions** for detailed information on application methods.
Potato	Root-knot nematode (*Meloidogyne* spp.)	60 mL/100 m of row Followed by 60 mL/100 m of row	Apply as an in-furrow spray at planting. Mount the nozzle so the spray is directed into the furrow at the seed as a 15-20 cm band applied just before the seed is covered. Alternatively, use a two-nozzle system spraying the furrow prior to seed drop and after it has dropped just prior to covering. Apply TERVIGO® in 1 – 2 L of water/100 m of row. Ensure the water volume applied does not wash off any previously applied seed treatments. Apply a second application as a banded spray over the row between 14 to 30 days after planting, maximising contact with the soil. Apply sufficient overhead irrigation immediately after treatment (whilst avoiding surface runoff), to allow the product to reach the root zone. DO NOT apply TERVIGO® if conditions or poor seed quality favour seed piece breakdown (eg high soil temperature and moisture, high relative humidity, light textured soils, insect damaged seed) as the addition of any moisture to the seed may enhance the problem.

Crop	Pest	Rate	Critical Comments
Sweet potato	Root-knot nematode (*Meloidogyne* spp.)	120 mL/100 m of row Followed by 60 to 120 mL/100 m of row	Apply TERVIGO® via a drip/trickle irrigation system no later than 3-days after transplanting. Product placement is important, and the drip/trickle irrigation tape and drip points should be positioned near the transplanted runner. The TERVIGO® application through drip/trickle irrigation should occur in the second quarter of the irrigation cycle after the soil has been partially wetted up. Follow the initial transplant treatment with up to 4 applications via the drip/trickle irrigation system of 60 to 120 mL/100 meters of row at 14-day intervals. Use more applications and the high rate for longer growing cycles or under moderate to high root-knot nematode pressure where nematode counts, or paddock history indicate that the population is above economic threshold and yield loss is expected. DO NOT apply more than five (5) applications per crop. Refer to **Mixing and Application** under **General Instructions** for detailed information on application methods.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:
Tomato, Capsicum, Chilli, Eggplant, Cucurbits, Potato, Sweet potato:
NOT REQUIRED WHEN USED AS DIRECTED

GENERAL INSTRUCTIONS

Mixing

Calculate the amount of TERVIGO® required based on the area to be treated. Prepare the injection solution using a minimum dilution for injection of 500 mL of product in 1 L of water. The volume of the injection solution should not exceed what can be injected in ¼ of the total irrigation volume. DO NOT prepare more injection solution than required to dose the total area to be treated.

Shake contents well before mixing. Add the required amount of TERVIGO® directly to the spray tank and mix well. TERVIGO® mixes completely with water and should only be mixed with compatible fertilisers as it has been formulated with an iron chelate.

Application

Drip/trickle Irrigation

TERVIGO® should be applied through direct injection into drip/trickle irrigation systems. Injection equipment must be accurately calibrated to achieve the required application rates prior to treatment. TERVIGO® should be incorporated into the normal irrigation schedule at the recommended timing within the irrigation cycle. Even distribution throughout the paddock is essential. Irrigation systems should be monitored periodically to ensure all delivery points are in working order and there are no clogs or blockages. TERVIGO® is most effective when drippers are located directly beside seedlings.

The TERVIGO® injection solution (see Mixing) should be injected into irrigation systems during the second quarter of the irrigation cycle, after the soil has been partially wetted up. The duration of injection should be long enough to distribute TERVIGO® uniformly. This will vary depending on irrigation system setup and design, area of crop treated, emitter type, irrigation flow rates etc (refer to the manufacturer specifications for the irrigation system).

The final half of the irrigation cycle is used to flush product from the system and aid in the distribution of product into the plant root zone. DO NOT over irrigate. Soil moisture levels should be at or slightly below field capacity at the end of the application and irrigation cycle.

Planting Hole Drench Application
TERVIGO® should be applied in a minimum total diluted volume of 50 mL per seedling directly at the base of transplanted seedlings or into the planting hole immediately before transplanting. Actual total volume will vary depending on equipment used, pre-existing soil moisture levels and soil type but should be sufficient to ensure uniform application and distribution.

Compatibility
As formulations of other manufacturer's products are beyond the control of Syngenta Australia Pty Ltd, and water quality varies with location, all mixtures should be tested prior to mixing commercial quantities. Check compatibility prior to use and contact your local Syngenta representative for further information.

Insecticide Resistance Warning **GROUP 6 INSECTICIDE**
For insecticide resistance management TERVIGO® Nematicide is a Group 6 insecticide. Some naturally occurring nematode biotypes resistant to TERVIGO® and other Group 6 insecticides may exist through normal genetic variability in any nematode population. The resistant individuals can eventually dominate the nematode population if TERVIGO® and other Group 6 insecticides are used repeatedly. The effectiveness of TERVIGO® on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Syngenta Australia Pty Ltd accepts no liability for any losses that may result from failure of TERVIGO® to control resistant nematodes. TERVIGO® may be subject to specific resistance management strategies. For further information contact your supplier, Syngenta representative or local agricultural department agronomist.

PRECAUTION
Re-entry Period
DO NOT manually transplant fruiting vegetable and cucurbit seedlings or sweet potato runners after initial treatment unless wearing chemical resistant gloves.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT
Very toxic to aquatic life. DO NOT contaminate streams, rivers, or watercourses with the chemical or used containers.

PROTECTION OF HONEY BEES AND OTHER INSECT POLLINATORS
Highly toxic to bees. DO NOT use TERVIGO® while bees are actively foraging.

STORAGE AND DISPOSAL
Store in the closed original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.
Triple rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal at an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation, and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. May irritate the eyes. Avoid contact with the eyes. Do not inhale spray mist. When opening the container and preparing the product for use wear:

- **cotton overalls buttoned to the neck and wrist (or equivalent clothing)**
- **chemical resistant gloves**
- **half face-piece respirator with organic vapour/gas cartridge or canister**

When using the prepared product wear:

- **cotton overalls buttoned to the neck and wrist (or equivalent clothing)**

Wash hands after use. After each day's use wash gloves, respirator (if rubber wash with detergent and warm water) and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.

SAFETY DATA SHEET

If additional hazard information is required refer to the Safety Data Sheet. For a copy phone 1800 067 108 or visit our website at www.syngenta.com.au

DISCLAIMER

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability.

Product names marked ® or ™, the ALLIANCE FRAME the SYNGENTA Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company

May cause damage to organs (nervous system) through prolonged or repeated exposure

Batch No	
Date of Manufacture	

Scan here for more information including SDS and label