Product Name: FLUTE 50 EW FUNGICIDE
APVMA Approval No: 66970 / 123183

RLP
APPROVED

Label Name:	FLUTE 50 EW FUNGICIDE

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	50 g/L CYFLUFENAMID

Mode of Action:	GROUP **U6** FUNGICIDE

Statement of Claims:	A Fungicide for the control of powdery mildew in grapevines, cucurbits, strawberries and strawberry runners and plug plants as per the Directions for Use Table.

Net Contents:	1L, 2.5L, 5L, 10L, 20L

Restraints:	DO NOT apply by air as uneven coverage is likely to result. DO NOT apply when wind speed is less than 3 or more than 20 km/hr at the application site. Except when applying with vineyard airblast equipment, DO NOT apply with spray droplets smaller than a MEDIUM spray droplet size category according to nozzle manufacturer specifications that refer to the ASAE S572 Standard or the British Crop Protection Council guidelines. DO NOT apply during surface temperature inversion conditions at the application site. Users of this product MUST make an accurate written record of the details of each spray application within 24 hours following application and KEEP this record for a minimum of 2 years. The spray application details that must be recorded are: 1. date with start and finish times of application; 2. location address and paddock/s sprayed; 3. full name of this product; 4. amount of product used per hectare and number of hectares applied to; 5. crop/situation and weed/pest; 6. wind speed and direction during application; 7. air temperature and humidity; 8. nozzle brand, type, spray angle, nozzle capacity and spray system pressure measured during application; 9. name and address of person applying this product. (Additional record details may be required by the state or territory where this product is used.)

Directions for Use:	This section contains file attachment.

Other Limitations:	

Withholding Periods:	GRAPES (DOMESTIC): DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION GRAPES (FOR WINE INTENDED FOR EXPORT): CONTACT AGNOVA TECHNOLOGIES PTY LTD OR CONSULT YOUR WINERY OR THE AUSTRALIAN WINE RESEARCH INSTITUTE (AWRI) FOR THE RECOMMENDED WITHHOLDING PERIOD. CUCURBITS: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION STRAWBERRIES: NIL

Trade Advice:	EXPORT OF TREATED PRODUCE Growers should note that MRLs or import tolerances do not exist in all markets for fruit treated with Flute 50 EW Fungicide. Fruit: If growing fruit for export (either fresh or dried) consult your peak industry body or AgNova Technologies Pty Ltd for the latest information on MRLs and import tolerances. Wine: Flute 50 EW Fungicide should not be applied later than EL31 [berries pea-size] when grapes are to be used to make wine for export. Contact your winery or the Australian Wine Research Institute (AWRI) at www.awri.com.au for the updated information on MRLs and overseas import tolerances BEFORE using Flute 50 EW Fungicide.

General Instructions:	This section contains file attachment.

Resistance Warning:	FUNGICIDE RESISTANCE WARNING Flute 50 EW Fungicide is a member of the phenyl-acetamide group of fungicides. For fungicide resistance management, the product is a Group U6 fungicide. Some naturally occurring individual fungi resistant to Flute 50 EW Fungicide and other Group U6 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Flute 50 EW Fungicide or other Group U6 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, AgNova Technologies Pty Ltd accepts no liability for any losses that may result from the failure of Flute 50 EW Fungicide to control resistant fungi. Refer to Directions for Use table and published CropLife Resistance Management Strategies www.croplifeaustralia.org.au

Precautions:	RE-ENTRY PERIOD Do not allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. Clothing must be laundered after each day's use. For glasshouses and other enclosed areas, do not allow re-entry until spray had dried or dissipated and enclosed area has been thoroughly ventilated.

Protections:	PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

	DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Very toxic to aquatic life. DO NOT contaminate wetlands or water course with this product or used containers.

Storage and Disposal:	Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Safety Directions:	Will irritate the skin. May irritate the eyes. Avoid contact with the eyes and skin. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist [or equivalent clothing] and elbow-length chemical resistant gloves. If product on skin, immediately wash area with soap and water. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

First Aid Instructions:	If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 13 11 26.

First Aid Warnings:	

CROPS	DISEASE	RATE	WHP	CRITICAL COMMENTS
Grapevines	Powdery mildew (*Erysiphe necator*)	*Dilute spraying*: 25 to 35 mL/100 L water *Concentrate spraying*: Refer to the Mixing/ Application section	5 weeks	Apply before first sign of disease or when conditions (varietal and climatic) are conducive to disease development as part of a disease management program. Apply one to two applications from E-L 18 (caps changing colour) to E-L 31 (berries pea-size) based on the full dilute spray volume matched with the size of the target canopy. Use a spray interval of 10 to 14 days in a program of effective powdery mildew fungicides to ensure that new growth is protected. Use the shorter interval during periods of rapid vine growth. A rate of 25 mL/100 L is applicable in most situations. If conditions favour high levels of infection or if powdery mildew levels were significant in the previous season, use the higher rate. A rate of 35 mL/100 L may be used with a spray interval up to 21 days provided disease pressure is low and before rapid vine growth or once vine growth has slowed. Apply by dilute or concentrate spraying equipment. Apply in sufficient water volume to obtain thorough coverage of the crop. The required dilute spray volume will increase and the sprayer setup and operation may also need to be changed, as the crop grows.  As an example, a 1 m x 1 m canopy (refer diagram) with a 3 m row spacing requires a true dilute water volume of around 1000 L/ha when using dilute application. In this case the dose will be 350 mL/ha when using a rate of 35 mL/100 L. For the same canopy, if using concentrate application equipment achieving good coverage applying 500 L/ha, then the concentrate factor is 2X. This means a rate of 70 mL/100 L. The dose will remain at 350 mL/ha. For further advice on application, contact AgNova. Refer to the section on concentrate spraying if low spray volumes are to be used. Apply the same amount of product to the target crop whether applying by dilute or concentrated spray volumes. Do not apply at concentrations greater than 3X. Do not apply more than 2 (two) Flute 50 EW sprays per season, and follow the CropLife Australia resistance management guidelines.

cont'd

Cucurbits	Powdery mildew (*Podosphaera xanthii*)	250 mL/ha or *Glasshouse:* 25 mL/100 L	1 day	Apply before first sign of disease or when conditions are conducive to disease as part of a disease management program. Apply sprays at 7 to 10 day intervals. Use the shorter intervals under conditions favouring disease infection. Glasshouse: To ensure thorough spray coverage of the target crop, apply to the point of run-off. The required dilute spray volume will change and the sprayer setup and operation may also need to be changed, as the crop grows. For resistance management: Do not apply Flute 50 EW Fungicide as consecutive sprays or apply more than 2 sprays per crop.
Strawberries (field and protected cropping)	Powdery mildew (*Podosphaera aphanis*)	*Dilute spraying*: 35 mL/100 L Use when spray volume exceeds 1000 L/ha OR 350 mL/ha Use when spray volume is less than 1000 L/ha Do not apply volumes less than 500 L/ha	Nil	Reduce background levels of disease by removing infected plant debris and/or rotted fruit throughout the season. A fungicide program for the control of powdery mildew should commence early before disease appears. Apply Flute 50 EW before first sign of disease or when conditions are conducive to disease development as part of a disease management program. Apply sprays at 7 to 10 day intervals. Use the shorter intervals under conditions favouring disease infection. To ensure thorough spray coverage of the crop, apply to the point of run-off targeting foliage, flowers and fruit. The required dilute spray volume will change and the sprayer setup and operation may also need to be changed, as the crop grows. Do not apply Flute 50 EW as consecutive sprays or apply more than 2 sprays per fruiting season, and follow the CropLife Australia resistance management guidelines.
Strawberry runners and plug plants		*Dilute spraying:* 350 mL/ha Do not apply volumes less than 500 L/ha.		Apply a maximum of 2 sprays per cropping season with at least 2 sprays of products from different Mode of Action Groups between them. In addition, a minimum of 2 sprays from different Mode of Action Groups should be made between the last Flute 50 EW spray and digging. Apply in a spray volume of 600 - 800 L water per hectare.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

MIXING/APPLICATION

Add the required quantity of Flute 50 EW Fungicide to a partly filled spray tank and agitate. Complete filling while agitating.

Ensure the spray mixture is properly agitated before restarting after stoppage.

Grapevines

Dilute Spraying

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of runoff. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer setup and operation may also need to be changed as the crop grows.

Concentrate Spraying

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of runoff) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:

EXAMPLE ONLY

1. Dilute spray volume as determined above: For example 1000 L/ha
2. Your chosen concentrate spray volume: For example 500 L/ha
3. The concentration factor in this example is: 2 X (i.e.1000 L ÷ 500 L = 2)
4. If the dilute label rate is 35 mL/100 L, then the concentrate rate becomes 2 x 35, that is 70 mL/100 L of concentrate spray.

- The chosen spray volume, amount of product /100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.
- Do not use a concentration rate higher that that specified in the Critical Comments
- When concentrate spraying, apply using a minimum spray volume of 300L/ha or a maximum concentration factor of 3X, whichever is the greater spray volume.

Example 1. The true dilute volume is 1200 L/ha, therefore the minimum spray volume is 400 L/ha (3X)

Example 2. The true dilute is 750 L/ha, therefore the minimum spray volume is 300 L/ha (2.5X)

- For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.