RLP APPROVED

Product Name: Campbell Fludy 230SC Fungicide

APVMA Approval No.: 83843 / 124061

Label Name:	Campbell Fludy 230SC Fungicide

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	230 g/L FLUDIOXONIL

Mode of Action:	GROUP 12 FUNGICIDE

Statement of Claims:	Controls certain post-harvest diseases in citrus, stone fruit, pome fruit, kiwi fruit, pomegranates, chestnuts, pineapple and mangoes.

Net Contents:	10L 1L 5L

Restraints:	

Directions for Use:	This section contains file attachment.

Other Limitations:	

Withholding Periods:	WITHHOLDING PERIODS: NOT REQUIRED WHEN USED AS DIRECTED

Trade Advice:	Growers should note that Maximum Residue Limits (MRLs) or import tolerances may not exist in all markets for edible produce treated with Campbell Fludy 230SC Fungicide. If you are growing edible produce

	for export, please check with Colin Campbell (Chemicals) Pty Ltd for the label information on MRLs and import tolerances before using this product.

General Instructions:	GENERAL INSTRUCTIONS: Campbell Fludy 230SC Fungicide is a protective fungicide used as a postharvest treatment to control certain postharvest diseases. Campbell Fludy 230SC Fungicide may be applied as a post-harvest dip, drench, flood or spray in citrus and mangoes, or as a dip or drench in pome fruit, pomegranates, stone fruit and kiwi fruit. Note: Campbell Fludy 230SC Fungicide may be degraded by exposure to direct sunlight. Treated fruit should not be stored in direct sunlight. MIXING: Clean mixing and spraying equipment before use. Prepare no more treatment solution than is needed for the immediate operation. Shake the container before use to ensure uniform dispersion of contents before measuring quantity required. Half-fill the spray or mixing tank with clean water or wax/oil emulsion (or aqueous dilution of a wax/oil emulsion) and start agitation. While filling the remainder of the spray tank add the required amount of Campbell Fludy 230SC Fungicide. Begin application of the solution after Campbell Fludy 230SC Fungicide has completely and uniformly dispersed into the mix carrier. Maintain agitation throughout the application operation. DO NOT let the treatment mixture stand overnight in the tank. Flush the spray equipment thoroughly after each use. APPLICATION: Dip and High Volume Drench Application: Citrus, kiwi fruit, pome fruit, pomegranates and stone fruit: Dip fruit in prepared solution for 30 to 60 seconds and allow fruit to drain. Ensure fruit is in complete contact with the dipping solution. Mangoes: Apply by hot dip or hot flood spray. Time of exposure: 5 minutes at 52°C. Dip and Low Volume Concentrate spray application. Chestnuts: Dip in prepared solution or spray with an application time one (1) minute and allow produce to drain and dry thoroughly prior to storage. Ensure thorough and even coverage of chestnuts in the dip or by spray application. High Volume Application: Mix appropriate volume of Campbell Fludy 230SC Fungicide (see Directions for Use) in water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion for the fruit being treated. Use T-jet, flooders, or similar application system. Fruit should be treated for approximately 30 seconds. Ensure all parts of fruit are well covered by mixture. Low Volume Application: Citrus: Apply by low volume application. Use properly calibrated low volume application equipment. Ensure all parts of fruit are well covered by the spray solution and application time is at least 30 seconds. Mix the required amount of Campbell Fludy 230SC Fungicide in 5 to 15 L of water or wax per 10,000kg fruit. Mangoes: Apply by low volume non-recirculating ambient spray, exposing fruit for 30 seconds. The use of brush rollers may improve efficacy. COMPATIBILITY: Campbell Fludy 230SC Fungicide Fungicide can be mixed with Vorlon, imazalil (e.g. Magnate), calcium hypochlorite and DPA. A reduction in efficacy may occur when Campbell Fludy 230SC Fungicide is applied with DPA at DPA rates above 500ppm. If tank mixes are to be used observe all directions, precautions and limitations on all products to be used. As formulations of other manufacturers' products are beyond the control of Colin Campbell Chemicals, and the quality of water may vary with location, all mixtures should be tested prior to mixing commercial quantities. CLEANING EQUIPMENT: Rinse application equipment with water after use. Spread rinsings over flat land away from desirable vegetation, waterways and drainage, and follow Dip Disposal instructions.
Resistance Warning:	GROUP 12 FUNGICIDE Campbell Fludy 230SC Fungicide Fungicide is a member of the phenylpyrrole group of fungicides. For fungicide resistance management. Campbell Fludy 230SC Fungicide is a Group 12 fungicide. Some naturally occurring individual fungi resistant to Campbell Fludy 230SC Fungicide and other Group 12 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Campbell Fludy 230SC Fungicide and other Group 12 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to

	use, Colin Campbell (Chemicals) Pty Ltd accepts no liability for any losses that may result from the failure of Campbell Fludy 230SC Fungicide to control resistant fungi.

Precautions:	PRECAUTIONS: Re-handling: DO NOT rehandle treated produce until the product has been allowed to dry, unless wearing chemical resistant gloves.

Protections:	PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT: Dangerous to fish and other aquatic organisms. This product is very toxic to aquatic invertebrates. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

Storage and Disposal:	STORAGE AND DISPOSAL Keep out of reach of children. Store in the closed original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray or dip tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product. DIP DISPOSAL Unused or spent dip should be disposed of carefully to avoid contamination of streams, rivers or waterways. Dispose of dip in an authorised dip disposal facility.

Safety Directions:	SAFETY DIRECTIONS: May irritate the eyes. Harmful if inhaled. Repeated exposure may cause allergic disorders. Avoid contact with the eyes. DO NOT inhale spray mist. When opening the container and preparing the product for use, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). If applying by hand wear half facepiece respirator with organic vapour/gas cartridge or canister. Change canister if odour or taste of pesticide is noticed (a maximum of 8 hours of use is recommended). Wash hands after use. After each days use, wash respirator and, if rubber, wash with detergent and warm water, and contaminated clothing.

First Aid Instructions:	FIRST AID: If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

First Aid Warnings:	

CROP	PEST	RATE		CRITICAL COMMENTS
		Dip and high volume	Low volume	
Citrus	Blue mould *(Penicillium italicum)* Green mould *(Penicillium digitatum)* Diplodia Stem End Rot	260 to 520mL in 100L water	87 to 174mL per 10,000kg fruit applied in 5 to 15L water	Raising the temperature of the prepared aqueous solution to a maximum of 50°C for up to 1 minute may improve the efficacy of this product. Avoid long heat exposure on fruit with a sensitive rind. Applied alone Fludy should be used in alternation with other products for easily protected fruit and when disease pressure is expected to be low to moderate. Under high disease pressure, Fludy should only be used with other postharvest fungicides (see Compatibility section). Refer to application section below for application instructions.
Kiwi fruit	Grey mould *(Botrytis cinerea)*	130 to 260mL in 100L water	-	Refer to application section below for application instructions.
Pome fruit	Blue mould (*Penicillium expansum* and *P. solitum*) Grey mould *(Botrytis cinerea)*	130 to 260mL in 100L water	-	Refer to application section below for application instructions.
Stone fruit – Except apricots and Peaches	Brown rot *(Monolinia* spp), Grey mould *(Botrytis cinerea)*, Rhizopus rot *(Rhizopus stolonifer)*	130 to 260mL in 100L water	-	Refer to application section below for application instructions.
Apricots and Peaches		130mL in 100L water	-	
Mangoes	Anthracnose *(Colletotrichum gleosporoides)* Stem end rot and Dendritic spot *(Botrosphaeria* family*)* and includes *Fusicoccum parvum, Dothiorella dominicana, Lasiodiplodia theobromae* and *Phomopsis mangiferae)*	60 to 120mL in 100L water (52°C for 5 minutes)	-	Hot dip or flood spray. Use the low rate for low disease pressure and the high rate for high disease pressure. Refer to application section below for application instructions.
	Anthracnose *(Colletotrichum gleosporoides)*	-	260mL /100L	Low volume non-recirculating spray only.
Pomegranates	Botrytis fruit rot *(Botrytis cinerea)*	260mL in 100L water	-	Dip or flood spray. Fruit should be dipped or drenched under high volume application for 30 seconds. The dip or flood spray may be heated to 49°C to further increase efficacy.
Chestnuts	Suppression of surface moulds and fungal rots	-	Dip or low volume spray 130 mL / 100 L water	Product may be applied as a post-harvest dip or low volume concentrate spray in chestnuts Ensure that nuts are clean of soil debris prior to treatment.
Pineapple	Postharvest moulds (*Penicillium* spp.)	230 mL product / 100 L water	-	Apply treatment immediately after harvest. Ensure thorough and even coverage.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.