POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Access®

HERBICIDE

ACTIVE CONSTITUENT: 240 g/L TRICLOPYR present as the butoxyethyl ester
120 g/L PICLORAM present as the isooctyl ester
SOLVENT: 389 g/L LIQUID HYDROCARBON

GROUP	4	HERBICIDE

For selective control of a wide range of woody and noxious weeds in commercial and industrial areas, public lands, fence lines and pastures, by basal bark and cut stump applications as specified in the Directions for Use.

Pack Sizes: 1 L, 2L, 5L, 10L, 20L, 100L

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre.
Phone: *Australia* 13 11 26.
If in eyes, hold eyes open, flood with water for at least 15 minutes and see a Doctor.

EMERGENCY RESPONSE (ALL HOURS)
RING FROM ANYWHERE IN AUSTRALIA
1800 370 754
(LOCAL CALL FEE ONLY)

IN A TRANSPORT EMERGENCY ONLY
DIAL 000
FOR POLICE OR FIRE BRIGADE

SAFETY DIRECTIONS

- Harmful if swallowed.
- Will irritate the eyes, nose, throat and skin.
- Avoid contact with eyes and skin.
- Repeated exposure may cause allergic disorders.
- Avoid inhaling vapour or spray mist.
- When opening the container and preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrists, a washable hat, and elbow-length neoprene gloves and a face shield or goggles.
- If product on skin, immediately wash area with soap and water.
- If product in eyes, wash it out immediately with water.
- After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.
- After each day's use wash gloves, face shield or goggles and contaminated clothing.

SAFETY DATA SHEET

Additional information is listed on the Safety Data Sheet for **ACCESS® HERBICIDE** which is available from Corteva Agriscience on request. Call Customer Service Toll Free on 1-800 700 096 or visit www.corteva.com.au

CORTEVA™ agriscience

™ ® Trademarks of Corteva Agriscience and its affiliated companies.

Visit us at Corteva.com.au

RESTRAINTS

- **DO NOT** add water to Access® Herbicide.
- **DO NOT** apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition, presence of disease, or previous herbicide treatment as reduced levels of control may result.
- **DO NOT** apply to charcoal coated or wet stems as this can repel the diesel mixture.

DIRECTIONS FOR USE

See GENERAL INSTRUCTIONS - APPLICATION section for application method details**.**

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FENCE LINES, FORESTRY, PASTURES AND RIGHTS-OF-WAY.			
WEEDS CONTROLLED	**WEED GROWTH STAGE**	**RATE / 60 L DIESEL[2]**	**CRITICAL COMMENTS**
Acacia spp. including: Black wattle[1] Brigalow[1] Hickory wattle[1] Mimosa bush Mulga[1] Sally wattle[1] Silver wattle[1]	**Basal bark method:** Plants with stems up to 5 cm basal diameter.	1 L	Ensure all stems on multi-stemmed varieties are treated. Delay treatment until all regrowth on bulldozed or ploughed land has had time to emerge to a height of at least one metre. Root suckering wattles will need follow-up treatment.
African boxthorn	**Cut Stump method:** Plants up to and in excess of basal bark size. Apply immediately after cut is made.		Treat all stems on multi-stem plants.
Algaroba (*Prosopis* spp.)			
Angophora spp. regrowth			
Australian blackthorn			This species may become dormant so should not be treated during winter.
Bitterbark			Regrowth from roots may occur.
Broadleaf privet		2 L	
Brush box			Treat single stemmed seedling plants only.
Cacti including: Common pest pear Harrisia cactus Prickly pear Snake cactus Tiger pear Tree pear		1 L	Apply as an overall spray, wetting all areas of the plant to ground level.
Calotrope			Species with rough corky bark require more thorough wetting than smooth barked species. Some regrowth may still occur.
Cockspur thorn[1]			
Corkwood wattle			Basal bark application only. Species with old, rough bark require more thorough wetting than smooth barked species. Some regrowth may still occur.
Corymbia spp. [1]			
Ellangowan poison bush			
Eucalyptus spp. regrowth			
False sandalwood			
Green cestrum			
Grevillea spp.			
Groundsel bush			
Guava		2 L	Regrowth may occur from roots of the main tree.
Hard milkwood		1 L	
Hawthorn			
Heartleaf poison bush			
Honey locust			

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FENCE LINES, FORESTRY, PASTURES AND RIGHTS-OF-WAY.

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE / 60 L DIESEL[2]	CRITICAL COMMENTS
Kiteleaf poison	**Basal bark method:** Plants with stems up to 5 cm basal diameter. **Cut Stump method:** Plants up to and in excess of basal bark size. Apply immediately after cut is made.	1 L	
Leucaena[1]			
Limebush			Treat single stem plants only.
Lantana			
Narrow leaf hop bush			
Needlewood			
Oleander			
Parkinsonia[1]			
Punty bush			
Red ash (white myrtle)			
Rubbervine			
Swamp box		2 L	Treat single stemmed seedling plants only.
Sweet briar		1 L	Do not treat during winter.
Tea trees			
Turpentine			
Turpentine bush			
Whitewood			
Wild orange			Basal bark treatment only.
Wild peach			
Wild Rosemary			
Wild tobacco tree			Cut stump treatment only.
Wilga			
Yellow bells (*Tecoma* spp.)			Some root suckering may occur.
Camphor laurel	**Basal bark method:** Plants with stems up to 10 cm basal diameter. **Cut stump method:** Plants in excess of basal bark size. Apply immediately after cut is made.	1 L	Species that become dormant should not be treated during winter.
Peppercorn tree (*Schinus molle*)			Basal bark method only.
Prickly acacia			
Sisal hemp			Basal bark method only.
Tree of Heaven			Species that become dormant should not be treated during winter.
Willow		4 L	Cut stump method only. Complete control may not occur due to the multi-stem growth of plant and difficulty treating all stems.
Chinee apple	**Basal bark method:** Plants with stems up to 15 cm basal diameter. **Cut stump method**: Plants up to and in excess of basal bark size. Apply immediately after cut is made.	1 L	

[1] The smooth stemmed species *Acacia* spp. (excluding Mimosa bush), Cockspur thorn, *Corymbia* spp., Leucaena and Parkinsonia can be treated using the alternative Access® ThinLine method for basal bark application. Please refer to the following table.

[2] Biosafe™ may be used as an alternative carrier to diesel. For alternative biodiesel products, first seek advice from Corteva Agriscience.

ACCESS® THINLINE BASAL BARK APPLICATION FOR SMOOTH STEMMED SPECIES.

See GENERAL INSTRUCTIONS - APPLICATION section for application method details**.**

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FENCE LINES, FORESTRY, PASTURES AND RIGHTS-OF-WAY.			
WEEDS CONTROLLED	**WEED GROWTH STAGE**	**RATE / 10 L DIESEL[2]**	**CRITICAL COMMENTS**
Acacia spp. including: Black wattle Brigalow Hickory wattle Mulga Sally wattle Silver wattle	**Basal bark method:** Plants with stems up to 5 cm basal diameter.	1 L	**Spray the bark around the stem from ground level up to 5 cm high.** Ensure all stems on multi-stemmed varieties are treated. Delay treatment until all regrowth on bulldozed or ploughed land has had time to emerge to a height of at least one metre. Root suckering wattles will need follow-up treatment.
Cockspur thorn			
Corymbia spp.			
Leucaena			
Parkinsonia			
[2] Biosafe™[1] may be used as an alternative carrier to diesel. For alternative biodiesel products, first seek advice from Corteva Agriscience.			

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

The use pattern of this product is such that no withholding periods are required.

GENERAL INSTRUCTIONS

MIXING

Access® herbicide is designed for use with diesel distillate only. It will NOT mix with water.

Quarter-fill the spray unit or mixing container with diesel and add the required amount of Access®. Add the remaining diesel and shake or agitate thoroughly to mix the contents. Only mix sufficient chemical for each day's work, mixing periodically.

Dilution Table:

Rate	**Volume of Access® Herbicide**			
1 L / 60 L	17 mL	83 mL	167 mL	250 mL
2 L / 60 L	33 mL	167 mL	333 mL	500 mL
4 L / 60 L	67 mL	333 mL	667 mL	1 L
1 L / 10 L	100 mL	500 mL	1 L	1.5 L
Volume of diesel added after Access®	**fill to 1 L**	**fill to 5 L**	**fill to 10 L**	**fill to15 L**

COMPATIBILITY:

- Access® is compatible with diesel and Biosafe™[1] and should not be used or diluted with any other chemicals. For alternative biodiesel products, first seek advice from Corteva Agriscience.

GENERAL APPLICATION:

- Apply only with hand-directed equipment such as a pneumatic sprayer. The use of diesel as a herbicide carrier may affect the rubber seals in some sprayers. To avoid this, use sprayers which use Viton®[1] seals and fittings.
- Spray equipment should be used at low pressures, up to 200 kPa, to avoid splashing.
- Ensure all stems on multi-stemmed varieties are treated. Delay treatment until all regrowth on bulldozed or ploughed land has had time to emerge to a height of at least 1 metre.
- Species that lose their leaves e.g. sweet briar; or become dormant, e.g. Australian blackthorn or camphor laurel, should not be treated during winter.
- Species with old, rough bark require more thorough wetting than the smooth barked species.
- **DO NOT** apply to charcoal coated or wet stems as this can repel the diesel mixture.

Basal Bark Application Method

- Dilute the indicated volume of Access® Herbicide with an appropriate carrier up to a total volume of 60 L. Liberally spray the bark around the stem from ground level to 30 cm high, wetting thoroughly to the point of runoff (unless otherwise stated).
- Access® ThinLine basal bark application method:

  Alternatively, the application band can be reduced to 5 cm high but with the concentration increased proportionally (one part Access® diluted with 9 parts diesel) for the treatment of the smooth stemmed woody weed species: *Acacia* spp. (excluding Mimosa bush), Cockspur thorn, *Corymbia* spp., Leucaena and Parkinsonia, as noted in the Directions for Use tables.

Cut Stump Method

- Stems should be cut less than 15 cm above ground level.
- Immediately apply Access®/carrier mixture liberally to the freshly cut stump by spraying the cut surface and sides of the stem.

CLEANING SPRAY EQUIPMENT

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.

- After completion of spraying, use a degreaser such as Caltex Kwik-D-Grease®[1] to remove traces of the carrier from the sprayer. Rinse tank and spray through nozzles thoroughly with water to remove degreaser.

  After the above, quarter fill the tank with clean water and add an alkali detergent, (e.g. liquid SURF®[1], OMO®[1], DRIVE®[1] at 500 mL / 100L of water or the powder equivalent at 500 g / 100L of water). If using a concentrated laundry detergent use 250 g (or mL) / 100 L water. Do not use chlorine based cleaners.
- Shake the water around the sprayer and spray the solution through the nozzles to thoroughly clean the lines and nozzles. Rinse well with clean water to remove detergent.
- To clean brushes and container, spray liberally with degreaser. Hose off thoroughly with water and repeat using detergents (see above).
- **DO NOT** use this equipment for any other purpose.

RESISTANT WEEDS WARNING

GROUP	4	HERBICIDE

Access® Herbicide is a member of the pyridines group of herbicides. The product has the disrupters of plant cell growth mode of action. For weed resistance management the product is a Group 4 Herbicide. Some naturally occurring weed biotypes resistant to the product and other disrupters of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group 4 Herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Corteva Agriscience Australia Pty Ltd accepts no liability for any losses that may result from the failure of Access® Herbicide to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local Corteva Agriscience representative.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

• DO NOT contaminate streams, rivers or waterways with the chemical or used containers.
• Alongside waterways treat only noxious weeds and poisonous plants.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

- Treat target weeds only. DO NOT allow mixture to contact non-target species.
- DO NOT use on land to be cultivated for growing sensitive plants. Legumes, vines, vegetables, cotton, tomatoes, ornamentals, coniferous species and many other broadleaf plants are highly susceptible to Access® during both growing and dormant periods.
- Picloram, one of the active ingredients in this product, can remain active in the soil for extended periods depending on soil type, rainfall, temperature, humidity, soil moisture and soil organic matter.
- DO NOT move soil, which has been sprayed, to areas where desirable plants are to be grown.
- DO NOT apply when the treated soil may be washed to areas growing, or to be planted with desirable plants.
- DO NOT apply close to, or on areas containing roots of desirable plants, or sites where surface water from heavy rain can be expected to run off to areas containing, or to be planted to, susceptible crops or plants.

FOR NATIVE VEGETATION

Use of Access® on native vegetation must be done in accordance with STATE and/or LOCAL legislation.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Store locked up.

DO NOT store for prolonged periods in direct sunlight.

DO NOT store near food, feedstuffs, fertilisers or seed.

<u>1L pack size:</u>

Triple-rinse containers with diesel before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. The cap should not be replaced but may be taken separately.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury

the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

5 L, 10 L, 20 L and 100 L pack sizes:

This container can be recycled if it is clean, dry, free of visible residues and has the ***drum****MUSTER* logo visible. Triple rinse container before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any ***drum****MUSTER* collection or similar container management programme site. The cap should not be replaced but may be taken separately.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

SPILL AND LEAK MANAGEMENT

Do not touch or walk through spilled material. Wear a face shield or goggles, overalls buttoned to neck and wrist, chemical resistant gloves and footwear. Stop leak when safe to do so. Dam area and prevent entry into waterways and drains.

Small spills/leaks: Absorb with material such as sand, soil or sawdust. Collect spilled product and place in sealable container for disposal. Spill residues may be cleaned using water and detergent. Contain and absorb wash water for disposal. Absorb and collect washings and place in the same sealable container for disposal. Dam the area of large spills and report them to Corteva Agriscience Emergency Services at 1-800 370 754.

APVMA Approval No.: 46640/147012

Corteva Agriscience Australia Pty Ltd A.B.N. 24 003 771 659
67 Albert Avenue, Chatswood, NSW 2067
www.corteva.com.au
CUSTOMER SERVICE TOLL FREE 1-800 700 096

® Trademarks of Corteva Agriscience and its affiliated companies. © 2024 Corteva.
®1 / TM1 Registered Trademark of Other Companies.

Barcode
for stock
identification

Hazard and precautionary statements according to classification under GHS (Globally Harmonised System of Classification and Labelling)

Store locked up. Combustible liquid. Keep away from flames, hot surfaces – no smoking. In case of fire use water fog or fine spray, dry chemical, carbon dioxide, foam or alcohol resistant foams (ATC type). Suspected of causing cancer*. If exposed or concerned: Get medical advice/attention. May cause damage to organs through prolonged or repeated exposure. Contaminated clothing should not be allowed out of the workplace. May be fatal if swallowed and enters airways. If swallowed immediately call a POISONS CENTER or doctor. DO NOT induce vomiting.

*this classification is derived from naphthalene, an ingredient in the hydrocarbon solvent.