Product Name: Ubeniq Rinskor active Herbicide

APVMA Approval No: 86450/130952

RLP
APPROVED

Label Name:	Ubeniq Rinskor active Herbicide

Signal Headings:	READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	300 g/L FLORPYRAUXIFEN BENZYL

Mode of Action:	GROUP 4 HERBICIDE

Statement of Claims:	A suspension concentrate formulation for in-flood control of sedge and broadleaf weeds in rice, caltrop in sorghum and fallow and Navua sedge in pasture as specified in the DIRECTIONS FOR USE.

Net Contents:	500 mL - 20 L

Restraints:	This section contains file attachment.

Directions for Use:	This section contains file attachment.

Other Limitations:	

Withholding Periods:	PASTURE

	DO NOT GRAZE OR CUT TREATED AREAS FOR STOCK FEED FOR 3 DAYS AFTER APPLICATION. FALLOW DO NOT GRAZE OR CUT TREATED AREAS FOR STOCK FEED FOR 1 DAY AFTER APPLICATION. RICE HARVEST: NOT REQUIRED WHEN USED AS DIRECTED. GRAZING: DO NOT GRAZE OR CUT TREATED CROPS FOR STOCK FEED FOR 7 DAYS AFTER APPLICATION. SORGHUM HARVEST: NOT REQUIRED WHEN USED AS DIRECTED. GRAZING: DO NOT GRAZE OR CUT TREATED CROPS FOR STOCK FEED FOR 1 DAY AFTER APPLICATION.

Trade Advice:	LIVESTOCK DESTINED FOR EXPORT MARKETS When Ubeniq® is used as directed and the above WITHHOLDING PERIOD is observed, livestock commodities are considered acceptable for export. When using Ubeniq® in a tank mix with another product, observe whichever HARVEST or GRAZING/STOCKFOOD WITHHOLDING PERIOD that is the longer of the products used.

General Instructions:	This section contains file attachment.

Resistance Warning:	GROUP 4 HERBICIDE Ubeniq® Rinskor® active Herbicide contains a member of the pyridine-carboxylate (arylpicolinate) group of herbicides. The product has the disrupters of plant cell growth mode-of-action. For weed resistance management, the product is a Group 4 herbicide. Some naturally occurring weed biotypes resistant to Ubeniq® and other Group 4 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds may not be controlled by Ubeniq® or other Group 4 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Corteva Agriscience Australia Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds. Strategies to minimize the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, Rice Crop Protection Guide (NSW DPI) or the CropLife website (www.croplifeaustralia.org.au). In rice, a follow-up treatment of Basagran®1 M60 for broadleaf weeds or Barnstorm® for grass weeds is recommended where needed to give a second mode of action, to limit the likelihood of weed survival and onset of resistance. To reduce the risk of weed resistance, it is recommended to use two (2) or more mode of action treatments for rice weed management in any season and follow Integrated Weed Management techniques, which include use of a foundation treatment at planting, as outlined in the Rice Crop Protection Guide (NSW DPI). Foundation treatment at planting is essential for effective rice weed management in either water or drill seeded rice.

Precautions:	RE-ENTRY PERIOD Not required.

Protections:	PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto non-target vegetation. Crops highly susceptible to spray drift include, but are not limited to, grain legumes (summer or winter), millets (Echinochloa spp), lucerne, pasture legumes, cotton, citrus and other fruit, nut trees, hops, ornamentals, potatoes, safflower, beets, sunflower, tobacco, tomatoes, poppies, peppercorn trees, all vegetables and vines. Contact your Corteva Agriscience representative for more information. Refer to CROP ROTATION RECOMMENDATIONS for crop rotation information and the list of highly susceptible crops above. PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Ubeniq® is very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers. PROTECTION OF LIVESTOCK • DO NOT graze or cut treated crops or plants for stock food except as specified under withholding periods. • Poisonous plants may become more palatable after spraying and stock should be kept away from these plants until they have died down.

Storage and Disposal:	• KEEP OUT OF REACH OF CHILDREN. • Store in the closed, original container in a securely locked, dry, cool, well-ventilated place, out of direct sunlight. • DO NOT store near food, feedstuffs, fertilisers or seed. Less than 1 L Do not re-use containers. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product. 1 L - 20 L Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product. This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank. Do not dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be taken separately. SPILL AND LEAK MANAGEMENT

	Do not touch or walk through spilled material. Wear a face shield or goggles, overalls buttoned to neck and wrist, chemical resistant gloves and footwear. Stop leak when safe to do so. Dam area and prevent entry into waterways, and drains. Small spills/leaks: Absorb with material such as sand, soil or sawdust. Collect spilled product and place in sealable container for disposal. Spill residues may be cleaned using water and detergent. Contain and absorb wash water for disposal. Absorb and collect washings and place in the same sealable container for disposal. Dam the area of large spills and report them to Corteva Agriscience Emergency Services at 1-800 370 754. .

Safety Directions:	May irritate the eyes, avoid contact with eyes. When using together with other products, consult their safety directions. Wash hands after use.

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone: Australia 13 11 26.

First Aid Warnings:	

RESTRAINTS

DO NOT apply to crops or weeds which may be stressed (not actively growing) due to a range of factors including, but not limited to prolonged periods of extreme cold, sustained high temperatures, moisture stress (water-logged or drought affected), poor nutrition (including deficiency and trace element toxicity), presence of disease, damage or previous herbicide treatment as reduced weed control and / or increased crop injury may result.

In rice **DO NOT** treat weeds if mud, cracks or firm soil have appeared through fields prior to treatment, as poor control will result due to moisture stress.

DO NOT treat weeds of high density or growth stage larger than specified in the **DIRECTIONS FOR USE** as poor control will result.

DO NOT apply if rain is likely within one (1) hour as weed control may be reduced.

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of field runoff for at least 3 days after application.

DO NOT apply outside of the recommended application window with regards to the crop growth stage (see Tables 3 and 4).

DO NOT apply more than one (1) application per season in rice, fallow and sorghum.

DO NOT apply more than two (2) applications per season in tropical grass pastures, commercial & industrial areas and rights-of-way.

In rice **DO NOT** apply by application methods other than SCWIIRT (Soluble Chemical Water Injection In Rice Technique) or Bickley boom.

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone tables below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between three (3) and twenty (20) kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by aircraft unless by SCWIIRT or Bickley boom in rice.

SCWIIRT or Bickley boom

DO NOT apply by a SCWIIRT or Bickley boom, ground or aerial, unless minimum distances between the application site and downwind sensitive areas are observed (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for SCWIIRT or Bickley boom').

Table 1. Buffer zones for SCWIIRT or Bickley boom (in-flood application)						
Application rate	Application technique	Mandatory downwind buffer zones				
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
150 mL/ha	Ground SCWIIRT (in-flood)	0 m	0 m	0 m	0 m	0 m
	Helicopter* SCWIIRT or Bickley boom (in-flood)		5 m		60 m	
	Fixed-wing SCWIIRT or Bickley boom (in-flood)		20 m		300 m	

* For helicopter operation at speeds greater than 50 knots, users must observe the fixed-wing buffer zones when applying in-flood applications via SCWIIRT or Bickley boom.

Boom sprayers

DO NOT apply by a boom sprayer unless the following requirements are met:

- Spray droplets are not smaller than a COARSE spray droplet size category.
- Minimum distances between the application site and downwind sensitive areas are observed (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for boom sprayers').

Table 2. Buffer zones for boom sprayers						
Application rate	Boom height above the target canopy	Mandatory downwind buffer zones				
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
Up to 150 mL/ha	0.5 m or lower	0 m	10 m	0 m	90 m	0 m
	1.0 m or lower		35 m		350 m	
25 mL/ha	0.5 m or lower		0 m		25 m	
	1.0 m or lower		10 m		55 m	

DIRECTIONS FOR USE

Table 3. ***WEEDS CONTROLLED OR SUPPRESSED IN RICE – SCWIIRT or Bickley boom (in-flood) application.*** **DO NOT** apply prior to three leaf growth stage of rice (BBCH 13). **DO NOT** apply any later than panicle initiation growth stage of rice (BBCH 30). Always use Ubeniq® in sequence after foundation herbicide treatments that have a different mode of action.

Always apply with Uptake® Spraying Oil, Hasten™[1] Spray Adjuvant or Loveland Products™[1] MSO®[1] with Leci-Tech®[1] at 2 - 4 L/ha in total spray volume of 5 - 20 L/ha. Pre-mix Ubeniq® thoroughly in oil, then once fully dispersed, add remaining water to make up total spray mix volume.			
WEEDS CONTROLLED	**WEED GROWTH STAGE**	**RATE (mL/ha)**	**CRITICAL COMMENTS**
Arrowhead (*Sagittaria calycina)*	Up to 4 leaf and not more than 5 cm across	150	See WATER MANAGEMENT in GENERAL INSTRUCTIONS. A follow up treatment with a herbicide having a different mode of action may be required for complete control of weeds larger than label size or growth stage.
Suppression Dirty dora (*Cyperus difformis*)	Up to 2 leaf stage and not more than 3 cm across		
Jerry Jerry (*Ammania multiflora*)	Up to 4 leaf and not more than 5 cm across		
Starfruit (*Damasonium minus*)			
Water plantain (*Alisma plantago-aquatica*)			

Table 4. ***WEEDS CONTROLLED IN SORGHUM – Post emergence ground boom application.*** **DO NOT** apply before the secondary roots are present and before four fully expanded leaves (BBCH 14). **DO NOT** apply any later than at eight fully expanded leaves of sorghum (BBCH 18). Use Ubeniq® + Starane Advanced in tank mixture with atrazine for broadspectrum weed control.

Always apply with Hasten™[1] Spray Adjuvant at 500 mL/100 L water.			
WEEDS CONTROLLED	**WEED GROWTH STAGE**	**RATE (mL/ha)**	**CRITICAL COMMENTS**
Caltrop (Yellow vine) (*Tribulus terrestris*)	Seedling plants up to 15 cm diameter	25 + 300 Starane Advanced	Only treat small seedling weeds that are actively growing.

Table 5. ***WEEDS CONTROLLED IN FALLOW – Post emergence ground boom application.***
Always use Ubeniq® in tank mixture with glyphosate for broadspectrum weed control.

WEEDS CONTROLLED	WEED GROWTH STAGE	RATE (mL/ha)	CRITICAL COMMENTS
Caltrop (Yellow vine) (*Tribulus terrestris*)	Seedling plants up to 15 cm diameter	25	Only treat small seedling weeds that are actively growing. When Ubeniq® is applied without a tank mix partner, add Hasten™[1] Spray Adjuvant at 500 mL/100 L water.

Table 6. ***WEEDS CONTROLLED IN TROPICAL GRASS PASTURES (GRAZING ONLY), COMMERCIAL & INDUSTRIAL AREAS, RIGHTS-OF-WAY – Post emergence ground boom and spot application.***

Always apply with Uptake® Spraying Oil.			
WEEDS CONTROLLED	**WEED GROWTH STAGE**	**RATE**	**CRITICAL COMMENTS**
Navua sedge (*Cyperus aromaticus*)	Seedlings prior to flowering	100 – 150 mL/ha	Apply to grass pastures 2 weeks after grazing in spring/summer. A second application 10 - 12 weeks later may be required to control the next germination of Navua sedge. Use the higher rate on denser infestations. Add Uptake® Spraying Oil at 1 L/100 L water.
		15 mL/100 L water	A second application 10 - 12 weeks later may be required to control the next germination of Navua sedge. Add Uptake® Spraying Oil at 500 mL/100 L water.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

GENERAL INSTRUCTIONS

Ubeniq® is an arylpicolinate herbicide in the form of suspension concentrate for in-flood use in rice and for ground broadcast applications in sorghum, pastures (grazing only), commercial & industrial areas, rights-of-way and fallow. It will not reliably control weeds that emerge after treatment. Best results are achieved under good growing conditions. Treatment of crop or weeds that are stressed must be avoided. Ubeniq® should only be used to treat small seedling weeds. Treatment of weeds larger than on the label will result in reduced control or possible treatment failure.

CROP SAFETY

Minor, transient crop effects may be observed following an application of Ubeniq®. Grain yield is unaffected. Crop effects may be slight leaf crinkling, height or minor growth retardation. Crops that are stressed due to one or more factors may be more likely to show crop effects and will be slower to recover. **DO NOT** apply to crops or weeds which may be stressed due to a range of factors including, but not limited to prolonged cold, sustained high temperatures, poor nutrition (including deficiency and trace element toxicity), root diseases or previous herbicide treatment as reduced weed control and / or increased crop injury may result. **DO NOT** apply outside of the recommended crop growth stages as per the Direction for Use tables.

WATER MANAGEMENT IN RICE

Ubeniq® may be directly applied to floodwater by SCWIIRT rig, using a minimum of 5–20 L spray mix/ha by steel wheeled tractor, all-wheel drive motorbike or aircraft (helicopter or plane). Apply to seedling weeds, more than 12 hours after water movement has stopped and is clear of sediment, with flood depth sufficient to allow even distribution throughout the bay and enable full water coverage of soil for a lock-up period of five (5) days. Ensure flood gates are secured to avoid water movement. Treated bays may be topped up for normal water management after the five (5) day lock-up period. Reduced weed control will result where there is muddy water, too shallow to allow effective mixing or where soil is exposed.

CROP ROTATION RECOMMENDATIONS

Safe re-cropping periods apply for all crops following Ubeniq® application. Susceptible crops include, but are not limited to, those listed in the table below. Also see PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS.

Crop to be sown	Application rate (mL/ha)	Minimum time from application to planting	Minimum Rainfall Requirement
Barley, oats, triticale, wheat, ryegrass	25	4 weeks	75 mm
Canola, Winter legumes (including chickpeas & field peas)		2 months	
Barley, oats, triticale, wheat, ryegrass	150	4 weeks	Not applicable (refers to rice uses where floodwater is present)
Legumes (grain or pasture)		6 months	

Ubeniq® is broken down in soil by microbial activity. Quickest breakdown occurs in warm, moist soil that favours highest microbial activity. Breakdown may be slow in very dry seasons, or in cold, waterlogged soils, extending the safe plant back interval for susceptible crops.

- Rotational crops may only be planted after both the time and rainfall requirement have been met.
- Plantback periods for other crops have not yet been established. Contact your Corteva Agriscience representative.

AVOID double overlaps to reduce risk of injury to rotational crops the following season.

Before using Ubeniq® in tank mixes with other herbicides, check the plant-back information on all product labels. The time between spraying and planting will be determined by the most residual product, i.e. the product with the longest plantback period.

WEED DENSITY
Control may be reduced where weed density is very high and limits movement in floodwater in rice.

WEED GROWTH STAGE
Best results are achieved when applied to small, seedling weeds up to four (4) leaf growth stage.

APPLICATION
SCWIIRT application – in-flood (rice)
Ubeniq® may be directly applied to floodwater by SCWIIRT rig, using 5–20 L spray mix/ha by steel wheeled tractor, all-wheel drive motorbike or aircraft (plane or helicopter).

Boom application (sorghum, pasture, commercial & industrial areas, rights-of-way, fallow): Apply Ubeniq® with an accurately calibrated boom sprayer, in at least 80 L water/ha for sorghum and fallow and at least 150 L water/ha for Navua sedge control.

Spot spraying (pasture, commercial & industrial areas, rights-of-way): To obtain good coverage, a spray volume of 1,000 L water/ha is required per treated hectare. Spray to ensure thorough coverage of all foliage, including stems, to the point of runoff.

MIXING
Measure the required quantity of liquid and pre-mix with 2-4 L crop oil e.g. Uptake® Spraying Oil. Ensure that Ubeniq® is thoroughly dispersed in crop oil, then add to water to make up the remaining spray volume. Apply in total spray mix volume of 5-20 L/ha. Apply immediately after mixing Ubeniq® plus crop oil with water. DO NOT leave to stand in pre-mix vat or plane. Maintain continuous agitation until mix applied.

COMPATIBILITY
Herbicides: atrazine granules, glyphosate, Starane® Advanced
Adjuvants:
Uptake® Spraying Oil at 2 – 4 L/ha is the recommended adjuvant in rice. Hasten™[1] Spray Oil or Loveland Products™[1] MSO®[1] with Leci-Tech®[1] Spray Adjuvant may be used as an alternative at 2 - 4 L/ha. Use the higher rate where weeds are large or dense at spray time.
Hasten™[1] Spray Oil at 500 mL/100 L is the recommended adjuvant for use in sorghum.
Hasten™[1] Spray Oil at 500 mL/100 L is the recommended adjuvant for use in fallow when Ubeniq® is applied without a tank mix partner.
Uptake® Spraying Oil is the recommended adjuvant for Navua sedge control: 1 L/100 L water for broadcast applications and 500 mL/100 L for spot applications.

CLEANING SPRAY EQUIPMENT

After using Ubeniq®, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose, drain the tank and clean tank, pump, line and nozzle filters.

Partial Cleaning - Rinse only - before using sprayer to treat wheat, barley, oats or triticale:

After cleaning the tank as above, quarter fill the tank with clean water and circulate through the pump, line, hoses and nozzles. Drain and repeat procedure twice.

Complete Cleaning - **Decontamination** - before using sprayer to treat crops that are sensitive to Ubeniq®:

Wash the tank and rinse as above. Then quarter fill the tank and add a standard alkali-based laundry detergent at 500 g (or mL)/100 L water and circulate throughout the system for at least 15 minutes. If using a concentrated laundry detergent use 250 g (or mL)/100 L water. Do not use chlorine-based cleaners.

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land <u>away from</u> desirable plants and their roots and watercourses.