Product Name: Verpixo Adavelt active Fungicide

APVMA Approval No: 91871/142508

RLP
APPROVED

Label Name:	Verpixo Adavelt active Fungicide

Signal Headings:	READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	100 g/L FLORYLPICOXAMID

Mode of Action:	GROUP 21 FUNGICIDE

Statement of Claims:	For the control of various diseases in cucurbits, fruiting vegetables, head and leafy lettuce, strawberry and grapevines as specified in the Directions for Use.

Net Contents:	1 - 100 L

Restraints:	This section contains file attachment.

Directions for Use:	This section contains file attachment.

Other Limitations:	

Withholding Periods:	Cucurbits, fruiting vegetables, leafy vegetables and strawberries: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION Grapes: DO NOT HARVEST FOR 10 DAYS AFTER APPLICATION. DO NOT graze treated vineyards.

Trade Advice:	TRADE ADVICE: Verpixo® may leave detectable chemical residues in harvested produce. Overseas markets may not have appropriate residue tolerances in place or may have established tolerances which are lower than Australian maximum residue limits. Some crops for export to these destinations may require a longer harvesting withholding period. If you are using this product on crops destined for export, please contact your exporter for advice.

General Instructions:	This section contains file attachment.

Resistance Warning:	For fungicide resistance management Verpixo® Adavelt® active Fungicide is a group 21 fungicide, with the Quinone inside Inhibitor (QiI) mode of action. Some naturally occurring individual fungi resistant to the product and other group 21 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by the product or other Group 21 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Corteva Agriscience Australia Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi. Verpixo® may be subject to specific industry resistance management strategies which may recommend limits to the number of sprays, constraints regarding consecutive sprays or sprays following in-furrow or seed treatments, minimum spray intervals and no-spray periods for specific fungicide activity groups. For further information refer to the CropLife Australia website.

Precautions:	RE-ENTRY No re-entry period is required.

Protections:	PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

Storage and Disposal:	Store in the closed, original container in a cool well-ventilated area. Do not store for prolonged periods in direct sunlight. Keep out of reach of children. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank. Do not dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be taken separately. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Safety Directions:	May irritate the eyes. Avoid contact with the eyes. When opening the container, preparing the spray, and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). Wash hands after use.

First Aid Instructions:	First aid is not generally required. If in doubt, contact a Poisons Information Centre (Phone Australia 13 11 26) or a doctor.

First Aid Warnings:	

RESTRAINTS

DO NOT apply if rain is likely within 3 hours (i.e. Verpixo will be rainfast in 3 hours).
DO NOT apply if heavy rains or storms are forecast within 3 days.
DO NOT irrigate to the point of field runoff for at least 3 days after application.

DO NOT apply more than two consecutive sprays or more than three times per crop, or 33% (1/3) of total sprays, whichever is lower. Apply up to five (5) applications per calendar year in case of multiple cropping cycles.

DO NOT apply by aircraft.

DO NOT apply in water only chemigation hydroponics e.g. with reticulation.

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between three (3) and twenty (20) kilometres per hour at the application site during the time of application.
DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

Boom sprayers
DO NOT apply by a boom sprayer unless the following requirements are met:

- Spray droplets are not smaller than a MEDIUM spray droplet size category.
- Minimum distances between the application site and downwind sensitive areas are observed (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for boom sprayers').

Buffer zones for boom sprayers

Application rate	Boom height above the target canopy	Mandatory downwind buffer zones				
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
1.5 L/ha	0.5 m or lower	0 metres	55 metres	0 metres	0 metres	0 metres
	1.0 m or lower	0 metres	150 metres	0 metres	10 metres	0 metres

Vertical sprayers
DO NOT apply by a vertical sprayer unless the following requirements are met:

- Spray is not directed above the target canopy.
- The outside of the sprayer is turned off when turning at the end of rows and when spraying the outer row on each side of the application site.
- For dilute water rates up to the maximum listed for each type of canopy specified, minimum distances between the application site and downwind sensitive areas are observed (see the following table titled 'Buffer zones for vertical sprayers').

Buffer zones for vertical sprayers

Application rate	Mandatory downwind buffer zones				
	Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
1.5 L/ha	0 metres	20 metres	0 metres	0 metres	0 metres

DIRECTIONS FOR USE

Always use as part of a CropLife Australia Fungicide Resistance Management Strategy (FRMS). Apply as part of an Integrated Disease Management (IDM) program. Thorough coverage of crop is essential to obtain good control. Good crop hygiene will aid disease control.			
CROP	**DISEASE**	**RATE**	**CRITICAL COMMENTS**
Cucurbits, including but not limited to cucumber, melons (e.g. honeydew and rockmelon), pumpkin, squash, watermelon and zucchini (field and protected cropping)	*Powdery mildew* *Podosphaera xanthii*	1.25 – 1.5 L/ha or 125 – 150 mL/100L	Use as a protectant spray. Apply prior to disease infection or when conditions favour disease development (wet and cloudy conditions). A second application can be made 7-10 days later. Use a shorter interval when disease pressure is high. **DO NOT** apply more than two consecutive sprays or more than three sprays per crop. See **General Instructions** - Fungicide Resistance Management Powdery mildew: Use the high rate under high disease pressure. In broadcast situations ensure 1.25 - 1.5 L Verpixo is applied per hectare. DO NOT apply in greater than 1000 L water per ha. Protected cropping situations is permitted with a robust FRMS in place.
	Gummy stem blight (*Stagonosporopsis cucurbitacearum*, *Stagonosporopsis citrulli* and *Stagonosporopsis caricae)* known as *Didymella bryoniae* *Suppression only*	1.5 L/ha or 150 mL/100L	
Capsicum, chilli, eggplant, okra and tomato (field and protected cropping)	Powdery mildew of capsicum (*Leveillula taurica*) Powdery mildew of tomato *(Oidium lycopersici)* Target Spot (Early Blight) *(Alternaria Solani)* Anthracnose (*Colletotrichum capsica*) Grey mould *(Botrytis cinerea)*	1.5 L/ha or 150 mL/100L	Use as a protectant spray. Apply prior to disease infection or when conditions favour disease development (wet and cloudy conditions). A second application can be made 7-10 days later. Use a shorter interval when disease pressure is high. **DO NOT** apply more than two consecutive sprays or more than three sprays per crop. See **General Instructions** - Fungicide Resistance Management In broadcast situations ensure 1.5 L Verpixo is applied per hectare. DO NOT apply in greater than 1000 L water per ha. Protected cropping situations is permitted with a robust FRMS in place.

Head lettuce, Leaf lettuce (field crops only)	Sclerotinia rot *(Sclerotinia minor, Sclerotinia sclerotiorum)* Grey mould *(Botrytis cinerea)* Septoria spot or late blight *(Septoria lactucae)* Anthracnose *(Microdochium panattonianum)*	1.5 L/ha	Use as a protectant spray. Apply before first sign of disease or when conditions are conducive to disease as part of a disease management program and continue with a 7-10 day interval between sprays. Use the shorter interval when disease pressure is high. **DO NOT** apply more than two consecutive sprays or more than three sprays per crop. See **General Instructions** - Fungicide Resistance Management
Strawberry (field and protected cropping)	Grey mould (*Botrytis cinerea*) Powdery mildew (*Sphaerotheca macularis*)	1.5L/ha or 150 mL/100L	Use a program of protectant fungicides during flowering (blossom) when conditions favour disease. Repeat applications of Verpixo® on a 7 - 10 day interval. Use the shorter interval when conditions favour disease development. **DO NOT** apply more than two consecutive sprays or more than three sprays per crop. See **General Instructions** - Fungicide Resistance Management Reduce background levels of disease by removing infected plant debris and rotten fruit. Thorough coverage of the crop is essential. Ensure 1.5 L Verpixo is applied per hectare whether broadcast or dilute spraying is used. If less than 1000 L/ha water volume is applied, use concentrate spraying. Refer to **General Instructions - Application** section. DO NOT apply in greater than 1000 L water per ha. **Adjuvants:** Always apply Verpixo® with an adjuvant at the recommended rate e.g. Agral at 10 mL/100L. See **General Instructions** - Adjuvants.

Grapevine	Powdery mildew (*Erysiphe necator*) Grey mould (*Botrytis cinerea*)	150 mL/100L To give good coverage adjust water rate to growth stage and size of foliage.	Use as a protectant spray only, and rotate with fungicides of different mode of action for resistance management. See **General Instructions** - Fungicide Resistance Management Apply as a preventative program when conditions favour disease. Apply a maximum of two applications per season using a minimum 10-day spray interval. Verpixo is not systemically mobile. Thorough coverage is required to ensure control. Apply as part of a botrytis control spray program, where critical application timings are 5% capfall, 80% capfall, pre-bunch closure or veraison. DO NOT apply in greater than 1000 L water per ha. **Adjuvants:** Always apply Verpixo® with an adjuvant at the recommended rate e.g. Agral at 10 mL/100L. See **General Instructions** - Adjuvants.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

MIXING
Verpixo® is a suspension concentrate (SC) formulation.
Agitate or shake the container immediately prior to use. Half-fill the spray tank with water, add the appropriate amount of accurately measured Verpixo®, then complete filling the tank. Ensure thorough agitation at all times during mixing and application, by either mechanical or hydraulic action. Only use clean water within the range pH 5-9 to dilute Verpixo®.

STORAGE OF DILUTED SPRAY MIX
Whenever possible the spray mix should be used immediately after it is prepared. However, if weather conditions or mechanical breakdown prevent immediate use, the spray mix may be stored for up to 24 hours without loss of activity. The spray mix should be agitated thoroughly by mechanical or hydraulic action at regular intervals during storage to prevent sedimentation. Ensure that the stored spray mix is thoroughly agitated at least once every 8 hours. The spray mix must be stored out of direct sunlight.

ADJUVANT
Strawberry and grapevines: Apply with a high quality non-ionic surfactant such as Agral®[1], or other compatible adjuvant, at the recommended label rate. **Other crops**: Addition of a wetting agent may improve control under less than ideal application conditions. Use the wetter according to its label directions. Apply with a non-ionic surfactant such as Agral®[1] at the recommended label rate.

APPLICATION
Apply by ground spraying equipment only. **DO NOT** apply with spray droplets smaller than a MEDIUM spray droplet size category.

Thorough coverage of the crop is essential. Ensure this by increasing water volume with plant growth stage. Do not apply when conditions are unsuitable for water-based spray applications. Avoid high temperature, strong winds, inversion conditions, imminent rain or any conditions that may reduce the quality of spray coverage or result in drift from the target area. Techniques to minimise drift should be employed at all times when applying sprays to, or near, sensitive areas.

If using high volume (dilute) sprayer or concentrate sprayer, ensure that the correct amount of Verpixo® is applied per hectare irrespective of the water application rate per hectare. Apply the same total amount of Verpixo® to the target crop whether applying via dilute or concentrate spraying equipment.

Dilute spraying
Use a sprayer designed to apply high spray volumes, up to the point of run-off and matched to the crop
being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off. The required spray volume may be determined by applying different test volumes, using different settings on the sprayer, or from industry guidelines or expert advice. Add the amount of product specified in the Direction for Use table for each 100 L of water. Spray to the point of run-off. If volume to be applied is <1000 L/ha then use the low volume (concentrate) application method for calculation of chemical rate. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying
Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies spray volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen
spray volume. Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way:
EXAMPLE ONLY
1. Dilute spray volume as determined above: For example 1000 L/ha

2. Your chosen concentrate spray volume: For example 500 L/ha
3. The concentration factor in this example is 2X (i.e. 1000 L ÷ 500 L = 2)
4. If the dilute label rate is 150 mL/100 L, then the concentrate rate becomes 2 x 150, that is, 300 mL/100 L of concentrate spray.
The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation
may need to be changed as the crop grows.
For further information on concentrate spraying users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry best practice.

RAINFASTNESS
Rain can wash Verpixo® from treated plant surfaces and result in reduced disease control. Avoid making spray applications if rain is expected before the spray can dry completely. Verpixo® is rainfast after 3 hours of drying time. Where Verpixo® is applied in tankmix, use the longer of the partner products rainfast periods to determine when to apply to avoid rain.

CLEANING SPRAY EQUIPMENT
After using Verpixo® empty the tank and completely drain the system. Rinse the tank, pumps, lines, hoses, filters and nozzles by circulating clean water through the system. Drain and repeat the rinsing procedure twice.

FUNGICIDE RESISTANCE MANAGEMENT
Verpixo® use is subject to a CropLife Australia resistance management strategy. Please refer to this strategy at http://www.croplife.org.au/industry-stewardship/resistance-management/ before use.

The following restrictions also apply to the use of Verpixo®:

- Apply Verpixo® as a protectant spray only.
- DO NOT apply more than two consecutive sprays or more than three times per crop with maximum of five applications per calendar year in case of multiple cropping cycles.
- When conditions favour disease development, DO NOT wait for the disease to appear, but apply Verpixo® at the spray interval recommended in the Directions for Use table. Then resume the spray program with treatments that have a different mode-of-action to Group 21.
- Exposure to Verpixo® (or any other Group-21 containing product) should not exceed thirty-three percent (33%) of the total period of protection needed per crop.
- Verpixo® applications are to be made preventatively and no more than two (2) times in sequence before applying a fungicide with a different mode of action.
- Where a fungicide with a different mode of action follows Verpixo® (or any other Group 21-containing product) application(s), this fungicide preferably should have curative activity.
- Continue alternation of fungicides between successive crops. There will be no more than five (5) applications of Verpixo® (or any other Group 21 containing product) per year on the same area.

A disease management program that includes rotation and/or tank mixing with fungicides with a different mode of action is essential to reduce the risk of fungicide resistance development. For guidance on a particular crop and disease control situation, contact your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, local Corteva Agriscience Representative or CropLife Australia.