Product Name: ZORVEC ENICADE FUNGICIDE

APVMA Approval No.: 68375/142021

RLP
APPROVED

Label Name:	ZORVEC ENICADE FUNGICIDE

Signal Headings:	READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	100 g/L OXATHIAPIPROLIN

Mode of Action:	GROUP 49 FUNGICIDE

Statement of Claims:	For the control of certain fungal disease species in oilseed poppies and vegetable crops as per the Directions for Use.

Net Contents:	1 L 5 L

Restraints:	DO NOT apply if rainfall is expected within 20 minutes of spray residues drying. DO NOT use on hydroponic crops. DO NOT apply in nursery production of transplanted crops. DO NOT apply by air. SPRAY DRIFT RESTRAINTS DO NOT apply with spray droplets smaller than a MEDIUM spray droplet size category according to nozzle manufacturer specifications that refer to the ASAE S572 Standard or the BCPC Guideline. DO NOT apply during surface temperature inversion conditions at the application site. DO NOT apply when wind speed is less than 3 or more than 20 km per hour are measured at the application site. Users of this product MUST make an accurate written record of the details of each spray application within 24 hours following application and KEEP this record for a minimum of 2 years. The spray application details that must be recorded are: 1. date with start and finish times of application; 2. location address and paddock/s sprayed; 3. full name of this product; 4. amount of product used per hectare and number of hectares applied to;

	5. crop/situation and weed/pest; 6. wind speed and direction during application; 7. air temperature and relative humidity during application; 8. nozzle brand, type, spray angle, nozzle capacity and spray system pressure measured during application; 9. name and address of person applying this product. (Additional record details may be required by the state or territory where this product is used.)

Directions for Use:	This section contains file attachment.

Other Limitations:	

Withholding Periods:	HARVEST BRASSICAS: DO NOT HARVEST FOR 0 DAYS AFTER APPLICATION BRASSICA LEAFY VEGETABLES, LEAFY VEGETABLES: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION. CUCURBIT VEGETABLES: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION. CHIVES, FENNEL BULB, GARLIC, LEEKS, ONIONS, SHALLOTS, SPRING ONIONS: DO NOT HARVEST FOR 10 DAYS AFTER APPLICATION. POPPIES: DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION. GRAZING DO NOT GRAZE OR CUT FOR STOCK FOOD.

Trade Advice:	EXPORT STATEMENT: Growers should note that suitable Maximum Residue Levels (MRLs) or import tolerances may not be established in all markets for produce treated with Zorvec Enicade. If you are growing produce for export, please check with Corteva Agriscience for the latest information on MRLs and export tolerances before using this product.

General Instructions:	GENERAL INSTRUCTIONS Zorvec Enicade fungicide is recommended for control of certain foliar plant diseases, and has preventive, and locally systemic activity. Zorvec Enicade must be applied in a regularly scheduled protective spray program in rotation with other fungicides. See the Directions for Use table for specific crop/ disease recommendations. MIXING INSTRUCTIONS Fill spray tank to ¼ to ½ full of water. Measure the amount of Zorvec Enicade required for the area to be sprayed. Add Zorvec Enicade directly to the spray tank with the agitation engaged. Mix thoroughly to disperse the fungicide. Once dispersed, the material must be kept in suspension at all times by continuous agitation. Use mechanical or hydraulic means, DO NOT use air agitation, premix or slurry. If spray solution is left standing, ensure thorough re-agitation of the spray mix until fully resuspended. DO NOT allow spray mix to sit overnight, as resuspension may be difficult. MIXING ORDER The mixing sequence recommended is: water soluble bags, dry flowable or water dispersible granules, wettable powders, water based suspension concentrates, water soluble concentrates, oil based suspension concentrates, emulsifiable concentrates, adjuvants and surfactants, soluble fertilisers.

	SURFACTANT/WETTING AGENT Use of a surfactant/wetting agent is not required. APPLICATION Use a sprayer fitted with high flow rate nozzles to apply the highest practical spray volume. Use sufficient water to obtain thorough coverage of plants, with a minimum 250 L/ha. Nozzles with higher rated flows produce larger droplets. Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size, DOES NOT improve canopy penetration and may increase drift potential. WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE. Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. COMPATIBILITY Zorvec Enicade is an oil-based formulation. Avoid applying Zorvec Enicade in tank mixture with other agricultural products that have a history of causing crop injury when applied with oils unless specifically recommended by Corteva Agriscience. Zorvec Enicade is compatible with many commonly used fungicides, liquid fertilisers, herbicides, insecticides, plant growth regulators and biological control products. However, since the formulations of products are always changing, it is advisable to test the physical compatibility of desired tank mixes and check for adverse effects like settling out or flocculation. To determine the physical compatibility, add the recommended proportions of the tank mix products to water, mix thoroughly and allow to stand for 20 minutes. If the combination remains mixed, or can be re-mixed readily, it is considered physically compatible. Zorvec Enicade is compatible with Antracol, Avatar, Benevia, Coragen, Fontelis, imidacloprid, mancozeb, Movento, Octave, Phosphorous acid, Polyram, Rovral, Success Neo, Sumagic and Ultimate. The crop safety of all potential tank-mixes, including additives and other pesticides, on all crops has not been tested. Before applying any tank-mix not specifically recommended on this label or other Corteva Agriscience supplemental labelling, the safety to the target crop must be confirmed. To test for crop safety, apply the combination to a small area of the target crop in accordance with the label instructions to ensure that a phytotoxic response will not occur. SPRAY EQUIPMENT CLEANOUT Prior to application, start with clean, well-maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove. Drain spray equipment. Thoroughly rinse sprayer and flush hoses, boom, and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. DO NOT clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

Resistance Warning:	FUNGICIDE RESISTANCE WARNING GROUP 49 FUNGICIDE Zorvec Enicade is a member of the oxysterol binding protein homologue inhibition (OSBPI) group of fungicides. For fungicide resistance management Zorvec Enicade is a Group 49 fungicide. Oxysterol binding proteins (OSBP) are implicated in the movement of lipids between membranes, among other processes. Inhibiting OSBP may disrupt other processes in the fungal cell, such as signalling, maintaining cell membranes, and the formation of more complex lipids that are essential for the cell to survive. Some naturally occurring fungal biotypes resistant to Zorvec Enicade and other Group 49 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if Zorvec Enicade and other Group 49 fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 49 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant individuals is difficult to detect prior to use Production Agriscience (Australia) Pty Ltd* accepts no liability for any losses that may result from the failure of Zorvec Enicade to control resistant fungi.

	* Production Agriscience (Australia) Pty Ltd is a member of Corteva Agriscience group of companies DISEASE RESISTANCE MANAGEMENT Zorvec Enicade may be subject to specific resistance management strategies. To help prevent the development of resistance to Zorvec Enicade, use Zorvec Enicade in accordance with the current Fungicide Resistance Management (FRM) strategy for your region. For the latest information refer to the CropLife Australia website (https://www.croplife.org.au/resources/programs/resistance-management/). To delay fungicide resistance; • Exposure to Zorvec Enicade (or any other Group 49-containing product) should not exceed thirty-three percent (33%) of the total period of protection needed per crop. • For crops when three (3) or more Downy mildew fungicide applications are made, use Zorvec Enicade (or any other Group 49-containing product) in no more than 33% of the total number of fungicide applications, with a maximum number of sprays as specified in the Critical Comments section of the Directions for Use table. Where less than three (3) Downy mildew fungicide applications are made, make no more than one (1) application of Zorvec Enicade (or any other Group 49-containing product). • Zorvec Enicade applications are to be made preventatively and no more than two (2) times in sequence before applying a fungicide with a different mode of action. • Where a fungicide with a different mode of action follows Zorvec Enicade (or any other Group 49-containing product) application, this fungicide preferably should have curative activity. • Very few foliar fungicides have activity on soil-borne pathogens and the resistance risk associated with these diseases • There will be no more than six (6) applications of Zorvec Enicade (or any other Group 49 containing product) per year on the same acreage, targeting the same pathogen. A disease management program that includes rotation and/or tank mixing with fungicides with a different mode of action is essential to reduce the risk of fungicide resistance development. For guidance on a particular crop and disease control situation, contact your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, or local Corteva Agriscience representative.

Precautions:	RE-ENTRY No re-entry restriction apply to this product.

Protections:	PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Very toxic to aquatic life with long lasting effect. DO NOT contaminate wetlands or watercourses with this product or used containers.

Storage and Disposal:	KEEP OUT OF THE REACH OF CHILDREN. Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. 1 L and 5 L containers Do not re-use 1 L containers. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packagingto an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in

	compliance with relevant local, state or territory government regulations. Do not burn empty containers or product. This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank. Do not dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be taken separately.

Safety Directions:	• May irritate the skin. • Avoid contact with the skin. • Repeated exposure may cause allergic disorders. • Sensitive workers should use protective clothing. • When opening the container and preparing spray, wear chemical resistant gloves. • Wash hands after use. • After each day's use wash gloves and contaminated clothing.

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

First Aid Warnings:	

DIRECTIONS FOR USE
For use in all States where appropriate for the crop and/or disease.

CROP	DISEASE	RATE	WHP	CRITICAL COMMENTS	
Bulb vegetables (excluding Onions) including: Chives, Fennel bulb, Florence fennel, Garlic, Leeks, Shallots, Spring onions.	Downy mildew (*Peronospora destructor*)	350 mL/ha Plus a registered Downy mildew protectant fungicide at registered rates.	10 days	Apply up to two consecutive sprays of Zorvec® Enicade®, 10 days apart and then change to a fungicide from another chemical group. **DO NOT** apply more than 3 sprays of Zorvec® Enicade® to each crop as a precaution against development of disease resistance.	Maintain a regular protectant spray programme. Apply when conditions favour disease development but before the disease is evident. Apply in 250 to 500 L of water per hectare.
Onions				Apply up to two consecutive sprays of Zorvec® Enicade®, 10 to 14 days apart. **DO NOT** apply more than 2 sprays of Zorvec® Enicade® to each crop as a precaution against development of disease resistance.	
Brassica vegetables including: Broccoli, Brussels sprout, Cabbage, Cauliflower.	Downy mildew (*Hyaloperonospora parasitica*)		0 days	Maintain a regular protectant spray programme. Apply when conditions favour disease development but before the disease is evident. Apply up to two consecutive sprays of Zorvec® Enicade®, 7 to 10 days apart and then change to a fungicide from another chemical group. Use the shorter interval when conditions favouring infection are creating a high risk. Apply in 250 to 500 L of water per hectare. **DO NOT** apply more than 3 sprays of Zorvec® Enicade® to each crop as a precaution against development of disease resistance.	
Cucurbit vegetables including (field and protected crops): Bitter melon, Chokos, Cucumber, Gherkin, Marrow, Melons, Pumpkin, Rockmelon, Squash, Zucchini	Downy mildew (*Pseudoperonospora cubensis*)		1 day	Maintain a regular protectant spray programme. Apply when conditions favour disease development but before the disease is evident. Apply up to two consecutive sprays of Zorvec® Enicade® 7 to 10 days apart and then change to a fungicide from another chemical group. Use the shorter interval when conditions favouring infection are creating a high risk. Apply in 250 to 500 L of water per hectare. **DO NOT** apply more than 3 sprays of Zorvec® Enicade® to each crop as a precaution against development of disease resistance.	

CROP	DISEASE	RATE	WHP	CRITICAL COMMENTS
Leafy vegetables and Brassica leafy vegetables including: Amaranth, leafy Arugula, Buk choy, Cardoon, Chard, Chervil, Chinese broccoli (Gai lum/ Gai lan/ Kai lan), Chinese cabbage (Pet sai/ Wombok/ Haksukai), Choy sum (Gai choy/ Am soy), Corn salad, Cress, Dandelion leaves, Dock, Endive, Kai choy, Kale, Leafy mustard including Indian mustard and Mustard spinach (Komatsuma), Lettuce (Head and Leafy), Mibuna, New Zealand spinach, Orach, Pak choy, Purslane (Garden and Winter), Radicchio, Rocket, Spinach, Swiss chard, Tat soy	Downy mildew (*Bremia lactucae, Peronospora farinose)*	350 mL/ha Plus a registered Downy mildew protectant fungicide at registered rates.	3 days	Maintain a regular protectant spray programme. Apply when conditions favour disease development but before the disease is evident. Apply up to two consecutive sprays of Zorvec® Enicade®, 7 to 10 days apart and then change to a fungicide from another chemical group. Use the shorter interval when conditions favouring infection are creating a high risk. Apply in 250 to 500 L of water per hectare. **DO NOT** apply more than 3 sprays of Zorvec® Enicade® to each crop as a precaution against development of disease resistance.
Poppies	Downy mildew (*Peronspora cristata*)	350 mL/ha Plus a registered Downy mildew protectant fungicide at registered rates	6 weeks	Commence spraying early (i.e. before the main disease infection period) with registered non Group 49 fungicides. Apply Zorvec® Enicade® when conditions favour disease development during the late run-up to hook stage. Apply in 250 to 500 L water per hectare. **DO NOT** apply later than hook stage. **DO NOT** apply more than 1 application of Zorvec® Enicade® to each crop, as a precaution against development of disease resistance.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.