CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Delegate®

Jemvelva® active

INSECTICIDE

ACTIVE CONSTITUENT: 250 g/kg SPINETORAM

GROUP	5	INSECTICIDE

For the control of various caterpillar pests, such as codling moth and lightbrown apple moth in citrus, grapes, pome and stone fruit, western flower thrips in pome and stone fruit and carob moth in almonds as specified in the Directions for Use.

Net contents: 500g – 5kg

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone: *Australia* 13 11 26. *New Zealand* 0800 764 766.

EMERGENCY RESPONSE (ALL HOURS)
RING FROM ANYWHERE IN AUSTRALIA
1800 370 754
(LOCAL CALL FEE ONLY)

SAFETY DIRECTIONS

- Will irritate the eyes.
- Avoid contact with eyes.
- Wash hands after use.

IN A TRANSPORT EMERGENCY ONLY
DIAL 000
FOR POLICE OR FIRE BRIGADE

SAFETY DATA SHEET

Additional information is listed on the Safety Data Sheet for **DELEGATE® JEMVELVA® ACTIVE INSECTICIDE** which is available from Corteva Agriscience on request. Call Customer Service Toll Free on 1-800 700 096 or visit www.corteva.com.au

CORTEVA™ agriscience

™ ® Trademarks of Corteva Agriscience and its affiliated companies.

Visit us at Corteva.com.au

RESTRAINTS

DO NOT make more than 4 applications to any crop in any one season, except where otherwise indicated in the CRITICAL COMMENTS (also see the RESISTANCE statement).

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at: www.apvma.gov.au/Spraydrift

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone tables below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply when the wind speed is less than 3 or more than 20 kilometres per hour, as measured at the application site.

DO NOT apply if there are surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply from aircraft.

Except when applying with orchard / vineyard airblast equipment, **DO NOT** apply with spray droplets smaller than a **MEDIUM** spray droplet size category.

DO NOT apply by a boom sprayer.

DO NOT apply by a vertical sprayer unless the following requirements are met OR the conditions provided in a relevant output of the Spray Drift Management Tool are met:

- spray is not directed above the target canopy;
- the outside of the sprayer is turned off when turning at the end of rows and when spraying the outer row on each side of the application site;
- for dilute water rates up to the maximum listed for each type of canopy specified, minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for vertical sprayers') are observed.

MANDATORY NO-SPRAY ZONES

DO NOT apply if there are aquatic and wetland areas including aquacultural ponds, surface streams and rivers downwind from the application area and within the **mandatory no-spray zones** shown in Table 1 below.

DO NOT apply if there are livestock, pasture or any land that is producing feed for livestock downwind from the application area and within the mandatory no-spray zones shown in the table below.

DECLARED SPRAY DRIFT RISK AREA—Spray drift deposits that can cause residue violations in livestock feeding on pasture and forage, can result from typical applications out to distances downwind from the application area as shown in Table 1 below:

Table 1: Buffer zones for vertical sprayers

Type of target canopy and dilute water rate	Mandatory downwind buffer zones				
	Bystander areas	**Natural aquatic areas**	**Pollinator areas**	**Vegetation areas**	**Livestock areas**
2 metres tall and shorter, maximum dilute water rate of 1,000 L/ha	0 metres	10 metres	0 metres	0 metres	0 metres
Taller than 2 metres (not fully-foliated), maximum dilute water rate of 4,000 L/ha	0 metres	40 metres	25 metres	0 metres	30 metres
Taller than 2 metres (fully-foliated), maximum dilute water rate of 4,000 L/ha	0 metres	30 metres	15 metres	0 metres	20 metres

DIRECTIONS FOR USE

SPRAYING TREE & VINE CROPS: *In the following table, all rates are given for dilute spraying where spray volumes may vary in order to obtain good coverage to the point of run-off. For concentrate spraying refer to the "CONCENTRATE SPRAYING" section on this label.*

CROP	PEST	RATE	CRITICAL COMMENTS
FOR ALL TREE & VINE CROPS: Carefully monitor crops for eggs and larvae of pest species by regular field scouting. Target sprays against mature eggs and newly-hatched larvae. Apply repeat applications at 14 day intervals as egg hatch continues or as new infestations occur unless otherwise directed in the CRITICAL COMMENTS. ***RESISTANCE MANAGEMENT***: DO NOT make more than 4 applications to any crop in any one season.			
Citrus Fruits; including Grapefruit, Lemons, Limes, Mandarins Oranges	Citrus leafminer, Lightbrown apple moth	5-10 g /100 L❶	***Do not*** *apply by air.* *Citrus leafminer*: Best results will be achieved when horticultural oil is used in place of a wetting agent. **Only** use oils when applying to non-bearing trees due to the risk of fruit phytotoxicity. For the other pests, use higher rates for heavy infestations.
	Helicoverpa (Corn earworm and native budworm)	10- 20 g/100 L❶	
Grapes	Lightbrown apple moth	10 g/100 L❶	See comments under **"*FOR ALL TREE & VINE CROPS*"** above.
	Grapevine moth	3 g/100 L❶*	

Pome fruit, *including*: apples, pears and nashi pears	Codling moth	15 – 20 g/ 100 L❶	Use higher rates when adverse weather conditions may cause the spray interval to become greater than 14 days, or in high pressure conditions, in orchards with a history of damage or when mating disruption does not provide adequate protection.
	Lightbrown apple moth and Oriental fruit moth	10 or 15 g/ 100 L❶	
	Loopers, pear and cherry slug, Helicoverpa	10 g/100 L ❶	
	Western flower thrips	20 g/100 L❶	Use this product as part of the Western flower thrips (WFT) Resistance Management strategy (see end of table for details).
Stone fruit, including Apricots, Cherries Nectarines, Peaches Plums	Pear and cherry slug	5 g/100 L ❶	Use the higher rate under high pest pressure or when adverse weather conditions may cause the spray interval to become greater than 14 days.
	Lightbrown apple moth and Oriental fruit moth	10 or 15 g/ 100 L❶	
	Western flower thrips	20 g/100 L❶	Use this product as part of the WFT Resistance Management strategy (see end of table for details).
Almonds	Carob moth	20 g/100 L ❶	Use as part of an IPM system, ensure good orchard hygiene with the removal of mummified fruit in winter. Application timing is critical. Early post-flower application: At the beginning of the season (Aug-Sept), carefully monitor pheromone traps for peak adult pest incidence of 1st generation. Once regular field scouting indicates egg laying and newly hatched larvae apply Delegate® as required. Apply to complete coverage. Hull-split application: Delegate® should be applied at the start of hull split (2 – 5% of the almond hull sutures open). Carefully monitor pheromone traps for peak activity of 2nd generation moths. Continue field scouting for egg-laying activity, and apply Delegate® to coincide with the initiation of egg hatch. Apply to complete coverage. Apply repeat applications at a minimum of 7 day intervals. Ensure thorough spray coverage of all leaf and nut surfaces using an airblast sprayer or equivalent. Add a non-ionic wetting agent at the recommended rate❶.
Western Flower Thrips Resistance Management Strategy: Make 3 consecutive applications at either 3-5 day intervals when temperatures are greater than 20°C or at 6-12 day intervals when temperatures are less than 20°C. For any further sprays required, use an approved product from another chemical group.			

Do not make more than 3 consecutive applications of Delegate® before switching to an approved product from another chemical group.
❶Addition of a non-ionic wetting agent at its label rate, such as Agral® at 10 mL/100 L, may improve control under less than ideal application conditions. **DO NOT** add non-ionic wetting agent if tank-mixing Delegate® with any other products.
PEST NAMES: Carob moth: *Ectomyelois ceratoniae*; Citrus leafminer: *Phyllocnistis citrella*; Codling moth: *Cydia pomonella*; Grapevine moth: *Phalaenoides glycinae*; Helicoverpa caterpillars, corn earworm, native budworm: *Helicoverpa* spp.; Lightbrown apple moth: *Epiphyas postvittana*; Loopers: *Phrissigonus* spp., *Ectropis* spp., *Chloroclystis* spp.; Oriental fruit moth: *Grapholita molesta;* Pear and cherry slug: *Caliroa cerasi*; Western flower thrips: *Frankliniella occidentalis.*

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

Almond: **DO NOT APPLY WITHIN 7 DAYS OF HARVEST OR NUT FALL.**

Citrus Fruit: **DO NOT HARVEST FOR 1 DAY AFTER THE LAST APPLICATION**

Grapes and Pome Fruit: **DO NOT HARVEST FOR 7 DAYS AFTER THE LAST APPLICATION.**

Stone Fruit: **DO NOT HARVEST FOR 3 DAYS AFTER THE LAST APPLICATION.**

DO NOT GRAZE ANY TREATED ORCHARD OR VINEYARD OR CUT THE ORCHARD OR VINEYARD FLOOR FOR STOCKFOOD.

The grazing limitations above apply specifically to stock slaughtered for the domestic market. Export markets may apply different standards. To ensure compliance with residue standards in export markets, comply with the above grazing withholding periods and ensure that the **Export Slaughter Interval (ESI)** is observed before stock are sold or slaughtered.

Export Slaughter Interval: 14 Days. This means that livestock that have grazed on or were fed treated crops or byproducts should be placed on clean feed for at least 14 days prior to slaughter.

PRECAUTION: Delegate® Jemvelva® active Insecticide may leave detectable chemical residues in harvested produce. Overseas markets may not have appropriate residue tolerances in place or may have established tolerances which are lower than Australian maximum residue limits. Some crops for export to these destinations may require a longer harvesting withholding period. If you are using this product on crops destined for export, please contact your exporter for advice.

WINE GRAPES: If you are growing grapes for export wine production, please check with your exporter or the AWRI for the latest information on MRLs and import tolerances before using Delegate®.

GENERAL INSTRUCTIONS
Insecticide Resistance Warning

GROUP	5	INSECTICIDE

For insecticide resistance management, Delegate® Jemvelva® active Insecticide is a Group 5 insecticide. Some naturally occurring insect biotypes resistant to Delegate® and other Group 5 insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Delegate® and other Group 5 insecticides are used repeatedly. The effectiveness of Delegate® on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Corteva Agriscience Australia Pty Ltd accepts no liability for any losses that may result from the failure of Delegate® to control resistant insects. Delegate® may be subject to specific resistance management strategies. For further information contact your local supplier, Corteva Agriscience representative or local agricultural department agronomist.

MIXING
Half-fill the spray tank with water, add the appropriate amount of accurately measured Delegate®, then complete filling the tank.
Ensure thorough agitation by mechanical or hydraulic action at all times during mixing and application.
Use only clean water within the range pH 5-9 to dilute Delegate®.
DO NOT add non-ionic wetting agent if tank-mixing Delegate® with any other products.

APPLICATION
Thorough coverage of the crop is essential. Do not apply when conditions are unsuitable for water-based spray applications. Avoid high temperature, strong winds, inversion conditions, imminent rain or any conditions that may reduce the quality of spray coverage or result in drift from the target area. Techniques to minimise drift should be employed at all times when applying sprays to, or near, sensitive areas.

DILUTE SPRAYING
Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of first run-off. Avoid excessive run-off. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of runoff. If volume to be applied is < 1,000 L/ha then use the low volume (concentrate) application method for calculation of chemical rate. For volumes > 1,000 L/ha use dilute spray rate.

CONCENTRATE SPRAYING
Use a sprayer designed and set up for concentrate spraying (that is, a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (see DILUTE SPRAYING above) for the crop canopy. Consult your local adviser, agronomist or Department of Agriculture to determine this volume. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way:

Concentrate Spraying Example
1. Dilute spray volume as determined above: e.g. 1,500 L/ha
2. Your chosen concentrate spray volume: e.g. 500 L/ha
3. The concentration factor is 3 X (1,500 / 500)

4. If the dilute label rate is 10g/100 L, then the concentrate rate becomes 3 X 10, *i.e*. 30g/100 L of concentrate spray

The chosen spray volume, amount of product per 100 L of water and the sprayer set up and operation may need to be changed as the crop grows. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training. Always follow Industry Best Practices.

RAINFASTNESS
Rain can wash Delegate® from treated plant surfaces and result in reduced insect control. Avoid making spray applications if rain is expected before the spray can dry completely.

CLEANING SPRAY EQUIPMENT
After using Delegate® empty the tank and completely drain the system. Rinse the tank, pumps, lines, hoses, filters and nozzles by circulating clean water through the system. Drain and repeat the rinsing procedure twice.

RE-ENTRY

Do not allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF LIVESTOCK
Bee Safety: This product is highly toxic to bees; will kill bees foraging in the crop to be treated or in hives which are over-sprayed or reached by spray drift. Residues may remain toxic to bees for several days after application. Once the spray deposit has dried, foraging bees should not be affected when using spray volumes of 2,000 L/ha or less. However, if using spray volumes greater than 2000 L/ha, it is possible that foraging bees may be affected for some days after spraying.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT
Very Toxic to aquatic life. Do not allow the product or used containers to enter dams, ponds, waterways or drains. Do not allow irrigation water from treated paddocks to enter adjacent pastures, crops or water supplies. Do not apply in strong winds, inversion conditions or any other conditions that may result in drift onto adjacent pastures, crops or water supplies.

PROTECTION OF NON-TARGET INSECTS
Risk to non-target insects. Delegate® may have adverse effects on some non-target beneficials, such as foliage dwelling predators, particularly where IPM is practiced.

STORAGE AND DISPOSAL
Store in the closed, original container in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT store near food, feedstuffs, fertilisers or seed.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.
If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

DrumMUSTER containers
This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinsate or any

undiluted chemical according to state legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be taken separately.

SMALL SPILL MANAGEMENT
Sweep up material and contain in a refuse vessel for disposal in the same manner as for the container (see STORAGE AND DISPOSAL section).

APVMA Approval No.: 61717/133429

D.O.M.
Batch No.

Corteva Agriscience Australia Pty Ltd A.B.N. 24 003 771 659
67 Albert Avenue, Chatswood, NSW 2067
www.corteva.com.au
CUSTOMER SERVICE TOLL FREE **1-800 700 096**

™ ® Trademarks of Corteva Agriscience and its affiliated companies. © 2021 Corteva.

Barcode
for stock
identification

This product is GHS compliant. No additional GHS hazard and precautionary statements are required under the Safe Work Australia exemptions for AgVet products.