READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Fontelis®

FUNGICIDE

ACTIVE CONSTITUENT: 200 g/L PENTHIOPYRAD

GROUP 7 FUNGICIDE

For the control of certain fungal diseases in fruit, nut and vegetable crops as per the Directions for Use.

Pack Sizes: 1 L, 5 L

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone: *Australia* 13 11 26.

EMERGENCY RESPONSE (ALL HOURS)
RING FROM ANYWHERE IN AUSTRALIA
1800 370 754
(LOCAL CALL FEE ONLY)

IN A TRANSPORT EMERGENCY ONLY
DIAL 000
FOR POLICE OR FIRE BRIGADE

SAFETY DIRECTIONS

- When opening the container, preparing and using the product wear cotton overalls (or equivalent clothing) buttoned to the neck and wrist and a washable hat.
- Wash hands after use.
- After each day's use wash contaminated clothing.

SAFETY DATA SHEET

Additional information is listed on the Safety Data Sheet for **FONTELIS® FUNGICIDE** which is available from Corteva Agriscience on request. Call Customer Service Toll Free on 1-800 700 096 or visit www.corteva.com.au

CORTEVA™ agriscience

™ ® Trademarks of Corteva Agriscience and its affiliated companies.

Visit us at Corteva.com.au

RESTRAINTS
DO NOT apply if rainfall is expected within 1 hour of application.
DO NOT use on hydroponic crops.
DO NOT apply by air.

SPRAY DRIFT RESTRAINTS
Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.

DO NOT allow bystanders to come into contact with the spray cloud.
DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.
DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.
DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings 1 to 2 hours before sunset and persist until 1 to 2 hours after sunrise.

Boom sprayers
DO NOT apply by a boom sprayer unless the following requirements are met:
- Spray droplets not smaller than a MEDIUM spray droplet size category.
- Minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for boom sprayers') are observed.

Buffer zones for boom sprayers

Application rate	**Mandatory downwind buffer zones**
	Natural aquatic areas
Up to maximum label rate	10 metres

Vertical sprayers
DO NOT apply by a vertical sprayer unless the following requirements are met:
- Spray is not directed above the target canopy.
- The outside of the sprayer is turned off when turning at the end of rows and when spraying the outer row on each side of the application site.
- For dilute water rates up to the maximum listed for each type of canopy specified, minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for vertical sprayers') are observed.

Buffer zones for vertical sprayers

Crop	**Mandatory downwind buffer zones**
	Natural aquatic areas
Pome/Stone fruit orchards	80 metres
Tree nuts, Macadamias	100 metres

DIRECTIONS FOR USE

For use in all States where appropriate for the crop and/or disease.

TREE AND FRUIT CROPS

CROP	DISEASE	RATE	WHP	CRITICAL COMMENTS
Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Refer to Application section of the label. Thorough fruit coverage is essential. DO NOT use a concentration factor greater than 3X, for concentrate application. Use in accordance with CropLife Fungicide Resistance Management Strategy guidelines – see Disease Resistance Management Strategy. General Instructions section.				
Pome fruit including: Apples, Nashi pears, Pears	Apple black spot (Scab) (*Venturia inaequalis*) Apple powdery mildew (*Podosphaera leucotricha)* Pear scab (*Venturia pirina*)	**Dilute spraying:** 75 mL/100 L **Concentrate spraying:** Refer to **Mixing/ Application** section	28 days	**Protectant program:** Begin applications prior to disease development and continue on a seven (7) to twenty-one (21) day interval depending on the targeted diseases. Application interval for Apple black spot (Scab) and Pear scab is seven (7) to ten (10) days. Application interval for Powdery mildew is fourteen (14) to twenty-one (21) days. Application intervals for Alternaria leaf blotch and Alternaria fruit spot are ten (10) to fourteen (14) days. Application for control of this disease should begin after blossom and during early fruit development to prevent establishment of the disease on the leaves. Use the shorter spray interval when weather conditions favour disease development.
	Alternaria leaf blotch and Alternaria fruit spot of apples (*Alternaria* spp.) (*Suppression only*)			
Strawberry	Grey mould (Botrytis) (*Botrytis cinerea*) Powdery mildew (*Sphaerotheca* spp.)	175 mL/100 L 1.75 L/ha	Nil	Begin applications prior to disease development and continue on a seven (7) to ten (10) day interval. Use shorter spray interval when disease pressure is high. Maximum seasonal use rate is 5.25 L/ha.
Stone fruits including: Apricots, Cherries, Nectarines, Peaches, Plums, Prunes (fresh)	Brown rot (Blossom blight) (*Monilinia* spp.) Scab (Freckle) (*Cladosporium carpophilum, Venturia carpophila)*	150 mL/100L		Begin applications prior to disease development and continue on a seven (7) to fourteen (14) day interval. Use higher rate and shorter spray interval when disease pressure is high.

TREE AND FRUIT CROPS *continued*

CROP	DISEASE	RATE	WHP	CRITICAL COMMENTS
Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Refer to Application section of the label. Thorough fruit coverage is essential. DO NOT use a concentration factor greater than 3X, for concentrate application. Use in accordance with CropLife Fungicide Resistance Management Strategy guidelines – see Disease Resistance Management Strategy. General Instructions section.				
Tree nuts including: Almond, Chestnut, Filbert (Hazelnut)	Brown rot (Blossom blight) (*Monilinia* spp.)	150 mL/100L	14 days	Begin applications prior to disease development and continue on a seven (7) to fourteen (14) day interval. Use shorter spray interval when disease pressure is high.
Macadamias	Flower blight (dry flower, raceme blight) including but not limited to: (*Pestalotiopsis* sp.) (*Neopestalotiopsis* sp.)			Apply the first application at bud break and the second application fourteen (14) to twenty-one (21) days later when conditions are conducive to disease development, to protect flower formation and retention. Under high disease pressure use the shorter spray interval. Apply sufficient water volume to enable thorough coverage of flower racemes – see Application section.
	Husk spot (*Pseudocercospora macadamiae*)			Apply the first application at the match head stage of the crop and apply the second application fourteen (14) to twenty-eight (28) days later when conditions are conducive to disease development. Under high disease pressure use the shorter spray interval. Apply sufficient water volume to enable thorough coverage of flower racemes – see Application section.

VEGETABLE CROPS

CROP	DISEASE	RATE	WHP	CRITICAL COMMENTS	
Brassica vegetables including: Broccoli, Brussels sprout, Cabbage, Cauliflower	White mould (Sclerotinia stem rot) (*Sclerotinia* spp.)	1.75 L/ha	Nil	Begin applications prior to disease development and continue on a seven (7) to fourteen (14) day interval. Maximum seasonal use rate is 5.25 L/ha.	
Brassica leafy vegetables including: Buk choy, Chinese broccoli (Gai lum/Gai lan/Kai lan), Chinese cabbage (Pet sai / Wombok / Haksukai), Choy sum Gai choy / Am soy, Kai choy, Kale, Mibuna, Leafy mustard including Indian mustard and Mustard spinach (Komatsuma), Pak choy, Tat soy	White mould (Sclerotinia stem rot) (*Sclerotinia* spp.)				
Cucurbit vegetables (field and protected crops) including: Bitter melon, Chokos, Cucumber, Gherkin, Marrow, Melons, Pumpkin, Squash, Zucchini	Grey mould (*Botrytis cinerea*) Powdery mildew (*Sphaerotheca fuliginea*, *Erysiphe cichoracearum*)		1 day	Begin applications prior to disease development and continue on a seven (7) to fourteen (14) day interval. Use shorter spray interval when disease pressure is high.	Maximum seasonal use rate is 3.5 L/ha.
	Gummy stem blight (*Didymella bryoniae*)			Begin applications prior to disease development and continue on a seven (7) to ten (10) day interval. Use shorter spray interval when disease pressure is high.	

CROP	DISEASE	RATE	WHP	CRITICAL COMMENTS
Fruiting vegetables (field and protected crops) including: Egg plants, Capsicums (Pepper), Chillies, Tomatillo, Tomatoes	Early blight (*Alternaria solani*) Grey mould (*Botrytis cinerea*) Powdery mildew *(Leveillula taurica)*	1.75 L/ha	Nil	Begin applications prior to disease development and continue on a seven (7) to ten (10) day interval. Use shorter spray interval when disease pressure is high. Maximum seasonal use rate is 5.25 L/ha.
Leafy vegetables including: Chinese cabbage, Endive, Fennel, Kale, Cress, Lettuce, Mustard, Silverbeet, Spinach	Sclerotina rot (Lettuce drop) *(Sclerotinia* spp.*)* Grey mould (*Botrytis cinerea*) Powdery mildew (*Erysiphe cichoracearum*)		3 days	
Onions, Shallots, Spring onions	Botrytis blight and Neck rot (*Botrytis* spp.) Purple blotch (*Alternaria porri*)			Begin applications prior to disease development and continue on a seven (7) to fourteen (14) day interval. Maximum seasonal use rate is 5.25 L/ha.
Root and Tuber vegetables including: Beetroot, Carrots, Celeriac, Galangal, Parsnips, Potatoes, Radish, Swedes, Sweet potato, Turnips	Early blight (Target spot, Leaf spot) (*Alternaria* spp.) Powdery mildew (*Erysiphe* spp.)		7 days	Begin applications prior to disease development and continue on a seven (7) to fourteen (14) day interval. Use shorter spray interval when disease pressure is high. Maximum seasonal use rate is 3.5 L/ha.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

HARVEST WITHHOLDING PERIODS

BRASSICA VEGETABLES, BRASSICA LEAFY VEGETABLES, FRUITING VEGETABLES (OTHER THAN CUCURBIT VEGETABLES), STONE FRUIT, STRAWBERRIES: NOT REQUIRED WHEN USED AS DIRECTED.

LEAFY VEGETABLES (EXCEPT BRASSICA LEAFY VEGETABLES), ONIONS, SHALLOTS, SPRING ONIONS: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION.

CUCURBIT VEGETABLES: DO NOT HARVEST FOR 1 DAYS AFTER APPLICATION.

TREE NUTS: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

POME FRUIT: DO NOT HARVEST FOR 28 DAYS AFTER APPLICATION.

ROOT AND TUBER VEGETABLES: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION.

GRAZING

DO NOT GRAZE OR CUT FOR STOCK FOOD.

EXPORT STATEMENT: Growers should note that suitable Maximum Residue Levels (MRLs) or import tolerances may not be established in all markets for produce treated with Fontelis® fungicide. If you are growing produce for export, please check with Corteva Agriscience for the latest information on MRLs and export tolerances before using this product.

GENERAL INSTRUCTIONS

Fontelis® fungicide is a broad-spectrum fungicide, formulated as an oil based suspension concentrate. Fontelis® is recommended for control of foliar and soil-borne plant diseases, and has preventive, and locally systemic activity. Fontelis® must be applied in a regularly scheduled **protective** spray program in rotation with other fungicides that have a different mode of action.

FUNGICIDE RESISTANCE WARNING

GROUP 7 FUNGICIDE

For fungicide resistance management Fontelis® fungicide is a Group 7 fungicide.

Some naturally occurring fungal biotypes resistant to Fontelis® fungicide and other Group 7 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if Fontelis® fungicide and other Group 7 fungicides are used repeatedly. The effectiveness of Fontelis® fungicide on resistant individuals could be significantly reduced. Since the occurrence of resistant individuals is difficult to detect prior to use Corteva Agriscience accepts no liability for any losses that may result from the failure of Fontelis® fungicide to control resistant fungi.

Disease Resistance Management Strategy

Fontelis® fungicide may be subject to specific resistance management strategies. To help prevent the development of resistance to Fontelis®, use Fontelis® in accordance with the current CropLife Australia Fungicide Resistance Management strategy for your crop/disease.

For further information contact your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, local Corteva Agriscience Representative or refer to CropLife Australia Fungicide Resistance Management strategies at https://www.croplife.org.au/resources/programs/resistance-management/fungicide-resistance-management-strategies1-draft-draft-3.

MIXING

Fill spray tank to ¼ to ½ full of water. Measure the amount of Fontelis® fungicide required for the area to be sprayed. Add Fontelis® fungicide directly to the spray tank with the agitation engaged. Mix thoroughly to disperse the fungicide. Once dispersed, the material must be kept in suspension at all times by continuous agitation. Use mechanical or hydraulic means, **DO NOT** use air agitation, premix or slurry.

If spray solution is left standing, ensure thorough re-agitation of the spray mix until fully resuspended. **DO NOT** allow spray mix to sit overnight, as resuspension may be difficult.

SURFACTANT/WETTING AGENT

Use of a surfactant/wetting agent is not required.

APPLICATION

Use a sprayer fitted with high flow rate nozzles to apply the highest practical spray volume. Use sufficient water to obtain thorough coverage of plants, with a minimum 110 L/ha.

Nozzles with higher rated flows produce larger droplets. Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size, DOES NOT improve canopy penetration and may increase drift potential. WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE. Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. For orchard/vineyard sprayers avoid directing spray above trees and always turn-off outward pointing nozzles at row ends and outer rows.

Dilute Spraying

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of runoff. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.
- Always apply sufficient water to cover the crop to the point of run-off, otherwise under dosing will occur and disease control may be inadequate.

Concentrate Spraying

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.

- Determine an appropriate dilute spray volume (see Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:

Example Only

1. Dilute spray volume as determined above: For example 1500 L/ha
2. Your chosen concentrate spray volume: For example 500 L/ha
3. The concentration factor in this example is : 3 times (i.e. 1500 L divided by 500 L = 3). DO NOT use a concentration factor greater than 3X, for concentrate application.
4. If the dilute label rate is 150 mL/100 L, then the concentrate rate becomes 3 x 150, that is, 450 mL/100 L of concentrate spray.

- The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows.
- For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

COMPATIBILITY

Fontelis® is compatible with many commonly used fungicides, liquid fertilisers, herbicides, insecticides, and biological control products. However, since the formulations of products are always changing, it is advisable to test the physical compatibility of desired tank mixes and check for adverse effects like settling out or flocculation. To determine the physical compatibility, add the recommended proportions of the tank mix products to water, mix thoroughly and allow to stand for 20 minutes. If the combination remains mixed, or can be re-mixed readily, it is considered physically compatible. Fontelis® is compatible with Altacor, Avatar, Dithane®, Polyram and Systhane®.

The crop safety of all potential tank-mixes, including additives and other pesticides, on all crops has not been tested. Before applying any tank-mix not specifically recommended on this label or other Corteva Agriscience supplemental labelling, the safety to the target crop must be confirmed. To test for crop safety, apply the combination to a small area of the target crop in accordance with the label instructions to ensure that a phytotoxic response will not occur.

Fontelis® contains a percentage of mineral oil in the formulation which may predispose oil sensitive products such as Captan to crop damage if they are not applied according to the restrictions referring to summer oils on their label.

The mixing sequence recommended is: water soluble bags, dry flowable or water dispersible granules, wettable powders, water based suspension concentrates, water soluble concentrates, oil based suspension concentrates (Fontelis®), emulsifiable concentrates, adjuvants and surfactants, soluble fertiliser.

Spray Equipment Cleanout

Prior to application, start with clean, well-maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove. Drain spray equipment. Thoroughly rinse sprayer and flush hoses, boom, and nozzles with clean water.
Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. **DO NOT** clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

RE-ENTRY PERIOD

DO NOT allow entry into treated areas until the spray has dried. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), chemical resistance gloves and footwear. Clothing must be washed after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

IMPORTANT: Not all crops within a crop group, and not all varieties, cultivars or hybrids of crops, have been individually tested for crop safety. To test for crop safety, apply the product in accordance with the label instructions to a small area of the target crop to ensure that a phytotoxic response will not occur, especially where the application is a new use of the product by the applicator.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. **DO NOT** contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank. Do not dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be taken separately.

If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

NOTICE TO BUYER

To the extent permitted by the Competition and Consumer Act (2010) or any relevant legislation of any State or Territory (the "Legislation") all conditions and warranties and statutory or other rights of action, whether arising in contract or tort or whether due to the negligence of Corteva Agriscience or Seller, which buyer or any other user may have against Corteva Agriscience or Seller are hereby excluded provided however that any rights of the buyer pursuant to non-excludable conditions or warranties of the Legislation are expressly preserved. Corteva Agriscience hereby gives notice to buyer and other users that to the extent permitted by the Legislation it will not accept responsibility for any indirect or consequential loss of whatsoever nature arising from the storage, handling or use of this Product. Where permitted by the Legislation Corteva Agriscience 's liability shall in all circumstances be limited to the replacement of the product, or a refund of the purchase price paid therefore.

The Product must be used and applied strictly in accordance with the label instructions and other directions for use. It is impossible to eliminate all risks associated with the use of this product. Such risks may arise from factors such as weather conditions, soil factors, off target movement, unconventional technique, presence of other materials, the manner

of use or application, or other unknown factors, all of which are beyond the control of Corteva Agriscience or the Seller. Buyer accepts these risks.

APVMA Approval No. : 65510/133739

Corteva Agriscience Australia Pty Ltd A.B.N. 003 771 659 769
67 Albert Avenue, Chatswood, NSW 2067
www.corteva.com.au
CUSTOMER SERVICE TOLL FREE **1-800 700 096**

™ ® Trademarks of Corteva Agriscience and its affiliated companies. © Corteva.

Barcode
for stock
identification

Hazard and precautionary statements according to classification under GHS (Globally Harmonised System of Classification and Labelling)
May cause an allergic skin reaction. Very toxic to aquatic life with long lasting effects. Avoid breathing dust / fumes / spray. Wear protective gloves. Contaminated clothing should not be allowed out of the workplace. Wash contaminated clothing before re-use.