READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Prodigy®

INSECTICIDE

ACTIVE CONSTITUENT: 240 g/L METHOXYFENOZIDE

GROUP 18 INSECTICIDE

For the control of lightbrown apple moth on apples & pears (pome fruit), blueberries, citrus, grapevines and kiwifruit; and for the control of various other lepidopteran pests of almonds, apples & pears (pome fruit), avocado, coffee, custard apple, eggplant, longan, lychee, peppers (capsicum and chilli), okra, macadamia and tomatoes as specified in the Directions for Use.

READ THE ATTACHED BOOKLET BEFORE USE
SHAKE WELL BEFORE USE

Pack Sizes: 1L, 5L

EMERGENCY RESPONSE (ALL HOURS)
RING FROM ANYWHERE IN AUSTRALIA
1800 370 754
(LOCAL CALL FEE ONLY)

IN A TRANSPORT EMERGENCY ONLY
DIAL 000
FOR POLICE OR FIRE BRIGADE

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

SAFETY DIRECTIONS

- When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves.
- After each day's use wash gloves and contaminated clothing.

SAFETY DATA SHEET

Additional information is listed on the Safety Data Sheet for **PRODIGY® INSECTICIDE** which is available from Corteva Agriscience on request. Call Customer Service Toll Free on 1-800 700 096 or visit www.corteva.com.au

CORTEVA™ agriscience

™ ® Trademarks of Corteva Agriscience and its affiliated companies.

Visit us at Corteva.com.au

RESTRAINTS:
DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto adjacent areas, particularly wetlands, waterbodies or watercourses.
DO NOT spray across open bodies of water.
DO NOT apply if rain or overhead irrigation is expected within 6 hours. Retain the first flush of irrigation tailwater / storm water in the tailwater dam after application.
DO NOT use more than 3 sprays per season in any single crop.
DO NOT use in covered or protected situations such as glasshouses, greenhouses, shade houses or plastic tunnels.
DO NOT apply by air.
DO NOT apply more than three times to any Almond crop.
DO NOT apply more than once at the hull split stage.

DIRECTION FOR USE

USE ON TREE AND VINE CROPS

In the following table, all rates given are for **dilute spraying**. For **concentrate spraying**, refer to the **MIXING/APPLICATION** section. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. For concentrate spraying, **DO NOT** use at rates greater than 5 times the dilute spraying rate.

CROP	PEST	RATE/ 100 L	CRITICAL COMMENTS
Almonds	Carob moth (1st generation)	80mL + Wetter	Practice good orchard hygiene with the removal of mummified fruit in winter. Early post-flower application: At the beginning of the season (Aug-Sept), carefully monitor pheromone traps for peak adult pest incidence of 1st generation. Once regular field scouting indicates egg laying and newly hatched larvae apply a maximum of 3 sprays of Prodigy® at a minimum of 10-day intervals between sprays. Apply to complete coverage. Refer to WETTING AGENTS in the GENERAL INSTRUCTIONS section for more information
	Carob moth (2nd generation)		Hull-split application: Prodigy® should be applied at the start of hull split (2 – 5% of the almond hull sutures open). Carefully monitor pheromone traps for peak activity of 2nd generation moths. Continue field scouting for egg-laying activity and apply Prodigy® to coincide with the initiation of egg hatch. Apply to ensure complete coverage of all foliage and fruit (hull/shell) surfaces. Concentrate spraying is not appropriate for this pest. Do not apply less than 2000 L/ha water.
Apples and Pears (and other pome fruit)	Lightbrown apple moth	25 mL	For control of the **spring generation** of lightbrown apple moth, commence spraying at petal fall and apply in a series of 3 applications at 14-day intervals. Alternatively, in areas where a later **summer generation** occurs, spray when moth activity is indicated by pheromone traps or lure pots. Apply a minimum of 2 sprays with a 21-day interval. Ensure thorough coverage of developing fruit clusters. When insect pressure is high in blocks with large, vase-shaped trees and with short-stemmed, bunchy varieties, Prodigy® must be applied by dilute spraying to ensure run-off into fruit clusters.
	Loopers		Control of these pests is best achieved using a schedule of 3 sprays of Prodigy® in spring (see above).
Avocado	Avocado leafroller	25 mL	Crop monitoring is recommended from pre-bloom. Apply at first sign of pest incidence and target eggs and newly hatched larvae. Additional sprays may be required if re-infestation occurs.
Blueberry	Lightbrown apple moth	25 mL	Thorough coverage of the plant is essential. Target eggs and newly hatched larvae. Apply when locally determined thresholds are exceeded.

CROP	PEST	RATE/ 100 L	CRITICAL COMMENTS
Citrus	Lightbrown apple moth	25 mL + a wetting agent at 10 mL/100L	***Use with high volume spraying equipment only.*** Thorough coverage is essential. Spray to runoff. Apply when eggs and very small larvae are first seen in flower clusters or developing fruitlets. A second spray may be required 2-3 weeks later if larvae hatch over an extended period.
Coffee	Avocado leafroller	25 mL	Crop monitoring for pest incidence is recommended from pre-bloom. Apply at first sign of pest incidence, targeting eggs and very small larvae before they are protected in retreats. Additional sprays may be required if re-infestation occurs.
Custard apple	Yellow peach moth	25 mL	Apply when locally determined pest thresholds are exceeded as indicated by field checks. Additional sprays may be required if re-infestation occurs. Spray to thoroughly cover fruit.
Grapevines	Lightbrown apple moth	25 mL	Thorough leaf tip and bunch coverage is essential. ***Spring generation*:** Apply from pre-flowering onwards targeting eggs and small (5-6 mm long) larvae (up to 3rd instar). Apply a second spray 14 - 21 days later. ***Summer generation*:** Apply at first sign of activity targeting eggs and larvae up to 5-6 mm long (up to 3rd instar). The summer generation can only be detected by monitoring.
Kiwifruit	Lightbrown apple moth	25 mL	Monitor pest activity from bud burst onwards. Target sprays at eggs and first instar larvae.
Longan	Macadamia nutborer	40 mL	Spray thoroughly when pest numbers reach economic threshold levels according to field checks. Target sprays against eggs and early instar larvae.
Lychee			
Macadamia	Macadamia flower caterpillar	25 mL	Monitor for eggs and very small larvae on flowers and apply at a threshold of 50-80% of racemes infested.
	Macadamia nutborer	40 mL	Spray to thoroughly cover nuts when pest numbers reach economic threshold levels according to field checks. Target sprays against eggs and early instar larvae.

USE ON NON-TREE AND VINE CROPS

Note: Prodigy® is a moult-accelerating insecticide that requires ingestion for control. Feeding ceases almost immediately after ingestion. Larvae in protected feeding sites (*eg*. flowers) will not be controlled. Mortality of larvae will not be evident until 4-6 days after application.			
CROP	**PEST**	**RATE**	**CRITICAL COMMENTS**
Tomatoes, Peppers (capsicum and chilli), Eggplant and Okra	Native budworm Tomato grub Cluster caterpillar	125 or 170 mL/100 L or 1.25 or 1.7 L/ha	Use the higher rate under heavy egg pressure. Apply Prodigy® to brown eggs or at egg hatch when pest numbers reach treatment threshold levels as determined by field checks. Maintain field checks and reapply after 7 days if necessary. Ensure thorough coverage of plants.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

Tomato, Peppers (Capsicum and Chilli), Eggplant and Okra: NOT REQUIRED WHEN USED AS DIRECTED

Citrus fruits: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

Longan: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION.

Almond: DO NOT APPLY WITHIN 7 DAYS OF HARVEST.

Blueberry: DO NOT APPLY WITHIN 7 DAYS OF HARVEST.

Avocado, Coffee, Custard apple, Kiwifruit, Lychee and Pome fruit: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

Grapes (for domestic consumption): DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION.

Grapes (for export or wine production): Consult your winery, industry spray diary or peak industry body for the recommended withholding period (export harvest interval) required to meet export residue requirements.

Macadamia: DO NOT HARVEST FOR 28 DAYS AFTER APPLICATION.

GRAZING WITHHOLDING PERIOD

All crops, orchards, plantations and vineyards:
DO NOT graze treated crops
DO NOT allow livestock to graze within treated orchards, plantations or vineyards.

Grapes, Blueberries and Coffee:
do not apply when there are livestock, pasture or any land that is producing feed for livestock within 5 m downwind of the application area.

Tomato, Peppers, Eggplants and Okra:
DO NOT apply when there are livestock, pasture or any land that is producing feed for livestock within 15 m downwind of the application area.

TRADE ADVICE
NOTE – Crops intended for export: Prodigy® insecticide may leave detectable chemical residues in harvested produce. Overseas markets may not have appropriate residue tolerances in place or may have established tolerances which are lower than Australian maximum residue limits. Some crops for export to these destinations may require a longer harvest withholding period. If you plan to use this product on crops destined for export, please check with your exporter before using Prodigy® insecticide.

GENERAL INSTRUCTIONS
Prodigy® Insecticide is very suitable for use in orchards and vineyards using integrated pest management (IPM) programs.

MIXING and APPLICATION
MIXING
Prior to pouring, shake container vigorously, then add the required quantity of Prodigy® to water in the spray vat while stirring or with agitators in motion.

COMPATIBILITY
Prodigy® is incompatible with mineral spray oils. **DO NOT** mix concentrates together but add each to the spray tank separately. As formulations of other manufacturers' products are beyond the control of Corteva Agriscience, all mixtures should be tested prior to mixing commercial quantities. As changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays Corteva Agriscience cannot be responsible for the behaviour of such mixtures.

WETTING AGENTS
Not all surfactants or crop oils are of equal quality. Corteva Agriscience does not support the use of alternative products other than those listed below.
Agral® Spray adjuvant, Nufarm Chemwet 1000.
If intending to use other wetting agents consult Corteva Agriscience.

DILUTE SPRAYING
- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the DIRECTIONS FOR USE table for each 100 L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changes as the crop grows.

CONCENTRATE SPRAYING
- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (see **Dilute Spraying** above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way.

EXAMPLE ONLY

1. Dilute spray volume as determined above: For example, 1500 L/ha
2. Your chosen concentrate spray volume: For example, 500 L/ha
3. The concentration factor in this example is 3X (i.e. 1500 L ÷ 500L = 3)
4. If the dilute label rate is 25 mL/100 L, then the concentrate rate becomes 3 x 25, that is; 75 mL/100 L of concentrate spray.

- The chosen spray volume and the sprayer set up and operation may need to be changed as the crop grows.
- DO NOT use a concentrate rate higher than that specified in the CRITICAL COMMENTS.
- For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

PRECAUTION

Re-entry

DO NOT allow entry into treated areas until the spray has dried. If prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

INSECTICIDE RESISTANCE WARNING

GROUP	18	INSECTICIDE

For insecticide resistance management Prodigy® Insecticide is a Group 18 insecticide. Some naturally occurring insect biotypes resistant to Prodigy® Insecticide and other Group 18 insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Prodigy® or other Group 18 insecticides are used repeatedly. The effectiveness of Prodigy® on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Corteva Agriscience Australia Pty Ltd accepts no liability for any losses that may result from the failure of Prodigy® to control resistant insects. Prodigy® may be subject to specific resistance management strategies. For further information contact your local supplier or Corteva Agriscience representative or local agricultural department agronomist.

PRECAUTIONS:

DO NOT apply using spray equipment carried on the back of the user.

RE-ENTRY

DO NOT allow entry into treated areas until the spray has dried. If prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day’s use.

PROTECTION OF LIVESTOCK

Low hazard to bees. May be applied on plants at any time. Keep livestock out of operational areas during treatment. Avoid spray drift onto adjoining properties or stock areas.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS, AND ENVIRONMENT

- TOXIC to aquatic life with long lasting effects.
- **DO NOT** contaminate streams, rivers or waterways with the chemical, used containers or containers used for mixing or holding treated seed.

STORAGE AND DISPOSAL

Keep Out of Reach of Children.
Store in the closed, original container in a cool, well ventilated area.
DO NOT store for prolonged periods in direct sunlight.
DO NOT store near food, feedstuffs, fertilisers or seed.

The method of disposal of the container depends on the container type. Read the STORAGE AND DISPOSAL instructions on the label that is attached to the container.

1L container
Rinse container before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. Dispose of at a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty container and product should not be burnt.

5L container
This container can be recycled if it is clean, dry free of visible residues and hos the drumMUSTER logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate by adding to the spray tank. DO NOT dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management site. The cap should not be replaced but may be taken separately. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SPILL AND LEAK MANAGEMENT
Do not touch or walk through spilled material. Wear a face shield or goggles, overalls buttoned to neck and wrist, chemical resistant gloves and footwear. Stop leak when safe to do so. Dam area and prevent entry into waterways, and drains.
Small spills/leaks: Absorb with material such as sand, soil or sawdust. Collect spilled product and place in sealable container for disposal. Spill residues may be cleaned using water and detergent. Contain and absorb wash water for disposal. Absorb and collect washings and place in the same sealable container for disposal. Dam the area of large spills and report them to Corteva Agriscience Emergency Services at 1800-370-754.

APVMA Approval No. : 61605/100302

Corteva Agriscience **Australia Pty Ltd** A.B.N. 24 003 771 659
67 Albert Avenue, Chatswood, NSW 2067
www.corteva.com.au
CUSTOMER SERVICE TOLL FREE **1-800 700 096**

™ ® Trademarks of Corteva Agriscience and its affiliated companies. © 2021 Corteva.

Barcode
for stock
identification