CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Success® Neo

Jemvelva® active

INSECTICIDE

ACTIVE CONSTITUENT: 120 g/L SPINETORAM

GROUP 5 INSECTICIDE

For the control of certain insect pests in fruit, tree nuts, herbs, ornamentals, vegetables, canola, cotton, pulses, soybeans, fodder beets, forage brassicas and forestry (*Eucalyptus* spp. and Tea Tree) as specified in the Directions for Use.

Contents: 1 L, 5 L &10 Litres

FIRST AID
If poisoning occurs, contact a doctor or Poison Information Centre.
Phone: *Australia* 13 11 26.

EMERGENCY RESPONSE (ALL HOURS)
RING FROM ANYWHERE IN AUSTRALIA
1800 370 754
(LOCAL CALL FEE ONLY)

IN A TRANSPORT EMERGENCY ONLY
DIAL 000
FOR POLICE OR FIRE BRIGADE

SAFETY DIRECTIONS

- May irritate the eyes. Avoid contact with the eyes.
- When using together with other products consult their safety directions.
- When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves.
- Wash hands after use. After each day's use, wash gloves and contaminated clothing.

SAFETY DATA SHEET
Additional information is listed on the Safety Data Sheet for **SUCCESS® NEO JEMVELVA® ACTIVE INSECTICIDE** which is available from Corteva Agriscience on request. Call Customer Service Toll Free on 1-800 700 096 or visit www.corteva.com.au

CORTEVA™ agriscience

™ ® Trademarks of Corteva Agriscience and its affiliated companies.

Visit us at Corteva.com.au

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.
DO NOT allow bystanders to come into contact with the spray cloud.
DO NOT allow irrigation water from treated paddocks to enter adjacent pastures, crops or water supplies.
DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone tables below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.
DO NOT apply unless the wind speed is between three (3) and twenty (20) kilometres per hour at the application site during the time of application.
DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

Boom sprayers

DO NOT apply by a boom sprayer unless the following requirements are met:

- Spray droplets are not smaller than a MEDIUM spray droplet size category.
- Minimum distances between the application site and downwind sensitive areas are observed (see ‘Mandatory buffer zones’ section of the following table titled ‘Buffer zones for boom sprayers’).

Buffer zones for boom sprayers

Application rate	Boom height above the target canopy	Mandatory downwind buffer zones	
		Natural aquatic areas	Pollinator areas
Up to 500 mL/ha	0.5 m or lower	20 metres	Not required
	1.0 m or lower	60 metres	25 metres
Up to 300 mL/ha	0.5 m or lower	10 metres	Not required
	1.0 m or lower	40 metres	20 metres

Vertical sprayers

DO NOT apply by a vertical sprayer unless the following requirements are met:

- Spray is not directed above the target canopy.
- The outside of the sprayer is turned off when turning at the end of the rows and when spraying the outer row on each side of the application site.

- For dilute water rates up to the maximum listed for each type of canopy specified, minimum distances between the application site and downwind sensitive areas are observed (see the following table titled 'Buffer zones for vertical sprayers').

Buffer zones for vertical sprayers

Application rate	Type of target canopy and dilute water rate	Mandatory downwind buffer zones	
		Natural aquatic areas	Pollinator areas
Up to 40 mL / 100 L	2 metres tall and shorter, maximum dilute water rate of 1,000 L/ha	10 metres	Not required
	Taller than 2 metres (not fully foliated), maximum dilute water rate of 4,000 L/ha	40 metres	25 metres
	Taller than 2 metres (fully foliated), maximum dilute water rate of 4,000 L/ha	30 metres	20 metres
	Taller than 2 metres (not fully foliated), maximum dilute water rate of 10,000 L/ha	60 metres	40 metres
	Taller than 2 metres (fully foliated), maximum dilute water rate of 10,000 L/ha	50 metres	30 metres

Aircraft

DO NOT apply by aircraft unless the following requirements are met:

- Spray droplets not smaller than a MEDIUM spray droplet size category.
- For maximum release height above the target canopy of 3 metres or 25 per cent of wingspan or 25 per cent of rotor diameter, whichever is the greatest, minimum distances between the application site and downwind sensitive areas are observed (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for aircraft').

Buffers zones for aircraft

Application rate	Type of aircraft	Mandatory downwind buffer zones	
		Natural aquatic areas	Pollinator areas
Up to 500 mL/ha	Fixed-wing	250 metres	100 metres
	Helicopter	170 metres	80 metres
Up to 250 mL/ha	Fixed-wing	150 metres	60 metres
	Helicopter	110 metres	50 metres

DIRECTIONS FOR USE:
Table 1 - BROADACRE CROPS: Canola, Chickpeas, Cotton and Pulses

RESTRAINTS (specific to broadacre crops)

DO NOT apply using Ultra Low Volume methods.

DO NOT apply by aircraft, except for applications to canola, cotton and pulses.

DO NOT apply more than two (2) applications to each of the following crops in any one season: Canola, chickpea, pulses and soybean.

DO NOT apply more than three (3) applications to each of the following crops in any one season: Cotton.

ALL CROPS: Carefully monitor crops for eggs and larvae of pest species by regular field scouting. Target sprays against mature eggs and newly-hatched larvae when numbers exceed local spray threshold. Apply repeat applications at 7-14 day intervals as new infestations occur or as specified under Critical Comments. Any subsequent sprays to control insects in that crop should be made with a product from a different chemical group.

CROP	PEST	RATE	CRITICAL COMMENTS
Canola	Diamondback moth (cabbage moth)	100 mL/ha + Uptake® spraying oil at 100 mL/ha or a non-ionic wetting agent❶	Success® Neo can be applied **once** at any time up to 14 days before harvest (windrowing). If initially applied at any time up to early pod formation, then a **second** application can be made from 7 days after the 1st application or at any time up to 14 days before harvest. Apply in a minimum of 50 L/ha water by ground or 30 L/ha by air. Larvae that are entrenched (hidden in leaves, stems or pods) will not be controlled.
	Cabbage cluster caterpillar, Cabbage white butterfly, Centre grub, Corn earworm (*heliothis*), Native budworm	150 mL/ha + Uptake® spraying oil at 100 mL/ha or a non-ionic wetting agent❶	
	Lucerne flea (springtails)	20 mL/ha + Uptake spraying oil at 100 mL/ha or a non-ionic wetting agent❶	For applications to seedling canola up to 6 leaf. Assess for presence of springtails. Use an accepted monitor method, such as by placing an A4 sheet of white paper on the ground and vigorously banging the ground on either side, once or twice, with a large hammer or similar object. If present, springtails will jump onto the paper.
Chickpeas	Heliothis (*Helicoverpa* spp.)	150 – 200 mL/ha	Product can be applied either twice to the crop during flowering or once at flowering and once up to 14 days before harvest. Use the low rate against light infestations of newly emerged larvae and higher rates when infestation is heavy and/or larvae are more advanced. Note: Entrenched larvae, or those not actively feeding, will not be controlled.

Cotton	Bollworm (*Helicoverpa armigera*) Native budworm (*Helicoverpa punctigera*)	300 – 400 mL/ha	Use the low rate against light infestations and higher rates when infestation is heavy (see Guidelines below). Carefully monitor eggs and larvae of Helicoverpa species by regular field scouting. Target sprays against brown eggs and newly hatched very small larvae. **Guidelines:** **Light infestation:** Use 300 mL/ha when infestation of Helicoverpa species is less than 10 eggs and 2 larvae/m of row. **Heavy infestation:** Use 400 mL/ha when infestation of Helicoverpa species exceeds 10 eggs and/or 2 larvae per metre of row. Note: Larvae larger than 8mm in length, and larvae feeding within bolls and squares may not be controlled.
	Western flower thrip (*Frankliniella occidentalis*)	400 mL/ha + non-ionic wetting agent❶	Product can be applied either twice to the crop during flowering or once at flowering and once up to 14 days before harvest. Use this product as part of the WFT Resistance Management strategy (see end of table 4 for details).
Pulses (including but not limited to adzuki beans, cowpeas, faba beans, field peas, lentils, lupins, kidney beans, mungbeans and navy beans)	Heliothis (*Helicoverpa* spp.) Loopers (*Chrysodeixis* spp.) Soybean looper (*Thysanoplusia orichalcea*)	200 – 300 mL/ha	Product can be applied either twice to the crop during flowering or once at flowering and once up to 14 days before harvest. Use the low rate against light infestations and higher rates when infestation is heavy (see Guidelines below). Carefully monitor eggs and larvae of pests by regular field scouting. To achieve best results target sprays against brown eggs and newly hatched, very small larvae. **Light infestation:** Use 200 mL/ha when infestation of Helicoverpa species is less than 10 eggs and 2 larvae/m of row (pulse crops). **Heavy infestation:** Use 300 mL/ha when infestation of Helicoverpa species exceeds 10 eggs and/or 2 larvae per metre of row (pulse crops). **For Southern Australia:** If determining insect thresholds by sweep netting, follow the recommendations of your local Department of Agriculture or equivalent advice. **Note:** Entrenched larvae, or those not actively feeding, will not be controlled.
Soybean			

Table 2 - FORAGE CROPS: Fodder Beets and Forage Brassicas

RESTRAINTS (specific to forage and fodder crops)

DO NOT apply using Ultra Low Volume methods.

DO NOT apply more than two (2) applications to any one crop in any one season.

ALL CROPS: Carefully monitor crops for eggs and larvae of pest species by regular field scouting. Target sprays against mature eggs and newly-hatched larvae when numbers exceed local spray threshold. Any subsequent sprays to control insects in that crop should be made with a product from a different chemical group.

CROP	PEST	RATE	CRITICAL COMMENTS
Fodder beets	Beet webworm (*Spoladea recurvalis*)	200 mL/ha + Uptake spraying oil 100 mL/ha (suppression only)	Target mature eggs and young larva when pest numbers exceed threshold. This treatment will provide knockdown of light infestations of early instar larvae in crops where good spray coverage has been achieved. Larvae that are entrenched (hidden in leaves, stems, bulbs or heads) may not be controlled. Minimum retreatment interval 7 to 10 days.
Forage Brassicas including but not limited to choumoellier, kale, fodder rape, swedes, turnips.	Diamondback moth (cabbage moth), Cabbage white butterfly, Cabbage cluster caterpillar, Centre grub, Corn earworm (*heliothis*), Native budworm, Soybean looper	100 mL/ha + Uptake® spraying oil 100 mL/ha or a non-ionic wetting agent❶	Apply in a minimum of 50 L/ha water. This treatment will provide knockdown of light infestations of early instar larvae in crops where good spray coverage has been attained. Larvae that are entrenched (hidden in leaves, stems, bulbs or heads) may not be controlled.
	Lucerne flea (springtails)	20 mL/ha + Uptake spraying oil at 100 mL/ha or a non-ionic a wetting agent❶	For applications to seedling canola up to 6 leaf. Assess for presence of springtails. Use an accepted monitor method such as by placing an A4 sheet of white paper on the ground and vigorously banging the ground on either side, once or twice, with a large hammer or similar object. If present, springtails will jump onto the paper.

Table 3 - VEGETABLE CROPS, ORNAMENTALS and HERBS

Brassica vegetables, bulb vegetables, cucurbits, culinary herbs, fruiting vegetables, leafy vegetables, legume vegetables, root and tuber vegetables, stalk and stem vegetables, sweet corn and ornamentals

RESTRAINTS (specific to vegetable crops, ornamentals and herbs)

The use of Success Neo® in protected cropping is permissible when used in conjunction with a proven Insect Resistance Management Strategy (IRMS).
DO NOT apply to seedlings of edible crops for transplanting i.e. seedling intended for food production. **DO NOT** apply to vegetable or herb seedlings in a production nursery system.

DO NOT apply using Ultra Low Volume methods.

DO NOT apply by aircraft.

DO NOT make more than four (4) applications to each of the following crops in any one season (also see the **RESISTANCE** statement): Brassica vegetables, cucurbits, culinary herbs, fruiting vegetables, leafy vegetables, root and tuber vegetables, stalk and stem vegetables, and sweet corn.

DO NOT apply more than three (3) applications to each of the following crops in any one season: Bulb vegetables and legume vegetables.

ALL VEGETABLES: Carefully monitor crops for eggs and larvae of pest species by regular field scouting. Target sprays against mature eggs and newly-hatched larvae when numbers exceed spray threshold. Apply repeat applications at 7-14 day intervals as new infestations occur or as specified under Critical Comments. As part of IPM programs for potato moth, Helicoverpa and diamondback moth, it is important to plough crops in immediately after harvest.

CROP	PEST	RATE	CRITICAL COMMENTS
Brassica vegetables; Including but not limited to Broccoli Brussels sprouts Cabbage Cauliflower Brassica Leafy vegetables	Diamondback moth, Cabbage white butterfly, Cabbage cluster caterpillar, Cabbage centre grub, Loopers	200 mL/ha + non-ionic wetting agent❶	Use a minimum spray volume of 250 L/ha and ensure thorough crop coverage by increasing water volume with plant growth stage.

Table 3 continued

CROP	PEST	RATE	CRITICAL COMMENTS
(**see list at end of table* 3*)* Radishes[1] Swedes[1] Turnips[1] [1]*See also under Root & Tuber Vegetables below)*	*Helicoverpa*	200 – 400 mL/ha + non-ionic wetting agent❶	Use the lower rate when good coverage can be achieved and the high rate in maturing crops if crop canopies prevent good coverage.
	Cluster caterpillar	300 mL/ha + non-ionic wetting agent❶	
	Western flower thrips	400 mL/ha + non-ionic wetting agent❶	Use this product as part of the WFT Resistance Management strategy (*see end of table 4 for details*).
Bulb vegetables; Including but not limited to bulb onion, spring onion and leek	Western flower thrips	400 mL/ha + non-ionic wetting agent❶	Use this product as part of a WFT Resistance Management strategy (*see end of table 4 for details*).
	Onion thrips (*Thrips tabaci*)	500 mL/ha + non-ionic wetting agent❶	Use as part of a season-long program utilizing all IPM techniques. Apply up to three “back to back” applications of Success Neo at 7 day intervals. Always apply Success Neo in sufficient water to ensure thorough coverage of foliage.
Cucurbits; including but not limited to cucumbers, melons, squash & zucchini	Cucumber moth, *Helicoverpa*	200 - 400 mL/ha	Use higher rates during periods of high insect pressure or when crop coverage is difficult.
	Western flower thrips	400 mL/ha	Use this product as part of the WFT Resistance Management strategy (*see end of table 4 for details*).
Culinary Herbs (***see list at end of table* 3)	Diamondback moth (cabbage moth), Loopers, Lightbrown apple moth	200 mL/ha + non-ionic wetting agent❶	Use a maximum spray volume of 250 L/ha. Ensure thorough coverage of the target area by increasing water volume with plant growth stage.
	Helicoverpa	200 - 400 mL/ha + non-ionic wetting agent❶	As above, plus use the lower rate when good coverage can be achieved and the high rate in maturing crops if crop canopies prevent good coverage.

Table 3 continued

CROP	PEST	RATE	CRITICAL COMMENTS
Fruiting vegetables; including but not limited to Eggplant Okra Peppers (Capsicums & Chillies), Sweet corn (see also under separate listing below) Tomatoes	Potato moth (tomato leaf miner) Helicoverpa	200 - 400 mL/ha or Dilute 20 - 40 mL/100L	Use the per hectare rate when applying to bush tomatoes and the dilute rate (per 100 L) in trellised crops (see the "DILUTE SPRAYING" Section in this booklet). Use the lower rate as part of an IPM program when Helicoverpa is the dominant pest and good crop coverage is possible. Use higher rates during periods of high insect pressure or when crop coverage is difficult. Addition of a non-ionic wetting agent may improve control.
	Western flower thrips	400 mL/ha or Dilute 40 mL/100L	Use this product as part of the WFT Resistance Management strategy (*see end of table 4 for details*).
	Tomato potato psyllid (*Bactericera cockerelli*)	400 mL/ha (suppression only) + non-ionic wetting agent[1]	Use as part of a season long program utilising all IPM techniques. Apply up to four "back to back" applications of Success Neo at 7 day intervals. Always apply Success Neo in sufficient water to ensure thorough coverage of foliage.
Leafy vegetables; including but not limited to Lettuce, Endive Silverbeet, Spinach & Brassica leafy vegetables (*see list at end of table 3)	Loopers	200 mL/ha	See above under "***ALL VEGETABLES***" Use the lower rate as part of an IPM program when Helicoverpa is the dominant pest and good crop coverage is possible. Use higher rates during periods of high insect pressure or when crop coverage is difficult.
	Helicoverpa	200 - 400 mL/ha	
	Western flower thrips	400 mL/ha	Use this product as part of the WFT Resistance Management strategy (*see end of table* 4 *for details*).
Legume vegetables (succulent seeds and immature pods only); including but not limited to beans, peas, snow peas and sugar snap peas.	Loopers	200 mL/ha	Use higher rates during periods of high insect pressure or when crop coverage is difficult. **Note:** Entrenched larvae will not be controlled.
	Helicoverpa	200 - 400 mL/ha	
	Western flower thrips	400 mL/ha	Use this product as part of the WFT Resistance Management strategy (see end of table 4 for details).

Table 3 continued

CROP	PEST	RATE	CRITICAL COMMENTS
Root and tuber vegetables; including but not limited to Beetroot, Carrots Celeriac, Galangal Parsnips Potatoes Radishes (incl. Daikon) Sweet potato Swedes, Turnips	Lightbrown apple moth, Loopers	200 mL/ha	See above under "**ALL VEGETABLES**" Use the lower rate when good coverage can be achieved and the high rate in maturing crops if crop canopies prevent good coverage. Entrenched larvae will not be controlled. Only target foliar infestations of potato moth. Potato moth larvae within stems or below the soil will not be controlled.
	Helicoverpa	200 – 400 mL/ha	
	Potato moth	200 - 400 mL/ha + non-ionic wetting agent❶	
	Tomato potato psyllid (*Bactericera cockerelli*)	400 mL/ha (suppression only) + non-ionic wetting agent❶	Use as part of a season long program utilizing all IPM techniques. Apply up to four "back to back" applications of Success® Neo at 7 day intervals. Always apply Success® Neo in sufficient water to ensure thorough coverage of foliage.
Stalk & Stem vegetables; including but not limited to Celery and Rhubarb	*Helicoverpa*	400 mL/ha	See comments under "**ALL VEGETABLES**" above
Sweet corn (*see also under Fruiting Vegetables above*)	*Helicoverpa*	200 – 400 mL/ha	Use higher rates during periods of high insect pressure or when crop coverage is difficult.
Ornamentals	Pear and cherry slug	10 mL/100L	Apply when infestation first identified. Repeat applications at no less than 10 day intervals. Caterpillars feeding in entrenched sites may not be controlled.
	Caterpillars	20 mL/100L	
	Western flower thrips	40 mL/100L	Use this product as part of the WFT Resistance Management strategy (*see end of table* 4 *for details*).
***Brassica Leafy Vegetables:** Includes Pak choi, Bok choi, Choi sum, Chinese broccoli (Gai lum/Gai lan/Kai lan), Chinese cabbage (Pet sai/Wong bok/Haksukai), Mibuna, Mustard spinach (Komatsuma), Kale, Indian mustard, Kai choi, Gai choi/Am soi, Tat soi and Leafy mustard.			
** **Culinary Herbs:** Includes Basil, Bay leaves, Borage, Chervil, Chives, Coriander, Dill, Fennel, Galangal, Lemon balm, Lemon grass, Lemon verbena, Kaffir lime leaves, Marigold flowers, Marjoram (Oregano), Mints, Mizuna, Nasturtium leaves, Parsley, Rosemary, Sage, Salad Burnett, Sorrel, Tarragon, Thyme, Tumeric, Savory.			

Table 4 - TREE, VINE and BERRYFRUIT CROPS

Avocados, bananas, berryfruit, carob, cocoa, citrus fruit, coffee, kiwifruit, mango, tropical and subtropical fruits, macadamia nuts, pistachio nuts and forestry

RESTRAINTS (specific to tree, vine and berryfruit crops)

DO NOT apply using Ultra Low Volume methods.

DO NOT apply by aircraft except for applications to mango and forestry uses.

DO NOT apply more than four (4) applications to each of the following crops in any one season: Berryfruit carob, citrus, cocoa, coffee, kiwifruit, macadamia nuts, mango or tropical and sub-tropical fruits.
DO NOT apply more than three (3) applications to each of the following crops in any one season: Forestry (Eucalyptus plantations and Tea tree).
DO NOT apply more than two (2) applications to each of the following crops in any one season: Banana or pistachio nuts.
DO NOT apply more than twice during flowering or twice after the completion of flowering to any tree nut crop.
DO NOT apply while bees are actively foraging.

In the following table, all rates (except in FORESTRY) are given for dilute spraying. For concentrate spraying refer to the "Concentrate Spraying" Section in this booklet. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
FOR ALL TREE, VINE & BERRYFRUIT CROPS: Carefully monitor crops for eggs and larvae of pest species by regular field scouting. Target sprays against mature eggs and newly-hatched larvae when numbers exceed spray threshold. Apply repeat applications at 7 - 14 day intervals as new infestations occur unless otherwise directed in the Critical Comments.

CROP	PEST	RATE	CRITICAL COMMENTS
Avocados *(see also under Tropical & Sub-Tropical Fruit Crops below)*	Leafrollers (including Avocado leafroller, Ivy leafroller and Lightbrown apple moth), Loopers (including *Ectropis* looper)	20 mL/100 L+ non-ionic wetting agent[Φ]	See comments under "**FOR ALL TREE, VINE & BERRYFRUIT CROPS**" above

Table 4 continued

CROP	PEST	RATE	CRITICAL COMMENTS
Bananas	Banana rust thrips, Sugarcane bud moth	20 mL/10 L	Bunch spray: Apply as a fine spray to point of run-off (50 - 60 mL of solution) ensuring complete coverage of the bunch. Application should be made no later than 2 weeks after bunch emergence. Application should be made immediately after placement of the bunch cover. Good coverage of the bunch is essential.
Berryfruit; including but not limited to Blackberries Blueberries Boysenberries Cranberries Currants Gooseberries Raspberries Strawberries	Loopers	20 mL/100 L	See comments under "**FOR ALL TREE, VINE & BERRYFRUIT CROPS**" above
	Lightbrown apple moth, *Helicoverpa*	20 – 40 mL/100 L	Use the higher rate in dense canopies and when larvae have begun webbing leaves and fruit. Use the lower rate under an IPM system or where good coverage is assured.
	Western flower thrips	40 mL/100 L	Use this product as part of the WFT Resistance Management strategy (see end of table 4 for details).
Carob	Carob moth	40 mL/100 L Minimum 1 L/ha + non-ionic wetting agent❶	Use as part of an IPM system, ensure good orchard hygiene with the removal of mummified fruit in winter. Application timing is critical. Early post-flower application: at the beginning of the season (Aug-Sep), carefully monitor pheromone traps for peak adult pest incidence of 1st generation. Once regular field scouting indicates egg laying and newly hatched larvae apply Success Neo as required. Apply to complete coverage. Carefully monitor pheromone traps for peak activity of 2nd generation moths. Continue field scouting for egg-laying activity and apply Success Neo to coincide with the initiation of egg hatch. Apply to complete coverage. Apply repeat applications at a minimum of 7 day intervals. Ensure thorough spray coverage of all leaf and pod surfaces with a minimum rate of 1000 L/ha water using an airblast sprayer or equivalent.
Cocao			
Citrus Fruits; including but not limited to Grapefruit, Lemons, Limes, Mandarins Oranges	Citrus leafminer, Lightbrown apple moth	10 - 20 mL/100 L+ non-ionic wetting agent❶	*Citrus leafminer*: Best results will be achieved when horticultural oil is used in place of a wetting agent. Only use oils when applying to non-bearing trees due to the risk of fruit phytotoxicity. For the other pests, use higher rates for heavy infestations
	Helicoverpa (Corn earworm and native budworm)	20 - 40 mL/100 L+ non-ionic wetting agent❶	

Coffee	Avocado leaf roller	20-40 mL/100 L+ non-ionic wetting agent❶	Use higher rates for heavy infestations
Kiwifruit (see also under Tropical & Sub-Tropical Fruit Crops below)	Lightbrown apple moth	20 mL/100 L	See comments under "**FOR ALL TREE, VINE & BERRYFRUIT CROPS**" above
Mango (see also under Tropical & Sub-Tropical Fruit Crops below)	Flower-eating caterpillars, Small mango tip borer	20 mL/100 L+ non-ionic wetting agent❶	
	Large mango tip borer	5 mL/100 L + non-ionic wetting agent❶	
Mango – Aerial Application ONLY	Flower-eating caterpillars, Small mango tip borer, Large mango tip borer	200 mL/ha + non-ionic wetting agent❶	AERIAL APPLICATION ONLY
Tropical and sub-tropical fruit crops (inedible peel); including but not limited to avocado[2], cherimoya, custard apple, durian, feijoa, guava, jackfruit, kiwifruit[2], longan, lychee, mango[2], mangosteen, papaya, passionfruit, persimmon, rambutan and star apple [2](See separate listings above also for these crops)	Flower-eating caterpillars, Leafrollers and loopers, Yellow peach moth	20 mL/100 L	See comments under "**FOR ALL TREE, VINE & BERRYFRUIT CROPS**" above Addition of a non-ionic wetting agent at its recommended rate may improve control on difficult to wet foliage & fruit.
	Red-banded thrips, Sorghum head caterpillar	40 mL/100 L	

Table 4 continued

CROP	PEST	RATE	CRITICAL COMMENTS
Macadamias	Macadamia nut borer Thrips including: Red-banded thrips	40 mL/100 L	Nut borer: Ensure complete spray coverage of nuts and nutlets. Commence applications when pest numbers reach local thresholds and repeat 10-14 days later if pests are still active. Use the shorter interval during wet weather or severe infestations. Addition of a non-ionic wetting agent at its recommended rate may improve control on difficult-to-wet foliage and fruit[1].
	Flower-eating caterpillars, Twig girdler, Yellow peach moth	20 mL/100 L	
Pistachios	Carob moth	1 L/ha + non-ionic wetting agent❶	Application timing is critical – the first application of an insecticide should be when 1-5% of the pistachio hulls are opening. Target sprays against mature eggs and newly hatched larvae. Ensure thorough spray coverage of all leaf and nut surfaces with a minimum rate of 1000 L/ha water using an airblast sprayer or equivalent. Make a follow up application if required.
FORESTRY:			
Eucalyptus Plantations	Larvae of Eucalyptus chrysomelid leaf beetle	25 – 50 mL/ha + sticker or non-ionic wetting agent❶	Use higher concentration for larger larvae and older trees. Larval mortality will not occur for at least four (4) days after spraying. Note that Success® Neo is not effective against adult beetles. **DO NOT** spray if rain expected in the following 24 hr. Follow code of practice for aerial spraying for relevant state, including appropriate buffers.
Tea tree (*Melaleuca* spp.)	Pyrgo beetle	100 - 250 mL/ha + non-ionic wetting agent❶	Closely monitor plantation for egg, larval numbers and age of larvae. Use the higher rate for heavy infestations and for larger tea trees. Apply by ground based application equipment only in a minimum of 100 L/ha water. Use sufficient spray volume to ensure thorough coverage of flush leaf, and adjust spray volumes to stage of crop growth. **For 1st- 2nd instar larvae**, apply 100 mL/ha. **For 3rd- 4th instar larvae**, apply 100-150 mL/ha. **For control of adults** apply 150-250 mL/ha.
[1]DO NOT add an adjuvant if tank-mixing Success® Neo with other products and applying to tree nuts without first consulting Corteva Agriscience.			

[2]See separate listing above also for these crops.
Western Flower Thrip (WFT) Resistance Management Strategy Make 3 consecutive applications at either 3-5 day intervals when temperatures are greater than 20°C or at 6-12 day intervals when temperatures are less than 20°C. For any further sprays required, use an approved product from another chemical group. **DO NOT** make more than 3 consecutive applications of Success® Neo before switching to an approved product from another chemical group.
PEST NAMES: Avocado leafroller: *Homona spargotis;* Banana rust thrips: *Chaetanaphothrips signipennis*; Beet webworm (*Spoladea recurvalis*); Cabbage cluster caterpillar: *Crocidolomia pavonana*; Cabbage centre grub: *Hellula hydralis*; Cabbage white butterfly: *Pieris rapae*; Carob moth: *Ectomyelois ceratoniae;* Citrus leafminer: *Phyllocnistis citrella;* Cluster caterpillar: *Spodoptera litura*; Codling moth: *Cydia pomonella*; Cucumber moth: *Diaphania indica*; Diamondback moth: *Plutella xylostella*; Elm Leaf Beetle: *Pyrrhalta luteola;* Eucalyptus chrysomelid leaf beetle: *Chrysophtharta bimaculata* and *C. Agricola;* Helicoverpa caterpillars, corn earworm, native budworm: *Helicoverpa* spp.; Ivy leafroller: *Cryptoptila immersana;* Large mango tipborer: *Penicillaria jocosatrix;* Lightbrown apple moth: *Epiphyas postvittana*; Loopers: *Chrysodeixis* spp., *Thysanoplusia* spp., and Geometrid loopers, Ectropis looper: *Ectropis savulosa*; Lucerne flea: *Sminthurus viridis;* Macadamia nutborer: *Cryptophlebia ombrodelta;* Onion thrips: *Thrips* tabaci; Potato moth/tomato leaf miner: *Phthorimaea operculella*; Pyrgo beetle: *Paropsistema tigrina;* Red-banded thrips: *Selenothrips rubrocinctus*; Small mango tipborer: *Chlumetia euthysticha;* Sorghum head caterpillar: *Cryptoblabes adoceta*; Soybean looper: *Thysanoplusia orichalcea;* Sugarcane bud moth: *Opogona glycyphaga*; Tomato potato psyllid (*Bactericera cockerelli*); Twig girdler: *Xyloricta luteotactella;* Western flower thrips: *Frankliniella occidentalis*; Yellow peach moth: *Conogethes punctiferalis.*

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

HARVESTING WITHHOLDING PERIODS (WHP)

Bananas, Subtropical and Tropical Fruit (excluding Kiwi Fruit): NOT REQUIRED WHEN USED AS DIRECTED.

Berry Fruit, Citrus Fruit and Fruiting Vegetables (except Sweet Corn), Stalk and Stem Vegetables: DO NOT HARVEST FOR 1 DAY AFTER THE LAST APPLICATION.

Bulb vegetables, Brassicas, Cucurbits, Culinary Herbs, Leafy Vegetables, Legume Vegetables, Root and Tuber Vegetables, Stone Fruit and Sweet Corn: DO NOT HARVEST FOR 3 DAYS AFTER THE LAST APPLICATION.

Coffee, Kiwi Fruit: DO NOT HARVEST FOR 7 DAYS AFTER THE LAST APPLICATION.

Tree nuts (except almonds): DO NOT APPLY WITHIN 7 DAYS OF HARVEST OR NUT FALL.

Canola, Chickpeas, Cocao, other Pulses and Soybeans: DO NOT HARVEST FOR 14 DAYS AFTER THE LAST APPLICATION.

Carob: DO NOT HARVEST FOR 21 DAYS AFTER THE LAST APPLICATION.

Cotton: DO NOT HARVEST FOR 28 DAYS AFTER APPLICATION.

GRAZING AND STOCKFOOD WITHHOLDING PERIOD (WHP)
Also note the instructions in the section LIVESTOCK DESTINED FOR EXPORT MARKETS below

DO NOT GRAZE ANY TREATED ORCHARD OR CUT THE ORCHARD FLOOR FOR STOCKFOOD.

DO NOT GRAZE ANY TREATED FOREST OR CUT THE FOREST FLOOR FOR STOCKFOOD.

Brassica and Leafy Vegetables: DO NOT USE ON BRASSICA VEGETABLES GROWN FOR FORAGE OR FODDER. DO NOT GRAZE TREATED BRASSICA or LEAFY VEGETABLE CROPS.

Cotton: DO NOT GRAZE OR CUT TREATED COTTON CROPs, STUBBLE OR GIN TRASH.

Sweet Corn: DO NOT CUT OR GRAZE FOR STOCKFEED FOR 3 DAYS AFTER APPLICATION. DO NOT FEED SWEET CORN FORAGE OR FODDER TO LACTATING DAIRY ANIMALS PRODUCING MILK FOR HUMAN CONSUMPTION.

Canola, Fodder beets and Forage Brassicas: DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 7 DAYS AFTER APPLICATION.

Pulses and legume vegetables (including chickpeas and soybeans):

Grazing or cutting for meat production: DO NO GRAZE OR CUT FOR STOCKFEED FOR 14 DAYS AFTER APPLICATION.

Grazing or cutting for milk production (forage): DO NOT FEED OR ALLOW LACTATING DAIRY ANIMALS PRODUCING MILK FOR HUMAN CONSUMPTION TO GRAZE TREATED PULSE OR LEGUME VEGETABLE FORAGE.

Grazing or cutting for milk production (fodder, i.e. hay or straw): DO NOT GRAZE OR CUT FODDER FOR STOCKFEED FOR LACTATING DAIRY ANIMALS FOR 14 DAYS AFTER APPLICATION.

TREATED CROPS FOR EXPORT
Some crops for export to particular destinations outside of Australia may require a longer interval before harvest to comply with the residue standards of importing countries, please check with your exporter.

LIVESTOCK DESTINED FOR EXPORT MARKETS
To ensure compliance with residue standards in export markets, comply with the above grazing withholding periods **and then** ensure that the **Export Slaughter Interval (ESI)** is observed before stock are sold or slaughtered.

Export Slaughter Interval:
Forage brassicas: **7 days (OR see EAFI below)**
Chickpeas, other pulses, Soybeans and Legume vegetables: **7 days**
Forage and fodder of sweet corn: **21 days**

This means that livestock that have grazed on or were fed treated crops should be placed on clean feed for at least the periods stated above prior to slaughter.

Export Animal Feed Interval (EAFI) For Forage Brassicas:

For situations where a grazing WHP and ESI are not practical after application to forage brassicas, the following EAFI can be observed:

Do not graze or cut treated forage brassicas for 28 days (4 weeks) after application if animals are intended to be sent directly for slaughter for export.

Sweet corn cannery waste:
When Success® Neo Insecticide is used as directed and the above WHPs are observed, sweet corn cannery waste can be fed to livestock. Animals fed sweet corn cannery waste are considered acceptable to slaughter for export, provided it makes up no more than 10% of the animals' diet for periods exceeding 7 days. If animals are fed exclusively on sweet corn cannery waste there could be a risk of animal residues exceeding export requirements. In this situation it is advisable to transfer stock to untreated feed for at least 14 days before sending to slaughter.

GENERAL INSTRUCTIONS

Success® Neo is a spinosyn insecticide, in the form of a suspension concentration.

INSECTICIDE RESISTANCE WARNING

GROUP	5	INSECTICIDE

For insecticide resistance management, Success® Neo Jemvelva® active Insecticide is a Group 5 insecticide. Some naturally occurring insect biotypes resistant to Success® Neo and other Group 5 insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Success® Neo and other Group 5 insecticides are used repeatedly. The effectiveness of Success® Neo on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Corteva Agriscience Australia Proprietary Limited accepts no liability for any losses that may result from the failure of Success® Neo to control resistant insects. Success® Neo may be subject to specific resistance management strategies. For further information contact your local supplier, Corteva Agriscience representative or local agricultural department agronomist.

MIXING

- Agitate or shake the container immediately prior to use.
- Half fill the spray tank with water, add the appropriate amount of accurately measured Success® Neo Insecticide (suspension concentrate), then complete filling the tank.
- Ensure thorough agitation by mechanical or hydraulic action at all times during mixing and application.
- Use only clean water within the range pH 5 - 9 to dilute Success® Neo Insecticide.

COMPATIBILITY

Success® Neo is compatible with most herbicides, fungicides and insecticides. Always conduct a bucket test to confirm physical compatibility. Always follow the instructions on both products and, if in any doubt, consult Corteva Agriscience, your agronomist or consultant or the maker of the other product.

ADJUVANT

[1] Where required in the Directions for Use tables, if not using Uptake® spraying oil, apply with a 100% non-ionic wetting agent such as BS-1000®[1] Chemwet®[1] 1000 or Spreadwet®[1] 1000 at the manufacturer's recommended rate.

STORAGE OF DILUTED SPRAY MIX

- Whenever possible the spray mix should be used immediately after it is prepared. However, if weather conditions or mechanical breakdown prevent immediate use, the spray mix may be stored for up to 72 hours without loss of activity.
- The spray mix should be agitated thoroughly by mechanical or hydraulic action at regular intervals during storage to prevent sedimentation. Ensure that the stored spray mix is thoroughly agitated at least once every 8 hours.
- The spray mix must be stored out of direct sunlight.

APPLICATION

Thorough coverage of the crop is essential. Ensure this by increasing water volume with plant growth stage. **DO NOT** apply when conditions are unsuitable for water-based spray applications. Avoid high temperature, strong winds, inversion conditions, imminent rain or any conditions that may reduce the quality of spray coverage or result in drift from the target area. Techniques to minimise drift should be employed at all times when aerially applying sprays to, or near, sensitive areas.
For optimum results follow the application specifications listed below:

- Ground Spraying: Apply in a minimum of 250 L/ha of water. Increase spray volumes as the crop grows.
- Ground Spraying forage brassicas and canola, cotton, chickpeas, other pulses and soybean ONLY: Apply in a minimum of 50L/ha of water.
- Aerial Spraying: Success® Neo must only be applied by aircraft fitted with accurately calibrated equipment. Apply a minimum total spray volume of 30 L/ha with nozzles (e.g. Micronnaire rotary atomisers, CP nozzles or conventional hydraulic nozzles) set to medium spray quality according to ASAE S572 definition for standard nozzles. A spray drift minimisation strategy should be employed at all times when applying this product.

DO NOT apply Success® NEO using Ultra Low Volume (ULV) methods.

DILUTE SPRAYING

- Use a sprayer designed to apply high volumes of water up to the point of run-off and match to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of first run-off. Avoid excessive run-off.

- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the DIRECTIONS FOR USE table for each 100 L of water. Spray to the point of runoff. If volume to be applied is <1,000 L/ha then use the low volume (concentrate) application method for calculation of chemical rate. For volumes >1,000 L/ha use dilute spray rate.

CONCENTRATE SPRAYING

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (see **DILUTE SPRAYING** above) for the crop canopy. Consult your local advisor, agronomist or Department of Agriculture to determine this volume. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:

Concentrate Spraying Example

1. Dilute spray volume as determined above: e.g. 1,000 L/ha
2. Your chosen concentrate spray volume: e.g. 500 L/ha
3. The concentration factor is 2 X (1,000 / 500)
4. If the dilute label rate is 40 mL/100 L, then the concentrate rate becomes 2 X 40, i.e. 80 mL/100 L of concentrate spray

The chosen spray volume, amount of product per 100 L of water and the sprayer set up and operation may need to be changed as the crop grows. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training. Always follow Industry Best Practices.

RAINFASTNESS

Rain can wash Success® Neo Insecticide from treated plant surfaces and result in reduced efficacy. Avoid making spray applications if rain is expected before the spray can dry completely.

CLEANING SPRAY EQUIPMENT

After using Success® Neo Insecticide empty the tank and completely drain the system. Rinse the tank, pumps, lines, hoses, filters and nozzles by circulating clean water through the system. Drain and repeat the rinsing procedure twice.

RE-ENTRY

DO NOT allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF HONEY-BEES AND OTHER INSECT POLLINATORS
Highly toxic to bees; will kill bees foraging in the crop to be treated or in hives which are over-sprayed or reached by spray drift. **DO NOT** spray while bees are actively foraging. Residues may remain toxic to bees for three days after application.
Beekeepers who are known to have hives in, or nearby, the area to be sprayed should be notified no less than 48 hours prior to the time of the planned application so that bees can be removed or otherwise protected prior to spraying.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT
Very toxic to aquatic life. **DO NOT** contaminate wetlands or watercourses with this product or used containers.

PROTECTION OF OTHER NON-TARGET INSECTS
Risk to non-target insects. This product may have adverse effects on some non-target beneficials, such as foliage dwelling predators, particularly where IPM is practiced.

STORAGE AND DISPOSAL
Store in the closed, original container in a cool well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. **DO NOT** store near food, feedstuffs, fertilisers or seed.

This container can be recycled if it is clean, dry, free of visible residues and has the ***drum****MUSTER* logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate by adding to the spray tank. **DO NOT** dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any ***drum****MUSTER* collection or similar container management site. The cap should not be replaced but may be taken separately.
If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

SPILL AND LEAK MANAGEMENT
DO NOT touch or walk through spilled material. Wear a face shield or goggles, overalls buttoned to neck and wrist, chemical resistant gloves and footwear. Stop leak when safe to do so. Dam area and prevent entry into waterways, and drains.
Small spills/leaks: Absorb with material such as sand, soil or sawdust. Collect spilled product and place in sealable container for disposal. Spill residues may be cleaned using water and detergent. Contain and absorb wash water for disposal. Absorb and collect washings and place in the same sealable container for disposal. Dam the area of large spills and report them to Corteva Agriscience Emergency Services at 1-800 370 754.

APVMA Approval No.: 64109/147372

Corteva Agriscience Australia Proprietary Limited
A.B.N. 24 003 771 659
67 Albert Avenue, Chatswood, NSW 2067
www.corteva.com.au
CUSTOMER SERVICE TOLL FREE **1-800 700 096**

®,TM Trademarks of Corteva Agriscience and its affiliated companies. ® 2026 Corteva.

Barcode
for stock
identification