CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
READ THE ATTACHED BOOKLET BEFORE USE

Transform®

Isoclast® active

INSECTICIDE

ACTIVE CONSTITUENT: 240 g/L SULFOXAFLOR

GROUP 4C INSECTICIDE

For the control of aphids and other insect pests in canola, cereals, cotton, forage brassicas, herbs, lucerne, nursery stock, pulses, tree nuts, soybeans and various fruit and vegetable crops as specified in the Directions for Use.

SHAKE WELL BEFORE USE **NET CONTENTS:** 1 L, 5 L

FIRST AID
If poisoning occurs, contact a doctor or Poisons Information Centre. Phone: *Australia* 13 11 26.

EMERGENCY RESPONSE (ALL HOURS)
RING FROM ANYWHERE IN AUSTRALIA
1800 370 754
(LOCAL CALL FEE ONLY)

IN A TRANSPORT EMERGENCY ONLY
DIAL 000
FOR POLICE OR FIRE BRIGADE

SAFETY DIRECTIONS
- May irritate the eyes. Avoid contact with the eyes.
- When opening the container and preparing the product for use by ground-boom and aerial application, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow length chemical resistant gloves and face shield or goggles.
- If applying by spraying equipment carried on the back of the user, wear cotton overalls over normal clothing, buttoned to the neck and wrist, elbow length chemical resistant gloves and face shield or goggles.
- Wash hands after each use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

SAFETY DATA SHEET
Additional information is listed on the Safety Data Sheet for **TRANSFORM® ISOCLAST® ACTIVE INSECTICIDE** which is available from Corteva Agriscience on request. Call Customer Service Toll Free on 1-800 700 096 or visit www.corteva.com.au

CORTEVA™ agriscience

™ ® Trademarks of Corteva Agriscience and its affiliated companies.

Visit us at Corteva.com.au

SPRAY DRIFT RESTRAINTS:

Please note SPRAY DRIFT RESTRAINTS apply to all uses.

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between three and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

- spray droplets not smaller than a **MEDIUM** spray droplet size category
- minimum distances between the application site and downwind sensitive aquatic and wetland areas including aquacultural ponds, surface streams and rivers (see 'Mandatory No-Spray Zones' below) are observed.

Buffer zones for boom sprayers

Application rate	Boom height above the target canopy	Mandatory downwind buffer zones				
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
400 mL/ha or lower	0.5 m or lower	0 metres	0 metres	0 metres	0 metres	0 metres
	1.0 m or lower	0 metres	0 metres	10 metres	0 metres	0 metres
200 mL/ha or lower	1.0 m or lower	0 metres	0 metres	0 metres	0 metres	0 metres

DO NOT apply by a vertical sprayer unless the following requirements are met:

- spray is not directed above the target canopy
- the outside of the sprayer is turned off when turning at the end of rows and when spraying the outer row on each side of the application site
- minimum distances between the application site and downwind sensitive aquatic and wetland areas including aquacultural ponds, surface streams and rivers (see 'Mandatory No-Spray Zones' below) are observed.

Buffer zones for vertical sprayers

Application rate	Maximum dilute water rate	Type of target canopy	Mandatory downwind buffer zones				
			Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
40 mL/ 100 L or lower	1000 L/ha	2 metres tall and shorter	0 metres	0 metres	0 metres	0 metres	0 metres
	2000 L/ha	Taller than 2 metres (not fully-foliated)	0 metres	0 metres	10 metres	0 metres	0 metres
		Taller than 2 metres (fully-foliated)	0 metres	0 metres	5 metres	0 metres	0 metres

DO NOT apply by aircraft unless the following requirements are met:

- spray droplets not smaller than a **MEDIUM** spray droplet size category
- for maximum release height above the target canopy of 3 metres or 25 per cent of wingspan or 25 per cent of rotor diameter, whichever is the greatest, minimum distances between the application site and downwind sensitive aquatic and wetland areas including aquacultural ponds, surface streams and rivers (see ‘Mandatory No-Spray Zones’ below) are observed.

Buffer zones for aircraft

Application rate	Type of aircraft	Mandatory downwind buffer zones				
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
400 mL/ha or lower	Fixed-wing	0 metres	5 metres	25 metres	0 metres	0 metres
	Helicopter	0 metres	10 metres	30 metres	0 metres	0 metres

DIRECTIONS FOR USE

Broadacre, Berry, Nursery stock, Herbs, Vegetable and Fruit Crops (refer to individual Tables 1 to 6 below for specific directions).

TABLE 1 BROADACRE CROPS

Canola, Cereals, Cotton, Forage brassicas, Lucerne, Pulses and Soybeans

DIRECTIONS FOR USE:

RESTRAINTS (specific to Broadacre Crops):

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of field runoff for at least 3 days after application.

DO NOT apply more than two (2) times to canola, cereals, forage brassicas, lucerne and pulses or four (4) times to cotton and soybean in any one (1) season.

DO NOT make more than two (2) consecutive applications per crop for all pests other than aphids.

DO NOT apply consecutive applications of Group 4C insecticides for control of aphids.

DO NOT use rotary atomisers when applying aerially.

Note: Monitor crops for pest species by regular field scouting. Target sprays against insect populations when they exceed threshold levels. Make repeated applications at 14-21 day intervals as new infestations occur unless otherwise directed in the CRITICAL COMMENTS.

CAUTION: this product is highly toxic to bees. If bees are present in the target area during flowering, see the PROTECTION OF LIVESTOCK section in this booklet before use.

CROP	PEST	RATE (mL/ha)	CRITICAL COMMENTS
Canola	Aphids (including cabbage aphid, green peach aphid and turnip aphid)	100 - 200 + wetting agent ❷	**DO NOT** make more than two (2) applications per crop.
Cereals <u>Wheat, similar Grains, and pseudocereals without husks</u> Rye, triticale, wheat, chia, quinoa, psyllium, Amaranth grain, Cañihua, cram-cram, huauzontle	Aphids (including cereal aphid vectors of barley yellow dwarf virus; corn aphid and oat aphid), grain aphid and rose-grain aphid	50 - 100	**DO NOT** apply after dough development (GS80). **DO NOT** make more than two (2) applications per crop. Use higher rate under heavy aphid infestations and/or when water volume is reduced, such as with aerial application❶. Some species of aphids tend to infest cereal plants at the base of the plant, often inside the leaf sheath and below the soil surface. These entrenched aphids at the base of the plant may not be adequately controlled by Transform®.

Table 1 continued

CROP	PEST	RATE (mL/ha)	CRITICAL COMMENTS
Barley similar grains and pseudocereals with husks (Except oats) Barley, buckwheat, buckwheat (tartary), canarygrass (annual) Sorghum grain and millet Sorghum grain, millet, Hungry rice, Job's tears, Teff or tef	Russian wheat aphid	100	**DO NOT** apply after dough development (GS80). **DO NOT** make more than two (2) applications per crop. Begin applications when action thresholds are reached. Monitor crops and make further insecticide applications as necessary. Entrenched aphids make not be controlled. For best results, use Transform® early in the aphid infestation, early in crops development. Use as part of an IPM program where beneficials are preserved. Thorough coverage is essential. Ensure water volume and spray quality is adequate to thoroughly cover all of the crop canopy.
	Aphids (including cereal aphid vectors of barley yellow dwarf virus; corn aphid and oat aphid), grain aphid and rose-grain aphid	50 - 100	**DO NOT** apply after flag leaf stage (GS39). **DO NOT** make more than two (2) applications per crop. Use higher rate under heavy aphid infestations and/or when water volume is reduced, such as with aerial application❶. Some species of aphids tend to infest cereal plants at the base of the plant, often inside the leaf sheath and below the soil surface. These entrenched aphids at the base of the plant may not be adequately controlled by Transform®.
Cereals Oats	Russian wheat aphid	100	**DO NOT** apply after flag leaf stage (GS39). **DO NOT** make more than two (2) applications per crop. Begin applications when action thresholds are reached. Monitor crops and make further insecticide applications as necessary. Entrenched aphids make not be controlled. For best results; use Transform® early in the aphid infestation, early in crops development. Use as part of an IPM program where beneficials are preserved. Thorough coverage is essential. Ensure water volume and spray quality is adequate to thoroughly cover all of the crop canopy.

Table 1 continued

CROP	PEST	RATE (mL/ha)	CRITICAL COMMENTS
Cotton	Aphids (including cotton aphid, cowpea aphid and green peach aphid)	200 - 300	Use higher rate under heavy aphid infestations and/or when water volume is reduced, such as with aerial application❶.
	Green mirid		Use the lower rate when infestation is predominantly nymphs. Use higher rate when control of adults and/or residual control is desired.
	Greenhouse whitefly	400	Ensure accurate species identification. Target early stage crawlers and nymphs. Apply when the pest reaches threshold levels. Good coverage is essential.
	Rutherglen bug	300 - 400 (Suppression only)	Use the lower rate when infestation is predominately nymphs. Use higher rate when control of adults and/or residual control is desired. Due to the highly mobile nature of Rutherglen bugs and their tendency to rapidly reinvade treated areas, Transform® should be expected to give residual control of 3-7 days.
	Solenopsis mealybug (Cotton mealybug)	400 (Suppression only)	Conservation of beneficials is essential for the control of this pest. Apply as part of a season long spray programme targeting pests early in the crop. The use of Transform® will conserve beneficials when used as part of an IPM system. Thorough spray coverage is essential. Addition of an adjuvant, although not critical, may improve control.
Forage brassicas	Aphids (including cabbage aphid and green peach aphid)	100 + wetting agent ❷	**DO NOT** make more than two (2) applications per crop. Apply at any stage of crop when pest builds to a damaging threshold.

Table 1 continued

CROP	PEST	RATE (mL/ha)	CRITICAL COMMENTS
Lucerne	Mirids (including brown mirid, green mirid, and yellow mirid (apple dimpling bug))	200 - 300	Use the lower rate when infestation is predominantly nymphs. Use higher rate when control of adults and/or residual control is desired.
	Aphids (including blue-green aphid, pea aphid and spotted alfalfa aphid)	100	Apply when the pest reaches threshold levels.
Pulses (adzuki beans, mung beans and navy beans only)	Aphids (including blue green aphid, cowpea aphid and green peach aphid)	100	**DO NOT** make more than two (2) applications per crop.
Soybeans	Soybean aphid	100 - 200	Use higher rate when canopy closure may adversely affect application coverage.
	Green mirid	200 - 300	Use the lower rate when infestation is predominantly nymphs. Use higher rate when control of adults and/or residual control is desired.
	Greenhouse whitefly	400	Ensure accurate species identification. Target early stage crawlers and nymphs. Apply when the pest reaches threshold levels. Good coverage is essential.
❶Apply by air using a minimum water volume of 30 L/ha			
❷Addition of a wetting agent may improve control under less than ideal application conditions. Use the wetter according to its label directions. See **WETTING AGENTS** section below for recommended products.			

TABLE 2 VEGETABLE CROPS

Cucurbits, Fruiting vegetables, Leafy vegetables, Root and tuber vegetables and Vegetable brassicas.

DIRECTIONS FOR USE:

RESTRAINTS (specific to Vegetable Crops):

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of field runoff for at least 3 days after application.

DO NOT apply more than four (4) times to any of these crops in any one (1) season, except where otherwise indicated.

DO NOT make more than two (2) consecutive applications per crop for all pests other than aphids.

DO NOT apply consecutive applications of Group 4C insecticides for control of aphids.

DO NOT use rotary atomisers when applying aerially.

The use of Transform® in protected cropping is permissible when used in conjunction with a proven Insect Resistance Management Strategy (IRMS).
DO NOT apply to seedlings of edible crops for transplanting i.e. seedling intended for food production.
DO NOT apply to vegetable or herb seedlings in a production nursery system.
Note: Monitor crops for pest species by regular field scouting. Target sprays against insect populations when they exceed threshold levels. Make repeated applications at 7 – 10 day intervals as new infestations occur unless otherwise directed in the CRITICAL COMMENTS.

CAUTION: this product is highly toxic to bees. If bees are present in the target area during flowering, see the PROTECTION OF LIVESTOCK section in this booklet before use.

CROP	PEST	RATE (mL/ha)	CRITICAL COMMENTS
Cucurbits, including but not limited to cucumbers, melons, pumpkin, and squash	Green peach aphid and Melon (cotton) aphid	200 - 300	Use higher rate under heavy aphid infestations or if longer residual control (> 7 days) is required.
	Greenhouse whitefly	400 (field) (40 mL/100 L (protected cropping))	Ensure accurate species identification. Target early stage crawlers and nymphs. Apply when the pest reaches threshold levels. Good coverage is essential.
	Rutherglen bug	300 - 400 (Suppression only)	Use the lower rate when infestation is predominately nymphs. Use higher rate when control of adults and/or residual control is desired. Due to the highly mobile nature of Rutherglen bugs and their tendency to rapidly reinvade treated areas, Transform® should be expected to give residual control of 3-7 days.

Table 2 continued

CROP	PEST	RATE (mL/ha)	CRITICAL COMMENTS
Fruiting vegetables, including but not limited to capsicum, chilli, eggplant, okra and tomatoes [excluding sweet corn and mushrooms]	Green peach aphid	200 – 300	Use higher rate under heavy aphid infestations or if longer residual control (>7 days) is required.
	Greenhouse whitefly	400 (field) (40 mL/100 L (protected cropping))	Ensure accurate species identification. Target early stage crawlers and nymphs. Apply when the pest reaches threshold levels. Good coverage is essential.
	Rutherglen bug	300 - 400 (Suppression only)	Use the lower rate when infestation is predominately nymphs. Use higher rate when control of adults and/or residual control is desired. Due to the highly mobile nature of Rutherglen bugs and their tendency to rapidly reinvade treated areas, Transform® should be expected to give residual control of 3-7 days.
	Tomato-potato psyllid	300 - 400 (Suppression only)	Apply as part of a season long spray programme targeting pests when active in the crop. Apply immediately psyllids are detected. Monitor crops and make further insecticide applications as necessary. The use of Transform® early in the fruiting/flowering stage of the crop will conserve beneficials when used as part of an IPM system. The use of Transform® for control of TPP may still result in the transmission of various plant diseases. Thorough coverage is essential.
Fruiting vegetables, sweet corn	Green peach aphid	200	**DO NOT** make more than two (2) applications per crop. If a second application is required, apply after a 14 day interval.
Leafy vegetables, including but not limited to Asian greens, lettuce (all varieties), silver beet and spinach	Brown sowthistle aphid and green peach aphid	200–300	Use higher rate under heavy aphid infestations or if longer residual control (>7 days) is required.
	Greenhouse whitefly	400	Ensure accurate species identification.
	Rutherglen bug	150 - 200 (Suppression only)	Use the lower rate when infestation is predominately nymphs. Use higher rate when control of adults and/or residual control is desired. Due to the highly mobile nature of Rutherglen bugs and their tendency to rapidly reinvade treated areas, Transform® should be expected to give residual control of 3-7 days.

Table 2 continued

CROP	PEST	RATE (mL/ha)	CRITICAL COMMENTS
Root and tuber vegetables, including but not limited to carrots, potatoes and turnips	Green peach aphid	200–300	Use higher rate under heavy aphid infestations or if longer residual control (>7 days) is required.
	Tomato-potato psyllid	300 - 400 (Suppression only)	Apply as part of a season long spray programme targeting pests when active in the crop. Apply immediately psyllids are detected. Monitor crops and make further insecticide applications as necessary. The use of Transform® early in the fruiting/flowering stage of the crop will conserve beneficials when used as part of an IPM system. The use of Transform® for control of TPP may still result in the transmission of various plant diseases. Thorough coverage is essential.
	Rutherglen bug	300 - 400 (Suppression only)	Use the lower rate when infestation is predominately nymphs. Use higher rate when control of adults and/or residual control is desired. Due to the highly mobile nature of Rutherglen bugs and their tendency to rapidly reinvade treated areas, Transform® should be expected to give residual control of 3-7 days.
Vegetable brassicas, including but not limited to Asian greens, broccoli, Brussels sprouts, cabbage and cauliflower	Aphids, including cabbage aphid, green peach aphid and turnip aphid	200 – 300 (+ wetting agent)❶	Use higher rate under heavy aphid infestations or if longer residual control (>7 days) is required.
	Greenhouse whitefly	400 (+ wetting agent)❶	Ensure accurate species identification. Target early stage crawlers and nymphs. Apply when the pest reaches threshold levels. Good coverage is essential.
	Rutherglen bug	300 - 400 (Suppression only)	Use the lower rate when infestation is predominately nymphs. Use higher rate when control of adults and/or residual control is desired. Due to the highly mobile nature of Rutherglen bugs and their tendency to rapidly reinvade treated areas, Transform® should be expected to give residual control of 3-7 days.
❶ Addition of a wetting agent may improve control under less than ideal application conditions. Use the wetter according to its label directions. See **WETTING AGENTS** section below for recommended products.			

TABLE 3 BERRY CROPS

Strawberries and Cane berries

DIRECTIONS FOR USE:
RESTRAINTS (specific to Berry Crop):

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of field runoff for at least 3 days after application.

DO NOT apply more than four (4) times to strawberries and two (2) times to caneberries in any one (1) season.

DO NOT make more than two (2) consecutive applications per crop for all pests other than aphids.

DO NOT apply consecutive applications of Group 4C insecticides for control of aphids.

DO NOT use rotary atomisers when applying aerially.

Carefully monitor crops for pest species by regular field scouting. Repeat applications at a 7 day interval if a new infestation occurs unless otherwise directed in the **CRITICAL COMMENTS.**

CAUTION: this product is highly toxic to bees. If honeybees are present in the target area during flowering, see the PROTECTION OF LIVESTOCK section in this booklet before use.

CROP	PEST	RATE (mL/ha)	CRITICAL COMMENTS
Cane berries (including but not limited to blackberries and raspberries)	Green peach aphid	300	Apply as a foliar cover spray following first signs of pest infestation.
	Greenhouse whitefly	400	Ensure accurate species identification. Target early stage crawlers and nymphs. Apply when the pest reaches threshold levels. Good coverage is essential.
	Mirids (brown mirid and green mirid)	300	Apply when the pest reaches threshold levels.
	Yellow mirid (apple dimpling bug)	300 – 400	Use the lower rate when infestation is predominately nymphs. Use higher rate when control of adults and/or residual control is desired.
	Rutherglen bug	300 – 400 (suppression only)	Use the lower rate when infestation is predominately nymphs. Use higher rate when control of adults and/or residual control is desired. Due to the highly mobile nature of Rutherglen bugs and their tendency to rapidly reinvade treated areas, Transform® should be expected to give residual control of 3-7 days.
	Scale insects (*Coccidae* spp. and *Diaspididae* spp.)	400 (suppression only)	Apply as part of a spray programme targeting crawlers when they are exposed. Make two (2) applications 14 - 21 days apart when crawlers are active. Complete spray coverage is essential. Addition of an adjuvant may improve control.

Table 3 continued

CROP	PEST	RATE (mL/ha)	CRITICAL COMMENTS
Strawberries	Green peach aphid	200 - 300	Use higher rate under heavy aphid infestations and/or when water volume is reduced.
	Green mirid		Use the lower rate when infestation is predominately nymphs. Use higher rate when control of adults and/or residual control is desired.
	Rutherglen bug	300 – 400 (suppression only)	Use the lower rate when infestation is predominately nymphs. Use higher rate when control of adults and/or residual control is desired. Due to the highly mobile nature of Rutherglen bugs and their tendency to rapidly reinvade treated areas, Transform® should be expected to give residual control of 3-7 days.

TABLE 4 TREE and VINE CROPS

Assorted tropical and sub-tropical fruits (inedible peel) (except banana), Avocado, Carob, Citrus, Grapes, Pome fruit, Macadamia nut, Tree nuts and Stone fruit.

DIRECTIONS FOR USE:

RESTRAINTS (specific to Tree and Vine Crops*)*:

DO NOT apply with aircraft.

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of field runoff for at least 3 days after application.

DO NOT apply more than twice per crop per season for all situations except for use in avocado and aphid control on stone fruit.

DO NOT apply more than four times per season for use in avocado and aphid control on stone fruit.

DO NOT apply consecutive applications of Group 4C insecticides for control of aphids excluding woolly apple aphid.

Carefully monitor crops for pest species by regular field scouting. Repeat applications at a 14 day interval if a new infestation occurs unless otherwise directed in the CRITICAL COMMENTS**.**

CAUTION: this product is highly toxic to bees. If honeybees are present in the target area during flowering, see the PROTECTION OF LIVESTOCK section in this booklet before use.

SPRAYING TREE and VINE CROPS: In the following table, all rates are given for dilute spraying where spray volumes may vary in order to obtain good coverage to the point of run-off. For concentrate spraying refer to the "**CONCENTRATE SPRAYING**" section of this label.			
CROP	**PEST**	**RATE (mL/100L)**	**CRITICAL COMMENTS**
Assorted tropical and sub-tropical fruits (inedible peel) (including but not limited to carob, custard apple, lychee, papaya, passionfruit and mango) (except bananas and pineapple (see separate claim))	banana-spotting bug and fruit spotting bug	40	**DO NOT** make more than two (2) applications to any one (1) crop in any one (1) season. If making repeat applications, **DO NOT** reapply before 14 days after the first application. Apply as part of a season long spray programme targeting pests when active in the crop. The use of Transform® early in the fruiting/flowering stage of the crop will conserve beneficials when used as part of an IPM system. Complete spray coverage is essential - concentrate sprays are not suitable for this pest. Addition of an adjuvant, although not critical, may improve control. Apply to the point of run-off.

Table 4 continued

CROP	PEST	RATE (mL/100L)	CRITICAL COMMENTS
Avocado	Banana-spotting bug and fruit-spotting bug	40	**DO NOT** make more than four (4) applications to any one (1) crop in any one (1) season. If making repeat applications, **DO NOT** reapply before 7 days after each application. Apply as part of a season long spray programme targeting pests when active in the crop. The use of Transform® early in the fruiting/flowering stage of the crop will conserve beneficials when used as part of an IPM system. Complete spray coverage is essential - concentrate sprays are not suitable for this pest. Addition of an adjuvant, although not critical, may improve control. Apply to the point of run-off.
Carob	Citrus mealybug	40	Apply as part of spray programme targeting crawlers when they are exposed and before they are protected under the fruit calyces or established between touching fruit. Mealybugs in protected feeding sites or protected by dense foliage may not be controlled. Make two (2) applications 14 - 21 days apart when crawlers are active. Addition of an adjuvant may improve control. Apply to the point of run-off.

Table 4 continued

CROP	PEST	RATE (mL/100L)	CRITICAL COMMENTS
Citrus, including but not limited to grapefruit, lemons, limes, mandarins, oranges and tangerines	Citrophilous mealybug Citrus mealybug and Longtailed mealybug	40	Apply as part of spray programme targeting crawlers when they are exposed and before they are protected under the fruit calyces or established between touching fruit. Mealybugs in protected feeding sites or protected by dense foliage may not be controlled. Make two (2) applications 14 - 21 days apart when crawlers are active. **DO NOT** apply more than 800 mL of product per hectare in a single application. Addition of an adjuvant may improve control.
	Citricola scale, citrus snow (white louse) scale, pink wax scale and red scale		Apply as part of a spray programme targeting crawlers when they are exposed. Make two (2) applications 14 - 21 days apart when crawlers are active. Complete spray coverage is essential - concentrate sprays are not suitable for this pest. Addition of an adjuvant may improve control.
	Kelly's citrus thrips		Apply as part of a spray programme. Monitor crops from flowering onwards and commence applications once local pest thresholds are reached. Continue to monitor crops and make further applications as required. Addition of an adjuvant may improve control. Thrips entrenched under fruit calyces will not be controlled.
	Banana-spotting bug and fruit spotting bug		**DO NOT** make more than two (2) applications to any one (1) crop in any one (1) season. Apply as part of a season long spray programme targeting pests when active in the crop. The use of Transform® early in the fruiting/flowering stage of the crop will conserve beneficials when used as part of an IPM system. **DO NOT** apply more than 800 mL of product per hectare in a single application. Complete spray coverage is essential - concentrate sprays are not suitable for this pest. Addition of an adjuvant, although not critical, may improve control.

Table 4 continued

CROP	PEST	RATE (mL/100L)	CRITICAL COMMENTS
Custard apple (see also Assorted tropical and sub-tropical fruits (inedible peel))	Mealybugs (including *pseudococcidae* spp. and Citrus mealybug)	40 (suppression only)	Apply as part of spray programme targeting crawlers when they are exposed and before they are protected under the fruit calyces or established between touching fruit. Mealybugs in protected feeding sites or protected by dense foliage may not be controlled. Make two (2) applications 14 - 21 days apart when crawlers are active. Addition of an adjuvant may improve control. Apply to the point of run-off.
Grapes (table grapes)	Longtailed mealybug	40	Apply as part of a spray programme targeting crawlers when they are exposed and before they become entrenched. Best control will be achieved by making two (2) applications 14 - 21 days apart when crawlers are active early in the season when good coverage can be achieved. Use 40 mL/100 L in up to 2000 L/ha of water. **DO NOT apply later than 80% capfall.** Complete spray coverage is essential – concentrate sprays are not suitable for this pest. Addition of an adjuvant may improve control. Apply to the point of run-off. See GENERAL INSTRUCTIONS.
Grapes (wine grapes)	Longtailed mealybug	30	Apply as part of a spray programme targeting crawlers when they are exposed and before they become entrenched. Best control will be achieved by making two (2) applications 14 - 21 days apart when crawlers are active early in the season when good coverage can be achieved. Use 30 mL/100 L rate in up to 2000 L of water. **DO NOT apply later than 80% capfall.** Complete spray coverage is essential – concentrate sprays are not suitable for this pest. Addition of an adjuvant may improve control. Apply to the point of run-off.

Table 4 continued

CROP	PEST	RATE (mL/100L)	CRITICAL COMMENTS
Lychee (see also Assorted tropical and sub-tropical fruits (inedible peel))	Citrus snow (white louse) scale and soft green scale	40 (suppression only)	Apply as part of a spray programme targeting crawlers when they are exposed. Make two (2) applications 14 - 21 days apart when crawlers are active. Complete spray coverage is essential. Addition of an adjuvant may improve control. Apply to the point of run-off.
	Mealybugs (*planococcus* spp.)		Apply as part of spray programme targeting crawlers when they are exposed and before they are protected under the fruit calyces or established between touching fruit. Mealybugs in protected feeding sites or protected by dense foliage may not be controlled. Make two (2) applications 14 - 21 days apart when crawlers are active. Apply to the point of run-off.
Macadamia	Banana-spotting bug and fruit-spotting bug	40	**DO NOT** make more than two (2) applications to any one (1) crop in any one (1) season. If making repeat applications, **DO NOT** reapply before 7 days after the first application. Apply as part of a season long spray programme targeting pests when active in the crop. The use of Transform® early in the fruiting/flowering stage of the crop will conserve beneficials when used as part of an IPM system. Complete spray coverage is essential - concentrate sprays are not suitable for this pest. Addition of an adjuvant, although not critical, may improve control. Apply to the point of run-off.
	Lace bugs		**DO NOT** make more than two (2) applications to any one (1) crop in any one (1) season. Apply immediately when the pest is detected. Make a second application as needed. The use of Transform® early in the fruiting/flowering stage of the crop will conserve beneficials when used as part of an IPM system. Complete spray coverage is essential. Addition of an adjuvant, although not critical, may improve control. Apply to the point of run-off.

Table 4 continued

CROP	PEST	RATE (mL/100L)	CRITICAL COMMENTS
Macadamia *continued*	Macadamia felted coccid		**DO NOT** make more than two (2) applications to any one (1) crop in any one (1) season. Apply immediately when the pest is detected. Make a second application as needed. The use of Transform® early in the fruiting/flowering stage of the crop will conserve beneficials when used as part of an IPM system. Complete spray coverage is essential. Addition of an adjuvant, although not critical, may improve control. Apply to the point of run-off.
Mango (see also Assorted tropical and sub-tropical fruits (inedible peel))	Apple dimpling bug (yellow mirid)	30	Apply when the pest reaches threshold levels in the lead up to flowering. As Apple dimpling bugs are a highly mobile pest and can rapidly reinvade crops, further sprays of Transform® (or another product) 14 days after application may be necessary. Apply to the point of run-off.
	Mango shield scale	40	Apply as part of a spray programme targeting crawlers when they are exposed. Make two (2) applications 14 - 21 days apart when crawlers are active. Complete spray coverage is essential. Addition of an adjuvant may improve control. Apply to the point of run-off.
	Pink wax scale		
	White mango scale		
Papaya (see also Assorted tropical and sub-tropical fruits (inedible peel))	Scale (*Coccidae* spp.)	40 (suppression only)	Apply as part of a spray programme targeting crawlers when they are exposed. Make two (2) applications 14 - 21 days apart when crawlers are active. Complete spray coverage is essential. Addition of an adjuvant may improve control. Apply to the point of run-off.
	Mealybugs (*Planococcus* spp.)		Apply as part of spray programme targeting crawlers when they are exposed and before they are protected under the fruit calyces or established between touching fruit. Mealybugs in protected feeding sites or protected by dense foliage may not be controlled. Make two (2) applications 14 - 21 days apart when crawlers are active. Addition of an adjuvant may improve control. Apply to the point of run-off.

Table 4 continued

CROP	PEST	RATE (mL/100L)	CRITICAL COMMENTS
Passionfruit (see also Assorted tropical and sub-tropical fruits (inedible peel))	Black scale	40 (suppression only)	Apply as part of a spray programme targeting crawlers when they are exposed. Make two (2) applications 14 - 21 days apart when crawlers are active. Complete spray coverage is essential. Addition of an adjuvant may improve control. Apply to the point of run-off.
	Citrus snow (white louse) scale		
	Red scale		
	Citrus mealybug		Apply as part of spray programme targeting crawlers when they are exposed and before they are protected under the fruit calyces or established between touching fruit. Mealybugs in protected feeding sites or protected by dense foliage may not be controlled. Make two (2) applications 14 - 21 days apart when crawlers are active. Addition of an adjuvant may improve control. Apply to the point of run-off.
	Longtailed mealybug		
	Passionvine mealybug		

Table 4 continued

CROP	PEST	RATE (mL/100L)	CRITICAL COMMENTS
Persimmon (see also Pome fruit)	Mealybugs (citrus mealybug, citrophilous mealybug and longtail mealybug)	40 (suppression only)	Apply as part of spray programme targeting crawlers when they are exposed and before they are protected under the fruit calyces or established between touching fruit. Mealybugs in protected feeding sites or protected by dense foliage may not be controlled. Make two (2) applications 14 - 21 days apart when crawlers are active. Addition of an adjuvant may improve control. **DO NOT** apply more than 800 mL of product per hectare in a single application.
	Banana-spotting bug and fruit spotting bug	40	**DO NOT** make more than four (2) applications to any one (1) crop in any one (1) season. If making repeat applications, **DO NOT** reapply before 7 days after each application. Apply as part of a season long spray programme targeting pests when active in the crop. The use of Transform® early in the fruiting/flowering stage of the crop will conserve beneficials when used as part of an IPM system. Complete spray coverage is essential - concentrate sprays are not suitable for this pest. Addition of an adjuvant, although not critical, may improve control. **DO NOT** apply more than 800 mL of product per hectare in a single application.
	Scale insects (*Coccidae* spp., *Diaspididae* spp., *Eriococcidae* spp.)	40 (suppression only)	Apply as part of a spray programme targeting crawlers when they are exposed. Make two (2) applications 14 – 21 days apart when crawlers are active. Complete spray coverage is essential. Addition of an adjuvant may improve control. **DO NOT** apply more than 800 mL of product per hectare in a single application.
Pineapple (see also Assorted tropical and sub-tropical fruits (inedible peel))	Pineapple mealybug		Apply as part of spray programme targeting crawlers when they are exposed and before they are protected under the fruit calyces or established between touching fruit. Mealybugs in protected feeding sites or protected by dense foliage may not be controlled. Make two (2) applications 14 - 21 days apart when crawlers are active. Ensure good coverage. **DO NOT** apply in a spray volume of more than 1000 L/ha. Addition of an adjuvant is advised.

Table 4 continued

CROP	PEST	RATE (mL/100L)	CRITICAL COMMENTS
Pome fruit, including but not limited to apples, pears and nashi	Apple dimpling bug	30	Apply when the pest reaches threshold levels in the lead up to flowering. As Apple dimpling bugs are a highly mobile pest and can rapidly reinvade crops, further sprays of Transform® (or another product) 14 days after application may be necessary. **DO NOT** apply more than 800 mL of product per hectare in a single application.
	Longtailed mealybug and tuber mealybug	40	Apply in spring as part of a spray programme targeting crawlers when they are exposed and before they become entrenched. Monitor for crawler emergence in spring and make two (2) applications of Transform® 14 days apart (or as indicated by monitoring). Good coverage is essential – concentrate sprays are not suitable for this pest. Addition of an adjuvant may improve control. **DO NOT** apply more than 800 mL of product per hectare in a single application. Monitoring for crawler emergence should continue and, if required, a product from a different mode of action group should be applied (e.g. Applaud®).
	Woolly (apple) aphid		Apply as part of a spray programme. Monitor for aphid activity and apply Transform® before colonies become clumped or heavily covered in wool. Consecutive applications of Transform® may be made if required. Repeat applications of Transform® 14-21 days apart will be required if targeting heavy, mature colonies. Continue monitoring throughout the season and apply further control measures as required. Addition of an adjuvant may improve wetting of plant surfaces and wool. Good coverage is essential - concentrate or low volume sprays are not suitable for control of this pest. **DO NOT** apply more than 800 mL of product per hectare in a single application.

Table 4 continued

CROP	PEST	RATE (mL/100L)	CRITICAL COMMENTS
Pome fruit, including but not limited to apples, pears and nashi *continued*	San Jose scale	40	**DO NOT** make more than two (2) applications per crop. Apply as part of a spray programme targeting crawlers when they are exposed. Make two (2) applications 14 - 21 days apart when crawlers are active. Crop scouting 7 - 10 days after the first application will help determine when the second application should be applied so as to target new crawler emergence. Complete spray coverage is essential - concentrate sprays arenot suitable for this pest. Addition of an adjuvant may improve control. **DO NOT** apply more than 800 mL of product per hectare in a single application.
Stone fruit, including but not limited to apricots, cherries, nectarines, peaches and plums	Apple dimpling bug	30	Apply when the pest reaches threshold levels in the lead up to flowering. As Apple dimpling bugs are a highly mobile pest and can rapidly reinvade crops, further sprays of Transform® (or another product) 14 days after application may be necessary. **DO NOT** apply more than 800 mL of product per hectare in a single application.
	Black peach aphid and cherry aphid	10	Apply when the pest reaches threshold levels. Good coverage is essential. Aphids that are protected inside curled leaves may not be controlled. **DO NOT** exceed 400 mL of product per hectare in a single application. If using more than 4000 L/ha water, dilute accordingly.
	Green peach aphid	10 – 20	Apply when the pest reaches threshold levels. Good coverage is essential. Aphids that are protected inside curled leaves may not be controlled. Use the higher rate under higher pest pressure. **DO NOT** apply more than 800 mL of product per hectare in a single application.
Tree nuts, including but not limited to almond, cashew, chestnut, filbert (hazelnut), pecan, walnuts and pistachios (except macadamia)	Aphids (including black peach aphid and green peach aphid)	10	Apply when the pest reaches threshold levels. Good coverage is essential. Aphids that are protected inside curled leaves may not be controlled. **DO NOT** apply more than 800 mL of product per hectare in a single application.

TABLE 5 NURSERY STOCK (NON-FOOD)

DIRECTIONS FOR USE:

RESTRAINTS (specific to Nursery Stock (Non-Food))

The use of Transform® in protected cropping is permissible when used in conjunction with a proven Insect Resistance Management Strategy (IRMS).

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of field runoff for at least 3 days after application.

DO NOT apply to seedling intended for food production.
DO NOT apply to vegetable or herb seedlings in a production nursery system.
DO NOT make more than two (2) applications per crop.
DO NOT make repeat applications within 7 days.
DO NOT apply if bees are foraging in the crop.

CAUTION: this product is highly toxic to bees. If honeybees are present in the target area during flowering, see the PROTECTION OF LIVESTOCK section in this booklet before use.

CROPS	PEST	RATE	CRITICAL COMMENTS
Nursery stock (non-food), including seedlings, tubes & plugs, potted colour, trees and shrubs, foliage plants, palms and grasses and fruiting plants (non-bearing); CUT FLOWERS, FOLIAGE and ORNAMENTALS	Aphid (*Aphis* spp.) Mirids (*Creontiades* spp.)	200 - 300 mL/ha OR 20 - 30 mL/100 L	Apply when the pest reaches threshold levels. Good coverage is essential. Aphids that are protected in sheltered feeding sites may not be controlled. **DO NOT** apply consecutive applications when aphids are present.
	Mealybugs (*Pseudococcus* spp.)	300 - 400 mL/ha OR 30 - 40 mL/100 L	Apply as part of a spray programme targeting crawlers when they are exposed. Make two (2) applications 14 - 21 days apart when crawlers are active. Complete spray coverage is essential - concentrate sprays are not suitable for this pest. Addition of an adjuvant may improve control.
	Scale (*Diaspididae* spp.)	400 mL/ ha OR 40 mL/100 L	Apply as part of a spray programme targeting crawlers when they are exposed. Make two (2) applications 14 - 21 days apart when crawlers are active. Complete spray coverage is essential - concentrate sprays are not suitable for this pest. Addition of an adjuvant may improve control.
	Greenhouse whitefly	400 mL/ha OR 40 mL/100 L	Ensure accurate species identification. Apply as part of a spray programme targeting crawlers when they are exposed. Complete spray coverage is essential. Addition of an adjuvant may improve control.
	Psyllids (including tomato potato psyllid) and stink bugs (including brown marmorated stink bug	400 mL/ha OR 40 mL/ 100L (suppression only)	Monitor crops and commence application when pests are first detected. Follow up treatment may be necessary to control new infestations 7 to 10 days later.

TABLE 6 HERBS

Herbs

DIRECTIONS FOR USE:
RESTRAINTS (specific to Herbs):
The use of Transform® in protected cropping is permissible when used in conjunction with a proven Insect Resistance Management Strategy (IRMS).

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of field runoff for at least 3 days after application.

DO NOT apply herb seedlings in a production nursery system.
DO NOT make more than two (2) applications per crop.

CAUTION: this product is highly toxic to bees. If honeybees are present in the target area during flowering, see the PROTECTION OF LIVESTOCK section in this booklet before use.

CROP	PEST	RATE (mL/100L)	CRITICAL COMMENTS
Herbs	Aphids (including cotton (melon) aphid and green peach aphid)	30	Apply when the pest reaches threshold levels. Good coverage is essential.
	Greenhouse whitefly	40	Ensure accurate species identification. Target early stage crawlers and nymphs. Apply when the pest reaches threshold levels. Good coverage is essential.
PEST NAMES: Apple dimpling bug (*Campylomma liebknechti*), Banana-spotting bug (*Amblypelta lutescens lutescens*), Black peach aphid (*Brachycaudus persica*), Black scale (*Saissetia oleae*), Bluegreen aphid (*Acyrthosiphon kondoi*), Brown marmorated stink bug *(Halyomorpha halys),* Brown sowthistle aphid (*Uroleucon sonchi)*, Cabbage aphid (*Brevicoryne brassicae*), Cereal aphids (*Rhopalosiphum* spp. – vectors of Barley Yellow Dwarf Virus), Cherry aphid (*Myzus cerasi*), Citricola scale (*Coccus pseudomagnoliarum*), Citrophilous mealybug (*Pseudococcus calceolariae*), Citrophilus mealybug (*Pseudococcus calceolariae*), Citrus mealybug (*Planococcus citri*), Citrus snow (white louse) scale (*Unaspis citri*) Corn aphid (*Rhopalosiphum maidis*), Cotton aphid (*Aphis gossypii*), Cotton mealybug *(Phenacoccus solenopsis),* Cowpea aphid (*Aphis craccivora*), Fruit-spotting bug (*Amblypelta nitida*), Grain aphid (*Sitobion miscanthi*), Green mirid (*Creontiades dilutus*), Green peach aphid (*Myzus persicae*), Greenhouse whitefly (*Trialeurodes vaporariorum*), Kelly's citrus thrips (*Pezothrips kellyanus*), Lace bug (*Ulonemia concave),* Longtailed mealybug (*Pseudococcus longispinus*), Macadamia felted coccid *(Acanthacoccus ironsidei),* Mango shield scale (*Milviscutulus mangiferae*), Mealybugs (*Planococcus* spp. and *pseudococcidae* spp.), Melon aphid (*Aphis gossypii*), Oat aphid (*Rhopalosiphum padi*), Passionvine mealybug (*Planococcus minor*), Pineapple mealybug (*Dysmicoccus brevipes*), Pink wax scale (*Ceroplastes rubens*), Red scale (*Aonidiella aurantii*), Rose-grain aphid (Metopolophium dirhodum), Russian wheat aphid, (*Diuraphis noxia*), Rutherglen bug (*Nysius vinitor*), San Jose scale (*Quadraspidiotus perniciosus*), Scale insects (*Coccidae* spp., *Diaspididae* spp. and *Eriococcidae* spp.), Spotted alfalfa aphid (*Therioaphis trifolii*), Soft green scale (*cuccus viridis*), Solenopsis mealybug *(Phenacoccus solenopsis),* Soybean aphid (*Aphis glycines),* Tomato-potato psyllid (*Bactericera cockerelli),* Tuber mealybug (*Pseudococcus viburni*), Turnip aphid (*Lipaphis pseudobrassicae*), White mango scale (*Aulacaspis tubercularis*) and Woolly (apple) aphid (*Eriosoma lanigerum*).			

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

HARVEST WITHHOLDING PERIODS (WHP)

Macadamia:
NIL

Oats, table grapes and wine grapes:
NOT REQUIRED WHEN USED AS DIRECTED.

Cane berries, carob, citrus fruit, cucurbits, custard apple, fruiting vegetables (except sweet corn), herbs, pineapple and strawberries:
DO NOT HARVEST FOR 1 DAY AFTER THE LAST APPLICATION.

Assorted tropical and sub-tropical fruits (inedible peel) (except avocado, banana, custard apple and pineapple), brassica vegetables, leafy vegetables and persimmon:
DO NOT HARVEST FOR 3 DAYS AFTER THE LAST APPLICATION.

Avocado, pome fruit (except persimmon), root and tuber vegetables and stone fruit, sweet corn and Tree nuts:
DO NOT HARVEST FOR 7 DAYS AFTER THE LAST APPLICATION.

Canola, cereals (except oats), cotton, pulses and soybeans:
DO NOT HARVEST FOR 14 DAYS AFTER THE LAST APPLICATION.

GRAZING AND STOCKFOOD WITHHOLDING PERIODS (WHP):

Pineapple:
DO NOT GRAZE OR CUT FOR STOCKFEED FOR 1 DAY AFTER APPLICATION.

Forage brassicas:
DO NOT GRAZE OR CUT FOR STOCKEED FOR 7 DAYS AFTER APPLICATION.

Canola forage, post-harvest straw and stubble, cereals and pulses:
DO NOT GRAZE OR CUT FOR STOCKFEED FOR 14 DAYS AFTER APPLICATION.

Lucerne:
DO NOT GRAZE OF CUT FOR STOCKFEED FOR 21 DAYS AFTER APPLICATION.

Cotton:
DO NOT FEED COTTON TRASH TO ANIMALS

Soybeans:
DO NOT GRAZE OR CUT FOR STOCKFEED FOR 7 DAYS AFTER APPLICATION.

Sweet corn:
DO NOT GRAZE OR CUT FOR STOCKFEED FOR 7 DAYS AFTER APPLICATION.

LIVESTOCK DESTINED FOR EXPORT MARKETS

The GRAZING AND STOCKFOOD WITHHOLDING PERIODS (above) only apply to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that in addition to complying with the GRAZING AND STOCKFOOD WITHHOLDING PERIODS, that the EXPORT SLAUGHTER INTERVAL, is observed before stock are sold or slaughtered.

EXPORT SLAUGHTER INTERVAL (ESI) – 14 days:

After observing the grazing withholding period, livestock that has been grazed on or fed treated crops should be placed on clean feed for 14 days prior to slaughter.

CROPS FOR EXPORT - **Before** using Transform® on crops destined for export it is essential to consult your exporter or Corteva Agriscience to ensure that an appropriate MRL is in place in the importing country.

GENERAL INSTRUCTIONS
Transform® is a sulfoximine insecticide, in the form of a suspension concentrate.

INSECTICIDE RESISTANCE WARNING

GROUP	4C	INSECTICIDE

For insecticide resistance management, Transform® Isoclast® active Insecticide is a Group 4C insecticide. Some naturally occurring insect biotypes resistant to Transform® and other Group 4C insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Transform® and other Group 4C insecticides are used repeatedly. The effectiveness of Transform® on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Corteva Agriscience Australia Pty Ltd accepts no liability for any losses that may result from the failure of Transform® to control resistant insects. Transform® may be subject to specific resistance management strategies. For further information contact your local supplier, Corteva Agriscience representative or local agricultural department agronomist.

MIXING
- Agitate or shake the container immediately prior to use.
- Half fill the spray tank with water, add the appropriate amount of accurately measured Transform®, then complete filling the tank.
- Ensure thorough agitation by mechanical or hydraulic action at all times during mixing and application.
- Use only clean water within the range pH 5-9 to dilute Transform®.

COMPATIBILITY
If intending to tank mix Transform® with other agricultural chemicals or plant nutrients consult Corteva Agriscience.

WETTING AGENTS
Not all surfactants or crop oils are of equal quality. Corteva Agriscience Australia Pty Ltd does not support the use of alternative products other than those listed below.
Agral® Spray Adjuvant, Nufarm Chemwet 1000 and Spreadwet 1000 Wetting Agent. If intending to use other wetting agents consult Corteva Agriscience.
Agral® Trademark of a Syngenta Group Company

STORAGE OF DILUTED SPRAY MIX
Whenever possible the spray mix should be used immediately after it is prepared. However, if weather conditions or mechanical breakdown prevent immediate use, the spray mix may be stored for up to 72 hours without loss of activity. The spray mix should be agitated thoroughly by mechanical or hydraulic action at regular intervals during storage to prevent sedimentation. Ensure that the stored spray mix is thoroughly agitated at least once every 8 hours. The spray mix must be stored out of direct sunlight.

APPLICATION
Thorough coverage of the crop is essential. Ensure this by increasing water volume with plant growth stage. Do not apply when conditions are unsuitable for water-based spray applications. Avoid high temperature, strong winds, inversion conditions, imminent rain or any conditions that may reduce the quality of spray coverage or result in drift from the target area. Techniques to minimise drift should be employed at all times when aerially applying sprays to, or near, sensitive areas (see RESTRAINTS).

For optimum results follow the application specifications listed below:
Ground Spraying (Broadacre crops): Apply in a minimum of 50 L/ha of water with spray droplets no smaller than medium category according to nozzle manufacturer specifications that refer to the ASAE S-572 Standard.
Increase spray volumes as the crop grows.

Ground Spraying (Vegetable crops): Apply in a minimum of 250 L/ha of water with spray droplets no smaller than medium category according to nozzle manufacturer specifications that refer to the ASAE S-572 Standard. Increase spray volumes as the crop grows.

Aerial Spraying (Broadacre arable and vegetable crops only): Apply in a minimum of 30 L/ha of water with spray droplets no smaller than a medium category according to nozzle manufacturer specifications that refer to the ASAE S-572 Standard.

PRECAUTIONS:

RE-ENTRY
DO NOT allow entry into treated areas until the product has dried

(Aerial Application):
DO NOT use human flaggers/markers unless they are protected by engineering controls such as enclosed cabs.

DILUTE SPRAYING
- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of first run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the DIRECTIONS FOR USE table for each 100 L of water. Spray to the point of runoff. If volume to be applied is <1000L/ha then use the low volume (concentrate) application method for calculation of chemical rate. For volumes > 1000 L/ha use dilute spray rate.

CONCENTRATE SPRAYING
Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
Determine an appropriate dilute spray volume (see **DILUTE SPRAYING** above) for the crop canopy. Consult your local advisor, agronomist or Department of Primary Industries to determine this volume. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way:

Concentrate Spraying Example
1. Dilute spray volume as determined above: e.g. 1000 L/ha
2. Your chosen concentrate spray volume: e.g. 500 L/ha
3. The concentration factor is 2X (1000 / 500)

4. If the dilute label rate is 40 mL/100 L, then the concentrate rate becomes 2 X 40, i.e. 80 mL/100 L of concentrate spray

The chosen spray volume, amount of product per 100 L of water and the sprayer set up and operation may need to be changed as the crop grows. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training. Always follow Industry Best Practices.

RAINFASTNESS

Rain can wash Transform® from treated plant surfaces and result in reduced insect control. Avoid making spray applications if rain is expected before the spray can dry completely.

CLEANING SPRAY EQUIPMENT

After using Transform® empty the tank and completely drain the system. Rinse the tank, pumps, lines, hoses, filters and nozzles by circulating clean water through the system. Drain and repeat the rinsing procedure twice.

PROTECTION OF HONEYBEES AND OTHER INSECT POLLINATORS

Hazard:

Highly toxic to bees. Harmful to bee brood. Sulfoxaflor has systemic action. Will kill foraging bees directly exposed through contact during spraying and while spray droplets are still wet. May harm bees in hives which are over-sprayed or reached by spray drift. **DO NOT** apply this product while bees are foraging in the crop to be treated.

Risk Management:

Treatments made to crops in flower or upwind of adjacent plants in flower that are likely to be visited by bees at the time of application, should not occur during the daytime if temperatures within an hour after the completion of spraying are expected to exceed 12 °C. It is recommended that orchard floors containing flowering plants be mown just prior to spraying. Beekeepers who are known to have hives in, or nearby, the area to be sprayed should be notified no less than 48 hours prior to the time of the planned application so that bees can be removed or otherwise protected prior to spraying.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. **DO NOT** contaminate wetlands or water courses with this product or used containers.

INTEGRATED PEST MANAGEMENT

This product may have adverse effects on parasitic wasps particularly where IPM is practiced.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT store near food, feedstuffs, fertilisers or seed.

This container can be recycled if it is clean, dry, free of visible residues and has the ***drum****MUSTER* logo visible. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any ***drum****MUSTER* collection or similar container management site. The cap should not be replaced but may be taken separately.

If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SPILL AND LEAK MANAGEMENT
Do not touch or walk through spilled material. Wear a face shield or goggles, overalls buttoned to neck and wrist, chemical resistant gloves and footwear. Stop leak when safe to do so. Dam area and prevent entry into waterways, and drains.
Small spills/leaks: Contain and absorb small spills with a proprietary absorbent suitable for chemical spills or inert materials such as sand, soil or sawdust. Collect spilled product and place in sealable container for disposal. Spill residues may be cleaned using water and detergent. Contain and absorb wash water for disposal. Absorb and collect washings and place in the same sealable container for disposal. Dam the area of large spills and report them to Corteva Agriscience Emergency Services at 1-800 370 754.

APVMA Approval No. : 64101/131569

Corteva Agriscience Australia Pty Ltd
A.B.N. 24 003 771 659
67 Albert Avenue, Chatswood, NSW 2067
www.corteva.com.au
CUSTOMER SERVICE TOLL FREE **1-800 700 096**

™ ® Trademarks of Corteva Agriscience and its affiliated companies. © 2022 Corteva.

Barcode
for stock
identification

This product is GHS compliant. No additional GHS hazard and precautionary statements are required under the Safe Work Australia exemptions for AgVet products.