Australian Pesticides & Veterinary Medicines Authority

Product Name: AzaMax Xtra Insecticide
APVMA Approval No: 90426/143138

Label Name:	AzaMax Xtra Insecticide

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	11.82 g/L AZADIRACHTIN A & B present as 29.55 g/L Azadirachta indica extract

Mode of Action:	GROUP **UN** INSECTICIDE

Statement of Claims:	For the control of various insect and mite pests in floriculture, ornamentals, nursery plants, turf and potting soil as shown in the Directions for Use

Net Contents:	1 - 1000L

Restraints:	This section contains file attachment.

Directions for Use:	This section contains file attachment.

Other Limitations:	

Withholding Periods:	WITHHOLDING PERIOD (WHP): DO NOT graze treated turf or lawn; or feed turf or lawn clippings from any treated area to poultry or livestock.

Trade Advice:	

General Instructions:	GENERAL INSTRUCTIONS: AzaMax Xtra is a biologically derived product that is active on the above sucking and chewing insect pests on a wide range of plants. Consistent applications are necessary during pest activity. MIXING: Half fill spray tank and commence agitation. Slowly add required amount of AzaMax Xtra. Top up with required quantity of water. Agitate well before commencing spray procedure.

Resistance Warning:	For insecticide resistance management AzaMax Xtra Insecticide is a Group UN insecticide. Some naturally occurring insect biotypes resistant to AzaMax Xtra Insecticide and other Group UN insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if AzaMax Xtra Insecticide or other Group UN insecticides are used repeatedly. The effectiveness of AzaMax Xtra Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Organic Crop Protectants accepts no liability for any losses that may result from the failure of AzaMax Xtra Insecticide to control resistant insects. AzaMax Xtra Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, Organic Crop Protectants representative or local agricultural department agronomist.

Precautions:	PRECAUTIONS Plant Safety: Trials on sensitive plants such as ferns, poinsettias and African violets have resulted in crop damage. Minor phytotoxic effects have occurred in other species. Testing in a small area is therefore recommended before spraying to determine sensitive species and varieties. Re-entry: DO NOT enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

Protections:	INTEGRATED PEST MANAGEMENT Toxic to beneficial arthropods in turf situations. Not compatible with integrated pest management (IPM) programs utilising beneficial arthropods. Minimise spray drift to reduce harmful effects on beneficial arthropods in non-crop areas. PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers. PROTECTION OF HONEY BEES AND OTHER INSECT POLLINATORS May be harmful to bees in turf situations. DO NOT allow spray drift to flowering plants in the vicinity of the treatment area. Before spraying, notify beekeepers to move hives to a safe location with an untreated source of nectar and pollen, if there is potential for managed hives to be affected by the spray or spray drift.

Storage and Disposal:	STORAGE AND DISPOSAL: Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in

	compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Safety Directions:	SAFETY DIRECTIONS: May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and mixing and loading wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length chemical resistant gloves. When applying wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). When using the prepared spray or drench, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). In addition, if applying by mechanical pressure hand wand/gun wear half facepiece respirator. Wash hands after use. After each day's use, wash gloves, respirator and if rubber wash with detergent and warm water, and contaminated clothing.

First Aid Instructions:	FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

First Aid Warnings:	

RESTRAINTS

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of field runoff for at least 3 days after application.

DO NOT apply where the slope exceeds 7%.

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply by aircraft.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between three and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

• Spray droplets are not smaller than a MEDIUM spray droplet size category.

• Minimum distances between the application site and downwind sensitive areas are observed (see the following table titled 'Buffer zones for boom sprayers').

Buffer zones for boom sprayers

Application rate	Boom height above the target canopy	Mandatory downwind buffer zones			
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas
Up to 20 L/ha	0.5 m or lower	0 m	10 m	0 m	0 m
	1.0 m or lower	10 m	40 m	0 m	25 m
Up to 3.0 L/ha	0.5 m or lower	0 m	0 m	0 m	0 m
	1.0 m or lower	0 m	10 m	0 m	0 m

DO NOT apply by a vertical sprayer unless the following requirements are met:

- Spray is not directed above the target canopy.
- The outside of the sprayer is turned off when turning at the end of rows and when spraying the outer row on each side of the application site.
- For dilute water rates up to the maximum listed for each type of canopy specified, minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for vertical sprayers') are observed.

Buffer zones for vertical sprayers

Type of target canopy and dilute water rate	Mandatory downwind buffer zones			
	Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas
2 metres tall and shorter, maximum dilute water rate of 1000 L/ha	0 m	0 m	0 m	0 m
taller than 2 metres (not fully-foliated), maximum dilute water rate of 2000 L/ha	0 m	20 m	0 m	10 m
taller than 2 metres (fully-foliated), maximum dilute water rate of 2000 L/ha	0 m	10 m	0 m	5 m

DIRECTIONS FOR USE

Crop/ Situation	Pest	Rate	Critical Comments
Floriculture, Ornamentals and Nursery Plants (including non-bearing fruit trees)	Two-Spotted Mites, Whitefly, Aphids, Western flower thrips Caterpillars	200 mL - 300 mL / 100 L water	Regularly monitor plants for presence of pests. Use a 7 to 10 day spray regime when pests are present. Multiple applications (3-5) may be required. Use the higher rate under higher pest pressure. Spray volume will vary according to the crop and location. Apply in a spray volume of 250 to 500 L/ha. Addition of a botanical oil concentrate adjuvant is recommended. Note plant safety PRECAUTION below. Control of Western flower thrips may require repeated applications. A single application might not give effective control. Add a botanical oil concentrate for best effects against Western flower thrips. **Caterpillars:** Use of an emulsified botanical oil spray additive is recommended.
	Mealybugs	300-600mL/100L	**Mealybugs**: use the higher rate when pest numbers are high.
	Grasshoppers	300 mL - 600 mL / 100 L water	Apply in sufficient water to achieve good coverage of foliage. Repeat application after 7 days. AzaMax Xtra will inhibit feeding by grasshoppers but does not provide rapid knockdown of grasshoppers.
Ornamental citrus	Citrus leaf miner	250ml / 100L	Begin applying when new flush growth is 4 cm long. Apply every 5 to 14 days with presence of new flush growth. Ensure that new flush leaves are sprayed on top and underneath.
Turf	Lawn Armyworm African black beetle	Mix 1 - 2 L in 200 - 500 L water to treat 1000 m^2	Apply at the first sign of pest infestation and before pest populations build up to damaging levels. Ensure thorough coverage of treated areas. AzaMax Xtra will suppress African black beetle populations but might not provide complete control.
	Sod webworm	5-10L /ha in 400-500 L of water	Monitor for moth activity and apply as soon after egg laying is observed. Apply the lower rate for low pest pressure. Apply every 7-10 days during infestation with HydraSoil soil wetter @ 250 ml/100m2 . Best applied early morning or late afternoon/night
	Ground pearl (suppression)	5-20L/ha in 400-500L	Apply very two weeks during egg laying period for a minimum of 6 weeks using a boom spray with HydraSoil @ 250m/100m2. Apply higher rate to high infestation areas. Apply to turf at min 80% field capacity and drench in with 10mm of irrigation.

Crop/ Situation	Pest	Rate	Critical Comments
Potting soil (for floriculture and ornamentals only)	Fungus Gnats African black beetle	30 mL / 10 L water	Apply as a soil drench using approximately 1 litre mixture per 8 litres potting soil.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION