CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm
2,4-DB 500

HERBICIDE

ACTIVE CONSTITUENT: 500 g/L 2,4-DB
present as the dimethylamine salt

GROUP 4 HERBICIDE

For the control of broadleaf weeds in seedling and established lucerne, medic and clover pastures, peanuts and cereal crops undersown with lucerne, medic or clover as specified in the DIRECTIONS FOR USE.

SL SOLUBLE CONCENTRATE

Nufarm

AUSTRALIAN THROUGH & THROUGH

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply if rain is expected within 4 hours.

DO NOT apply to crop or weeds under stress (not actively growing) due to periods of extreme cold or heat, moisture stress (water logged or drought affected), poor nutrition or previous herbicide treatments as increased levels of crop effect and/or reduced levels of weed control may result.

DO NOT use with hard water (>500 ppm) or water containing suspended matter.

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

SITUATION & CROP	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Lucerne, - seedling and established	Amaranths (*Amaranthus* spp) **Burrs** - Bathurst (*Xanthium spinosum)* - Californian (*Xanthium orientale)* - Noogoora (*Xanthium strumarium)* Capeweed (*Arctotheca calendula)* Charlock (*Sinapis arvensis)* Dock (seedlings) (*Rumex spp)* Evening primrose (*Oenothera stricta)* Fat hen (*Chenopodium album)* Fumitory (*Fumaria* spp) Marshmallow (*Malva parviflora)* Mexican poppy (*Argemone* spp) Mustards (*Sisymbrium* spp) Paterson's curse (Salvation Jane) *(Echium plantagineum)* Plantains (Flat weeds) (*Plantago* spp) Prickly lettuce (*Lactuca serriola)* Shepherd's purse (*Capsella bursapastoris)* Sowthistle (*Sonchus oleraceus)* Spiny emex (Doublegee) (*Emex australis)* Stinging nettle (*Urtica urens)* **Thistles** - Saffron (*Carthamus lanatus)* - Slender (*Carduus pycnocephalus)* - Soldier (*Picnomon acarna)* - Spear (*Cirsium vulgare)* - Variegated (*Silybum marianum)* Turnip weed (*Rapistrum rugosum)* Wild (Mediterranean) turnip (*Brassica tournefortii)* Wireweed (*Polygonum* aviculare)	1-3.2 L/ha	Spray seedling weeds at the 2-6 leaf stage when seedling lucerne has between 1-8 trifoliate leaves. Use lower end of rate range only for 2-4 leaf weeds under good growing conditions or for spray grazing. Small weeds which have been growing slowly under stress and are therefore physiologically older may not be well controlled. To avoid leaf distortion cut or graze established lucerne before spraying. Unless "Spray-Grazing" **DO NOT** cut, mow or graze for 4 weeks after application to maximise weed control.

SITUATION & CROP	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Clover and Medic Pastures - seedling and established Legumes showing acceptable tolerance Clover - Arrow leaf cvs Cefalu, Zulu II - Balansa cvs Bolta, Frontier, Paradana - Berseem cv Elite II - Persian cvs Nitro, Lazer, Leeton, Prolific, Shaftal - Rose cv Hykon - Red - Sub cvs Coolamon, Dalkeith, Gosse, Goulburn, Junee, Nungarin, Riverina, Seaton Park, Trikkala, Urana - White cv NuSiral etc Medic - Barrel cvs Caliph, Parabinga - Burr (polymorpha) cv Circle valley - Gama cvs Paraponto,Sapo - Murex cv Zodiac - Snail cvs Robinson, Sava Serradella - French cv Erica	Refer to Weeds Controlled in Lucerne	1-3.2 L/ha	Spray weeds at the 2-6 leaf stage when seedling medic and clover has between 1-8 trifoliate leaves. Use lower end of rate range only for 2-4 leaf weeds under good growing conditions or for spray grazing. **Medic and Clover safety** Nufarm 2,4-DB 500 may reduce initial vigour of most pasture legumes. Depending on inherent tolerance this may not result in reduced Spring growth or seed yield. Tolerance can vary between seasons, legume type and cultivars. Because of the continuing introduction of new pasture legumes and cultivars caution is advised before applying Nufarm 2,4-DB 500 to any new legume. Caution is especially advised if a treated crop is intended for seed production. Application beyond the 8 trifoliate leaf stage can increase the chance of reduced vigour and production. The use of higher rates can also increase vigour reduction. **DO NOT** exceed 3L/ha when spraying Robinson or Sava snail Medic. **DO NOT** exceed 2.5L/ha when spraying Sapo or Paraponto gama medic. **DO NOT** use on Berseem clover or on Red clover when it is being harvested for seed. Growth of Red clover may be retarded. Nufarm 2,4-DB 500 is unsafe to use on Biserrula, Gland clover, Popany vetch and Orion, Santiago & Serena medics due to the risk of excessive growth and seed. Nufarm 2,4-DB 500 is unsafe to use on chicory and plantain.
Sown Pasture containing Clover or Medics			Apply when clovers or medics are at the 1-8 trifoliate leaf stage. Grass seedlings should have 3 or more leaves. **Medic and Clover safety** refer to the Critical Comments above.
Sub. clover pasture	Doublegee (*Emex australis)*	1.2 L/ha	Spray when Doublegee has less than 14 leaves: best results are obtained when weed has 6-8 leaves. Clover should have 2-8 trifoliate leaves. **DO NOT** apply to pastures when one or more false breaks in the season occur, as uneven germination will result. Doublegee which emerge after spraying will not be controlled. **Medic and Clover safety** refer to the Critical Comments above.
Peanuts - Runner types - Virginia types - Spanish types	*Ipomoea* spp including - Bellvine (*Ipomoea plebeian)* - Morning glory (*Ipomoea purpurea)* - Red convolvulus (*Ipomoea hederifolia* Spiny *emex, Emex australis,* Thornapples (*Datura* spp)	1.3-1.7 L/ha	Apply between 1 and 12 weeks after seedlings have emerged. Use the lower end of the rate range for control of *Ipomoea* spp weeds only.
Wheat, Barley, Oats, Triticale & Cereal Rye undersown with lucerne, medic or clover	Refer to Weeds Controlled in Lucerne	2.1-3.2 L/ha	Spray weeds at the 2-6 leaf stage when seedling lucerne, medic and clover has between 1-8 trifoliate leaves. Use lower rate for 2-4 leaf weeds under good growing conditions. Apply to cereals after the "double ridge" stage which is normally through tillering after the main shoot has five leaves until booting (Z15-33). The time to "double ridge" stage will vary depending on growing conditions and variety. **Medic and Clover safety** refer to the Critical Comments above.
Spot spraying (All Situations)	Refer to Weeds Controlled in Lucerne	200-400 mL/100L water	Use lower rate for easy to kill weeds eg Fat hen or Variegated thistle. For harder to kill weeds eg Capeweed, Marshmallow, use higher rate. Where difficult to wet, hairy or very thick weed populations are present, add Nufarm BS1000® to the spray mixture at a rate of 65mL/100L water.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 7 DAYS AFTER APPLICATION

GENERAL INSTRUCTIONS

Spray applications and drift risk assessment
USE ONLY COARSE or larger spray quality according to the ASAE S572 definition for standard nozzles.

Mixing
Mix with the required amount of water in the spray tank or mixing vessel. Apply using a low-volume boom spray fitted with fan jets, or by aircraft. Use mains or rain water for mixing. If not available, water with a hardness of less than 500ppm (expressed as calcium carbonate equivalent) can be used but is not preferred. **DO NOT** tank mix Nufarm Liase with Nufarm 2,4-DB 500 in hard water for this may cause nozzle blockages.

Application
Apply using boom or spot spray equipment. Application must be uniform. Nozzle type, output, speed and pressure should be adjusted to deliver a Coarse spray quality (ASAE S527) in a minimum of 80L/ha by ground application and a minimum of 40L of water per hectare by aerial application.

COMPATIBILITY
This product can be tank mixed with 900 g/Kg diuron for use in lucerne only. This mix can be very damaging to clovers. Nufarm 2,4-DB 500 is compatible with 800 g/kg Flumetsulam but not for control of Paterson's curse or Doublegee – refer to 800 g/kg Flumetsulam product label. The effectiveness of Dimethoate, Le Mat* and Imidan* may be reduced by alkaline hydrolysis if tank mixed with this product. For these situations the addition of Nufarm Collide surfactant is recommended which should be added to the tank first. The addition of Nufarm CanDo Adjuvant can improve control of weeds under some conditions, for example moisture stress but may also increase early effects on legumes. When tank mixing add the required amount of compatible product (and surfactant if recommended) when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying. If left overnight, agitate thoroughly before spraying.

RESISTANT WEEDS WARNING

GROUP 4 HERBICIDE

Nufarm 2,4-DB 500 is a member of the Phenoxys group of herbicides. Nufarm 2,4-DB 500 has the disruptors of plant cell growth mode of action. For weed resistance management Nufarm 2,4-DB 500 is a Group 4 herbicide. Some naturally-occurring weed biotypes resistant to Nufarm 2,4-DB 500 and other disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Nufarm 2,4-DB 500 or other disruptors of plant cell growth herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of Nufarm 2,4-DB 500 to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS
Drift Warning: DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray drift onto nearby susceptible plants, adjacent crops, crop lands or pastures. Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

PROTECTION OF LIVESTOCK
SPRAY-GRAZING WARNING: Some pasture weeds can be controlled by Spray-Grazing, a combination of herbicide treatment followed by heavy grazing 7-10 days later. Sprayed weeds become more palatable to stock and a higher intake of some weed species after spraying can cause stock deaths and other problems. Increased quantities of poisonous plants may be eaten by stock when the Spray-Grazing technique is being used. Take care if pastures are dominated by Caltrop, Capeweed, Cape tulip, Paterson's curse (Salvation Jane) or Variegated thistle. With Paterson's curse preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. **DO NOT** graze horses or pigs in Paterson's curse (Salvation Jane) infested pasture.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT
DO NOT contaminate streams, rivers or waterways with this product or used containers.

STORAGE AND DISPOSAL
Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.
For non-refillable containers
Triple-rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. **DO NOT** burn empty containers or product.
For returnable containers
Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS
Poisonous if swallowed. May irritate the eyes and skin. This product is alkaline. Avoid contact with the eyes and skin. **DO NOT** inhale spray mist. When preparing spray, wear cotton overalls buttoned to the neck and wrist and washable hat, elbow-length PVC gloves and face shield. When using the prepared spray wear cotton overalls buttoned to the neck and wrist and washable hat. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID
If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

ADDITIONAL STATEMENTS (WHS REGULATIONS 2011)

Harmful if swallowed. Causes serious eye damage. Wash hands and exposed skin thoroughly after handling. **DO NOT** eat, drink or smoke when using this product. Wear protective gloves, clothing, eye and face protection. IF SWALLOWED: Call a POISON CENTER or doctor/physician if you feel unwell. Rinse mouth. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor/physician.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained by scanning the QR code, from your supplier or the Nufarm website.

In case of emergency: Phone 1800 033 498 (24 hrs) and ask for shift supervisor.

PRODUCT STEWARDSHIP INFORMATION AND TOOLS

nufarm.com.au/spraywise is a website which contains resources on spray drift management, recording keeping and application technologies.
spraywisedecisions.com.au is an online weather forecasting program and is recommended for use when planning your pesticide application.
When spraying in or near areas with sensitive crops such as cotton, check online at **satacrop.com.au** for the proximity of sensitive crops mapped by growers.
stewardshipfirst.com.au is a website with CropLife Australia's suite of world-leading product stewardship initiatives, programs and best-practice guides.

CONDITIONS OF SALE

Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute.

® All trade marks owned or used under licence by Nufarm Australia Limited
* Other registered trade marks

Nufarm AUSTRALIAN THROUGH & THROUGH

nufarm.com.au

NUFARM AUSTRALIA LIMITED
103-105 Pipe Rd, Laverton North VIC 3026
For product advice, or to learn more about complementary Nufarm solutions, reach out to your local Nufarm specialist or call **1800 NUFARM (1800 683 276)**.

APVMA Approval No.: 46043/142220 Date: 04-03-2024