POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm
Pyroxasulfone 850WG

HERBICIDE

ACTIVE CONSTITUENT:
850 g/kg PYROXASULFONE

GROUP 15 HERBICIDE

For the pre-emergence control of annual ryegrass, barley grass, annual phalaris, silver grass and toad rush and suppression of certain grass weeds in wheat (not durum wheat), triticale and certain winter legume crops as per the DIRECTIONS FOR USE.

WG WATER DISPERSIBLE GRANULE

Nufarm

AUSTRALIAN THROUGH & THROUGH

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply with aircraft.
DO NOT plant durum wheat (*Triticum durum*) after the application of Nufarm Pyroxasulfone 850WG (refer to **Crop Rotation Recommendations** for further advice).
DO NOT apply if heavy rain has been forecast within 48 hours.
DO NOT apply unless incorporation by sowing (IBS) can be performed within 3 days of application.
DO NOT apply to waterlogged soil.
DO NOT allow first irrigation tailwater from land treated with Nufarm Pyroxasulfone 850WG to enter aquatic and wetland areas including aquacultural ponds, surface streams and rivers.

Spray drift restraints

Specific definitions for terms used in this section of the label can be found at www.apvma.gov.au/spraydrift
DO NOT allow **bystanders** to come into contact with the spray cloud.
DO NOT apply in a manner that may cause an **unacceptable impact** to **native vegetation**, **agricultural crops**, **landscaped gardens** and **aquaculture production**, or cause contamination of plant or livestock commodities, outside the application site from **spray drift**. The buffer zones in the relevant buffer zone table below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.
DO NOT apply unless the **wind speed** is between 3 and 20 kilometres per hour at the **application site** during the time of application.
DO NOT apply if there are **hazardous surface temperature inversion** conditions present at the **application site** during the time of application. **Surface temperature inversion conditions** exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.
DO NOT apply by a boom sprayer unless the following requirements are met:

- spray droplets are not smaller than a COARSE **spray droplet size category**
- minimum distances between the **application site** and downwind **sensitive areas** are observed. See "**Mandatory Downwind Buffer Zones**" section of the following table titled "Buffer Zones for Boom Sprayers" –

Buffer Zones for Boom Sprayers

Application rate	Mandatory buffer zones
	Natural Aquatic Areas
118 g/ha	80 m

CROP	WEEDS	RATE	CRITICAL COMMENTS
Wheat (not durum wheat), triticale Chickpeas, field peas, lentils, lupins	Annual ryegrass (*Lolium rigidum*) Annual Phalaris or paradoxa grass (*Phalaris paradoxa* only) Barley grass (*Hordeum leporinum*) Silver grass (*Vulpia bromoides, Vulpia myuros*) Toad rush (*Juncus bufonius*) ***Suppression of:** Great brome (*Bromus diandrus*) Wild oat (*Avena fatua*) *Refer suppression of great brome and wild oat in general instructions for further details.	118 g/ha	Apply pre-sowing and incorporate by sowing (IBS) using knife points and press wheels or narrow points and harrows. For best results apply just before sowing (refer to **Interval between Application and Sowing section of General Instructions**). Avoid throwing treated soil into adjacent crop rows when sowing with knife points and press wheels. To reduce the risk of crop effects refer **to Crop Safety section of General Instructions**. **Cultivation:** To optimize weed control apply directly to uncultivated soil. Weed control may be greatly reduced where weed seeds have been buried by cultivation prior to sowing. **Rainfall soon after application:** • Weed control may be adversely affected by insufficient rainfall within 7 to 10 days after application. Adequate rainfall is necessary to facilitate uptake of the product by the germinating weed seeds, however the quantity of rainfall required will depend on many factors including stubble load, soil type, the existing soil moisture at sowing, the pattern of rainfall and other considerations. • In soils prone to leaching, rainfall which is sufficiently heavy to cause movement of the herbicide out of the weed seed zone may lead to reduced weed control. **Other factors which may adversely affect weed control include:** • uneven application • application to rigid or cloddy soil • stubble, plant residue or other ground cover particularly where this exceeds 50% • germinated and emerged weeds that are not controlled by a knockdown herbicide The factors above, when combined, may substantially reduce weed control. Competition provided by the crop can assist with the final weed control achieved by Nufarm Pyroxasulfone 850WG. Chickpea, field pea, lentil and lupin crops may provide less competition than cereal crops, hence weeds that survive the application of Nufarm Pyroxasulfone 850WG may grow taller (relative of the height of the crop), tiller more and generally give the appearance that weed control is poorer compared to weed control in wheat or triticale.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

Harvest:

All crops - NOT REQUIRED WHEN USED AS DIRECTED.

Grazing/Stockfood:

Wheat, Triticale - DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 6 WEEKS AFTER APPLICATION.

Chickpeas, Field peas, Lentils, Lupins - DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 8 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Nufarm Pyroxasulfone 850WG is a residual, soil applied, pre-emergence herbicide. It is absorbed by the roots and to a lesser extent by the shoots of germinating weeds, and works by inhibiting growth in the meristematic area. Weed control is optimised when Nufarm Pyroxasulfone 850WG is applied evenly to moist soil just prior to incorporation by sowing and there is sufficient rainfall soon after sowing to ensure uptake of the herbicide by germinating weeds. Weed control may be greatly reduced where weed seeds have been buried by cultivation prior to application. Weed control may also be reduced where there is insufficient soil moisture for herbicide uptake or in soils prone to leaching where rainfall is sufficiently heavy to cause movement of the herbicide out of the weed seed zone. Nufarm Pyroxasulfone 850WG will not reliably control emerged weeds. A knockdown herbicide should be used to control emerged weeds at sowing.

Crop Safety

Nufarm Pyroxasulfone 850WG generally shows good crop selectivity when used as directed. The following directions will help minimise the risk of crop effects.

- **DO NOT** plant durum wheat after the application of Nufarm Pyroxasulfone 850WG as it may be severely damaged. Refer to **Crop Rotation Recommendations** for further advice.
- When incorporation is by sowing with knife points and press wheels avoid throwing treated soil into adjacent crop rows.
- **DO NOT** use a combination of both press wheels and a covering device such as harrows or chains when sowing.

The potential for crop damage is increased when there is substantial rainfall after the application of Nufarm Pyroxasulfone 850WG, especially where this leads to temporary waterlogging. Situations which lead to concentration of herbicide in the planting row, or movement of herbicide to the depth of the crop seed, may also increase the potential for crop damage. This includes the following scenarios -

- Where deep furrows are formed by the sowing operation, soil movement into the crop row may occur due to wind or heavy rainfall soon after sowing resulting in concentration of herbicide in the crop row.
- Where soil has a high potential for leaching, heavy rainfall between application and crop emergence may result in movement of herbicide into the crop seed zone.

Other circumstances which may increase the potential for crop damage include where Nufarm Pyroxasulfone 850WG is applied in tank mixes with other herbicides, where crop vigour is reduced due to factors such as frosts, insect attack or crop disease, when weather damaged seed is used and/or with the use of some fungicide seed treatments especially in conjunction with crop varieties with short coleoptile length. A combination of individual factors which increase the potential for crop damage may increase the extent of crop damage.

Chickpeas, field peas, lentils and lupins

- Nufarm Pyroxasulfone 850WG may occasionally delay emergence or flowering of winter legume crops.
- Luxor variety of albus lupin has been identified as potentially more sensitive to Nufarm Pyroxasulfone 850WG than other lupin varieties, particularly in situations of late sowing and/or wet conditions around the time of sowing.

Incorporation by Sowing

Nufarm Pyroxasulfone 850WG should be applied prior to sowing, and incorporated by sowing using knife points and press wheels, or narrow points and harrows. When incorporation is by knife points and press wheels, weeds germinating in the seed row may not be controlled. Weeds germinating from depth, weeds just about to emerge, or weeds that have emerged which are not controlled by a knockdown herbicide at sowing may not be controlled by Nufarm Pyroxasulfone 850WG.

Interval between Application and Sowing

Incorporate by sowing as soon as practicable after the application of Nufarm Pyroxasulfone 850WG, but no later than 3 days after application.

Sandy Soils

Weed control may be reduced in soil prone to leaching where rainfall after application and sowing is sufficiently heavy to cause movement of the herbicide out of the weed seed zone.

Suppression of great brome and wild oat

Nufarm Pyroxasulfone 850WG is most effective when grass weed seeds are present on or very close to the soil surface at the time of application. For this reason, it is recommended that Nufarm Pyroxasulfone 850WG is applied to uncultivated soil. As the depth of weed seeds increases, control from Nufarm Pyroxasulfone 850WG tends to

decrease. It is rare that all great brome and wild oat weed seeds will be on the soil surface at the time of Nufarm Pyroxasulfone 850WG application, especially considering that these seeds may remain viable in the soil for several seasons. Plants may germinate from seeds buried by the sowing operation in previous seasons, by livestock or by weed seed self-burial mechanisms particularly in some soil types (e.g. cracking clays and sand). **Therefore only partial control or suppression of the great brome or wild oat population should generally be expected**. In these situations, a follow up application with a suitable post-emergence herbicide may be required to control remaining plants.

Mixing

Ensure sprayer and nozzle filters are clean before preparing the spray mixture. Half fill the spray tank with water and, with the agitators in motion, add the correct amount of Nufarm Pyroxasulfone 850WG directly to the spray tank. Complete filling the tank with agitators in motion. Agitation must continue before and during spraying. When other products are to be applied in addition to Nufarm Pyroxasulfone 850WG, always add Nufarm Pyroxasulfone 850WG to the spray tank first and ensure it is fully dispersed in the spray tank before adding other products.

Application

Ensure complete and even spray coverage of the soil is achieved. Poor spray coverage may result from application to ridged or excessively cloddy soil or in situations of high stubble, plant residue or other ground cover. A significant reduction in weed control may result where stubble, plant residue or other ground cover exceeds 50%, and in situations where a 'cold' or incomplete burn of stubble results in a mass of material which can act as a physical barrier between the herbicide and germinating weeds - this can be exacerbated in header trails where there may be greater weed seed numbers and higher levels of plant residue. Weed control can be particularly affected where Nufarm Pyroxasulfone 850WG is applied to a barrier of stubble, plant residue or other ground cover and there is insufficient following rainfall to transfer Nufarm Pyroxasulfone 850WG to the soil surface and the germinating weed seeds.

Equipment

Ground Sprayers – standard boom sprayers only are recommended and must be fitted with by-pass or mechanical agitation. It is recommended that 50 to 100 L water/ha is applied with spray droplets of a COARSE droplet size category. In some situations (e.g. high stubble loads) high water volumes may give higher levels of weed control.

Aircraft - **DO NOT** apply Nufarm Pyroxasulfone 850WG by aircraft.

Sprayer clean-up

Following the use of Nufarm Pyroxasulfone 850WG the spraying equipment should be thoroughly cleaned before it is used for application of other products. Cleaning should occur immediately following application of Nufarm Pyroxasulfone 850WG. The spray unit should first be completely emptied. The sprayer, including all filters and lines, should be thoroughly rinsed with water, to remove all traces of product. Ensure that the sprayer clean-up is carried out in an area that is clear of waterways, desirable vegetation and tree roots. If using Nufarm Pyroxasulfone 850WG with a tank-mix partner, refer to the sprayer clean-up instructions for the other product, which may be more rigorous than those for Nufarm Pyroxasulfone 850WG.

Crop Rotation Recommendations

Nufarm Pyroxasulfone 850WG breaks down by microbial degradation, which is favoured by warm, moist aerobic soil. Minimum recropping intervals (months after application) have been established for Nufarm Pyroxasulfone 850WG to minimise the risk of damage to following crops (see table below). However, environmental and agronomic factors make it impossible to eliminate all risk and therefore the potential for damage to following crops exists.

Rainfall of less than the minimum interim rainfall required (see table below) may result in extended recropping intervals. Interim rainfall is the total rainfall between the application of Nufarm Pyroxasulfone 850WG and planting of the particular following crop. For recropping with winter crops, where a minimum of 250 mm of interim rainfall is required, if rain from application to the end of spring is less than 125 mm and isolated heavy summer and autumn falls and break rains are required to achieve the 250 mm interim rainfall, then extended recropping intervals may apply.

Crops	Recropping recommendation	
	Minimum recropping interval	Minimum Interim Rainfall
Wheat (not durum wheat) and triticale	0 months	0 mm
Cotton, maize, mung beans, sorghum, soybeans and sunflowers	5 months	150 mm
Barley, canola*, chickpeas**, faba beans, field peas**, lentils**, lupins**, vetch and subterranean clover	9 months	250 mm
Durum wheat, oats, lucerne and medic	21 months	550 mm

*For canola sown the year after the application of Nufarm Pyroxasulfone 850WG there may occasionally be some crop stunting but no yield reductions have been measured.
**Chickpeas, field peas, lentils and lupins can be sown immediately after the application of Nufarm Pyroxasulfone 850WG where Nufarm Pyroxasulfone 850WG has not already been incorporated. However, where Nufarm Pyroxasulfone 850WG has been incorporated into the soil, for example, by a previous sowing operation for a subsequently failed crop, these legume crops should not be sown for at least 9 months after the application of Nufarm Pyroxasulfone 850WG.
For advice on crops and situations not listed above, contact Nufarm Australia Ltd.

RESISTANT WEEDS WARNING

GROUP 15 HERBICIDE

Nufarm Pyroxasulfone 850WG is a member of the isoxazoline group of herbicides and has the inhibitor of very long chain fatty acids (VLCFA inhibitors) mode of action. For weed resistance management Nufarm Pyroxasulfone 850WG is a Group 15 herbicide. Some naturally-occurring weed biotypes resistant to Nufarm Pyroxasulfone 850WG, and other Group 15 herbicides, may exist through normal genetic variability in any weed population. These resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Nufarm Pyroxasulfone 850WG or other Group 15 herbicides. **DO NOT** rely exclusively on Nufarm Pyroxasulfone 850WG for weed control. Use as part of an integrated weed management program involving herbicides with other modes of action and non-chemical methods of control. CropLife Australia resistance management strategies are available from your local agricultural chemical supplier or at the CropLife Australia website (www.croplife.org.au). Refer to these strategies for details on how to manage the build-up of resistant weeds on your farm.
Since occurrence of resistant weeds is difficult to detect prior to use Nufarm Australia Ltd accepts no liability for any losses that may result from the failure of Nufarm Pyroxasulfone 850WG to control resistant weeds.

COMPATIBILITY

Crop damage seen in adverse conditions, particularly wet or waterlogged conditions (refer **Crop Safety** above) may be exacerbated when Nufarm Pyroxasulfone 850WG is used in conjunction with other herbicides that may also cause crop damage in such conditions. **Always refer to the crop tolerance, plant back restrictions, rate recommendations and other directions for use on the label of the tank mix partner.**
Refer to **Mixing** section above for advice on preparing tank mixtures with Nufarm Pyroxasulfone 850WG. Mixtures with products containing paraquat require particular attention to these instructions, including ongoing agitation to ensure Nufarm Pyroxasulfone 850WG remains in suspension in the spray tank. For advice on compatibilities not listed below, contact Nufarm Australia Ltd.
For application prior to planting wheat or triticale
Nufarm Pyroxasulfone 850WG is compatible with any one of the following herbicides - Nufarm Associate®, Nufarm Avadex® Xtra, Nufarm Kamba® 750, diuron 900 WG formulations, 960 g/L S-metolachlor formulations, Nufarm 2,4-D Ester 680, Nufarm 2,4-D Amine 625, Glean®, Nufarm Crucial®, Nufarm Glyphosate 450, Nufarm weedmaster DST®, Nufarm weedmaster Duo®, Nufarm TriflurX®, Goal* EC, paraquat, Spray.Seed*, Nufarm Nail®, triasulfuron, B-Power*, Monza*, oxyfluorfen.
Nufarm Pyroxasulfone 850WG is compatible with mixtures of glyphosate with any one of the following herbicides - Nufarm Associate®, Nufarm Kamba® 750, Nufarm 2,4-D Ester 680, Goal* EC, Nufarm Nail®, B-Power*, Monza* and oxyfluorfen.
Nufarm Pyroxasulfone 850WG is compatible with any one of the following insecticides - Fastac® and Le-mat®.

For application prior to planting chickpeas, field peas, lentils or lupins

Knockdown herbicides, some "spike" herbicides and insecticides shown to be compatible with Nufarm Pyroxasulfone 850WG prior to planting cereals, should also be suitable prior to planting chickpeas, field peas, lentils or lupins e.g. Nufarm Glyphosate 450, weedmaster DST®, weedmaster Duo®, Goal® EC, paraquat, Spray.Seed*, Nufarm Nail®, Fastac® and Lemat®. Note that plantback restrictions may render some herbicides unsuitable for mixing with Nufarm Pyroxasulfone 850WG where legume crops are to be planted.

Limited studies have shown that the following residual herbicides appear to be compatible with Nufarm Pyroxasulfone 850WG when used according to label directions -

Chickpeas: Nufarm TriflurX®, simazine 900 WG formulations, Nufarm TriflurX® + simazine, Nufarm Igran®.

Field peas: Nufarm TriflurX®, Nufarm Pendimethalin 440, Bladex* 900 WG, Nufarm Igran®.

Lentils: Nufarm Pendimethalin 440, Nufarm Igran®.

Lupins: Nufarm TriflurX®, simazine 900 WG formulations, Nufarm TriflurX® + simazine, simazine + atrazine 900 WG formulations, Nufarm Pendimethalin 440, Nufarm Igran®.

PRECAUTIONS

Re-entry Period

DO NOT allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto non-target plants, cropping lands or pastures.

Undersown Pasture Species - DO NOT undersow with pasture species (legumes or grasses) following the application of Nufarm Pyroxasulfone 850WG.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate streams, rivers or watercourses with this product or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. **DO NOT** re-use container for any other purpose. **DO NOT** burn empty containers or product.

SAFETY DIRECTIONS

May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When using together with other products, consult their label safety directions. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length chemical-resistant gloves. Wash hands after use. After each day's use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

ADDITIONAL STATEMENTS (WHS REGULATIONS 2011)

May cause an allergic skin reaction. Causes serious eye irritation. Suspected of causing cancer. May cause damage to organs through prolonged or repeated exposure. Obtain special instructions before use. **DO NOT** handle until all safety precautions have been read and understood. **DO NOT** breathe dust or spray. Wash skin thoroughly after handling. Contaminated work clothing should not be allowed out of the workplace. Wear protective gloves/protective clothing/face protection. IF exposed or concerned, get medical advice/attention. IF IN EYES: rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do so. If eye irritation persists: get medical advice/attention. IF ON SKIN: Wash with plenty of water. If skin irritation or rash occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse. Get medical advice/attention if you feel unwell. Collect spillage.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained by scanning the QR code, from your supplier or the Nufarm website.

In case of emergency: Phone 1800 033 498 (24 hrs) and ask for shift supervisor. .

PRODUCT STEWARDSHIP INFORMATION AND TOOLS

nufarm.com.au/spraywise is a website which contains resources on spray drift management, recording keeping and application technologies.
spraywisedecisions.com.au is an online weather forecasting program and is recommended for use when planning your pesticide application.
When spraying in or near areas with sensitive crops such as cotton, check online at **satacrop.com.au** for the proximity of sensitive crops mapped by growers.
stewardshipfirst.com.au is a website with CropLife Australia's suite of world-leading product stewardship initiatives, programs and best-practice guides.

CONDITIONS OF SALE

Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute.

® All trade marks owned or used under licence by Nufarm Australia Limited
* Other registered trade marks

Nufarm AUSTRALIAN THROUGH & THROUGH

nufarm.com.au

NUFARM AUSTRALIA LIMITED
103-105 Pipe Rd, Laverton North VIC 3026
For product advice, or to learn more about complementary Nufarm solutions,
reach out to your local Nufarm specialist or call **1800 NUFARM (1800 683 267)**.

APVMA No.: 93314/138710 Date: 21-11-2023