CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm SENTRY®

HERBICIDE

ACTIVE CONSTITUENTS:
525 g/kg IMAZAPIC
175 g/kg IMAZAPYR

GROUP 2 HERBICIDE

For pre-emergence or post-emergence application in various imidazolinone herbicide tolerant crops for control of a range of weeds as per the DIRECTIONS FOR USE.

WG WATER DISPERSIBLE GRANULE

Nufarm

AUSTRALIAN THROUGH & THROUGH

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply to barley, wheat or canola varieties that lack imidazolinone tolerance or oats other than Kingbale® and Archer® varieties.
DO NOT apply post-emergence to imidazolinone tolerant barley or oats.
DO NOT apply to crops or weeds which are suffering moisture stress (waterlogged or drought affected) or stressed from previous herbicide or foliar fertiliser treatment.
DO NOT apply by aircraft.

Spray Drift Restraints

Specific definitions for terms used in this section of the label can be found at www.apvma.gov.au/spraydrift.
DO NOT allow bystanders to come into contact with the spray cloud.
DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table below provides guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.
DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.
DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.
DO NOT apply by a boom sprayer unless the following requirements are met:

- minimum distances between the application site and downwind sensitive areas (see 'Mandatory downwind buffer zones' section) are observed.

Buffer zones for boom sprayers

Application rate	Mandatory downwind buffer zones	
	Natural aquatic areas	Vegetation areas
Up to maximum label rate	50 m	0 m

SECTION 1 - PRE-EMERGENCE USE

SITUATION	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Imidazolinone herbicide tolerant Barley, Canola and Wheat varieties only Kingbale® or Archer® oats - **DO NOT** apply to other oat varieties.	Barley grass (*Hordeum* spp.) Brome grass (*Bromus* spp.) Climbing buckwheat (*Fallopia convolvulus*) Hedge mustard (*Sisymbrium officinale*) Indian hedge mustard (*S. orientale*) Wild radish (*Raphanus raphanistrum*) Wireweed (*Polygonum aviculare*) **Suppression of the following weeds:** Annual ryegrass (*Lolium rigidum*) Capeweed (*Arctotheca calendula*) Clover (*Trifolium* spp.) Fumitory (*Fumaria* spp.) Long storksbill (*Erodium botrys*) Paterson's curse (*Echium plantagineum*) Phalaris (*Phalaris* spp.) Volunteer barley (*Hordeum vulgare* - other than imidazolinone herbicide tolerant barley varieties) Volunteer canola (*Brassica napus* - other than imidazolinone herbicide tolerant canola varieties) Volunteer wheat (*Triticum aestivum* - other than imidazolinone herbicide tolerant wheat varieties) Volunteer oats (*Avena sativa* - other than imidazolinone herbicide tolerant oat varieties) Wild oats (*Avena* spp.)	40 - 50 g/ha Refer to Compatibility Section and Critical Comments for advice and rates of herbicide tank mix partners.	**DO NOT apply Nufarm Sentry® (to a crop) more than once in a growing season.** **Where Nufarm Sentry® is used pre-emergence followed by a post-emergent treatment with another imidazolinone-based herbicide or any other Group 2 herbicide to control grasses in cereals or brassicaceous weeds in canola, only sow an imidazolinone herbicide tolerant crop the following season.** **Ensure follow crop comments and restrictions on the label are consulted prior to use.** Nufarm Sentry® may be applied up to and immediately prior to planting with incorporation by the sowing process using knife/blade points and press wheels. Best weed control will be achieved when applied to weed free, moist, friable soil immediately prior to sowing. Nufarm Sentry® can be applied to dry soil but will not be active until follow up rain disperses the product to the root zone of germinating weeds. Applying Nufarm Sentry® to dry soil when weeds are germinating from depth can impair performance. A 15-20 mm rainfall event received within a fortnight of application will limit this risk. The low rate of Nufarm Sentry® may not provide adequate control when used in heavy stubble covers, on high weed density burdens and in heavier soil types. Tank mixing with a suitable pre-emergence grass herbicide is recommended (see Compatibility section). Applying Nufarm Sentry® in tank mix with pre-emergence grass herbicides will improve grass control, particularly Annual ryegrass, and the control of some broadleaf weed species. This is especially applicable when using Nufarm Sentry® at 40 g/ha. Choice of the mixing partner regime depends upon weed spectrum and site conditions. **DO NOT** apply a Nufarm Sentry® tank mix in a manner contrary to advice provided on the label of the mixing partner. **Wherever possible, an appropriate follow up post- emergence herbicide regime using alternative modes of action herbicides is recommended for weed seed set management.** Harvest Weed Seed Set Control (HWSSC) measures are recommended in all situations in order to limit the survival of Group 2 resistant seeds, but are essential where Group 2 based herbicides, including Nufarm Sentry®, have been applied pre- and post-emergence. See Best Management Practice. **Imidazolinone herbicide tolerant Oats** Kingbale® and Archer® are the only varieties approved for use with Nufarm Sentry®, **DO NOT** apply to other oat varieties. Kingbale® and Archer® are oaten hay varieties. Grain harvested from failed hay crops, screenings from seed crops or excess seed cannot be delivered to bulk handlers for export. Grain from these situations can be fed to livestock on the Australian domestic market but not used for the domestic human consumption market. If crops are stressed due to drought conditions, they should be grazed or harvested for grain for livestock on the domestic market and not cut for export oaten hay markets. **DO NOT** use Nufarm Sentry® at 40 – 55 g/ha on shallow, strongly duplex soils in Western Australia.

*Alternatively to Nufarm CanDo™ Adjuvant another registered high quality methylated seed oil adjuvant or paraffinic oil can be used according to their label instructions.

SECTION 2 - POST-EMERGENCE USE

CROP	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
POST-EMERGENCE USE Imidazolinone herbicide tolerant Canola varieties only	Hedge mustard (*Sisymbrium officinale*) Indian hedge mustard (*S. orientale*) London rocket (*S. irio*) Shepherd's purse (*Capsella bursa-pastoris)* Wild radish (*Raphanus raphanistrum*) Wild turnip (*Brassica tournefortii*) *Continued over*	20 g/ha + Nufarm CanDo™ Adjuvant* at 0.5 L/100L spray volume Refer to Compatibility Section and Critical Comments for advice and rates of Nufarm Archer® 750 and the 240 g/L clethodim formulation.	**DO NOT use a post emergence application if a pre- emergence Nufarm Sentry® application was applied.** **DO NOT apply more than once per season to any one crop.** **Ensure follow-crop comments and restrictions on the label are consulted prior to use.** Apply to crops in the 2-6 leaf stage. Apply early post-emergence to actively growing grass weeds in the 3-5 leaf stage (Z13-15) and broadleaf weeds in the 2 - 4 leaf stage (20 g/ha rate) or grass weeds in the 3 leaf to 2 tillers stage (Z13-22) and broadleaf weeds in the 2-6 leaf stage (40 or 55 g/ha rate). Where Nufarm Sentry® is used alone and annual ryegrass is a significant weed in the total weed population, use the 55 g/ha rate to achieve suppression. Weeds will either be killed or will be stunted and uncompetitive with the crop. Nufarm Sentry® is physically and biologically compatible with Nufarm Archer® 750 and 240 g/L clethodim formulations, both in two- way and three-way mixes. A 240 g/L clethodim formulation aids in the control of selected grass species (refer to the 240 g/L clethodim formulation label). When these weeds require control apply a tank mix of Nufarm Sentry® and a 240 g/L clethodim formulation using the recommended label rate of the 240 g/L clethodim formulation. 175 mL/ha of a 240 g/L clethodim formulation will normally be sufficient to achieve good control of light populations. Use higher rates when grasses are primary weeds in the paddock, and when their growth stage requires it, to ensure highest levels of control. When mixing with a 240 g/L clethodim formulation, use 0.5 L/100L spray volume of Nufarm CanDo™ Adjuvant*.

CROP	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
POST-EMERGENCE USE Imidazolinone herbicide tolerant Canola varieties only (*cont.*)	Amsinckia (*Amsinckia* spp.) Barley grass (*Hordeum leporinum*) Bedstraw (*Galium tricornutum*) Brome grass (*Bromus* spp.) Climbing buckwheat (*Fallopia convolvulus*) Clover (*Trifolium* spp.) Corn gromwell (*Buglossoides arvense*) Crassula (*Crassula* spp.) Dead nettle (*Lamium amplexicaule*) Double gee (*Emex australis*) Fumitory (*Fumaria* spp.) Hedge mustard (*Sisymbrium officinale*) Indian hedge mustard (*S. orientale*) London rocket (*S. irio*) Paterson's curse (*Echium plantagineum*) Phalaris (*Phalaris* spp.) Shepherd's purse (*Capsella bursa-pastoris)* Toadrush (*Juncus bufonius*) Volunteer barley (*Hordeum vulgare* - other than imidazolinone herbicide tolerant barley varieties) Volunteer canola (*Brassica napus* - other than imidazolinone herbicide tolerant canola varieties) Volunteer wheat (*Triticum aestivum* - other than imidazolinone herbicide tolerant wheat varieties) Volunteer oats (*Avena sativa* - other than imidazolinone herbicide tolerant oat varieties) Wild oats (*Avena* spp.) Wild radish (*Raphanus raphanistrum*) Wild turnip (*Brassica tournefortii*) Wireweed (*Polygonum aviculare*) **Suppression only:** Annual ryegrass# (*Lolium rigidum*) Medics (*Medicago* spp.) Silver grass (1-2 leaf only) (*Vulpia* spp.) Storksbill (*Erodium* spp.) Volunteer vetch (*Vicia* spp.)	40 or 55 g/ha + Nufarm CanDo™ Adjuvant* at 0.5 L/100L spray volume Refer to Compatibility Section and Critical Comments for advice and rates of Nufarm Archer® 750 and the 240 g/L clethodim formulation	Nufarm Archer® 750 aids in the control of legumes, prickly lettuce, capeweed and other thistle-like weed species. When these weeds require control apply a tank mix of Nufarm Sentry® and Nufarm Archer® 750 using the recommended label rate of Nufarm Archer® 750. Rates of Nufarm Archer® 750 above 60 mL/ha can impair grass control. The addition of a 240 g/L clethodim formulation at 175 mL/ha to the tank mix will overcome this antagonism. (Refer to the Compatibility section of this label and the 240 g/L clethodim formulation and Nufarm Archer® 750 labels for further details of use). #Where Group 1 and/or Group 2 – resistant ryegrass is known to be present or ryegrass populations in excess of 200 plants per m^2 are expected, an application of Nufarm Pendimethalin 440, Nufarm TriflurX® or Nufarm Avadex® Xtra should be made prior to sowing. **DO NOT** use Nufarm Sentry® at 40 - 55 g/ha on shallow, strongly duplex soils in Western Australia.

*Alternatively to Nufarm CanDo™ Adjuvant, another registered high quality methylated seed oil adjuvant or paraffinic oil can be used according to their label instructions.

CROP	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
POST-EMERGENCE USE Imidazolinone herbicide tolerant Wheat (single gene) varieties only	Hedge mustard (*Sisymbrium officinale*) Indian hedge mustard (*S. orientale*) London rocket (*S. irio*) Shepherd's purse (*Capsella bursa- pastoris*) Wild radish (*Raphanus raphanistrum*) Wild turnip (*Brassica tournefortii*)	20 g/ha + Nufarm CanDo™ Adjuvant* at 0.5 L/100L spray volume	**DO NOT use a post emergence application if a pre- emergence Nufarm Sentry® application was applied.** **DO NOT apply more than once per season to any one crop. Ensure follow-crop comments and restrictions on the label are consulted prior to use.** Apply to crops in the 4 leaf (Z14) to commencement of flag leaf (Z37) stage. Apply early post-emergence to actively growing grass weeds in the 3 - 5 leaf stage (Z13-15) and broadleaf weeds in the 2 - 4 leaf stage (20 g rate) or grass weeds in the 3 leaf to 2 tillers stage (Z13-22) and broadleaf weeds in the 2 - 6 leaf stage (40 g/ha rate). Weeds will either be killed or will be stunted and uncompetitive with the crop. #Where ALS resistant ryegrass is known to be present or ryegrass populations in excess of 200 plants per m^2 are expected, an application of a Group 3 herbicide, such as Nufarm Pendimethalin 440 or Nufarm TriflurX®, or a Group 15 herbicide such as Nufarm Avadex® Xtra should be made prior to sowing. **DO NOT** use Nufarm Sentry® at 40 g/ha on shallow, strongly duplex soils in Western Australia.
	Amsinckia (*Amsinckia* spp.) Barley grass (*Hordeum leporinum*) Bedstraw (*Galium tricornutum*) Brome grass (*Bromus* spp.) Climbing buckwheat (*Fallopia convolvulus*) Clover (*Trifolium* spp.) Corn gromwell (*Buglossoides arvense*) Crassula (*Crassula* spp.) Dead nettle (*Lamium amplexicaule*) Double gee (*Emex australis*) Fumitory (*Fumaria* spp.) Hedge mustard (*Sisymbrium officinale*) Indian hedge mustard (*S. orientale*) London rocket (*S. irio*) Paterson's curse (*Echium plantagineum*) Phalaris (*Phalaris* spp) Shepherd's purse (*Capsella bursa- pastoris*) Toadrush (*Juncus bufonius*) Volunteer barley (*Hordeum vulgare* - other than imidazolinone herbicide tolerant barley varieties) Volunteer canola (*Brassica napus* - other than imidazolinone herbicide tolerant canola varieties) Volunteer wheat (*Triticum aestivum* - other than imidazolinone herbicide tolerant wheat varieties) Volunteer oats (*Avena sativa* - other than imidazolinone herbicide tolerant oat varieties) Wild oats (*Avena* spp.) Wild radish (*Raphanus raphanistrum*) Wild turnip (*Brassica tournefortii*) Wireweed (*Polygonum aviculare*) **Suppression only:** Annual ryegrass# (*Lolium rigidum*) Medics (*Medicago* spp.) Silver grass (1-2 leaf only) (*Vulpia* spp.) Storksbill (*Erodium* spp.) Volunteer vetch (*Vicia* spp.)	40 g/ha + Nufarm CanDo™ Adjuvant* at 0.5 L/100L spray volume	

CROP	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
POST-EMERGENCE USE Imidazolinone herbicide tolerant Wheat (single gene) varieties only (*cont.*)	**Above weeds plus:** African turnip weed (*Sisymbrium thellungii*) Capeweed (*Arctotheca calendula*) Spear thistle (*Cirsium vulgare*) Spreading night phlox (*Zaluzianskya divaricata*) Turnip weed (*Rapistrum rugosum*) Variegated thistle (*Silybum marianum*) Volunteer canola (*Brassica napus*) **Suppression only:** Cat's-ear (*Hypochaeris radicata*) Common vetch (*Vicia* spp.) Cotula (*Cotula australis*) Mintweed (*Salvia reflexa*) Medics (*Medicago* spp.) Prickly lettuce (*Lactuca serriola*) Saffron thistle (*Carthamus lanatus*) Sowthistle (*Sonchus oleraceus*) Volunteer faba beans (*Vicia faba*)	40 g/ha + 455 mL/ha Nufarm MCPA LVE 570 + Nufarm CanDo™ Adjuvant* at 0.5 L/100L spray volume	**DO NOT use a post emergence application if a pre- emergence Nufarm Sentry® application was applied.** **DO NOT apply more than once per season to any one crop. Ensure follow-crop comments and restrictions on the label are consulted prior to use.** Apply to crops in the 4 leaf (Z14) to commencement of flag leaf (Z37) stage. Apply early post-emergence to actively growing grass weeds in the 3 - 5 leaf stage (Z13-15) and broadleaf weeds in the 2 - 4 leaf stage (20 g rate) or grass weeds in the 3 leaf to 2 tillers stage (Z13-22) and broadleaf weeds in the 2 - 6 leaf stage (40 g/ha rate). Weeds will either be killed or will be stunted and uncompetitive with the crop. #Where ALS resistant ryegrass is known to be present or ryegrass populations in excess of 200 plants per m^2 are expected, an application of a Group 3 herbicide, such as Nufarm Pendimethalin 440 or Nufarm TriflurX®, or a Group 15 herbicide such as Nufarm Avadex® Xtra should be made prior to sowing. **DO NOT** use Nufarm Sentry® at 40 g/ha on shallow, strongly duplex soils in Western Australia.

*Alternatively to Nufarm CanDo™ Adjuvant, another registered high quality methylated seed oil adjuvant or paraffinic oil can be used according to their label instructions.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

Imidazolinone herbicide tolerant Barley and Canola:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 6 WEEKS AFTER APPLICATION.
HARVEST FOR GRAIN: NOT REQUIRED WHEN USED AS DIRECTED.

Imidazolinone herbicide tolerant Wheat:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION.
HARVEST FOR GRAIN: NOT REQUIRED WHEN USED AS DIRECTED.

Imidazolinone herbicide tolerant Oat:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION.
HARVEST FOR GRAIN: NOT REQUIRED WHEN USED AS DIRECTED.

EXPORT OF TREATED PRODUCE

Growers should note that maximum residue limits (MRLs) or import tolerances may not exist in all markets for crops treated with Nufarm Sentry®. If you are growing produce for export, please check with Nufarm Australia Limited or your Industry Association for the latest information on MRLs and import tolerances before using Nufarm Sentry®.

GENERAL INSTRUCTIONS

Nufarm Sentry® is for use pre-emergence in imidazolinone herbicide tolerant barley, wheat oat and canola varieties only and for use post emergence in imidazolinone herbicide tolerant canola and wheat (single gene) varieties. Varieties with imidazolinone herbicide tolerance are those which have been bred specifically to tolerate imidazolinone herbicides. Nufarm Sentry® is an imidazolinone herbicide combination that provides both soil and foliar activity. Soil activity can be variable when dry soil conditions prevail following application because the product is not freely available to weeds. Best pre-emergence weed control will be achieved when soil moisture at or soon after application is good allowing the roots of germinating weed seedlings to take up the herbicide. Best post-emergence weed control will be achieved when application is made to young actively growing weeds. The pre-emergence use of Nufarm Sentry® will assist crop establishment and protect yield potential but may not preclude follow-up post-emergence weed control (in imidazolinone herbicide tolerant canola and wheat, single gene, only). When applied post-emergence Nufarm Sentry® must be mixed with Nufarm CanDo™ Adjuvant* as per the Directions for Use. Weeds treated post-emergence will either die or will remain stunted and will not compete with the crop. Vigorous crop growth will assist in suppressing weeds not completely killed or germinating later.

Crop Safety

This product may, in some circumstances, lead to transient crop yellowing and temporary slowing of growth but plants soon recover and yield is unaffected. This effect may be more pronounced when the product is used under poor growth conditions. **DO NOT** use this product on any canola or cereal varieties other than imidazolinone herbicide tolerant varieties. Severe crop damage will result to any other canola or cereal varieties. Spray overlapping can heighten crop injury and can be severe in CL STL and CL JNZ wheat varieties.

Mixing
Nufarm Sentry® is a water dispersible granule formulation. Part-fill the spray tank with water, then with agitator running, add the required amount of product. **Once completely dissolved** add the required quantities of recommended products, in the correct mixing order, whilst filling the remainder of the tank. For large tank loads requiring large amounts of Nufarm Sentry®, or in situations where agitation is insufficient, consider pre-mixing Nufarm Sentry® and then adding the pre-mix to the spray tank.

Application
Generally, apply in a minimum 70 L water per hectare. For post-emergence situations where grass weed populations exceed 200 plants per m^2, apply in a minimum of 100 L water per hectare. When used alone post-emergence, Nufarm Sentry® should be applied a minimum of two hours before rainfall or overhead irrigation. In mixtures with Nufarm Archer® 750 or a 240 g/L clethodim formulation, application should be made a minimum of three or six hours (as per advice for these products) before rainfall or overhead irrigation.

Equipment Clean-up
Thoroughly flush all spray equipment with water following the use of Nufarm Sentry® and before use with other products. After Nufarm Sentry® has been tank mixed with other products ensure that all spray equipment is additionally cleaned according to recommendations for the tank mix partners before using on sensitive crops.

Compatibility
Nufarm Sentry® is compatible with the herbicides, insecticides and adjuvants as listed in the following table. Other simultaneous mixes have not been tested.

HERBICIDES	INSECTICIDES	ADJUVANTS#
Nufarm Archer® 750 (clopyralid)	100 g/L alpha-cypermethrin formulations	Nufarm CanDo™ Adjuvant
Nufarm Avadex® Xtra (triallate)	Dimethoate	Hasten*
960 g/L S-metolachlor formulations	Le-Mat* (omethoate)	Kwickin*
Nufarm CRUCIAL® Herbicide		
900 g/kg diuron formulations		
Nufarm Factor® (butroxydim)		
Nufarm Glyphosate 450 Herbicide		
240 or 360 g/L clethodim formulations		
Spray-Seed* (diquat + paraquat)		
Nufarm Pendimethalin 440 Herbicide		
Nufarm Pyroxasulfone 850WG Herbicide		
Sakura* 850WG Herbicide (pyroxasulfone)		
Nufarm Shirquat® Herbicide		
Nufarm MCPA LVE 570 Herbicide		
Nufarm TriflurX® (trifluralin)		
Nufarm weedmaster® DST (glyphosate)		
Nufarm weedmaster® ARGO (glyphosate)		

When applying Nufarm Sentry® pre-emergence tank mixed with alternate mode of action pre-emergent herbicides, the following options are suggested for use in barley, canola and wheat based on weed profile and conditions:

i) Nufarm TriflurX® at up to 3 L/ha (**DO NOT** use on Oat above 2 L/ha),
ii) Nufarm Avadex® Xtra at 1.6 L/ha for wild oats control (**DO NOT** use on Oat) or
iii) Nufarm TriflurX® at up to 2 L/ha plus Nufarm Avadex® Xtra at up to 3.2 L/ha (**DO NOT** use on Oat)
iv) Sakura* at 118 g/ha (**DO NOT** use on Barley, Canola or Oat)

When applying Nufarm Sentry® pre-emergence on Imidazolinone herbicide tolerant Oat, the following options are suggested for use in conjunction with Nufarm Sentry® based on weed profile and conditions:

i) Nufarm TriflurX® at up to 1.5 - 2 L/ha in min-till/no-till pre-sowing or IBS (refer to Nufarm TriflurX® label for instructions)
ii) 960 g/L S-metolachlor formulations at 0.375 to 0.5 L/ha applied post-sowing pre-emergence,
iii) 900 g/kg diuron formulations at 275 to 550 g/ha (WA only), pre or post plant application (refer to label for instructions)

Tank mixes with grass herbicides are recommended as a sound herbicide resistance management strategy but are encouraged when a high grass weed burden is expected or when Group 2 resistance is suspected. These mixes can also assist broadleaf weed control. Tank mixtures of Nufarm Sentry® and these various grass herbicides can be applied with the knockdown herbicides and insecticides indicated but more complex mixes have not been tested.
When applying Nufarm Sentry® post-emergence tank mixed with Nufarm Archer® 750 or a 240 g/L clethodim formulation use 60 – 80 mL/ha Nufarm Archer® 750 (depending on legume and composite weed burden) and 175 – 500 mL/ha the 240 g/L clethodim formulation (depending on grass burden and Group 2 resistance status). When applying Nufarm Sentry® with a mix of Nufarm Archer® 750 and a 240 g/L clethodim formulation, note that Nufarm Archer® 750 can impact on grass control and should not be used at 80 mL/ha with low rates of a 240 g/L clethodim formulation.
DO NOT tank mix with foliar fertilisers.

Best Management Practice (BMP) Program

Sound agronomic practices including the practice of integrated weed management and Harvest Weed Seed Set Control (HWSSC) will optimise the performance of Imidazolinone herbicide tolerant crops and minimise the potential for the development of Group 2 herbicide resistance in weed populations. Whether using Nufarm Sentry® pre-emergence or post-emergence, or both pre- and post-emergence, in a crop, or whether integrating its use with another Group 2 herbicide, use herbicides with alternative mode of action, either in tank mix or sequentially, to alleviate selective pressures. Where two Group 2 treatments have been applied to a crop ensure HWSSC techniques are practiced. Avoid allowing surviving weeds to set seed. Consult a Nufarm representative in regard to BMP prior to using Nufarm Sentry®, especially when growing an Imidazolinone herbicide tolerant crop for the first time or when paddocks require an imidazolinone or Group 2 herbicide treatment following the use of Nufarm Sentry® pre-emergence. Implementation of the BMP is an essential part of herbicide resistance management.

Follow Crops

This product is broken down in the soil by microbes in wet, aerobic conditions. Under conditions which **DO NOT** favour breakdown, such as impoverished soils low in organic matter, non-wetting sands, anaerobic situations such as waterlogging and prolonged dry periods, soil residues will persist longer and may affect susceptible follow crops. Otherwise normally safe residue levels may still affect follow crops when soil nutrition is low or marginal, or when drought conditions or cold and very wet soil conditions prevail, or when soil pathogens or nematodes are present as these situations will add stress to the crops. As environmental and agronomic factors make it impossible to eliminate all risks associated with this product, rotational crop injury is always possible. Consult your local Nufarm representative for advice should you have any concerns.

Consult the table below for minimum re-cropping intervals for a variety of commonly sown crops. **Where Nufarm Sentry® is used pre-emergence and followed with a post-emergence treatment of another imidazolinone-based herbicide, or any other Group 2 herbicide, only sow an imidazolinone herbicide tolerant crop the following season.** Additional advice is provided below the table in regard to growing conventional (non-imidazolinone tolerant) cereals the winter after using Nufarm Sentry®. When planning to plant legumes the winter after using Nufarm Sentry®, advice should be sought from a Nufarm representative if rainfall is below 150 mm prior to autumn. In calculating rainfall, place greater emphasis on rain received from application up to the end of spring and lesser emphasis on break rains. If rainfall from application to the end of spring is less than 200 mm and if single isolated heavy summer and autumn falls and break rains are required to achieve rainfall targets, it may not be safe to sow non-Imidazolinone herbicide tolerant cereals within 10 months of application. Consult your local Nufarm representative for advice.

Note: When the intention is to grow cereals on Imidazolinone herbicide tolerant canola stubble (treated with Nufarm Sentry®) self-sown canola volunteers must be removed before they mature beyond 2-leaf, all macro and micro-nutrients must be maintained at levels necessary to grow the planned crops, and sulfonylureas must not be used.

The following minimum re-cropping intervals (months after application) should be observed.

Nufarm Sentry® rate used	Replant interval – Months after application	Follow Crops
20, 40, 50 and 55 g/ha	0	Imidazolinone herbicide tolerant crops only
20 and 40 g/ha	8	Chickpeas, Faba beans, Field peas, Lucerne, Lupins, Pasture legumes, vetch, **Oats[Ψ], Triticale[Ψ], Barley[Ψ], Wheat[Ψ]**
20 and 40 g/ha	22	Safflower
50 and 55 g/ha	8	Chickpeas, Faba beans, Field peas, **Triticale[Ψ], Barley[Ψ], Wheat[Ψ]**
50 and 55 g/ha	22	Lucerne, Lupins, Oats, Pasture legumes, Safflower, Vetch
20, 40, 50 and 55 g/ha	34	All other crops including conventional, TT and RR canola varieties

[Ψ] The following additional requirements apply if it is intended to sow conventional (non-imidazolinone herbicide tolerant) cereals the Winter season following the use of Nufarm Sentry®:

- **DO NOT** use Nufarm Sentry® at rates above 20 g/ha in areas where rainfall from spraying to sowing non-imidazolinone herbicide tolerant cereals is expected to be below 150 mm.
- **DO NOT** use Nufarm Sentry® at 40 g/ha or higher in areas where rainfall from spraying to sowing non-imidazolinone herbicide tolerant cereals is expected to be below 250 mm.
- **DO NOT** use Nufarm Sentry® at 50 – 55 g/ha in areas where rainfall from spraying to sowing non-imidazolinone herbicide tolerant cereals is expected to be below 350 mm.
- **DO NOT** apply Nufarm Sentry® later than the end of August (end of July in Western Australia) unless it is intended to grow an imidazolinone herbicide tolerant crop the following year.
- **DO NOT** use Nufarm Sentry® at 40-55 g/ha on shallow, strongly duplex soils unless Imidazolinone herbicide tolerant crops are sown the following year.
- If Nufarm Sentry® is applied pre-emergence followed by a post-emergent application of another Imidazolinone herbicide, sow only an Imidazolinone herbicide tolerant crop the following year regardless of the total load of Nufarm Sentry® applied.
- **DO NOT** use Nufarm Sentry® above 20 g/ha in the lower Great Southern area of Western Australia, nor rates above 40 g/ha anywhere in Western Australia unless it is intended to grow an Imidazolinone herbicide tolerant crop the following year.
- **If expected rainfall is not received following use of Nufarm Sentry®, or the rainfall pattern is inappropriate, consult a Nufarm representative before planting conventional (non-imidazolinone herbicide tolerant) cereal crops**.

RESISTANT WEEDS WARNING

GROUP 2 HERBICIDE

Nufarm Sentry® Herbicide is a member of the Imidazolinone group of herbicides. Nufarm Sentry® has the inhibitors-of-acetolactate-synthase (ALS) mode of action. For weed resistance management, Nufarm Sentry® is a Group 2 herbicide. Some naturally occurring weed biotypes resistant to Nufarm Sentry® and other Group 2 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Nufarm Sentry® or other Group 2 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from failure of Nufarm Sentry® to control resistant weeds.

RE-ENTRY PERIOD

For Nufarm Sentry® used alone - not required when used as directed.

For Nufarm Sentry® PLUS Nufarm Archer® 750 or a 240 g/L clethodim formulation - when tank-mixing Nufarm Sentry® with Nufarm Archer® 750 or a 240 g/L clethodim formulation refer to the labels of these products for re-entry advice. When tank-mixing with MCPA, **DO NOT** allow entry into treated areas for 24 hours. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist and a washable hat, goggles and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT allow spray overlapping as crop injury may occur.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container, in a dry, cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. **DO NOT** burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray, wear elbow-length chemical resistant gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After each day's use, wash gloves and face shield or goggles. Wash hands after use. When tank-mixing with other products, also consult the safety directions for those products.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

ADDITIONAL STATEMENTS (WHS REGULATIONS 2011)

Causes serious eye damage. Wear protective gloves, clothing, eye and face protection. Wash hands and exposed skin thoroughly after handling. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. IF ON SKIN: Wash with plenty of soap and water. If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash before reuse.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained by scanning the QR code, from your supplier or the Nufarm website.

In case of emergency: Phone 1800 033 498 (24 hrs) and ask for shift supervisor.

PRODUCT STEWARDSHIP INFORMATION AND TOOLS

nufarm.com.au/spraywise is a website which contains resources on spray drift management, recording keeping and application technologies.
spraywisedecisions.com.au is an online weather forecasting program and is recommended for use when planning your pesticide application.
When spraying in or near areas with sensitive crops such as cotton, check online at **satacrop.com.au** for the proximity of sensitive crops mapped by growers.
stewardshipfirst.com.au is a website with CropLife Australia's suite of world-leading product stewardship initiatives, programs and best-practice guides.

CONDITIONS OF SALE

Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute.

® All trade marks owned or used under licence by Nufarm Australia Limited.
* Other registered trade marks

Nufarm AUSTRALIAN THROUGH & THROUGH

nufarm.com.au

NUFARM AUSTRALIA LIMITED
103-105 Pipe Rd, Laverton North VIC 3026
For product advice, or to learn more about complementary Nufarm solutions, reach out to your local Nufarm specialist or call **1800 NUFARM (1800 683 267)**.

APVMA Approval No.: 67951/142681 Date: 05-02-2024