POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm CONQUEROR®

ACTIVE CONSTITUENTS: 100 g/L PICLORAM
present as the hexyloxypropylamine salt
300 g/L TRICLOPYR present as the butoxyethyl ester

HERBICIDE

GROUP 4 HERBICIDE

For the control of environmental and noxious woody and herbaceous weeds as specified in the DIRECTIONS FOR USE.

EC EMULSIFIABLE CONCENTRATE

Nufarm

AUSTRALIAN THROUGH & THROUGH

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition, presence of disease, damage or previous herbicide treatment, as reduced levels of control may result.
DO NOT spray if rain is likely within one hour or if foliage is wet from rain or dew.
DO NOT burn off, cut or clear blackberry or other woody weeds for at least six months after spraying.
DO NOT apply by aerial application at air temperatures above 35°C. In areas prone to flooding, treatment should commence after any annual flooding as such areas flooded within 9 months following application may have reduced results.

Spray Drift Restraints

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.
DO NOT allow bystanders to come into contact with the spray cloud.
DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.
DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.
DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings 1 to 2 hours before sunset and persist until 1 to 2 hours after sunrise.

1. WOODY WEED SITUATIONS

Table A: High Volume Spraying

See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS			
WEEDS CONTROLLED	**WEEDS GROWTH STAGE**	**RATE**	**CRITICAL COMMENTS**
African boxthorn	Less than 2m tall	500 mL/100L	Apply when bushes have good leaf cover, growth and no leaf fall.
Angophora spp.	1 - 3m tall	350 mL/100L	
Australian blackthorn	Less than 2m tall	500 mL/100L	Apply from late spring to early autumn.
Banksia spp.	1 - 3m tall	350 mL/100L	
Biddy bush (Chinese shrub), (Sifton bush)	Autumn when actively growing	500 mL/100L	Add Nufarm BS1000® at 125 mL/100L of water for best results.
Blackberry in association with: Docks, Ragwort, Smartweed Thistles	Late spring to autumn	350 or 500 mL/100L	Use the higher rate on plants that have been damaged by grazing stock or insects and on known difficult to kill blackberry. Where herbicides other than Group 4 herbicides have been used, allow two seasons regrowth to occur before respraying with Nufarm Conqueror®.
Blue heliotrope	Flowering	500 mL/100L	Apply in a minimum spray volume of 1250 L/ha.
Brooms: Cape, English, Flax leaf, Montpellier	Spring to mid-summer prior to pod formation	250 mL/100L	Apply as a thorough foliage spray.
	Autumn to winter	350 mL/100L	
Camphor laurel	Less than 2m tall		
	Above 2m tall	500 mL/100L	
Casuarina spp.	1 - 3m tall	350 mL/100L	
Chinee apple	Less than 2m tall	350 mL/100L	Add Nufarm BS1000® at 100 mL/100L of water for best results.
Cockspur thorn	Spring to Autumn	350 mL/100L	Apply as thorough foliage spray.
Common sensitive plant	Any time when actively growing	200 mL/100L	To avoid leaves closing during application, spray plants while moving forward. Add Nufarm BS1000® at 100 mL/100L of water for best results.
Crofton weed	Spring to Autumn	350 mL/100L	Apply as thorough foliage spray.
Eastern cotton bush (*Maireana microphylla*)		500 mL/100L	Add Nufarm CanDo® Adjuvant at 500 mL/100L of water. Some bushes may require a follow up spray to control regrowth.
Eucalyptus spp.	Seedling and regrowth from small lignotubers, 1 - 3m tall	350 or 500 mL/100L	Apply the high rate where difficult to control species of Eucalyptus regrowth is present. Addition of an adjuvant may improve results - contact Nufarm Australia Limited for details.
Galenia	Fresh growth in spring to summer	500 mL/100L	Use 200L of spray mixture/ha.
Giant bramble	Spring to autumn		Penetration of thick clumps may be difficult and respraying may be necessary. Add Nufarm BS1000® at 100 mL/100L of water for best results.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS			
WEEDS CONTROLLED	**WEEDS GROWTH STAGE**	**RATE**	**CRITICAL COMMENTS**
Gorse	1 - 1.5m tall	250 mL/100L	Spring and Summer treatment only. Add Nufarm BS1000® at 100 mL/100L of water for best results.
	Over 1.5m tall or autumn treatment	350 mL/100L	
	Winter treatment	500 mL/100L	Brownout may not be complete until summer. Add Nufarm BS1000® at 100 mL/100L of water for best results.
Groundsel bush (*Baccharis halimifolia*)	1 - 1.5m tall in spring to summer	250 mL/100L	Apply as a thorough foliar spray.
	Over 1.5m tall or autumn treatment	350 mL/100L	
Green cestrum	Late spring to early autumn	500 mL/100L	One application may give satisfactory control. Any subsequent regrowth and seedlings must be resprayed at approximately 1 metre high.
Hawthorn	Less than 2m tall		Apply from late spring to early autumn.
Horehound	Pre- flowering	350 mL/100L	Apply as a thorough foliar spray.
Japanese sunflower			
Lantana (*Lantana camara*) (*Lantana montevidensis*)	Up to 1m tall in summer to autumn		Add one of the following adjuvants, when using 350 mL rate: Nufarm CanDo™ Adjuvant at 0.5% v/v, Nufarm Pulse® Penetrant at 0.1% v/v.
	1 - 2 m tall in summer to autumn	500 or 750 mL/100L	Thoroughly wet foliage, stems and soil around the base of the plants. Use higher rate on known harder to kill varieties.
Lion Tail (*Leonotis nepetifolia*)	Pre-flowering	200 mL/100L	Apply as a thorough foliar spray. Add Nufarm BS1000® at 100 mL/100Lof water for best results.
Limebush	Any time of the year with good leaf cover and soil moisture	350 mL/100L	Penetration of thick clumps may be difficult and respraying may be required. Addition of an adjuvant may improve results - contact Nufarm Australia Limited for details.
Manuka	At flowering	500 mL/100L	For optimum results, add Nufarm Pulse* Penetrant at 200 mL/100 L of spray. Thoroughly wet foliage, stems and soil around the base of the plants.
Mesquite (*Prosopis* spp.) *Prosopis velutina*	Seedling, full leaf and flowering before podding	350 mL/100L	**DO NOT** spray plants bearing pods. Add Nufarm BS1000® at 100 mL/100L of water for best results.
		670 mL/100L	
Mistflower	Spring to autumn	350 mL/100L	Apply as a thorough foliar spray.
Mother-of-millions	Flowering	500 mL/100L	Add Nufarm BS1000® at 100 mL/100L of water.
Paddy's lucerne	Active growth		Plants that have been continually slashed or grazed over many seasons may be difficult to control and regrowth may occur.
Parkinsonia	Under 2m tall	350 mL/100L	Add Nufarm CanDo™ Adjuvant at 500 mL/100Lwater. Avoid spraying under dry conditions when plants are stressed or bearing pods. Thoroughly wet foliage.
Prickly pear (common) Smooth tree pear	Active phyllode growth	500 mL/100L	Apply as a thorough foliage spray. Regrowth may occur, so a follow-up application may be necessary.
Rubber vine (Not infected with rust)	Up to 1.5m tall at flowering	350 mL/100L	Spray all leaves and stems just to the point of runoff and thoroughly spray the base of the plant. With larger, more dense stands regrowth may appear. Subsequent control of any regrowth should be done by basal bark spraying.
	Dense stands greater than 1.5m tall at flowering	500 mL/100L	
Siam weed	Active growth	350 mL/100L	Add Nufarm BS1000® at 100 mL/100 L of water for best results.
Sicklepod	Up to flowering	200 mL/100L	**DO NOT** apply to podding plants. Add Nufarm BS1000® at 100 mL/100L of water for best results.
St. John's wort	From flowering to early seed set	500 mL/100L	Late spring to early summer.
Sweet briar	Up to 1.5m tall	350 mL/100L	Add Nufarm Associate® at 10 g/100L water to obtain more reliable results with the lower rate of Nufarm Conqueror®. Full leaf to ripe fruit prior to leaf fall. Thorough wetting including the crown is recommended.
		500 mL/100L	
Tobacco weed	Actively growing plants	300 mL/100L	Add Nufarm BS1000® at 100 mL/100L of water for best results.
Wattle (*Acacia* spp.) (except corkwood wattle	1 - 3m tall	350 mL/100L	
Wild rosemary (*Cassinia laevis*)	Active growth, 0.5 - 1m tall	350 – 500 mL/100L	Use lower rate on seedlings 0.5 m tall. Apply as a thorough foliar spray.
Wild tobacco tree	Spring to autumn up to 2m tall	350 mL/100L	

Table B: Aerial Application
See GENERAL INSTRUCTIONS - APPLICATION section for application method details

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS			
WEEDS CONTROLLED	**WEED GROWTH STAGE**	**RATE/ha**	**CRITICAL COMMENTS**
Blackberry	Summer to autumn	10 L/ha	Where herbicides other than Group 4 herbicides have been used, allow two seasons regrowth to occur before re spraying with Nufarm Conqueror®. WARNING: *Eucalyptus* species up to 4 m may be killed if sprayed during this treatment. Mature trees which are 15 - 20 m tall may be partially defoliated but are likely to recover.
Gorse			Apply by helicopter only in TAS.
Cockspur thorn, Crofton weed, Lantana, Mistflower	Late autumn	1.5 L/ha + 6 L/ha Nufarm 2,4-D Amine 625	Spray with calibrated equipment using the half overlap opposite pass technique applying a minimum spray volume of 150 L/ha. Follow-up respraying will be required. Apply by helicopter only in NSW, QLD and NT.
Lantana		10 L/ha	Helicopter application only.
Rubber vine (Not infected with rust)	When flowering	3 – 5 L/ha	Use rates will depend upon the density and height of the rubber vine stand. The higher rate should be used on dense stands. However, complete coverage and penetration may be difficult. Follow-up respraying will be required. Any regrowth should be sprayed with a suitable basal bark herbicide. Apply by helicopter only in QLD and NT
St. John's wort	Flowering to early seed set (Nov-Jan)	4 L/ha	Apply by helicopter only in NSW. Follow up spraying will be required in the following season.
AGRICULTURAL NON-CROP AREAS ON FLOOD PLAINS			
WEEDS CONTROLLED	**WEED GROWTH STAGE**	**RATE /ha**	**CRITICAL COMMENTS**
Parkinsonia	Seedlings 1 - 2m tall, or 12 – 24 months old	3 L/ha	Add Nufarm CanDo™ Adjuvant at 1 L/ha. Apply by helicopter only in QLD and NT.

Table C: Controlled Droplet Application (C.D.A.)
See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS			
WEEDS CONTROLLED	**WEED GROWTH STAGE**	**RATE**	**CRITICAL COMMENTS**
Blackberry in association with: Docks, Ragwort, St. John's wort, Thistles	Summer to autumn	Apply undiluted	One application may give satisfactory control but subsequent regrowth and seedlings should be resprayed after hardening off. Where herbicides other than Group 4 herbicides have been used, allow two seasons regrowth to occur before respraying with Nufarm Conqueror®.

Table D: Low Volume High Concentrate Application Techniques (GAS POWERED GUN, SPRINKLER SPRAYER)
See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS			
WEEDS CONTROLLED	**WEED GROWTH STAGE**	**RATE**	**CRITICAL COMMENTS**
Blackberry	Late Spring to autumn	335 mL/10L	Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high volume application technique is recommended.
Camphor laurel, Cockspur thorn, Crofton weed	Less than 1.5 m high	500 mL/10L	
Eucalyptus species	Seedlings up to 2 m tall	335 mL/10L	
Mistflower	Less than 1.5 m high	500 mL/10L	
Sweet briar	1.5 m tall, full leaf to ripe fruit	500 mL/10L	**Gas powered Gun only**: Apply to actively growing bushes not more than 1.5 m tall that have not more than 5 stems from the crown.
St. John's wort	During flowering to early seed set		**Gas powered Gun only:** One application should provide control. Minor regrowth and seedlings may be retreated the following summer.
Wild tobacco tree	Less than 1.5 m high		Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high volume application technique is recommended.

Table E: Boom Application
See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHT-OF-WAYS			
WEEDS CONTROLLED	**WEED GROWTH STAGE**	**RATE**	**CRITICAL COMMENTS**
Galenia	Fresh growth during spring to summer	5 L/ha	Rough mine sites will require adequate spray equipment such as boomless nozzles for effective coverage.
Sicklepod	Up to flowering	3 L/ha	**DO NOT** apply to podding plants. Add Nufarm BS1000® at 100 mL/100L of water.
St. John's wort	Flowering to early seed set (Nov-Jan)	2 – 4 L/ha	Use the higher rate on dense infestations and when longer residual control is required. Follow-up respraying will be required in the following season.

2. FALLOW SITUATIONS
Table F: Boom Application
See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

FALLOW			
WEEDS CONTROLLED	**WEED GROWTH STAGE**	**RATE**	**CRITICAL COMMENTS**
Blackberry nightshade- Suppression only	10 to 25 cm tall prior to flowering	200 – 400 mL/ha + 1.2 L Nufarm Glyphosate 450 + Nufarm Collide® 700	FOR USE BY GROUND EQUIPMENT ONLY. Plants must be actively growing. Use the lower rate on the smaller weeds, as specified in the weed growth stage (or up to 5 cm diameter for *Polymeria pusilla*). Refer to mixture product label for use of Nufarm Collide® 700. **DO NOT** plant susceptible crops for up to nine months after application, as specified in General Instructions - Minimum Recropping Periods - Black Cracking Clay Soils, Northern NSW & Qld. Dry conditions after application will increase the recropping interval.
Camel melon, Prickly paddy melon, Cucumber melon (*Cucumis melo*)	From 2 leaf to 50 cm diameter		
Common sowthistle	From 8 leaf to flowering		
Cow vine	From 2 - 5 leaf up to 15 cm diameter, prior to flowering		
Lucerne (established)	Active growth, 15 to 25 cm high, during spring	300 - 500 mL/ha + 1.2 L Nufarm Glyphosate 450 + Nufarm Collide® 700	
Polymeria pusilla	2 - 12 leaf up to 20 cm diameter, prior to flowering	200 – 400 mL/ha + 1.2 L Nufarm Glyphosate 450	

Table G: Blanket Wiper Application
See GENERAL INSTRUCTIONS - APPLICATION section for application method details.

FALLOW			
WEEDS CONTROLLED	**WEED GROWTH STAGE**	**RATE**	**CRITICAL COMMENTS**
Bitterbark (*Alstonia constricta*)	From summer to end of autumn	1:4 (1 part Nufarm Conqueror® to 4 parts water) 2% solution for spot spray (e.g. 100 mL Nufarm Conqueror® in 5L water)	For use with blanket wipers only. For best results apply in autumn to tall (>60 cm) plants using two opposite directional passes (up and back). Follow up "missed" plants with a spot spray application. These will be obvious after 6 weeks. Blanket wiper application can be made in summer when plants are smaller but follow up spot spraying may be necessary. **DO NOT** disturb (cultivate) the treated patches for at least 3 months. Best long term control is achieved when patches are left undisturbed for as long as possible after treatment (at least 6 months). Spot spraying "missed" plants: thoroughly wet all stems and leaves without producing any solution run off. Avoid any spray reaching the soil surface.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.
IN TASMANIA FOR BLACKBERRY: DO NOT treat bushes carrying mature or near mature fruit.
FOR NATIVE VEGETATION: Use of Nufarm Conqueror® on native vegetation must be done in accordance with STATE and/or LOCAL legislation.

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS
Fallow Situations
***Minimum Recropping Intervals* - Black Cracking Clay Soils, Northern NSW & Qld.**

Table F: Boom Application

Plant-back periods for crops following the application of Nufarm Conqueror® for rates up to 600 mL/ha				
RATE mL/ha	**200**	**300**	**400**	**600**
CROP	**MONTHS**			
Wheat	2	2	4	4
Barley	2	2	4	4
Canola	2	4	4	4
Faba Beans	4	4	6	6
Chickpeas	4	6	6	6
Lucerne	6	9	9	9

These plant back periods are based on a normal rainfall pattern. During drought conditions (or when the rainfall is less than 100 mm for a period of 4 months or greater) the plant back period may be significantly longer.

Table G: Blanket Wiper Application

Plant-back periods for crops following blanket wiper application.	
CROP	**MONTHS**
Broadleaf crops	18
Lucerne	6 This will allow any potential soil residues to dissipate, if any, and allow effective control.

NOTE: Before using Nufarm Conqueror® in tank mixes with other herbicides, check the plant-back information on all product labels. The most residual product with the longest plant-back period, will determine the time between spraying and planting.

Mixing
Mix only with water. Half fill the spray unit with water, and add the required amount of Nufarm Conqueror®. Add the remaining water with the agitator running. If required, then add spray oils or wetters (surfactants). Maintain mechanical or by-pass agitation in the spray tank during spraying. Only mix sufficient solution for immediate daily use and avoid storing.

Application
1. Woody Weed Situations
Weeds need to be actively growing for herbicides to have optimum effect. Delay treatment until all regrowth has had time to grow approximately 1 metre in situations which have been bulldozed, slashed, burnt, ploughed or areas having a previous chemical treatment.

Table A. High Volume Spraying
Thorough coverage of foliage to the point of run-off is essential, however, avoid excess spraying which is wasteful of chemical.
Hand Gun: Apply the recommended mix to give full coverage of leaves and stems through a No. 6- 8 tip at 700 – 1500 kPa (400 – 500 kPa for St John's wort). A spray volume of 3000 – 4000 L per infested hectare of 1 - 2 metre high blackberry (30 – 40 L/100m^2) should be used. Use 2000 L of spray mixture/ha of galenia infestation (i.e. 20 L/100m^2 infested area).
Knapsack: Apply the recommended spray mix to give full coverage of leaves and stems. The final volume of application should be similar to hand gun. A spray volume of 3 – 4 L/10m^2 infested area should be used. A spray volume of 2L/10m^2 should be used for an area infested with galenia.

Table B. Aerial Application
Apply in 200 L of water/ha using an aircraft to apply 100 L per pass on a double overlap pattern using nozzle configurations to produce droplets of 250 - 350 micron diameter. The potential for damage from drift can be greatly reduced by avoiding unsuitable spraying conditions and using spray pressure and nozzles to minimise the production of small droplets. **DO NOT** spray when wind exceeds 15 km/hr and /or air temperature reaches 35°C.

Table C. Controlled Droplet Application (C.D.A)
Results similar to high volume spraying can be obtained using Micron Herbi* or similar equipment. Select a nozzle to give a flow rate of 2 mL/sec and sweeping action of approximately 1 m/sec to ensure a droplet density of 20/cm^2. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage. Also, consult directions provided with C.D.A unit.

Table D. Low Volume High Concentrate Application Techniques
Good control will be achieved, similar to high volume application, where bush size enables good coverage of entire bush. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage.
Gas powered gun: Apply 50 mL shots to obtain uniform coverage of 4 – 5 m^2 of surface area of bush. This relates to 20 droplets/cm^2 of leaf surface.
Sprinkler sprayer: This technique involves using a micro sprinkler that is disconnected to a hollow fibre glass rod attached to a pneumatic knapsack sprayer. Use at low pressures (50 – 200 kPa) and apply with a slow sweeping action over the top of the plants ensuring even coverage on the leaves.

Table E. Boom Application

Application in a minimum spray volume of 200 L/ha for galenia and St. John's wort and 600 L of water/ha for sickle pod. Flat fan nozzles are recommended, using pressure in the range of 200 – 300 kPa. Boom height must be set to ensure double overlap of nozzle patterns.

2. Fallow Situations

Table F. Boom Application

Application of Nufarm Conqueror® in a minimum spray volume of 50 L/ha is recommended. Flat fan nozzles are recommended, using pressure in the range of 200 – 300 kPa. Boom height must be set to ensure double overlap of nozzle patterns.

Table G. Blanket Wiper Application

Blanket needs to be made from durable and wettable material with a rigid backing. Blanket should be rigidly mounted behind motorised vehicle (eg. tractor, 4-wheel drive vehicle) and set low but never touching the ground. The chemical solution should be fed to the blanket at a flow rate sufficient to keep the blanket wet but not dripping. In thick patches the blanket may require more frequent solution recharge (rewetting). Ideally, a scraper bar should be mounted at the front of the blanket in order to scrape or damage the bark (but not sever the stems) prior to the blanket wiping stems. This scraper may be mounted at the front of the vehicle. Two passes (in opposite direction) with the blanket increases the contact with the plant. Ground speeds of 10 - 15 kph are ideal for blanket wiping application.

Cleaning Spray Equipment

Rinsing

After using Nufarm Conqueror® empty the spray unit completely and drain the whole system. Thoroughly wash inside the unit using a pressure hose. Drain the spray unit, and clean any filters in the tank, pump, lines, hoses and nozzles. After cleaning the spray unit as above, quarter fill with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

Decontamination

Before spraying cotton and other sensitive crops with equipment that has been used to apply Nufarm Conqueror®, see PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section. Wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent or appropriate tank and equipment cleaner and circulate throughout the system for at least 15 minutes. Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.

RESISTANT WEEDS WARNING

GROUP 4 HERBICIDE

Nufarm Conqueror® Herbicide is a member of the pyridines group of herbicides. Nufarm Conqueror® has the disruptors of plant cell growth mode of action. For weed resistance management, Nufarm Conqueror® is a Group 4 herbicide. Some naturally-occurring weed biotypes resistant to Nufarm Conqueror® and other Group 4 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Nufarm Conqueror® or other Group 4 herbicides. Strategies to minimise the risk of herbicide resistance are available.

COMPATIBILITY

Follow any regional restrictions, and all directions and restrictions on the label, for any chemical mixed with Nufarm Conqueror® (e.g Nufarm 2,4,-D Amine 625).

Nufarm Conqueror® is compatible with the following herbicides: Nufarm Amicide® Advance 700, Nufarm 2,4-D Amine 625, Nufarm Associate®, Nufarm Glyphosate 450, Nufarm weedmaster®ARGO®, Nufarm weedmaster® DST®.

Nufarm Conqueror® is compatible with the following adjuvants, as per Directions for Use: Nufarm CanDo™ Adjuvant, Nufarm BS1000®, Nufarm Collide® 700 and Nufarm Pulse® Penetrant.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Crops susceptible to Nufarm Conqueror® include, but are not limited to: peas, lupins, lucerne, navy beans, peanuts, soybeans and other legumes, cotton, flowers, fruit, hops, ornamentals, shade trees and *Pinus* spp., potatoes, safflower, sugar beet, sunflowers, tobacco, tomatoes, vegetables and vines. Nufarm Conqueror® is damaging to susceptible crops during both growing and dormant periods. Grasses are normally unaffected and establish quickly after treatment. Picloram, one of the active constituents in this product can remain active in the soil for extended periods depending on soil type and application rate, rainfall, temperature, humidity, soil moisture and soil organic matter.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands, pastures, waterways or native vegetation.

DO NOT allow physical spray drift onto waterways, native vegetation and susceptible crops.

DO NOT apply close to, or in areas, containing roots of desirable vegetation, where treated soil may be washed onto areas growing (or areas to be planted with) desirable plants.

DO NOT apply on sites where surface water from heavy rain can be expected to run off to areas containing, or to be planted with susceptible crops or plants.

DO NOT move soil which may have been treated to areas where desirable plants are to be grown.

PROTECTION OF LIVESTOCK

Poisonous plants may become more palatable after spraying and stock should be kept away from these plants until they have died down. Many plants remain poisonous after death, and stock should not be allowed access, as there is a likelihood that they may graze the dead material. Such material should be burnt if possible.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or watercourses with the chemical or used containers.

STORAGE AND DISPOSAL
Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.
Non-refillable containers
Triple-rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. **DO NOT** burn empty containers or product.
Returnable containers
Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.
Small Spill Management
Wear protective equipment (See Safety Directions). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal. (See Storage and Disposal section). If necessary, wash the spill area with an alkali detergent and water and absorb as above the wash liquid for disposal.

SAFETY DIRECTIONS
Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length chemical resistant gloves and face shield or goggles. If the product is in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID
If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

ADDITIONAL STATEMENTS (WHS REGULATIONS 2011)
Harmful if swallowed. May cause an allergic skin reaction. Causes serious eye irritation. Combustible liquid. Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No smoking. Avoid breathing spray. Wash skin thoroughly after handling. **DO NOT** eat, drink or smoke when using this product. Contaminated work clothing should not be allowed out of the workplace. Wear protective gloves, clothing, eye and face protection. IF SWALLOWED: call a POISON CENTRE/doctor if you feel unwell. Rinse mouth. IF ON SKIN: wash with plenty of water. Take off contaminated clothing and wash it before reuse. If skin irritation or rash occurs: Get medical advice/attention. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.

SAFETY DATA SHEET
For further information refer to the Safety Data Sheet (SDS), which can be obtained by scanning the QR code, from your supplier or the Nufarm website.

In case of emergency: Phone 1800 033 498 (24 hrs) and ask for shift supervisor.

PRODUCT STEWARDSHIP INFORMATION AND TOOLS
When spraying in or near areas with sensitive crops such as cotton, check online at **satacrop.com.au** for the proximity of sensitive crops mapped by growers.
stewardshipfirst.com.au is a website with CropLife Australia's suite of world-leading product stewardship initiatives, programs and best-practice guides.

CONDITIONS OF SALE
Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute.

® All trade marks owned or used under licence by Nufarm Australia Limited
* Other registered trade marks

Nufarm AUSTRALIAN THROUGH & THROUGH

nufarm.com.au

NUFARM AUSTRALIA LIMITED
103-105 Pipe Rd, Laverton North VIC 3026
For product advice, or to learn more about complementary Nufarm solutions,
reach out to your local Nufarm specialist or call **1800 NUFARM (1800 683 276)**.

APVMA Approval No.: 57783/145021 Date: 15-11-2024