POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm
MCPA Amine 750

HERBICIDE

ACTIVE CONSTITUENT: 750 g/L MCPA
present as the dimethylamine salt

GROUP 4 HERBICIDE

For the selective control of broadleaf weeds in cereals, linseed, oilseed poppies, pastures, rice, sugar cane and turf as per the DIRECTIONS FOR USE.

SL SOLUBLE CONCENTRATE

Nufarm

AUSTRALIAN THROUGH & THROUGH

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.
DO NOT apply if rain is likely within 6 hours.

Spray Drift Restraints

DO NOT apply by a vertical sprayer.
Specific definitions for terms used in this section of the label can be found at www.apvma.gov.au/spraydrift
DO NOT allow **bystanders** to come into contact with the spray cloud.
DO NOT apply in a manner that may cause an **unacceptable impact** to **native vegetation**, **agricultural crops**, **landscaped gardens** and **aquaculture production**, or cause contamination of plant or livestock commodities, outside the application site from **spray drift**. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.
DO NOT apply unless the **wind speed** is between 3 and 20 kilometres per hour at the **application site** during the time of application.
DO NOT apply if there are **hazardous surface temperature inversion** conditions present at the **application site** during the time of application. **Surface temperature inversion conditions** exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

1. FIELD CROPS

SITUATION & CROP	WEEDS	STATE	RATE	CRITICAL COMMENTS
Wheat, Barley, Oats, Triticale	Refer to Weed Table	All States	2 - 4 leaf (Z12 - 14): 460 mL/ha 5 leaf to before booting (Z15 - 39): 460 mL - 1.45 L/ha	Apply at the 2 - 4 leaf stage when the weeds marked # in the weed table have reached 2 - 6 leaf stage, otherwise wait until the crop is at the 5 leaf stage and use higher rates as listed in the weed table.
Wheat, Barley, Triticale, Cereal rye	Capeweed, Double gee, Erodium, Mustard, Rapistrum, Wild turnip, Wild radish	WA only	270 mL/ha plus 200 g/ha of a 900WG diuron formulation at CROP: 3 - 4 leaf stage. (Z13 - 14)	Apply by aircraft or boom with 20 - 100 L/ha water, when most weeds have germinated and are in 2 - 5 leaf stage. Crop and weeds should be dry at time of application. Some temporary yellowing of crop may occur after application. Undersown sub-clovers may be slightly retarded. **DO NOT** apply to undersown medics.
			330 mL/ha plus 220 g/ha a 900WG diuron formulation at CROP: 4 - 5 leaf stage. (Z14 - 15)	
Wheat, Barley, Triticale, Cereal rye	Amsinckia (Yellow burr), Capeweed, Charlock, Dead nettle, Double gee (Spiny emex, Three-cornered Jack), Erodium spp., Mustard, Turnip weed, White ironweed (Corn gromwell, Sheepweed), Wild radish, Wild turnip Ward's weed, Common ice plant, Pimpernel	NSW, ACT, Vic and SA.	330 mL/ha (230 mL/ha in SA) plus 280 g/ha a 900WG diuron formulation	Apply by boom spray or aircraft when the crop is at the 3 - 5 leaf stage (NSW and SA only) and 5 leaf stage (Vic only). Weeds must be in the 2 - 5 leaf stage. **DO NOT** apply to undersown medic or lucerne. **DO NOT** use on Durati, Kamilaroi, Kite, Olympic or Shortim wheat. **DO NOT** use on very light sandy soils.
	Soldier thistle	SA only	460 mL/ha plus 80 mL/ha Nufarm Archer® 750 Herbicide	Apply to crop at tillering stage after the crop has 5 leaves and before the crop commences jointing.
	Skeleton weed	SA, NSW, VIC and WA.	660 mL/ha plus 200 mL/ha Nufarm Archer® 750 Herbicide	Apply to rosettes from 5 - 15cm diameter when the crop is between the 5 leaf and late tillering growth stages. It is essential that all the Skeleton weed is emerged and minimum of 5cm diameter which is usually not before August. This rate will provide control of Skeleton weed until harvest and will substantially reduce seed numbers the following season.
Wheat, Barley, Oats, Triticale, Cereal rye (not before the 3 leaf cereal stage)	Volunteer canola (*Brassica napus*) including Roundup Ready* varieties and canola hybrids with the Optimum GLY® herbicide tolerance trait	All States	WEED STAGE: 4 leaf: 330 mL/ha plus 85 mL/ha Unity 240 EW	**DO NOT** apply the tank mix of Unity 240 EW + Nufarm MCPA Amine 750 before the three leaf stage of cereals. **DO NOT** apply mixtures containing Unity to cereals by Aircraft.
			WEED STAGE: 6 leaf: 330 mL/ha plus 100 mL/ha Unity 240 EW	

SITUATION & CROP	WEEDS CONTROLLED	STATE	WEED STAGE	RATE	CRITICAL COMMENTS
Field peas	Prickly lettuce *(Lactuca serriola)*	NSW, Vic, Tas, SA only	4 leaf	80 mL/ha plus 125 mL/ha diflufenican 500 g/L as an SC	**FIELD PEAS:** Apply early post-emergence after the third node stage and before the start of flowering. **Warning:** Field peas grown on high pH soils in the presence of free lime may be less tolerant to diflufenican. **APPLICATION AND WEED CONTROL** Apply when weeds are actively growing. For optimum results apply 4 - 6 weeks post-sowing. Application beyond 8 weeks post-sowing may result in reduced levels of weed control. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as: • high crop and weed density, • late season germinations, • abnormal weed growth (including early flowering); higher rates of product (up to the maximum rate of application specified for that weed) may be required. Diflufenican will not effectively control: • regrowth of suppressed weeds, • transplanted weeds, • regrowth from rhizomes or roots, • weeds growing under stress from previous herbicide applications.
			6 leaf	100 mL/ha plus 150 mL/ha diflufenican 500 g/L as an SC	
	Charlock (Wild mustard) *(Sinapis arvensis)*		4 leaf	80 mL/ha plus 125 mL/ha diflufenican 500 g/L as an SC	
			6 leaf	100 mL/ha plus 150 mL/ha diflufenican 500 g/L as an SC	

SITUATION & CROP	WEEDS CONTROLLED	STATE	RATE	CRITICAL COMMENT
Linseed	Refer to Weed Table	Qld only	460 mL/ha - 1.45 L/ha	Apply when crop is 10 - 15 cm tall with at least 170 L/ha of water. **DO NOT** spray after buds appear. Some wilting effect may be seen following application but crop usually recovers quickly. **DO NOT** apply by aircraft.
		NSW, ACT, Vic, SA, Tas only	460 - 730 mL/ha	
Sugar Cane	Blue top, Chinese burr, Flannel weed, Gambia pea, Bell vine, Streaked rattle pod, Bindweed, Pink convolvulus, Cupid's flower, Merremia vine, Morning glory	Qld only	930 mL/ha	POST-EMERGENT: Apply as a directed inter-row spray. **DO NOT** exceed rates with phenoxy sensitive varieties.
	As above plus Fat hen, Noogoora burr		1.45 L/ha	
Oilseed Poppies *(Papaver somniferum)*	Fat hen (*Chenopodium album*), Spear thistle (*Cirsium vulgare*)	Tas only	1 L/ha	Use only in consultation with the poppy contracting company. Apply at least one week after petal fall after poppy seed has been set. Apply by aircraft in 30 - 40 L of water/ha. This treatment will control late germinating weeds if they are green and at the bud or early flowering stage. Spraying at this time will twist weeds below the crop canopy. **DO NOT** apply in horticultural areas where there is a risk of spray or vapour drift over susceptible crops.
Rice	Dirty Dora, Starfruit, Sedges, Buddah (Butter) pea, Docks	NSW, QLD and VIC.	485 - 970 mL/ha	Apply by aircraft at early/mid tillering stage of rice but not later than panicle initiation. Use 485 mL/ha at 35 days after aerial sowing (approx. 2 tillers per rice plant). Use 970 mL/ha at 45 days or more after cereal sowing (3 tillers or more per rice plant). The crop should be sprayed before flowering of the weeds. WATER MANAGEMENT: Before herbicide application, lower water level in the bays to at least two thirds of the weed growth for direct contact with Nufarm MCPA Amine 750. Weeds covered by water will not be controlled. Raise water depth to normal levels 24 hours following spraying.
Flax	Refer to Weed Table	Vic, Tas only	325 mL/ha	Apply to crop when 8 - 20 cm high.

2. PASTURES, SEED CROPS & TURF

SITUATION & CROP	WEEDS	STATE	RATE	CRITICAL COMMENT
Grass Pastures	Refer to Weed Table	All States	460 mL/ha - 2.7 L/ha	Apply to established grass based pastures. Damage may occur to legumes if present. See "Legume Tolerance" below. NOTE: Capeweed, Paterson's curse and Variegated thistle may be poisonous to stock after spraying.
	Soldier thistle	SA only	460 mL/ha - 1.45 L/ha	Use low rate on seedlings with no more than 2 true leaves. Use 100 L of water for best results. If clover present, **DO NOT** exceed 575 mL/ha.
	Seedling and established Horehound	NSW, ACT only	2 L/ha	Apply in Autumn when horehound plants are at the seedling stage. If sub clover or medic plants are present they will be damaged by this application.
Grass Seed Crops	Refer to Weed Table	Qld, SA, Tas, Vic, WA only	460 mL/ha - 2.7 L/ha	Apply to established grass seed crops.
Sub. clover seed crops		Vic, SA only	460 - 960 mL/ha	**DO NOT** exceed rate specified. Spray in Autumn and early Winter.
Pastures - Spray/Graze Technique	Caltrop, Capeweed, Charlock, Mustards, Paterson's curse, Shepherd's purse, Saffron, Slender, Spear or Variegated thistle, Turnip weed, Wild radish, Wild turnip	NSW, ACT, Vic only	230 - 930 mL/ha	Spray actively growing 6 - 8 week old weeds. Introduce stock 7 - 10 days after spraying, preferably sheep as cattle are less effective. Use stocking rate at least 5 times heavier than normal until weeds have been reduced but before survival of desirable pasture species is threatened. Lucerne and medics may be damaged and should be grazed short before spraying. Refer "Legume Tolerance" under General Instructions. PRECAUTION: An increased quantity of poisonous plants may be eaten by stock using Spray/Graze, eg. Caltrop, Capeweed, Paterson's curse and Variegated thistle and deaths could result from causes such as nitrate poisoning. With Paterson's curse preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. **DO NOT** graze horses or pigs on Paterson's curse. Observe Withholding Period of at least 7 days after spraying.
		Tas only	460 mL/ha	
	Refer to Weed Table	WA only	610 mL/ha - 1.25 L/ha	
	Capeweed, Dandelion, Seedling dock, Three-cornered Jack, Geranium, Erodium, Mustard, Annual thistles, Turnip weed, Wild radish, Wild turnip	SA only	330 mL/ha	
	Soldier Thistle		395 mL/ha	
Turf	Refer to Weed Table	All States	330 mL/ha or 2.8 mL/100 m^2	Newly laid - Delay until well established or at least two months old. Apply in high volume to actively growing weeds. See "Legume Tolerance". Buffalo grass is more easily damaged than other grasses.
			930 mL/ha - 1.8 L/ha or 10 - 19 mL/100 m^2	Established - Apply in high volume to actively growing weeds. **DO NOT** mow for 2 days before application. Some transitory damage may occur to fine turf grasses and strawberry clover. See "Legume Tolerance". Buffalo grass is more easily damaged than other grasses.

3. SPOT SPRAYING

SITUATION & CROP	STATE	MIXING RATES / CRITICAL COMMENTS
High Volume Spraying	All States	Use 130 mL/100 L of water for easy to kill weeds such as Wild turnip, Mustards and Wild radish. Use 270 mL/100 L of water for more difficult to control weeds such as Docks, Hoary cress and Bindweed.
Knapsack Application		Use 13 mL/10 L of water for easy to kill weeds such as Wild turnip, Mustards and Wild radish. Use 27 mL/10 L of water for more difficult to control weeds such as Docks, Hoary cress and Bindweed.

4. NON CROP AREAS, FALLOW LAND, INDUSTRIAL & COMMERCIAL AREAS, RIGHTS-OF-WAY

SITUATION & CROP	WEEDS	STATE	RATE	CRITICAL COMMENTS
Non Crop Areas, Fallow Land, Industrial & Commercial Areas, Rights of Way	Refer to Weed Table	All States	Refer to Weed Table	Apply at rates shown in Weed Table when weeds only are present and no crop or pasture is involved.

WEED TABLE

Where weeds are to be sprayed in a crop or pasture use only the rates given for particular crop or situation indicated under the Directions for Use table. In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed. The rates listed in the Weed Table below are for use where weeds only are present and no crop or pasture is involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage. This product should only be used in those states where a rate or range of rates is indicated for the particular weeds listed.

APPLICATION RATE PER HECTARE

#: refer to Critical Comments for Wheat, Barley, Oats, Triticale

WEEDS CONTROLLED	VIC	NSW, ACT	SA	TAS	QLD	WA	CRITICAL COMMENTS
Ball mustard	-	-	1.15 L	-	-	-	Apply at rosette stage.
Bathurst burr	1.35 L	660 mL - 1.35 L	1.35 L	1.35 L	1.35 L	1.35 L	Spray young seedlings only.
Black bindweed	-	970 mL - 1.35 L	-	-	930 mL - 1.35 L	-	Apply at young stage prior to flowering
California burr	-	660 mL - 1.35 L	-	1.45 L	-	-	Spray young seedlings only.
Cape tulip	-	-	-	-	-	1.35 L	Apply just prior to flowering. Gives mild suppression only.
Capeweed	-	1.45 L	1.45 L	1.45 L	1.45 L	-	Apply at seedling stage.
Carrot weed	1.35 L	-	-	-	1.35 L	-	Apply at seedling stage.
Charlock	460 mL	660 mL	460 mL	600 mL - 1.45 L	460 mL	-	Apply at rosette stage.
Cutleaf mignonette	-	-	1.45 L	-	-	-	Apply at young rosette stage.
Dandelion	1.35 L	1.35 L	1.35 L	1.35 L	1.35 L	1.35 L	Apply at young rosette stage.
Dead nettle	-	1.45 L	-	-	1.45 L	-	Moderately susceptible. Apply at seedling stage.
Docks	690 mL	-	1.45L	1.45 L	-	-	Apply at rosette stage and before flower heads emerge. Only the top growth of old established plants will be affected.
Fat hen	1.35 L	660 mL - 1.35 L	-	1.35 L	1.35 L	-	Spray up to rosette stage.
Fennel	1.8 L	1.8 L	1.8 L	1.8 L	1.8 L	-	Apply at young stage.
Field bindweed	1.35 L	-	1.35 L	1.35 L	1.35 L	1.35 L	Apply at bud stage, repeating over several seasons.
Field cress	-	-	-	1.45 L	-	-	Apply at young stage
Flatweed or Cat's-ear	-	-	-	1.45 L	-	930 mL	Apply when weeds are 10-15cm high.
Fumitory (red)	930 mL	930 mL	930 mL	-	930 mL	-	Apply at young stage prior to flowering.
Hoary cress	3 L	-	1.45 L	-	3 L	-	Apply before flowering
Horehound	-	2 L	1.45 L	-	-	-	Apply at rosette stage.
Hedge mustard#	730 mL	-	460 mL	460 mL	460 mL	825 mL	Apply at rosette stage.
Hexham scent or Melilotus	960 mL	800 mL	930 mL	-	930 mL	-	Apply at young seedling stage.
Lesser swinecress	-	-	-	1.45 L	-	-	Apply at young stage.
Lincoln weed	-	-	460 mL	-	-	-	Apply at rosette stage.
London rocket	-	-	-	-	-	825 mL	
Lupins	-	460 - 960 mL	-	-	-	825 mL	Spray up to 10cm high.
Mintweed	1.35 L	1.35 L	-	-	1.35 L	-	Spray young seedlings only.
Mustards#	730 mL	660 mL	730 mL	730 mL	-	825 mL	Spray up to rosette stage.
Noogoora burr	1.35 L	660 mL - 1.35 L	1.35 L	-	1.35 L	1.35 L	Spray young seedlings only.
Opium poppy	-	-	-	1.45 L	-	-	Apply at young stage.
Paterson's curse	-	660 - 960 mL	-	-	2.7 L	-	Apply early rosette stage.
Pimpernel	-	-	460 mL	-	-	-	Apply at rosette stage.
Plantains	-	1.35 L	-	1.35 L	1.35 L	-	Apply at young stage.
Rapeseed	-	660 - 960 mL	-	-	-	-	Spray up to rosette stage.
Rough poppy	-	460 - 960 mL	-	-	-	-	Spray up to rosette stage.
Safflower	-	460 - 660 mL	-	-	-	-	Spray up to rosette stage.
Shepherd's purse	-	-	-	1.45 L	-	-	Apply at rosette stage.
Skeleton weed	-	960 mL - 1.35 L	-	-	-	-	Spray rosettes before aerial growth commences.
Sorrel	-	-	1.45 L	-	-	-	Apply to rosette stage.
Stinkwort	1.35 L	660 - 960 mL	1.35 L	-	-	1.35 L	Apply when weeds are 10-15cm high.
Sunflower (seedlings)	-	690 mL - 1 L	-	-	-	-	Spray up to rosette stage.
Thistles: - Artichoke	1.05 L	-	1.45 L	-	1.35 L	460 mL	Apply at rosette stage.
- Nodding	-	1.65 L	-	-	-	-	Spray early rosettes. Repeat spray required in following Spring or Autumn.
- Saffron	960 mL - 1.65 L	660 mL - 1.35 L	930 mL - 1.65 L	930 mL - 1.65 L	930 mL - 1.65 L	930 mL - 1.65 L	Spray up to rosette stage. (September WA).
- Scotch or Cotton	1.45 L	-	460 - 930 mL	1.45 L	-	-	Apply at seedling stage.
- Slender	2 L	660 mL - 1.35 L	660 mL	2 L	-	2 L	Spray young seedlings only.
- Soldier	730 mL	-	730 mL	-	-	-	Spray young seedlings. Use higher rate for larger weeds.

WEEDS CONTROLLED	VIC	NSW, ACT	SA	TAS	QLD	WA	CRITICAL COMMENTS
- Spear	-	960 mL - 1.35 L	-	1.45 L	-	-	Apply at rosette stage.
- Star	-	-	730 - 930 mL	-	-	-	Apply at seedling stage. Use higher rate for larger weeds.
- Variegated	960 mL	660 mL - 1.35 L	460 - 930 mL	960 mL	960 mL	960 mL	Spray young seedlings only. Use 1.35L/ha (SA) for well developed weeds.
Turnip weed#	730 mL	660 mL	660 mL	-	-	-	Spray up to rosette stage.
Wild artichoke	-	-	1.45 L	-	-	-	Apply at rosette stage.
Wild radish#	930 mL	660 mL	930 mL	1.35 L	460 - 730 mL	800 mL	Spray up to rosette stage.
Wild turnip#	730 mL	660 mL	460 mL	1.35 L	460 - 730 mL	600 mL	Spray up to rosette stage.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

CEREALS PASTURE: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.
HARVEST: WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.
OILSEED POPPIES: DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavourable. The best conditions are when the soil is moist, weather fine and rain unlikely within 6 hours of application. **DO NOT** spray crops or weeds outside the stages indicated in "Critical Comments" as damage, loss of yield or inadequate weed control may result. **DO NOT** spray in high winds.

Application/Mixing Information

This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding the remaining water.
Boom spraying - Use 30 - 120 L water per hectare. - Use minimum of 170 L water per hectare for LINSEED.
Aerial spraying - Use 10 - 90 L water per hectare.

Equipment Maintenance and Usage

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants unless it has been cleaned with Tank & Equipment Cleaner or well washed out with a 2% solution of cloudy ammonia followed by several clear water rinses.

RESISTANT WEEDS WARNING

GROUP 4 HERBICIDE

Nufarm MCPA Amine 750 Herbicide is a member of the Phenoxy group of herbicides. Nufarm MCPA Amine 750 has the disruptors of plant cell growth mode of action. For weed resistance management Nufarm MCPA Amine 750 is a Group 4 herbicide. Some naturally-occurring weed biotypes resistant to Nufarm MCPA Amine 750 and other disruptors of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Nufarm MCPA Amine 750 or other disruptors of plant cell growth herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of Nufarm MCPA Amine 750 to control resistant weeds.

COMPATIBILITY

This product can be tank mixed with diflufenican 500g/L SC, Associate®, diuron 900 WG formulations, Nufarm Kamba® 750, Nufarm Bromicide 200, Nufarm Archer® 750 Herbicide, Revolver®, Igran 500 Flowable and Trooper®. The effectiveness of Imidan® and Dimethoate may be reduced by alkaline hydrolysis if tank mixed with this product. For these situations, the addition of COLLIDE 700® is recommended. When tank mixing, add the required amount of compatible product (and surfactant if recommended) when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying.

PRECAUTIONS

Re-entry period

Hand weeding should not be performed for 7 days after spray application unless workers wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), chemical resistant gloves and footwear.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT spray in high winds.

Drift Warning

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift onto susceptible crops such as cotton, tobacco, tomatoes, vines, fruit trees, vegetables, legume crops and pastures, oilseed crops and susceptible trees (eg. kurrajongs, belahs, eucalypts).

Legume Tolerance

The use of MCPA (Nufarm MCPA Amine 750) is preferred to 2,4-D Amine (Amicide) or Ester (Estercide) in situations where legumes are undersown in the crop.
NB: The tolerance of pasture legumes to applied product can vary with species, variety, growing conditions, stage of growth and companion crop cover. **DO NOT** spray cereal crops undersown with lucerne as damage may result.

PROTECTION OF HONEY BEES AND OTHER INSECT POLLINATORS

Low hazard to bees. May be applied at any time as recommended in the Directions for Use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT
DO NOT contaminate streams, rivers or watercourses with the chemical or used containers.

STORAGE AND DISPOSAL
Store in the closed, original container in a cool, well-ventilated area. **DO NOT** store for prolonged periods in direct sunlight.
Triple-rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. **DO NOT** burn empty containers or product.
Refillable containers
Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS
Harmful if swallowed. Attacks eyes. Will irritate the skin. When opening the container, preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product on skin, immediately wash area with soap and water. If product or spray in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID
If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

ADDITIONAL STATEMENTS (WHS REGULATIONS 2011)
DO NOT eat, drink or smoke when using this product. Wear protective gloves, protective clothing, eye and face protection. IF SWALLOWED: Rinse mouth. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. IF ON SKIN: Wash with plenty of soap and water. If skin irritation occurs: Get medical advice/attention.

SAFETY DATA SHEET
For further information refer to the Safety Data Sheet (SDS), which can be obtained by scanning the QR code, from your supplier or the Nufarm website.

In case of emergency: Phone 1800 033 498 (24 hrs) and ask for shift supervisor.

PRODUCT STEWARDSHIP INFORMATION AND TOOLS
nufarm.com.au/spraywise is a website which contains resources on spray drift management, recording keeping and application technologies.
spraywisedecisions.com.au is an online weather forecasting program and is recommended for use when planning your pesticide application.
When spraying in or near areas with sensitive crops such as cotton, check online at **satacrop.com.au** for the proximity of sensitive crops mapped by growers.
stewardshipfirst.com.au is a website with CropLife Australia's suite of world-leading product stewardship initiatives, programs and best-practice guides.

CONDITIONS OF SALE
Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute.

® All trade marks owned or used under licence by Nufarm Australia Limited
* Other registered trade marks

nufarm.com.au

NUFARM AUSTRALIA LIMITED
103-105 Pipe Rd, Laverton North VIC 3026
For product advice, or to learn more about complementary Nufarm solutions,
reach out to your local Nufarm specialist or call **1800 NUFARM (1800 683 267)**.

APVMA Approval No.: 60505/134311 Date: 05-04-2022