Product Name: NUFARM STRIKER HERBICIDE
APVMA Approval No: 56994/142772

RLP
APPROVED

Label Name:	NUFARM STRIKER HERBICIDE

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	ACTIVE CONSTITUENT: 240 g/L OXYFLUORFEN SOLVENTS: 626.4 g/L LIQUID HYDROCARBON 100 g/L N-METHYL-PYRROLIDONE

Mode of Action:	GROUP 14 HERBICIDE

Statement of Claims:	For the selective weed control of certain broadleaf and grass weeds as per the DIRECTIONS FOR USE.

Net Contents:	1 L - 1000 L

Restraints:	

Directions for Use:	This section contains file attachment.

Other Limitations:	

Withholding Periods:	WITHHOLDING PERIODS Grazing – DO NOT GRAZE TREATED WEEDS.

	Harvest – NOT REQUIRED WHEN USED AS DIRECTED.

Trade Advice:	

General Instructions:	GENERAL INSTRUCTIONS Nufarm Striker® is a selective herbicide for the control of certain annual grasses and broadleaf weeds in dormant apples, grapes, peaches, pears, plums, apricots, almonds, macadamia, dubosia, tobacco, coffee, pyrethrum, winter cereals, tropical/sub-tropical crops, brassicas, Eucalyptus and Pinus spp. and other forestry trees applied either to a weed free soil or to seedling weeds at the 4 to 6 true leaf stage. Nufarm Striker® applied to well-prepared weed free soil should not be disturbed or incorporated after application. Weed control for up to 6 months can be expected with high label rates, but spot treatment with knockdown herbicides of escape weeds and perennial grasses may be necessary. When Nufarm Striker® is applied to seedling weeds at the 4 to 6 true leaf stage a non-ionic surfactant such as Activator® Surfactant should be added at recommended rates to improve activity where suggested in the critical comments. Nufarm Striker® can also be utilised at low rates as a “spike” to improve the weed spectrum of knockdown herbicides. Duboisia seedlings and mature plants will tolerate “over the top” applications. Eucalypt and pine transplants and established trees can tolerate “over the top” applications. Timing For optimum residual weed control, Nufarm Striker® should be applied to the soil surface prior to weed emergence after all other agricultural operations have been completed, such as mechanical cultivation and reshaping of irrigation furrows. The area should be left undisturbed during the period of desired weed control. When applied to seedling weeds, they should be young and actively growing. Weed control for up to 6 months is expected but spot treatment, with knockdown herbicides, for escape weeds and perennial grasses may be necessary. Mixing Shake well before use. When using Nufarm Striker® alone, fill the spray tank, at least one-third full with clean water, add the recommended amount of Nufarm Striker® while the pump and agitator are running, then complete filling the spray tank. A non-ionic surfactant like Activator®, if required by label directions, should be added near the end of the filling process to minimise foaming. When tank mixing with Surflan* 500 or Devrinol*-C, add to one-third filled tank, then the Nufarm Striker® during the filling operation. Maintain agitation during mixing and until spraying is completed. When tank mixing with Nufarm weedmaster® DST, Nufarm weedmaster® DUO, Nufarm Glyphosate 450, 135 g/L paraquat + 115 g/L diquat, Spray•Seed* 250 or 250 g/L paraquat, add these after Nufarm Striker® during the filling operation. When using Nufarm Striker® in mixtures always refer to the appropriate label and read and follow all label directions. Application Nufarm Striker® should be applied as a ground spray directed at the base of the crop plants. The spray mixture should be applied evenly with a carefully calibrated, standard low pressure herbicide sprayer with a flat fan or hollow cone nozzle. When seedling weeds are present ensure thorough coverage of their foliage is required to obtain the maximum knockdown effect. Ensure both the weed foliage and the soil surface are sprayed. When applying to bare soil use 250 to 500 L of water/ha. For application to seedling weeds (4 to 6 leaf stage) use 250 to 1,350 L/ha. Use higher volumes for high weed density. Crop Safety Nufarm Striker® may be applied as directed around dormant peach, plum, apricot, almond, apple and pear trees and grape vines of all ages when applied at rates of less than 1 L/ha. When applied at 3 L/ha and above, the trees and grape vines should be at least 3 years of age. DO NOT apply Nufarm Striker® once bud swell has occurred when using rates greater than 1 L/ha. Duboisia seedlings and mature plants will tolerate "over-the-top" applications

	of Nufarm Striker®. When using the 75 mL/ha rate, Nufarm Striker® may be applied as directed prior to sowing winter cereals.

Resistance Warning:	RESISTANT WEEDS WARNING GROUP 14 HERBICIDE Nufarm Striker® is a member of the diphenyl ether group of herbicides. Nufarm Striker® has the inhibitor of protoporphyrinogen oxidase mode of action. For weed resistance management Nufarm Striker® is a Group 14 herbicide. Some naturally occurring weed biotypes resistant to Nufarm Striker® and other Group 14 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Nufarm Striker® or other Group 14 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of Nufarm Striker® to control resistant weeds.

Precautions:	

Protections:	PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby water bodies, susceptible plants/crops, cropping lands or pastures. PROTECTION OF LIVESTOCK Use with care when applying in areas frequented by stock. PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT This product is highly toxic to wildlife and fish. DO NOT contaminate streams, rivers or watercourses with the chemical or used containers. Use care when applying in areas frequented by wildlife or adjacent to any body of water. DO NOT apply when weather conditions favour drift from target areas.

Storage and Disposal:	STORAGE AND DISPOSAL Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Non-refillable containers Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product. Returnable containers Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

Safety Directions:	SAFETY DIRECTIONS Harmful if inhaled or swallowed. Will damage eyes. Will irritate the nose, throat and skin. Avoid contact with eyes and skin. DO NOT inhale vapour or spray mist. WHEN USING TOGETHER WITH OTHER PRODUCTS, CONSULT THEIR LABEL SAFETY DIRECTIONS. When opening the container, preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length butyl rubber gloves and face shield or goggles. If product on skin, immediately wash area with

	soap and water. Wash hands after use. After each day's use wash gloves, face shield or goggles and contaminated clothing.

First Aid Instructions:	FIRST AID If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, DO NOT induce vomiting. If in eyes wash out immediately with water.

First Aid Warnings:	

Nufarm Striker® Herbicide – Directions for Use

DIRECTIONS FOR USE

CROP	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Tree fruit, nuts and vines not less than 3 years old DORMANT APPLICATION including grapevines, olive trees, pome fruit (eg. apple, pear, nashi, quince), stone fruit (eg. apricot, cherry, nectarine, peach, plum) Tree nuts (eg. almond, macadamia, pecan, walnut)	Refer to Weeds Controlled list	Weed free soil	3 or 4 L/ha	**DO NOT** apply once budswell has occurred. Apply to freshly cultivated weed free soil. Use higher rate for longer residual control (up to 4 months). Where grass weeds are expected to be the major problem, or when control of a wider weed spectrum is needed, mix the lower rate with 4.5 L/ha Surflan* 500 (500 g/L) or 4.5 kg/ha Devrinol-C* (500 g/kg). Refer to individual product labels for crops, rates, and weeds controlled.
		Weeds at 4 - 6 true leaf stage		**DO NOT** apply once budswell has occurred. Use higher rate for longer residual control (up to 4 months). When young seedling grasses and/or broadleaf weeds are present, apply as a tank mix with 250 g/L paraquat, ,135 g/L paraquat + 115 g/L diquat, Reglone*, Nufarm weedmaster® DST, Nufarm weedmaster® DUO, or Nufarm Glyphosate 450 to give both knockdown and residual control. Refer to individual product labels for rates. Addition of a non-ionic surfactant such as Activator® Surfactant should be used in the spray mixture at 125 mL/100L. Mature, established weeds must be eliminated prior to application by mechanical or chemical means. **Macadamias:** Apply in 250 to 500 L water/ha. Apply after harvest to prevent spray contacting nuts. Avoid spray contact with macadamia tree foliage and stem. **DO NOT** apply to nuts on the ground.
Fruit and nut trees, vines including grapevines, olive trees, pome fruit (eg. apple, pear, nashi, quince), stone fruit (eg. apricot, cherry, nectarine, peach, plum) Tree nuts (eg. almond, macadamia, pecan, walnut)	Refer to 135 g/L paraquat + 115 g/L diquat product label	Emerged weeds present	250 mL/ha plus 135 g/L paraquat + 115 g/L diquat product at recommended label rates **OR** 25 – 30 mL/100L plus 135 g/L paraquat + 115 g/L diquat product at recommended label rates	The addition of Nufarm Striker® in a tank mix with a 135 g/L paraquat + 115 g/L diquat product will improve control of small flowered mallow, evening primrose and other weeds sensitive to Nufarm Striker®. Apply as a tank mix with 135 g/L paraquat + 115 g/L diquat - refer to the 135 g/L paraquat + 115 g/L diquat product label and follow all label directions.
	Refer to Nufarm weedmaster® DST, Nufarm weedmaster® DUO, or Nufarm Glyphosate 450 labels	Apply as a tank mix with glyphosate products	75 mL/ha plus Nufarm weedmaster® DST, Nufarm weedmaster® DUO * or Nufarm Glyphosate 450 at recommended label rates	Addition of Nufarm Striker® to glyphosate products will improve knockdown and increase the speed at which treated weeds develop visible symptoms of phytotoxicity (compared to results achieved with glyphosate applied alone) and give control of annual nettles *Urtica spp.*, and Marshmallow, *Malva parviflora*. For rates refer to the appropriate label. Read and follow all label directions. **DO NOT** apply the tank mix of glyphosate and Nufarm Striker® near trees or vines less than 3 years old unless they are effectively shielded from spray and spray drift.
Pyrethrum - as bare rooted transplants or seedlings	Refer to Weeds Controlled list (except chickweed)	Pre-plant incorporated into weed free soil worked to a fine tilth	4 or 6 L/ha	Apply prior to final soil preparation. The preferred implements for final soil preparation would be either a multiple tyne cultivator or rotary harrows. Use the 6 L/ha rate for heavy black clay soils only (as found in the Derwent and Coal River Valleys, Tasmania). Nufarm Striker® will not provide consistent control of chickweed.
Pyrethrum - greater than 4 leaf	Blackberry nightshade (*Solanum nigrum*) Fumitory (*Fumaria* spp.) Hogweed (*Polygonum aviculare*) Sorrel (*Rumex acetosella*) Volunteer potato (*Solanum tuberosum*) Cleavers (*Galium aparine*) Field Bindweed (*Convolvulus vulgaris*) Groundsel (*Senecio vulgaris*)	Emerged weeds present	100 – 150 mL/ha	Apply when pyrethrum growth stage is greater than 4 true leaf.
Pyrethrum - established crops, greater than 1 year old	As above plus Sowthistle (*Sonchus* spp.*)*, Spear thistle (*Cirsium vulgare*)	Emerged weeds present	200 mL/ha - 4 L/ha	Apply when pyrethrum is greater than 100 mm rosettes. Apply rates of greater than 1 L/ha ONLY between 1st of February and 31st of March. **DO NOT** apply later than 25 weeks before harvest.
Duboisia	Refer to Weeds Controlled list	Weed free soil	4 or 8 L/ha	Apply after transplantation of young seedlings or after harvest of mature plants. Use higher rate where longer residual control (up to 6 months) is required. "Over-the-top" application will be tolerated.
		Weeds at 4 - 6 true leaf stage		Best results are achieved when applied to moist soil free of weeds. However, some newly germinated small seedling grasses and broadleaf weeds (4-6 true leaves) will be controlled at these rates. Addition of Activator® Surfactant at recommended rates will improve activity. Established weeds must be eliminated prior to application by mechanical means.
Coffee	Refer to Weeds Controlled list	Weed free soil	2 – 4 L/ha	Apply after transplanting coffee plants or under established bushes. Use the higher rate where longer residual activity (up to 4 months) is required. Best results are achieved when applied to moist soil, free of weeds with follow-up irrigation.
		Weeds at 2 - 4 true leaf stage		When seedlings are present, apply as a tank mix with 250 g/L paraquat to produce both knockdown and residual control. Addition of Activator® Surfactant at 125 mL/100L should be used in the spray mixture. Apply as a directed spray to avoid contact with coffee plants. Mature established weeds must be eliminated by mechanical or chemical means prior to application.

CROP	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Tropical and subtropical fruit (inedible peel) including: avocado, cherimoya, custard apple, durian, feijoa, guava, jackfruit, kiwi fruit, longan, lychee, mango, mangosteen, papaya, passionfruit, pawpaw, persimmon, rambutan, star apple	Refer to Weeds Controlled list	Weed free soil	4 L/ha	Best results are obtained when applied to moist soil free of weeds and followed by rainfall or irrigation.
		Emerged weeds present		If weeds are present Nufarm Striker® should be applied as a tank mix with the recommended rates of Nufarm weedmaster® DST, Nufarm weedmaster® DUO, Nufarm Glyphosate 450, 135 g/L paraquat + 115 g/L diquat or 250 g/L paraquat.
Brassica crops, broccoli, cabbage, cauliflower	Refer to Weeds Controlled list	Weed free soil	1.5 - 2 L/ha	Apply Nufarm Striker® to prepared ground 4 - 7 days prior to transplanting. If soil is dry irrigation or rainfall is required prior to transplanting for activation of Nufarm Striker®. Utilise transplanting techniques which cause minimal soil disturbance. Excessive soil disturbance will lessen herbicide activity. Use the higher rate in situations where weed pressure is known to be heavy.
Forestry - *Eucalyptus* spp., *Pinus* spp. Nursery situations open rooted cuttings and seedlings	Refer to Weeds Controlled list	Cutting: May - Dec (4 weeks post setting) Seedlings: Jan - Jun (3 months post emergence in nursery)	2 L/ha	Use 2 L/ha/month, maximum of 3 applications. **DO NOT** use a surfactant. Apply to weed free soil either before or preferably soon after transplanting. Disturbance of the herbicidal barrier on the soil surface may reduce the length of weed control. If air temperature exceeds 20°C, application is not recommended. Application should be preferably on an overcast day if air temperatures exceed 20°C. **DO NOT** apply under hot or windy conditions.
Forestry - *Eucalyptus* spp., *Pinus* spp.	Refer to Weeds Controlled list	Weed free soil	3 or 4 L/ha	Apply to weed free soil either before or preferably soon after transplanting. Disturbance of the herbicidal barrier on the soil surface by the transplantation process may reduce the length of weed control. Use higher rate for longer residual control. Apply as a directed or over-the-top spray. Nufarm Striker® may be tank mixed with a 900 g/kg simazine product to extend the spectrum and length of weed control. **DO NOT** use this tank mix in Eucalyptus plantations grown on sands, with no clay or organic matter. Application to foliage that is soft and has not hardened off and/or if air temperatures exceed 20°C may increase the likelihood of foliar damage to trees, especially eucalypts. Nufarm Striker® has been found to be safe to commonly planted species used in forestry. Sensitivity of less common species should be checked prior to large scale application.
Forestry - *Eucalyptus* spp. *Pinus* spp. Cont.	Refer to Weeds Controlled list	Weeds at 4 - 6 true leaf stage	3 or 4 L/ha	Apply as an over-the-top spray to eucalypts and pines. Best results are achieved when applied to moist soil free of weeds. However, some newly germinated small seedling grasses and broadleaf weeds (4 to 6 true leaves) will be controlled at these rates. Addition of Activator® Surfactant at recommended rates will improve activity. Established weeds must be eliminated prior to application. Nufarm Striker® may be tank mixed with a 900 g/kg simazine product to extend the spectrum and length of weed control. **DO NOT** use this tank mix in Eucalyptus plantations grown on sands, with no clay or organic matter. Application to foliage that is soft and has not hardened off and/or if air temperatures exceed 20°C may increase the likelihood of foliar damage to trees, eucalypts especially. Nufarm Striker® has been found to be safe to commonly planted species used in forestry. Sensitivity of less common species should be checked prior to large scale application.
Forestry Trees	Broadleaf weeds and grasses	Weeds at 4 - 6 true leaf stage	4 L/ha or 4 mL/10m^2	For the establishment of trees for approved farm practices such as wind breaks, erosion control, woodlots and forestry plantings. When applying as a post-plant spray, ensure spray is directed to the base of seedlings, or that seedlings are protected. **DO NOT** apply under hot or windy conditions.
Prior to sowing winter cereals Prior to sowing cotton	Refer to Nufarm weedmaster® DST, Nufarm weedmaster® DUO, or Nufarm Glyphosate 450 labels	Apply as a tank mix with glyphosate products	75 mL/ha plus Nufarm weedmaster® DST, Nufarm weedmaster® DUO, or Nufarm Glyphosate 450 at recommended label rates	Addition of Nufarm Striker® to glyphosate products will improve knockdown and increase the speed at which treated weeds develop visible symptoms of phytotoxicity (compared to results achieved with glyphosate applied alone) and give control of annual nettles *Urtica spp.*, and Marshmallow, *Malva parviflora*. For rates refer to the appropriate label. Read and follow all label directions.

When using Nufarm Striker® in combination with other herbicides, refer to the appropriate label and read and follow all label directions.

WEEDS CONTROLLED	
BEFORE GERMINATION	**SEEDLINGS**
Amsinckia *Amsinckia* spp.	Amsinckia *Amsinckia* spp.
Barley grass *Hordeum leporinum*	Bellvine *Ipomoea* spp.
Barnyard grass *Echinochloa* spp.	Capeweed *Arctotheca calendula*
Blackberry nightshade *Solanum nigrum*	Common cotula *Cotula australis*
Bladder ketmia *Hibiscus trionum*	Crowsfoot grass *Eleusine indica*
Burrgrass *Cenchrus australis*	Dead nettle *Lamium amplexicaule*
Caltrop *Tribulus terrestris*	Groundsel *Senecio vulgaris*
Capeweed *Arctotheca calendula*	Liverseed grass *Urochloa panicoides*
Chickweed *Stellaria media*	Marshmallow *Malva parviflora*
Crowsfoot grass *Eleusine indica*	Pigweed *Portulaca oleracea*
Dead nettle *Lamium amplexicaule*	Potato weed *Galinsoga parviflora*
Fat hen *Chenopodium album*	Redshank *Amaranthus cruentus*
Giant pigweed *Trianthema portulacastrum*	Shepherd's purse *Capsella bursa-pastoris*
Liverseed grass *Urochloa panicoides*	Sowthistle *Sonchus oleraceus*

Lovegrass *Eragrostis* spp. Pigeon grass *Setaria* spp. Pigweed *Portulaca oleracea* Prickly lettuce *Lactuca* spp. Red natal grass *Rhynchelytrum repens* Redshank *Amaranthus cruentus* Ryegrass *Lolium* spp. Sesbania pea *Sesbania cannabina* Shepherd's purse *Capsella bursa-pastoris* Smallflower mallow *Malva parviflora* Soursob *Oxalis pes-caprae* Sowthistle *Sonchus oleraceus* Starburr *Acanthospermum hispidum* Stinkgrass *Eragrostis cilianensis* Summer grass *Digitaria* spp. Thornapple *Datura stramonium* White eye *Richardia brasiliensis* Wild mustard *Sisymbrium* spp. Wild radish *Raphanus raphanistrum* Wireweed *Polygonum aviculare*	Stinging nettle *Urtica urens* Stinkgrass *Eragrostis cilianensis* Wild radish *Raphanus raphanistrum*

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.