RLP APPROVED

Product Name: NUFARM SINKER FUNGICIDE

APVMA Approval No: 67732/133927

Label Name:	NUFARM SINKER FUNGICIDE

Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	500 g/L FLUTRIAFOL

Mode of Action:	GROUP 3 FUNGICIDE

Statement of Claims:	For control of sugarcane smut and pineapple disease in sugarcane planting material as per the directions for use table.

Net Contents:	1L - 20L

Restraints:	DO NOT apply if heavy rains or storms are forecast within 24 hours. DO NOT irrigate to the point of runoff for at least 24 hours after application.

Directions for Use:	This section contains file attachment.

Other Limitations:	

Withholding Periods:	Harvest: NOT REQUIRED WHEN USED AS DIRECTED. Grazing: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 32 WEEKS AFTER TREATMENT.

Trade Advice:	

General Instructions:	This section contains file attachment.

Resistance Warning:	NUFARM SINKER FUNGICIDE is a member of the DMI group of fungicides. For fungicide resistance management Nufarm Sinker is a Group 3 fungicide. Some naturally occurring individual fungi resistant to Nufarm Sinker and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Nufarm Sinker or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Nufarm Australia Limited accepts no liability for any losses that may result from the failure of Nufarm Sinker to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

Precautions:	RE-ENTRY/RE-HANDLING DO NOT enter treated areas until the spray has dried unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use. DO NOT re-handle treated sugarcane planting material unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

Protections:	PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Toxic to aquatic life with long lasting effects. DO NOT contaminate wetlands or watercourses with this product or used containers.

Storage and Disposal:	Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product. DIP DISPOSAL Disposal of used/excess solution should be through an authorized disposal facility. If such a facility is not available dispose of used dip solution and sludge at a rate not exceeding 20,000 L/ha over an area of dedicated and bunded (soil at least 15 cm high) flat land.

Safety Directions:	Harmful if swallowed. Harmful if absorbed by skin contact. Will irritate the eyes. May irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing the spray or dip, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length chemical resistant gloves and face shield or goggles. When using the prepared spray or dip, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

First Aid Warnings:	

GENERAL INSTRUCTIONS

MIXING

Spray-planter or planter mounted dip application: Half fill the clean spray or dip tank with clean water then add the required volume of Nufarm Sinker as listed in Table A and 50 mL/100 L Activator with the agitator in operation. Make up to final volume with water and continue agitation during spraying or dipping and after a stoppage.

Table A. Volume of Nufarm Sinker required for different water rates per 100 m row of cane when a spray planter or planter mounted dip is used

Volume of spray/dip mixture (L) per 100 m row of cane	Total Volume of spray/dip mixture (L/ha)	Mixing rate of Nufarm Sinker (mL/100L)
6	400	125
6.75	450	113
7.5	500	100
8.25	550	90
9	600	83
9.75	650	77
10.5	700	71
11.25	750	67
12	800	63
12.75	850	58
13.5	900	56
14.25	950	53
15	1000	50
DO NOT apply more than 15 L/100m row as Nufarm Sinker must be maintained at a level of at least 50 mL/100 L of spray mixture		

Static Dip application: Half fill the clean dip tank with clean water. Add the required volume of Nufarm Sinker and a wetting agent at the appropriate rates while filling the tank to the correct level. Agitate while filling using either a mechanical device or recirculating pump. Agitate between each batch of cane stalks/billets being dipped.

APPLICATION

Table B (single row planting) and Table C (dual row planting) below show the different rates of Nufarm Sinker to be used per hectare with different row or bed spacings.

Table B. Rate per hectare of Nufarm Sinker required for different row spacings in single row planting

Row spacing (single row planting)	Rate of Nufarm Sinker (mL/ha)
1.5	500
1.55	484
1.6	469
1.65	455
1.7	441
1.75	429
1.8	417

Table C. Rate per hectare of Nufarm Sinker required for different bed spacings in dual row planting

Bed spacing (dual row planting)	Rate of Nufarm Sinker (mL/ha)
1.8	833
1.9	789
2	750
2.1	714
2.2	682
2.3	652
2.4	625

DIRECTIONS FOR USE

Spray planters or planter mounted dip applicators

CROP	DISEASE	RATE	CRITICAL COMMENTS
Sugarcane planting material (setts/billets) - **SINGLE ROW PLANTING** (Planters with planting boards that result in wide planting furrows in single rows with a uniform spacing between each row)	Pineapple disease (*Ceratocystis paradoxa)* Sugarcane smut *(Sporisorium scitamineum)*	7.5 mL/100 m row + Activator at 50 mL/100 L spray mixture	For the prevention of primary infection of pineapple disease and sugarcane smut in sugarcane, apply as a spray onto setts/billets in the planting chute or with a planter mounted dip. The use of Activator® will enhance coverage of the fungicide on the planting material. Calibrate the spray application or planter mounted dip so that the total volume of Nufarm Sinker applied per 100 metre row of cane is 7.5 mL. This can be done by mixing Nufarm Sinker at 125 mL/100 L of water and applying 6L of this mixture per 100 metre row of cane. Wherever possible, calibrate the planter to ensure thorough coverage with this water volume. However, if coverage is not adequate higher water volumes can be used by reducing the rate of Nufarm Sinker per 100 L of water accordingly (refer to Table A in the GENERAL INSTRUCTIONS). For single row planting, **DO NOT** apply more than 500 mL/ha. For dual row planting, **DO NOT** apply more than 833 mL/ha. Split application is not recommended for dual row planting systems. Ensure the rate of 7.5 mL/100 m row is applied in both rows of a dual row planting system. **FOR SPRAY PLANTER APPLICATION** The spray should be applied with a minimum of 4 nozzles arranged in the planting chute to give thorough coverage of all surfaces of the setts/billets before they are planted in the furrow (up to the point of run-off). **FOR PLANTER MOUNTED DIP APPLICATION** Cane may or may not have previously been hot water treated for disinfection from other diseases. Check dip frequently for dirt and trash as these reduce the effectiveness of the fungicide. If solution colour changes from blue it should be discarded and a fresh dip made up after the dip tank has been cleaned out to extend the life of the dip. Ensure cane is dipped for at least 5-10 seconds for thorough coverage.
Sugarcane planting material (setts/billets) - **DUAL ROW PLANTING** (Cane planted in beds with dual rows with a non-uniform spacing between each row)		7.5 mL/100 m row + Activator at 50 mL/100 L spray mixture	

Static Dip Application (For treatment of planting material in a non-planter mounted dip for use in production of seed cane only)

CROP	DISEASE	RATE	CRITICAL COMMENTS
Sugarcane planting material (setts/billets, seed cane only)	Pineapple disease (*Ceratocystis paradoxa)* Sugarcane smut *(Sporisorium scitamineum)*	50-125 mL/100 L water + Activator at 50 mL/100 L spray mixture	For the prevention of primary infection of pineapple disease and sugarcane smut of sugarcane planting material apply utilising a static dip as a 5-10 minute ambient dip treatment prior to planting. For heavy disease pressure, or where superior control is required, use the higher rate and increase the duration of dipping (up to 12 hours). The use of Activator® will enhance coverage of the fungicide on the planting material. Cane may or may not have previously been hot water treated for disinfection from other diseases. Check dip frequently for dirt and trash as these reduce the effectiveness of the fungicide. If solution colour changes from blue it should be discarded.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.