RLP
APPROVED

Product Name: DURIVO INSECTICIDE
APVMA Approval No: 63936/127043

Label Name:	DURIVO INSECTICIDE

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	100 g/L CHLORANTRANILIPROLE 200 g/L THIAMETHOXAM

Mode of Action:	GROUP 4A 28 INSECTICIDE

Statement of Claims:	For the control of various insect pests in Brassicas, and Fruiting and Leafy Vegetables as per the Directions for Use

Net Contents:	1L 20L-100L 5L-10L

Restraints:	DO NOT apply more than 1 application of DURIVO or any other soil applied Group 4A Insecticide per crop DO NOT apply any other Group 4A Insecticide as a foliar spray after soil application of DURIVO in that crop DO NOT transplant seedlings treated by seedling drench into hydroponic production systems DO NOT apply DURIVO where planting density exceeds 56,000 plants/ha for brassicas, 83,000 plants/ha for leafy vegetables or 33,000 plants/ha for fruiting vegetables DO NOT plant or transplant more than four DURIVO treated crops in the same area in any one year

Directions for Use:	This field has been left intentionally blank by the applicant

Other Limitations:	

Withholding Periods:	WITHHOLDING PERIODS Brassicas: DO NOT HARVEST FOR 6 WEEKS AFTER APPLICATION Brassica Leafy Vegetables, Leafy Vegetables: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION Fruiting Vegetables: NOT REQUIRED WHEN USED AS DIRECTED

Trade Advice:	EXPORT TRADE ADVICE – TREATED CROPS: Treated crop commodities destined for export may require extra time being allowed between application and harvest to be accepted in some export markets. Before you use this product, you are advised to contact Syngenta and/or your industry body about any potential trade issues and their management.

General Instructions:	GENERAL INSTRUCTIONS DURIVO INSECTICIDE is a broad spectrum, systemic insecticide controlling key lepidopteron, sucking and chewing insect pests in a range of vegetable crops. When applied as a seedling application prior to transplant or as a plant hole drench at transplant, DURIVO is readily taken up by the roots of seedling transplants and is rapidly translocated throughout the plant to provide residual insect control. DURIVO, applied to seedlings as a seedling tray or plant hole drench, controls insects that feed on plant parts or fluids. It should have minimal impact on beneficial arthropods and therefore its use is compatible with integrated pest management programs. MIXING Add the required amount of DURIVO to clean water in half-filled spray tank with the agitator or by-pass in operation. Maintain agitation while filling tank with remainder of water. Agitate whilst spraying. APPLICATION DURIVO must be applied in a manner which allows the insecticide to be absorbed by plant roots. Sufficient water volumes to ensure uniform application and incorporation into seedling root zones are required. Seedling Tray Application: DURIVO should be applied to the seedling tray prior to transplanting through dedicated nursery spray equipment such as a calibrated hydraulic boom. Apply a single application as a slurry with water in a total volume of 2.5 L per 1000 seedlings. Some adjustment in water volume may be required depending on seedling cell size and moisture content. Final volume must be sufficient to wash product into the seedling root ball but not cause runoff or leaching from seedling cells. Failure to wash product into the root ball may result in reduced control. Watering of seedling trays after application may wash product from cells. If watering is required between application and transplanting, it should be done so sparingly and only as required. DO NOT allow run through from the cells. Where spray drift cannot be captured, calibrated shrouded sprayers or other methods to minimise spray drift must be used to deliver a uniform distribution to each transplant cell. Excess spray includes applied spray which misses the target, runs off or inadvertently runs through the cells. Any such excess must be captured and should be filtered and returned to the spray tank. DO NOT use watering cans, sprinklers or other imprecise methods of application. Losses from spray drift or through runoff can be minimised by ensuring there are no gaps between the seedling trays during application.

	Application should occur immediately before transplanting to a maximum of 24 hours before seedling dispatch. Some seedling damage may occur post application, but symptoms are generally transient and will not affect crop vigour. Seedlings should be transplanted within 48 hours of application. Planting Hole Drench Application: DURIVO should be applied with water in a minimum total volume of 50 mL per seedling directly at the base of transplanted seedlings or into the planting hole immediately before transplanting. Actual total volume will vary depending on equipment used, pre-existing soil moisture levels and soil type but should be to ensure uniform application and distribution. With applications made after transplanting DURIVO should be incorporated to seedling root depth using sufficient irrigation within 24 hours. COMPATIBILITY As formulations of other manufacturer's products are beyond the control of Syngenta, and water quality varies with location, all mixtures should be tested prior to mixing or using commercial quantities. For applications via drip/trickle irrigation, DURIVO is compatible with TERVIGOÂ® and selected fertilisers. For more information, contact your local Syngenta Representative.

Resistance Warning:	GROUP 4A 28 INSECTICIDE For insecticide resistance management DURIVO INSECTICIDE is a Group 4A and Group 28 insecticide. Some naturally occurring insect biotypes resistant to DURIVO and other Group 4A and/or Group 28 insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if DURIVO or other Group 4A and/or Group 28 insecticides are used repeatedly. The effectiveness of DURIVO on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Syngenta Australia Pty Ltd accepts no liability for any losses that may result from failure of DURIVO to control resistant insects. DURIVO may be subject to specific resistance management strategies. For further information contact your supplier, Syngenta representative or local agricultural department agronomist. DURIVO is a Group 4A and Group 28 insecticide and should be applied in accordance with CropLife Insecticide Resistance Management guidelines. In addition, DURIVO should, wherever possible, be incorporated into an Integrated Pest Management (IPM) program that includes cultural and biological control practices in association with the following IRM guidelines: • DO NOT apply more than 1 application of DURIVO or any other soil applied Group 4A insecticide per crop • Following the use of DURIVO rotate to alternative mode of action (MoA) insecticide class(es) for a period covering at least one generation of the target pest. For many pests this will require a minimum of 2 applications of alternate MoA insecticides • The total exposure period for all Group 28 insecticides applied through the crop cycle (from seedling to harvest) should not exceed 50% of the crop cycle • DO NOT apply other Group 4A insecticides as a foliar spray after soil application of DURIVO in that crop • Monitor insect populations for product effectiveness Additional information on Insecticide Resistance Management Strategies can be found on the CropLife Australia website: http://www.croplife.org.au/industry-stewardship/resistance-management.

Precautions:	PRECAUTIONS DO NOT apply as a seedling tray application using equipment carried on the back of the user. Rehandling: DO NOT handle treated seedlings until the spray has dried unless wearing cotton overalls buttoned to neck and wrist (or equivalent clothing) and chemical resistant gloves.

Protections:	PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Dangerous to aquatic life. DO NOT contaminate dams, waterways or drains with the chemical or used containers. DO NOT apply under weather conditions or from spraying equipment which could be expected to cause spray drift on adjacent areas, particularly wetlands, waterbodies or watercourses. PROTECTION OF HONEY BEES AND OTHER INSECT POLLINATORS Highly toxic to bees. Thiamethoxam has systemic action. The persistence of residues and expression of thiamethoxam in nectar and pollen suggests the potential for long term effects on honey bee larvae and colonies. Residues may remain at levels toxic to bees for several days after application. Where DURIVO has been applied in or near the treatment/ pollinator area, notify beekeepers prior to placing beehives in the treated field.

Storage and Disposal:	STORAGE AND DISPOSAL Store in the closed, original container in a cool, well ventilated area out of direct sunlight. Refillable Containers (20 – 100 L pack size range only) Empty contents fully into application equipment. Close all valves and return to point of sale for refill or storage. Other Containers (1 L and 5 - 10 L pack size only) Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

Safety Directions:	SAFETY DIRECTIONS Will irritate eyes. Avoid contact with eyes. When opening the container, preparing and using the product and transplanting treated seedlings wear: • cotton overalls buttoned to neck and wrist (or equivalent clothing) • elbow-length chemical resistant gloves. Wash hands after use. After each day's use wash gloves and contaminated clothing.

First Aid Instructions:	FIRST AID If poisoning occurs contact a doctor or Poisons Information Centre. Phone 131 126.

First Aid Warnings:	SAFETY DATA SHEET If additional hazard information is required refer to the Safety Data Sheet. For a copy phone 1800 067 108 or visit our website at www.syngenta.com.au

Direction for use:

Crop	Pest	Rate	Critical comments
Brassicas Including Broccoli, Brussels Sprouts, Cabbage, Cauliflower,	Diamondback Moth (*Plutella xylostella*), Cabbage White Butterfly (*Pieris rapae*), Corn Earworm (*Helicoverpa armigera*), Native Budworm (*Helicoverpa punctigera*), Cabbage Centre Grub (*Hellula hydralis*), Cabbage Cluster Caterpillar (*Crocidolomia pavonana*), Cluster Caterpillar (*Spodoptera litura*), Soybean Looper (*Thysanoplusia orichalcea*), Cabbage Aphid (*Brevicoryne brassicae*), Green Peach Aphid (*Myzus persicae*), Silverleaf Whitefly (*Bemisia tabaci*), Greenhouse Whitefly (*Trialeurodes vaporariorum*), Green Vegetable Bug (*Nezara viridula*), Western Flower Thrips (*Frankliniella occidentalis*), Onion Thrips (*Thrips tabaci*).	**Planting densities up to 41,000 plants/ha** 22 to 30 mL per 1000 seedlings	Apply DURIVO as a seedling tray application prior to transplant, or as a plant hole drench at transplant. Refer to Application section below for further instructions. **Where a rate range exists** Use lower rates when using the planting hole drench application method. Use lower rates when treating crops and varieties with growing seasons of less than 7 weeks duration. Use rates toward the upper end of the range when season conditions favour rapid crop development or high pest pressure is anticipated. Application should occur immediately before transplanting to a maximum of 24 hours before seedling dispatch. Seedlings should be transplanted within 48 hours of application. DO NOT apply the seedling drench within 48 hours of heavy rain being forecast or irrigate to the point of runoff for 48 hours after drenching. After application, continue to monitor crops and apply a foliar insecticide when target pests reach threshold levels. Refer to Resistance Management section below for resistance management strategy information. **Planting densities and rate selection** **The maximum recommended application rate per 1000 seedlings will depend on the proposed field planting density.**
Brassica Leafy Vegetables Including Chinese broccoli, Chinese cabbage, garden cress, kale and rocket		**Planting densities from 41,000 to 56,000 plants/ha** 22 mL per 1000 seedlings	
Leafy Vegetables Including Lettuce, Endive, Silverbeet, Spinach, Chard	Corn Earworm (*Helicoverpa armigera*), Native Budworm (*Helicoverpa punctigera*), Cluster Caterpillar (*Spodoptera litura*), Looper (*Chrysodeixis* spp.), Lettuce Aphid (*Nasonovia ribis-nigri*), Green Peach Aphid (*Myzus persicae*), Brown Sowthistle Aphid (*Uroleucon sonchi*), Silverleaf Whitefly (*Bemisia tabaci*), Western Flower Thrips (*Frankliniella occidentalis*), Vegetable Leafhopper (*Austroasca viridigrisea*), Lucerne Leafroller (*Merophyas divulsana*)	**Planting densities up to 41,000 plants/ha** 15 to 30 mL per 1000 seedlings	
		Planting densities from 41,000 to 83,000 plants/ha 15 mL per 1000 seedlings	
Fruiting vegetables Including Tomatoes, Capsicum, Eggplant	Corn Earworm (*Helicoverpa armigera*), Native Budworm (*Helicoverpa punctigera*), Potato Moth (*Phthorimaea operculella*), Cluster caterpillar (*Spodoptera litura*), Green peach aphid (*Myzus persicae*) Silverleaf Whitefly (*Bemisia tabaci*), Greenhouse whitefly (*Trialeurodes vaporariorum*), Western Flower Thrips (*Frankliniella occidentalis*)	**Planting densities up to 25,000 plants/ha** 37 to 50 mL per 1000 seedlings	

Crop	Pest	Rate	Critical comments
	Tomato Thrips (*Frankliniella schultzei*)	**Planting densities from 25,000 to 33,000 plants/ha** 37 mL per 1000 seedlings	Apply DURIVO at the rate corresponding to the desired planting density. DO NOT exceed the maximum rate of DURIVO based on anticipated planting density. DO NOT apply DURIVO where planting density exceeds 56,000 plants/ha for brassicas, 83,000 plants/ha for leafy vegetables 33,000 plants/ha for fruiting vegetables

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.