RLP
APPROVED

Product Name: MINECTO FORTE Insecticide

APVMA Approval No: 87610/127218

Label Name:	MINECTO FORTE Insecticide

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	80 g/L CYANTRANILIPROLE 400 g/L DIAFENTHIURON

Mode of Action:	GROUP 12A 28 INSECTICIDE

Statement of Claims:	For the control of various mite and insect pests in cucurbits and fruiting vegetables

Net Contents:	1 to 10 L

Restraints:	This section contains file attachment.

Directions for Use:	This section contains file attachment.

Other Limitations:	NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

Withholding Periods:	WITHHOLDING PERIODS: Cucurbits: DO NOT HARVEST TREATED CROPS WITHIN 1 DAY OF APPLICATION Fruiting vegetables (other than cucurbits): DO NOT HARVEST TREATED CROPS WITHIN 4 DAYS OF APPLICATION

Trade Advice:	

General Instructions:	This section contains file attachment.

Resistance Warning:	Insecticide Resistance Warning GROUP 12A 28 INSECTICIDE For insecticide resistance management MINECTO FORTE is a Group 12A and Group 28 insecticide. Some naturally occurring insect biotypes resistant to MINECTO FORTE and other Group 12A and Group 28 insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if MINECTO FORTE and other Group 12A and Group 28 insecticides are used repeatedly. The effectiveness of MINECTO FORTE on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Syngenta Australia Pty Ltd accepts no liability for any losses that may result from failure of MINECTO FORTE to control resistant insects. MINECTO FORTE may be subject to specific resistance management strategies. For further information contact your supplier, Syngenta representative or local agricultural department agronomist. Unless directed otherwise in the specific crop/insect sections of the label, the following practices are recommended to prevent or delay the development of insecticide resistance to MINECTO FORTE and to Group 28 insecticides: • Apply MINECTO FORTE and other Group 28 insecticides using a “treatment window” approach to avoid exposure of consecutive insect pest generations to the same mode of action. Multiple successive applications of Group 28 insecticides are acceptable if they are used to treat a single insect generation. • A “treatment window” is defined as the period of residual activity provided by single or sequential applications of products with the same mode of action. This “treatment window” should not exceed approximately the length of one generation of the target pest. • Following a “treatment window” of MINECTO FORTE or other Group 28 insecticides, rotate to a treatment window of effective insecticides with a different mode of action. • The total exposure period of all “Group 28-active windows” applied throughout the crop cycle (from seedling to harvest) should not exceed 50% of the crop cycle. • Incorporate IPM techniques into the overall pest management program. • Monitor insect populations for product effectiveness. Additional information on Insecticide Resistance Management Strategies can be found on the CropLife Australia website: http://www.croplife.org.au/industry-stewardship/resistance-management.

Precautions:	PRECAUTION Re-entry Period DO NOT undertake occupational re-entry activities until the specified day after treatment, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length chemical resistant gloves: Scouting, hand weeding, hand pruning of fruiting vegetables and cucurbits, and hand harvesting of cucurbits Day 0 Tying/training, hand harvesting of fruiting vegetables Day 6 Hand setting irrigation of fruiting vegetables Day 11 Hand setting irrigation of cucurbits Day 12

Protections:	PROTECTION OF HONEY BEES AND OTHER INSECT POLLINATORS

	Highly toxic to bees. DO NOT apply to crops from the onset of flowering until flowering is complete unless the application is made in the time period between 2 hours prior to sunset and 8 hours prior to sunrise. DO NOT allow spray drift to flowering weeds or flowering crops in the vicinity of the treatment area. Before spraying, notify beekeepers to move hives to a safe location with an untreated source of nectar and pollen, if there is potential for managed hives to be affected by the spray or spray drift. INTEGRATED PEST MANAGEMENT Toxic to beneficial arthropods. Not compatible with integrated pest management (IPM) programs utilising beneficial arthropods. Minimise spray drift to reduce harmful effects on beneficial arthropods in non-crop areas. PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

Storage and Disposal:	Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

Safety Directions:	SAFETY DIRECTIONS Harmful if inhaled. May irritate the eyes. Avoid contact with eyes. DO NOT inhale spray mist. When opening the container and preparing the spray, wear: - cotton overalls buttoned to the neck and wrist (or equivalent clothing) - elbow-length chemical resistant gloves If applying by low pressure hand wand, wear: - cotton overalls buttoned to the neck and wrist (or equivalent clothing) - a washable hat - elbow-length chemical resistant gloves Wash hands after use. After each day’s use wash gloves and contaminated clothing.

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.

First Aid Warnings:	

GENERAL INSTRUCTIONS

Mixing

Measure the required amount of MINECTO FORTE, add to the partly filled spray tank, and then add the remainder of the water. Add oil or non-ionic surfactant after the MINECTO FORTE is well mixed. Maintain continuous agitation while spraying.

Application

Ground Application Only: To be effective MINECTO FORTE requires thorough spray coverage. Ensure that equipment is properly calibrated to give an even distribution at the correct volume and delivering a MEDIUM spray quality according to nozzle manufacturer specifications that refer to the ASAE S572 Standard or the BCPC Guideline.

Compatibility

As formulations of other manufacturer's products are beyond the control of Syngenta Australia, and the quality of water may vary with location, all mixtures should be tested prior to mixing commercial quantities.
Wetting agents: Not all adjuvants or crop oils are of equal quality. Syngenta does not support the use of adjuvants/crop oils, other than those listed below:

- AGRAL® Spray Adjuvant
- Hasten* Spray Adjuvant
- SACOA BioPest* Paraffinic Oil (Fruiting Vegetables Other than Cucurbits only)

***Cucurbits*:** DO NOT apply SACOA BioPest Paraffinic Oil with MINECTO FORTE in cucurbit crops as phytotoxicity may occur.

Crop safety: Caution must be exercised when applying Hasten Spray Adjuvant with MINECTO FORTE on sensitive crops during periods when plants are subjected to drought stress, high temperatures or high humidity.

DIRECTIONS FOR USE

Crop	Pest	Rate	Critical Comments
Critical comments – all crops Monitor crops and apply when pest action thresholds are reached. Continue to monitor crops and make subsequent insecticide applications as necessary. Ensure spray volume is adequate to thoroughly cover all parts of the crop. DO NOT apply more than 2 applications of MINECTO FORTE per crop. Addition of a wetting agent may assist control under less than ideal conditions. Use the wetting agent according to its label directions. See ***Wetting Agents*** section below for recommended products.			
Cucurbits (including cucumber, zucchini, rockmelon, watermelon, pumpkin, Asian melons) **Open field only**	Silverleaf whitefly *(Bemisia tabaci)*	500 mL/ha	Target immature whitefly stages. Addition of AGRAL at 10 mL/100 L or Hasten at 200 mL/100 L may assist control of silverleaf whitefly in cucurbits. This use is subject to a **Croplife Australia Insecticide Resistance Management strategy.**
	Melon aphid *(Aphis gossypii)*	500 mL/ha	Addition of AGRAL at 10 mL/100 L may assist control of melon aphid. This use is subject to **Croplife Australia Insecticide Resistance Management strategy.**
	Cotton bollworm (*Helicoverpa armigera),* Native budworm *(Helicoverpa punctigera),* Cucumber moth *(Diaphania indica),* Cluster caterpillar *(Spodoptera litura)*	500 mL/ha	Target sprays against eggs and newly hatched larvae before they become entrenched. Addition of AGRAL at 10 mL/100 L may assist control of these pests in cucurbits. **This use is subject to a Resistance Management strategy for Group 28 insecticides.**
	Green peach aphid *(Myzus persicae)*	625 mL/ha	Addition of Hasten at 200 mL/100 L may assist control of green peach aphid. This use is subject to **Croplife Australia Insecticide Resistance Management strategy**
	Two-spotted mite *(Tetranychus urticae)*	750 mL/ha	Addition of Hasten at 200 mL/100 L may assist control of two-spotted mite in cucurbits.
	Suppression of: Western flower thrips *(Frankliniella occidentalis),* Tomato thrips *(Frankliniella schultzei),* Plague thrips *(Thrips imaginis)*	750 mL/ha	
Fruiting vegetables (including tomatoes, capsicums, eggplant, chillies) **Open field only**	Silverleaf whitefly *(Bemisia tabaci)*	500 mL/ha	Target immature whitefly stages. This use is subject to a **Croplife Australia Insecticide Resistance Management strategy**
	Cotton bollworm (*Helicoverpa armigera),* Native budworm *(Helicoverpa punctigera),* Potato Moth (*Phthorimaea operculella),* Cluster caterpillar *(Spodoptera litura)*	500 mL/ha	Target sprays against eggs and newly hatched larvae before they become entrenched. The use against *Helicoverpa* spp. is subject to **Croplife Australia Insecticide Resistance Management strategy. Refer also to note below regarding Group 28 insecticides.**
	Green peach aphid *(Myzus persicae)*	625 mL/ha	Addition of Hasten at 200 mL/100 L may assist control of green peach aphid. This use is subject to **Croplife Australia Insecticide Resistance Management strategy.**
	Two-spotted mite *(Tetranychus urticae)*	750 mL/ha	

Crop	Pest	Rate	Critical Comments
	Suppression of: Western flower thrips *(Frankliniella occidentalis),* Tomato thrips *(Frankliniella schultzei),* Plague thrips *(Thrips imaginis)*	750 mL/ha	Addition of Hasten at 200 mL/100 L, SACOA BioPest* Paraffinic Oil at 1000 mL/100 L or AGRAL at 10 mL/100 L may assist suppression of thrips in fruiting vegetables.

Restraints
DO NOT apply by aircraft.
DO NOT apply if heavy rains or storms are forecast within 3 days.
DO NOT irrigate to the point of runoff for at least 3 days after application.
DO NOT apply by backpack application.
DO NOT use in protected cropping situations.
DO NOT apply consecutive applications of MINECTO FORTE
DO NOT apply within 28 days of previous application of MINECTO FORTE

Spray Drift Restraints:
Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.
DO NOT allow bystanders to come into contact with the spray cloud.
DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.
DO NOT apply unless the wind speed is between three and 20 kilometres per hour at the application site during the time of application.
DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.
DO NOT apply by a boom sprayer unless the following requirements are met:

- Spray droplets are not smaller than a MEDIUM spray droplet size category.
- Minimum distances between the application site and downwind sensitive areas are observed (see '**Mandatory buffer zones**' section of the following table titled '**Buffer zones for boom sprayers'**).

Buffer zones for boom sprayers

Application rate	Boom height above the target canopy	Mandatory downwind buffer zones
		Natural aquatic areas
Up to maximum label rate	1.0 m or lower	5 metres