POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Pirimor® WG

Aphicide

syngenta®

ACTIVE CONSTITUENT: 500 g/kg PIRIMICARB
(an anticholinesterase compound)

GROUP	1A	INSECTICIDE

For control of certain Aphids on crops and pastures as per Directions for Use

Syngenta Australia Pty Ltd
Level 1, 2-4 Lyonpark Road, Macquarie Park NSW 2113

In a transport emergency dial 000, Police or Fire Brigade
For specialist advice in an emergency only, call 1800 033 111 (24 hours)

APVMA Approval No: 49835/128530
Item number

WG	Formulation type **Water-dispersible Granule**

DIRECTIONS FOR USE

Restraints

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application. DO NOT apply by equipment carried on the back of the user.

Rate	Critical Comments
In the following table: "Fruit Crops - Tree and Vine Crops" all rates given are for dilute spraying. For concentrate spraying, refer to the Mixing / Application Section.	For all uses in the table "Fruit Crops - Tree and Vine Crops": Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.

Crop	Pest	States	Rate		WHP	Critical Comments
			Per 100 L	Per ha		
FRUIT CROPS - TREE AND VINE CROPS						
Apples	Woolly Aphid	All States	50 g	1.1 to 1.7 kg	2 days	Use the higher rate when less than 2200 L of spray/ha. Add 18 mL Agral®/100 L.
Citrus	Citrus Aphid			1.1 kg		Apply when aphids appear. Use at least 2200 L spray/ha.
	Aphids		2.5 g	-		Apply only where pest management is being implemented and only when it is apparent that parasite predators will not control the infestation within a reasonable time.
Stone Fruit	Green Peach Aphid, Black Peach Aphid, Cherry Aphid		50 g	1.1 kg		Apply at pink bud or when aphids appear. Use at least 1100 L spray/ha.
FRUIT CROPS - NON-TREE AND VINE CROPS						
Blueberries	Aphids	All States	50 g	-	2 days	-
Pepinos						
Strawberries			35 g			Thorough coverage is necessary. If used in conjunction with predatory mites, apply before release or only after mites have fully established within the crop.
Blackberries (field and protected crops**)**	Aphids including Green Peach Aphid		35 g		7 days	Monitor crops and commence applications once local aphid thresholds are reached. Apply product to coincide with a susceptible life stage (adults and nymphs). DO NOT apply consecutive applications of Pirimor. If treatment is required, use an approved aphicide from a different chemical group before reapplying Pirimor. DO NOT re-apply within 10-14 days of the previous Pirimor spray. DO NOT apply more than 2 applications per year. Apply as a dilute foliar application by ground-based air blast sprayer, boom sprayer or equivalent spray equipment. DO NOT apply using a backpack or knapsack sprayer. Apply in a spray volume up to 900 L/ha. Ensure complete and thorough coverage of target crop. The sensitivity of some aphid species and blackberry varieties has not been fully evaluated. Treat a small number of plants to determine their sensitivity before treating the whole crop.

VEGETABLES						
Crop	**Pest**	**States**	**Rate**		**WHP**	**Critical Comments**
			Per 100 L	**Per ha**		
Artichokes (Globe)	Aphids	All States	50 g	-	2 days	-
Asparagus						
Beetroot	Green Peach Aphid		-	1 kg		Use at least 500 L spray/ha plus 18 mL/100 L Agral.
Beans	Cowpea Aphid (*Aphis craccivora*)			500 g		Apply as beans emerge and thereafter whilst aphid attack continues. Spray thoroughly.
Broad Beans, Lima Beans	Aphids		50 g	-		As above.
Broccoli, Brussels Sprouts, Cabbages, Cauliflowers	Cabbage Aphid	All States	50 g	500 g or 1 kg	2 days	Apply every 10 to 14 days commencing 1 week after planting. Use 500 L spray/ha. Add 18 mL Agral/100 L. If lower spray volumes per ha are used, increase the rate per 100 L to maintain the equivalent rate of product per ha. Use the higher rate if temperature falls below 20°C.
Brussels Sprouts, Cabbages, Cauliflowers	Green Peach Aphid			500 g to 1 kg		
Chinese cabbages	Cabbage Aphid, Green Peach Aphid					
Capsicums	Aphids					Spray when aphids are detected and repeat if necessary. Use the higher rates if applied in conditions below 20°C.
Cucurbits						Apply every 5 to 10 days during periods of rapid plant growth to delay virus buildup. Use the higher rates if applied in conditions below 20°C.
Endive, Garden Cress			50 g	-		-
Kale	Cabbage Aphid, Green Peach Aphid		100 to 200 g	500 g to 1 kg		Apply every 10 to 14 days commencing 1 week after planting. Use 500 L spray/ha. Add 18 mL Agral/100 L. If lower spray volumes per ha are used, increase the rate per 100 L to maintain the equivalent rate of product per ha. Use the higher rate if temperature falls below 20°C.
Kiwano	Aphids					Apply every 5 to 10 days during period of rapid plant growth to delay virus buildup.
Leeks			50 g	-		-
Lettuce			100 to 200 g	500 g to 1 kg		Spray when aphids are detected and repeat if necessary. Use the higher rates if applied in conditions below 20°C.

Crop	Pest	States	Rate		WHP	Critical Comments
			Per 100 L	Per ha		
Okra	Aphids	All States	50 g	-	2 days	For use in IPM programs where Helicoverpa resistance strategy is being practiced.
Peas			100 to 200 g	500 g to 1 kg		Spray when aphids are detected and repeat if necessary. Use the higher rates if applied in conditions below 20°C.
Potatoes	Potato Aphid		250 to 500 g	500 g or 1 kg		Apply in 200 L spray/ha when aphids first appear. If higher spray volumes per ha are used, decrease the rate per 100 L to maintain the equivalent rate of product per ha. Use the higher rate if temperature falls below 20°C.
Radishes	Cabbage Aphid, Green Peach Aphid	All States	100 to 200 g	500 g to 1 kg	2 days	Apply every 10 to 14 days commencing 1 week after planting. Use 500 L spray/ha. Add 18 mL Agral/100 L. If lower spray volumes per ha are used, increase the rate per 100 L to maintain the equivalent rate of product per ha. Use the higher rate if temperature falls below 20°C.
Shallots, Spinach	Aphids		50 g	-		-
Spring onion (field only**)**	Aphids including Green peach aphid (*Myzus persicae*) and Onion aphid (*Neotoxoptera formosana*) and Cabbage aphid		100 to 200 g	500 g to 1 kg		Monitor crops and apply Pirimor when local aphid thresholds are reached and before aphid populations reach high infestation levels. Use a spray volume of 200-500 L/ha, depending on crop density. If lower spray volumes per hectare are used, increase the product rate per 100 L to maintain the equivalent rate per hectare. Use the higher rate if the temperature falls below 20°C. Ensure thorough coverage of all plant surfaces, including the underside leaf surface. Always add 18 mL/100 L of a non-ionic surfactant, e.g Agral. DO NOT apply more than 2 spray applications per crop. DO NOT apply consecutive applications of Pirimor or an insecticide that has the same mode of action within and between seasons or exceed the recommended maximum number of applications in a crop. **DO NOT use on spring onions in protected cropping situations.**
Swedes, Turnips	Cabbage Aphid, Green Peach Aphid		100 to 200 g	500 g to 1 kg		Apply every 10 to 14 days commencing 1 week after planting. Use 500 L spray/ha. Add 18 mL Agral/100 L. If lower spray volumes per ha are used, increase the rate per 100 L to maintain the equivalent rate of product per ha. Use the higher rate if temperature falls below 20°C.

Crop	Pest	States	Rate		WHP	Critical Comments
			Per 100 L	Per ha		
Sweet potato	Aphids including Green peach aphid (*Myzus persicae*), Melon aphid (*Aphis gossypii*) and Cabbage aphid	All States	100 to 200 g	500 g to 1 kg	2 days	Monitor crops and apply Pirimor when local aphid thresholds are reached and before aphid populations reach high infestation levels. Use a spray volume of 200-500 L/ha, depending on crop density. If lower spray volumes per hectare are used, increase the product rate per 100 L to maintain the equivalent rate per hectare. Use the higher rate if the temperature falls below 20°C. Ensure thorough coverage of all plant surfaces, including the underside leaf surface. Always add 18 mL/100 L of non-ionic surfactant e.g. Agral. DO NOT apply more than 2 spray applications per crop. DO NOT apply consecutive applications of Pirimor or an insecticide that has the same mode of action within and between seasons or exceed the recommended maximum number of applications in a crop.
Tomatoes	Aphids					Spray when aphids are detected and repeat if necessary. Use the higher rates if applied in conditions below 20°C.
Watercress	Aphids		50 g	-		-
ORNAMENTALS						
Roses, Chrysanthe-mums	Rose Aphid, Chrysanth-emum Aphid	All States	50 g	-	-	Apply when aphids appear. Spray to give complete coverage of plants.
Ornamentals	Aphids	NSW, WA only		-		
		Qld, WA only	35 g			If applied in conjunction with predatory mites (*Phytoseiulus persimilis*) apply before release or only after mites have become fully established. DO NOT apply more than 2 sprays in quick succession where IPM is practiced. Where roses are grown in dense hedges and IPM is used, spray only the upper half of the hedge where the aphids are attacking the buds. Allow the lower half of the hedge to serve as a refuge for *P. persimilis*. In such situations PIRIMOR WG should NOT be mixed with a fungicide which requires very careful coverage. **Note:** Because of the wide range of crops and pests encompassed under Ornamentals, the phytotoxicity and efficacy of this compound cannot be guaranteed under all uses. Test the chemical for phytotoxicity on a small area of crop before widespread usage.

BROADACRE CROPS						
Crop	**Pests**	**States**	**Rate**		**WHP**	**Critical comments**
			Per 100 L	**Per ha**		
Adzuki bean, Mung bean, Soybean	Cowpea aphid (*Aphis craccivora*) Soya bean aphid (*Aphis glycines*)	All States		300 g	21 days harvest/ grazing	Use a minimum spray volume of 100 L/ha for ground applications and 30-50 L/ha for aerial applications. Use a Medium spray droplet size category. For best coverage with aerial application, apply when a slight crosswind with a minimum speed of 3 km/h is present. DO NOT apply more than 2 applications per season. DO NOT re-apply within 10 days of a previous Pirimor spray. DO NOT apply consecutive applications of Pirimor. If retreatment is required, use an approved aphicide from a different chemical group before reapplying Pirimor. If low relative humidity conditions prevail add 1% or 10% Summer Spray Oil (DC-Tron or Caltex) to the spray solution for ground and aerial application respectively. Soybeans: Monitor aphids during the vegetative stage, but unless crops are stressed and aphid populations are very high, delay spraying to give beneficial insects time to multiply and effect control. Spray if there are more than 250 aphids per plant during R1 - R4 (flowering to pod elongation). As a guideline, if soybean aphids are visible on the plant's main stem, there are likely to be greater than 400 aphids per plant present.
Canola and Mustard (oilseed cultivars) (Brassica juncea)	Cabbage Aphid (*Brevicoryne brassicae*), Green Peach Aphid (*Myzus persicae*)	All States	-	500 g or 1 kg	14 days harvest/ grazing	Apply when aphids first appear using 30 to 100 L water/ha (ground rig) or 20 to 50 L/ha (aerial). Use higher rate if temperature falls below 20 ^{0}C. Add 10 mL Agral/100 L spray.
Cotton	Cotton Aphid (*Aphis gossypii),* Green Peach Aphid *(Myzus persicae)*	All States	-	500 g or 750 g **Aerial Application** Spray volume of 30 to 50 L/ha **Ground Application** Spray volume of 100 L/ha minimum	21 days harvest/ grazing	Thorough spray coverage is essential to produce best results. Stage I - apply when aphids exceed 90% plants infested. Stage II - apply when aphids exceed 95% plants infested. Stage III - apply when aphids exceed 50% plants infested, or - if 1% bolls are open and honeydew is present apply when aphids exceed 10% of plants infested. If a spray is required for another pest, the aphid threshold for each stage should be reduced by half to determine the need to spray for aphids; eg 50% becomes 25% plants infested. Use PIRIMOR WG no more than twice per season. Adhere to the resistance management strategy and rotate chemistry. DO NOT use PIRIMOR WG as consecutive sprays.
Duboisia	Aphids		50 g	1.1 kg	-	Apply when populations warrant.
Crop	**Pests**	**States**	**Rate**		**WHP**	**Critical comments**

			Per 100 L	Per ha		
Lucerne, Medic pastures	Spotted Alfalfa Aphid, Blue Green Aphid	All States	-	100 to 150 g in 200 to 300 L water	3 days grazing	**Application** High rate to be used when there is a high aphid population density or with dense crop growth. Complete coverage and penetration of the crop is essential. **Timing** ***Queensland only***: On seedling lucerne, treatment is necessary **when 1 or 2 aphids are present per plant**. Established lucerne and medics (spring and summer) - spray when there are 20 to 40 aphids per stem. Spotted Alfalfa Aphid (NSW, Tas, SA, WA only): Apply when aphids begin to build up on stems and repeat as necessary. Spotted Alfalfa Aphid and Pea Aphid (Vic only): Seedling lucerne and medics - treat when 1 to 2 aphids per plant are present. Established lucerne and medics (spring and summer) - spray when there are 20 to 40 aphids per stem. ***Tasmania only***: A cold weather adapted species which may be found at most times of the year. Spraying should be carried out when aphids begin to increase and be repeated as necessary. For the control of both the above lucerne and medic pasture pests PIRIMOR WG at the recommended rate should be applied in 200 to 300 L of water/ha. Complete coverage and penetration of the crop is essential. **Blue Green Aphid** This aphid is cooler weather tolerant and will be found at all times of the year. Spraying should be carried out as aphids are observed. In some areas Spotted Alfalfa Aphid has developed tolerance to pirimicarb and in those areas PIRIMOR will not give satisfactory control. However, apparent tolerance should not be confused with poor application. Consult your local Syngenta or Department of Agriculture representative for local information.
Lucerne	Pea Aphid		-	100 to 150 g	3 days grazing	Apply when aphids begin to build up on stems. Repeat as necessary. Use 200 to 300 L/ha. Use the higher rate when there is a high aphid population density or dense crop or when temperature falls below 20°C.

Crop	Pests	States	Rate		WHP	Critical comments
			Per 100 L	Per ha		
Lupins	Green Peach Aphid	NSW, VIC, WA, only	-	250 g	6 weeks harvest/ grazing	Apply when aphids first appear using 20 to 100 L/ha (ground rig) or 4 to 15 L/ha (aerial). Repeat as necessary. If low relative humidity conditions prevail add 1% or 10% Summer Spray Oil (DC-Tron or Caltex) to spray solution for ground and aerial application respectively. Note: For best coverage and hence best results with aerial application, apply when a slight crosswind with a minimum speed of 3 km/hr is present.
	Cowpea Aphid					
	Green Peach Aphid	WA only		300 g		Apply when aphids first appear. Good coverage of the infested area is essential. Apply in 30 to 100 L/ha for aerial application. Repeat as necessary, if low relative humidity conditions prevail add Summer Spray Oil to the mix.
Vetch	Bean aphid (*Megoura crassicauda*)	All States		250 g		Apply as a foliar spray when aphids reach local thresholds, using a boom sprayer or similar equipment, in a sufficient water volume to ensure good coverage of the target crop. DO NOT apply more than 2 applications per crop. DO NOT re-apply within 15 days of a previous Pirimor spray.
Winter Cereals	Aphids (*Rhopalosip hum maidis*)	All States	-	150 g		Apply when 20 or more aphids are present per tiller and when the crop is between emergence of last leaf and flowering stages.
	Aphids (*R. maidis*, *R. padi*)	WA only		250 to 300 g		Use 20 to 100 L/ha (ground rig) or 4 to 15 L/ha (aerial). If low relative humidity conditions prevail add Summer Spray Oil as for lupins. Where there is concern for infection by Barley Yellow Dwarf virus, spraying should be carried out as soon as aphids are located in the crop.
	Russian wheat aphid *(Diuraphis noxia)*	All States		225 to 300 g		Begin applications when action thresholds are reached. The higher rate is recommended under conditions of higher pest pressure and/or lower temperatures. Thorough coverage is essential: ensure water volume and spray quality is adequate to thoroughly cover all parts of the crop canopy, including the undersides of, and in between leaves. Addition of a non-ionic surfactant (eg Agral, BS1000) or esterified crop oil (eg Hasten) may assist control. Continue to monitor crops and make further insecticide applications as necessary.

NURSERY STOCK						
Crop	**Pests**	**States**	**Rate**		**WHP**	**Critical comments**
			Per 100 L	**Per ha**		
Nursery stock [(non-food) – seedlings, plugs, potted colour, trees, shrubs, foliage plants, palms, grasses, fruit trees (non-bearing)], cut flowers and foliage	Aphids	-	Soil drench 50 g / 100 L			Apply at the first signs of infestation, or when pest thresholds are reached. Apply product as a targeted drench to soil surface. DO NOT use more than twice in quick succession where IPM is used or in consecutive treatments. The sensitivity of some species and varieties of the crops to be treated has not been fully evaluated. Treat a small number of plants to ascertain their reaction before treating the whole crop.
Nursery stock [(non-food) – seedlings, plugs, potted colour, trees, shrubs, foliage plants, palms, grasses, fruit trees (non-bearing)]		-	Foliar application 35 - 50 g / 100 L			Apply at the first signs of infestation, or when pest thresholds are reached. Ensure spray application provides thorough coverage of all plant foliage. Adjust water spray volume to crop density and growth stage. Use correctly calibrated ground application equipment. Use the higher rate when longer residual control is required, or during periods of high pest pressure or rapid crop growth, or when crops are well advanced, or when insect infestation is high, or when conditions favour high pest pressure. Continue to monitor crops and make a subsequent application as necessary. DO NOT use more than two (2) applications per crop with a minimum re-treatment interval of 14 days. The sensitivity of some species and varieties of the crops to be treated has not been fully evaluated. Treat a small number of plants to ascertain their reaction before treating the whole crop.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS

Adzuki bean, Mung bean,	**DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION.**
Soybean:	**DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION**
Blackberries:	**DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION**
***Fruit, Vegetables*:**	**DO NOT HARVEST FOR 2 DAYS AFTER APPLICATION**
Sweet potato:	**DO NOT GRAZE OR CUT TREATED CROPS FOR LIVESTOCK FEED**
***Canola, Mustard*:**	**DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION**
Cotton:	**DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION**
Lucerne, Medic Pastures:	**DO NOT GRAZE, CUT FOR STOCKFOOD OR FEED TO POULTRY FOR 3 DAYS AFTER APPLICATION**
***Lupins, Winter Cereals*:**	**DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 6 WEEKS AFTER APPLICATION**
***Vetch*:**	**DO NOT GRAZE OR CUT FOR LIVESTOCK WITHIN 6 WEEKS OF APPLICATION**
Nursery stock, cut flowers and foliage (when used as a soil drench):	***DO NOT GRAZE TREATED CROPS OR FEED MATERIAL CUT FROM TREATED CROPS TO LIVESTOCK, INCLUDING POULTRY***

GENERAL INSTRUCTIONS

Mixing
Fill tank 50 to 75% full then start agitation. Slowly pour granules into tank and maintain agitation throughout spraying. DO NOT pre-mix PIRIMOR granules.

Application
If this product is applied when the temperature is less than 15°C it is essential to have thorough spray coverage for quick and total aphid control.

Fruit Crops - Tree and Vine Crops
Dilute Spraying:
Use a sprayer designed to apply high volumes of water up to the point of runoff and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of runoff. Avoid excessive runoff. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of runoff. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying:
Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of runoff) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (See *Dilute Spraying*) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way:

Example Only:
1. Dilute spray volume as determined above: For example 1500 L/ha.
2. Your chosen concentrate spray volume: For example 500 L/ha.
3. The concentration factor in this example is: 3 X (ie 1500 L ÷ 500 L = 3)
4. If the dilute label rate is 10 mL / 100 L, then the concentrate rate becomes 3 x 10, that is 30 mL / 100 L of concentrate spray.

The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to change as the crop grows. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

Compatibility
PIRIMOR WG may be mixed in the spray vat with any one of the following products: Agral, BS1000, Hasten, Omite or Summer Spray Oil (DC-Tron or Caltex). As formulations of other manufacturer's products are beyond the control of Syngenta Australia, and the quality of water may vary with location, all mixtures should be tested prior to mixing commercial quantities. Tank mixtures of PIRIMOR WG with more than one other product are not recommended and may result in inadequate pest control or crop damage.

Insecticide Resistance Warning | **GROUP 1A INSECTICIDE**

For insecticide resistance management PIRIMOR WG Aphicide is a Group 1A insecticide. Some naturally occurring insect biotypes resistant to PIRIMOR WG and other Group 1A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if PIRIMOR WG or other Group 1A insecticides are used repeatedly. The effectiveness of PIRIMOR WG on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Syngenta Australia Pty Ltd accepts no liability for any losses that may result from the failure of PIRIMOR WG to control insects. PIRIMOR WG may be subject to specific resistance management strategies. For further information contact your local supplier, Syngenta representative or local agricultural department agronomist.

Resistance Management for Green Peach Aphid Control
Some Green Peach Aphids are resistant to all commonly used insecticides including PIRIMOR WG. If resistance is known or suspected, DO NOT use PIRIMOR WG and seek advice on resistance management. If there is no PIRIMOR WG resistance, use a maximum of 2 non-consecutive aphid sprays per season. If PIRIMOR WG was used as the last aphid spray for the season, DO NOT use it as the first aphid spray next season.

Trade Advice
Growers should note that maximum residue limits (MRLs) or import tolerances may not exist in all markets for edible produce treated with pirimicarb. If you are growing edible produce for export, please check with industry representative or Syngenta Australia Pty Ltd for the latest information on MRLs and import tolerances before using this product.

PRECAUTIONS:

Re-entry Period
DO NOT enter treated areas until the spray has dried. If prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). Clothing must be washed after each day's use.

PROTECTION OF LIVESTOCK

Animals and poultry should be kept away from orchards and other treated areas during the horticultural spraying season.
Keep animals and poultry away from spray and treated areas for a minimum of 3 days.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.
DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto adjacent areas, particularly wetlands, waterbodies or watercourses.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area out of direct sunlight.

Plastic containers
Triple rinse containers before disposal. Add rinsings to the mixing tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with Local, State or Territory government regulations. DO NOT burn empty containers or product.

Lined cardboard boxes
Single rinse or shake remainder into spray tank. DO NOT dispose of undiluted chemicals on site. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Harmful if absorbed by skin contact, inhaled or swallowed. May irritate the nose and throat and will irritate the eyes. Avoid contact with eyes and skin. DO NOT inhale dust or spray mist. When opening the container and preparing product for use, wear:

- **elbow length chemical resistant gloves, and**
- **goggles**

If applying by aerial spraying equipment wear gloves
If applying by high pressure hand wand, wear

- **elbow length chemical resistant gloves and**
- **googles**

If applying by low pressure hand wand, wear

- **cotton overalls, over normal clothing, buttoned to neck and wrist and a washable hat,**
- **elbow length chemical resistant gloves,**
- **goggles and**
- **a half face piece respirator with organic vapour/gas cartridge or canister.**

When opening the container, preparing the product for use and applying by one-cab boom spray equipment or open-cab airblast equipment, wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat, elbow-length chemical resistant gloves, goggles.

If clothing becomes contaminated with product, or wet with spray remove clothing immediately. If product or spray on skin, immediately wash area with soap and water. If product or spray in eyes wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles, respirator (if rubber wash with detergent and warm water) and contaminated clothing.

FIRST AID

If swallowed, splashed on skin or in eyes, or inhaled, contact the Poisons Information Centre (Phone 131 126) or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed.

SAFETY DATA SHEET

If additional hazard information is required refer to the Safety Data Sheet. For a copy phone 1800 067 108, or visit our website at www.syngenta.com.au

Suspected of causing cancer.

NOTE: A varying degree of consolidation of the contents of this container may occur during transit but the net weight of the contents is not changed and is as stated on the label.

DISCLAIMER

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability.

Product names marked ® or ™, the ALLIANCE FRAME the SYNGENTA Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company

Scan here for more information including SDS and label

Batch Number	
Date of Manufacture	

UN 2757
CARBAMATE PESTICIDES, SOLID, TOXIC, N.O.S.
(PIRIMICARB)
MARINE POLLUTANT
PACKING GROUP III
HAZCHEM 2X

In a transport emergency dial 000, Police or Fire Brigade
For specialist advice in an emergency only,
call 1800 033 111 (24 hours)