# POISON

**KEEP OUT OF REACH OF CHILDREN**
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# Simodis®

**PLINAZOLIN® technology Insecticide**

syngenta®

**ACTIVE CONSTITUENT: 100 g/L ISOCYCLOSERAM**

| GROUP | 30 | INSECTICIDE |
|---|---|---|

For control of certain insect and mite pests in Vegetables as per the Directions for Use

***Syngenta Australia Pty Ltd***
*Level 1, 2 Lyonpark Road, Macquarie Park NSW 2113*

***In a transport emergency dial 000, Police or Fire Brigade***
***For specialist advice in an emergency only, call 1800 033 111 (24 hours)***

***APVMA Approval No: 89460/124955***

| DC | Formulation Type **Dispersible Concentrate** |
|---|---|

## DIRECTIONS FOR USE

**Restraints:**

DO NOT apply by aircraft
DO NOT apply if heavy rains or storms that are likely to cause runoff are forecast within 3 days
DO NOT apply to bulb vegetables in Mackay/Whitsunday in August to end of December
DO NOT apply to leafy vegetables in Mackay/Whitsunday in October to end of November
DO NOT irrigate to the point of water runoff for at least 3 days after application
DO NOT apply more than 2 applications of SIMODIS® per crop
DO NOT apply to nursery crops

**Spray Drift Restraints**

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the buffer zone table below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

- Spray droplets are not smaller than a COARSE spray droplet size category.
- Minimum distances between the application site and downwind sensitive areas are observed (see 'Mandatory downwind buffer zones' section of the following table titled 'Buffer zones for boom sprayers').

***Buffer zones for boom sprayers***

| Application Rate | Boom Height Above the Target Canopy | Mandatory Downwind Buffer Zones | | | | |
|---|---|---|---|---|---|---|
| | | Natural Aquatic Areas | Vegetation Areas | Bystander Areas | Pollinator Areas | Livestock Areas |
| Up to 300 mL/ha | 0.5 m or lower | 350 metres | 0 metres | 0 metres | 0 metres | 0 metres |

DO NOT apply by a vertical sprayer unless the following requirements are met:

- Spray is not directed above the target canopy
- The outside of the sprayer is turned off when turning at the end of rows and when spraying the outer row on each side of the application site
- For dilute water rates up to the maximum listed for each type of canopy specified, minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for vertical sprayers') are observed.

***Buffer zones for vertical sprayers***

| Application Rate | Mandatory Downwind Buffer Zones | | | | |
|---|---|---|---|---|---|
| | Natural Aquatic Areas | Vegetation Areas | Bystander Areas | Pollinator Areas | Livestock Areas |
| Up to 300 mL/ha | 75 metres | 0 metres | 0 metres | 0 metres | 0 metres |

| Crop | Pest | Rate | Critical Comments |
|---|---|---|---|
| **Brassica Vegetables** (Broccoli, Broccolini, Brussels Sprouts, Cabbage, Cauliflower, Kohlrabi)<br><br>**Brassica Leafy Vegetables** including Bok Choy, Chinese Broccoli (Gai Lum / Gai Lan / Kai Lan), Chinese Cabbage (Pet Sai / Wombok / Haksukai), Choy Sum, Gai Choy / Am Soy / Kai Choy, Kale, Mibuna, Mustard (Leafy) including Indian Mustard and Mustard Spinach (Komatsuma), Pak Choy, Tat Soy | Diamondback Moth (*Plutella xylostella*), Cabbage White Butterfly (*Pieris rapae*)<br><br>**Suppression of** Heliothis (*Helicoverpa* spp.) | 300 mL/ha plus adjuvant [Δ] | Monitor crops and commence SIMODIS® applications immediately once local economic spray thresholds are reached. Apply at egg hatch or very soon after egg hatch to target young larvae. Avoid applying SIMODIS® to established populations dominated by large, late instar larvae.<br><br>Continue to monitor crops and make subsequent applications as required. DO NOT re-apply within 7 days of a previous SIMODIS® spray.<br><br>DO NOT apply more than 2 applications per crop.<br><br>Ensure thorough coverage of the target crop – refer to the Application section in GENERAL INSTRUCTIONS.<br><br>[Δ]Always add a specified non-ionic surfactant – refer to the Adjuvant section in GENERAL INSTRUCTIONS.<br><br>**Note:** This use is subject to a CropLife resistance management strategy. Refer to www.croplife.org.au for more information. |
| **Bulb Vegetables** (including Garlic, Onion Bulb, Spring Onion and Leeks) | **Suppression of** Onion Thrips (*Thrips tabaci*), Plague Thrips (*Thrips imaginis*) | 300 mL/ha plus adjuvant [Δ] | Monitor crops and commence SIMODIS® applications once local economic spray thresholds are reached.<br><br>Continue to monitor crops and make subsequent applications as required. DO NOT re-apply within 7 days of a previous SIMODIS® spray.<br><br>DO NOT apply more than 2 applications per crop.<br><br>Ensure thorough coverage of the target crop – refer to the Application section in GENERAL INSTRUCTIONS.<br><br>[Δ]Always add a specified adjuvant – refer to the Adjuvant section in GENERAL INSTRUCTIONS. |

| Crop | Pest | Rate | Critical Comments |
|---|---|---|---|
| **Cucurbits**<br>Open field only | Two-spotted Mite<br>(*Tetranychus urticae*)<br><br>**Suppression of**<br>Broad Mite<br>(*Polyphagotarsonemus latus*),<br>Bean Red Spider Mite<br>(*Tetranychus ludeni*) | 300 mL/ha plus adjuvant [Δ] | Monitor crops and commence SIMODIS® applications immediately once local economic spray thresholds are reached. Best results will be achieved when spray is applied to low mite populations. Avoid applying SIMODIS® to high established mite populations.<br>Use the higher application rate where extended residual control is required.<br><br>To manage insecticide resistance, DO NOT apply sequential applications of SIMODIS® for mite control. If retreatment is required, apply a miticide from a different chemical group, before using SIMODIS® again.<br><br>DO NOT apply more than 2 applications per crop.<br><br>Ensure thorough coverage of the target crop – refer to the Application section in GENERAL INSTRUCTIONS.<br><br>[Δ]Always add a specified non-ionic surfactant – refer to the Adjuvant section in GENERAL INSTRUCTIONS. |
| | Cucumber Moth<br>(*Diaphania indica*)<br><br>**Suppression of**<br>Western Flower Thrips<br>(*Frankliniella occidentalis*),<br>Tomato Thrips<br>(*Frankliniella schultzei*),<br>Melon Thrips<br>(*Thrips palmi*),<br>Plague Thrips<br>(*Thrips imaginis*),<br>Heliothis<br>(*Helicoverpa* spp.) | | Monitor crops and commence SIMODIS® applications once local economic spray thresholds are reached.<br><br>**Heliothis:** Apply at egg hatch or very soon after egg hatch to target young larvae.<br><br>Continue to monitor crops and make subsequent applications as necessary. DO NOT re-apply within 7 days of a previous SIMODIS® spray.<br><br>DO NOT apply more than 2 applications per crop.<br><br>Ensure thorough coverage of the target crop – refer to the Application section in GENERAL INSTRUCTIONS.<br><br>[Δ]Always add a specified non-ionic surfactant – refer to the Adjuvant section in GENERAL INSTRUCTIONS.<br><br>**Note:** This use is subject to a CropLife resistance management strategy. Refer to www.croplife.org.au for more information. |

| Crop | Pest | Rate | Critical Comments |
|---|---|---|---|
| **Fruiting Vegetables** (including Capsicum, Chilli, Eggplant and Tomato)<br><br>**Open field only** | Two-spotted Mite (*Tetranychus urticae*), Broad Mite (*Polyphagotarsonemus latus*)<br><br>**Suppression of** Tomato Russet Mite (*Aculops lycopersici*), Western Flower Thrips (*Frankliniella occidentalis*), Tomato Thrips (*Frankliniella schultzei*), Melon Thrips (*Thrips palmi*), Plague Thrips *(Thrips imaginis),* Heliothis (*Helicoverpa* spp.) | 300 mL/ha plus adjuvant [Δ] | Monitor crops and commence SIMODIS® applications once local economic spray thresholds are reached. Best results will be achieved when spray is applied to low mite populations. Avoid applying SIMODIS® to high, established mite populations.<br><br>**Heliothis:** Apply at egg hatch or very soon after egg hatch to target young larvae.<br><br>To manage insecticide resistance, DO NOT apply sequential applications of SIMODIS® for mite control. If retreatment is required, apply a miticide from a different chemical group, before using SIMODIS® again.<br><br>For thrips and Heliothis, continue to monitor crops and make subsequent applications as necessary. DO NOT re-apply within 7 days of a previous SIMODIS® spray.<br><br>DO NOT apply more than 2 applications per crop.<br><br>Ensure thorough coverage of the target crop – refer to the Application section in GENERAL INSTRUCTIONS.<br><br>[Δ]Always add a specified non-ionic surfactant – refer to the Adjuvant section in GENERAL INSTRUCTIONS.<br><br>**Note:** This use is subject to a CropLife resistance management strategy. Refer to www.croplife.org.au for more information. |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIODS:**
***Brassica Vegetables, Brassica Leafy Vegetables:***
**Harvest: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION**
**Grazing: DO NOT USE TREATED CROP, CROP WASTE OR PRODUCE FOR STOCK FOOD**
***Bulb Vegetables:*** **DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION**
***Cucurbits and Fruiting Vegetables:***
**Harvest: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION**
**Grazing: DO NOT USE TREATED CROP, CROP WASTE OR PRODUCE FOR STOCK FOOD**

## GENERAL INSTRUCTIONS

**Mixing**
SIMODIS® PLINAZOLIN® technology Insecticide is a Dispersible Concentrate (DC) formulation that mixes readily with water and is applied as a foliar spray.
Measure the required amount of SIMODIS®, add to the partly filled spray tank, and then add the remainder of the water.

**Adjuvant**
***Vegetables (except Bulb Vegetables):*** Apply SIMODIS® with a non-ionic adjuvant such as AGRAL® Spray Adjuvant[Δ] at 10 mL per 100 L of spray mixture.
***Bulb vegetables (including Bulb Onions, Garlic, Spring Onions and Leeks):*** Apply SIMODIS® with Maxx Organosilicone Surfactant*[Δ] at 60 mL per 100 L of spray mixture.
[Δ] or other specified adjuvant - refer to Syngenta Australia Pty Ltd representative for information

**Application**
Thorough coverage of the target area is essential. Apply in sufficient water and adjust water volumes according to the crop growth stage. Use suitable application parameters (nozzles, pressure, boom height, speed, etc.) to ensure thorough and even coverage. Use only COARSE spray droplets according to nozzle manufacturers specifications that refer to the ASAE S572 standard or the BCPC guideline.

**Compatibility**
SIMODIS® may be mixed with the following crop protection products: ORONDIS® Flexi Fungicide, AMISTAR® 250 SC Fungicide, AGRAL® Spray Adjuvant, ADIGOR® Spray Adjuvant and Maxx Organosilicone Surfactant*. For the latest information on the compatibility of SIMODIS® with other products, contact your local Syngenta representative.

**Insecticide Resistance Warning** | **GROUP** | **30** | **INSECTICIDE**

For insecticide resistance management, SIMODIS® PLINAZOLIN® technology Insecticide is a Group 30 insecticide. Some naturally occurring insect biotypes resistant to SIMODIS® and other Group 30 insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if SIMODIS® or other Group 30 insecticides are used repeatedly. The effectiveness of SIMODIS® on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Syngenta Australia Pty Ltd accepts no liability for any losses that may result from the failure SIMODIS® to control resistant insects. For further information contact your local supplier, Syngenta representative or local agricultural department agronomist.

In order to avoid or delay the selection of resistant insects, Group 30 insecticides should be used as part of an insecticide resistance management (IRM) strategy which incorporates the following:

- Insecticides from the same mode of action group should not be used to treat successive generations of the target pest.
- Multiple applications of SIMODIS® and other products containing Group 30 insecticides may be applied successively but only when targeting a single generation of the target insect.
- DO NOT apply sequential applications of SIMODIS® for mite control. If retreatment is required, apply a miticide from a different chemical group before using SIMODIS® again.
- If more than one application of an insecticide is required to control successive generations of the target pest then alternative insecticides with different modes of action should be utilised in rotation with SIMODIS®.
- Where possible incorporate alternative methods of pest control as part of an integrated pest management (IPM) approach.

## PRECAUTIONS
**Re-entry Period:** DO NOT enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

## PROTECTION OF HONEY BEES AND OTHER INSECT POLLINATORS

Toxic to bees and harmful to bee brood. DO NOT apply to crops from the onset of flowering until flowering is complete unless the application is made in the time period between 2 hours prior to sunset and 8 hours prior to sunrise. DO NOT allow spray drift to flowering weeds or flowering crops in the vicinity of the treatment area. Before spraying, notify beekeepers to move hives to a safe location with an untreated source of nectar and pollen, if there is potential for managed hives to be affected by the spray or spray drift.

## INTEGRATED PEST MANAGEMENT

Toxic to beneficial arthropods. Not compatible with integrated pest management (IPM) programs utilising beneficial arthropods. Minimise spray drift to reduce harmful effects on beneficial arthropods in non-crop areas.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

## STORAGE AND DISPOSAL

Store in the closed original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight.

Triple rinse containers before disposal. Add rinsings to treatment tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

## SAFETY DIRECTIONS

**Will irritate the eyes. Avoid contact with eyes and skin. When opening the container, mixing and preparing spray and using the prepared spray, wear**

- **cotton overalls buttoned to the neck and wrist (or equivalent clothing) and**
- **goggles or safety glasses.**

**Wash hands after use. After each day's use wash goggles or safety glasses and contaminated clothing.**

## FIRST AID

**If poisoning occurs contact a doctor or Poisons Information Centre. Phone 131 126.**

## WARNING - Suspected of damaging the unborn child.

## SAFETY DATA SHEET

If additional hazard information is required refer to the Safety Data Sheet. For a copy visit our website at www.syngenta.com.au or use the QR code on this label.

## DISCLAIMER

This product complies with the specifications in its statutory registration. Implied terms and warranties are excluded. Syngenta's liability for breach of the express or any non-excludable implied warranty is limited to product replacement or purchase price refund. The purchaser must determine suitability for intended purpose and take all proper precautions in the handling, storage and use of the product including those on the label and/or safety data sheet failing which Syngenta shall have no liability.

Product names marked ® or ™, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company

*Trademark

APVMA Approval No: 89460/124955

Scan here for more information including SDS and label