Australian Pesticides & Veterinary Medicines Authority

Product Name: SWITCH FUNGICIDE
APVMA Approval No: 51797/148814

| | |
|---|---|
| Label Name: | SWITCH FUNGICIDE |

| | |
|---|---|
| Signal Headings: | CAUTION<br>KEEP OUT OF REACH OF CHILDREN<br>READ SAFETY DIRECTIONS BEFORE OPENING OR USING |

| | |
|---|---|
| Constituent Statements: | ACTIVE CONSTITUENTS<br>375 g/kg CYPRODINIL<br>250 g/kg FLUDIOXONIL |

| | |
|---|---|
| Mode of Action: | GROUP 9 12 FUNGICIDE |

| | |
|---|---|
| Statement of Claims: | For the control of various fungal diseases in Alliums, Apples, Capsicum, Cucumber, Cut Flowers, Garlic, Grapes, Green Beans, Green Peas, Leafy Vegetables, Lettuce, Nursery Stock, Ornamentals, Pyrethrum and Strawberries as per the Directions for Use |

| | |
|---|---|
| Net Contents: | 1 kg - 5 kg |

| | |
|---|---|
| Restraints: | This section contains file attachment. |

| | |
|---|---|
| Directions for Use: | This section contains file attachment. |

| | |
|---|---|
| Other Limitations: | |

| | |
|---|---|
| Withholding Periods: | WITHHOLDING PERIODS<br><br>HARVEST<br>Strawberries, Cucumbers: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION |

| | |
|---|---|
| | Capsicums, Green Beans, Lettuce, Leafy Vegetables, Onions, Alliums: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION<br><br>Apples, Green Peas: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION. DO NOT PROCESS INTO STOCK FOOD FOR 14 DAYS AFTER APPLICATION.<br><br>Grapes: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION<br><br>Nursery stock (non-food, Ornamentals and Cut Flowers: NOT REQUIRED WHEN USED AS DIRECTED<br><br>GRAZING<br>Green Beans, Green Peas: DO NOT FEED GREEN BEAN OR GREEN PEA FORAGE OR FODDER TO LACTATING DAIRY ANIMALS PRODUCING MILK FOR HUMAN CONSUMPTION<br><br>Apple Orchards: DO NOT ALLOW LIVESTOCK TO GRAZE TREATED ORCHARDS OR CUT PASTURE WITHIN TREATED ORCHARDS FOR STOCK FOOD |

| | |
|---|---|
| Trade Advice: | EXPORT OF TREATED PRODUCE<br>Growers should note that some export markets have either no Maximum Residue Limits (MRLs) or different MRLs to those in Australia for crops treated with SWITCH fungicide. Treated crop commodities destined for export may require extra time being allowed between application and harvest. if you are growing edible crops for export, please check with Syngenta and/or your industry body for the latest information on MRLs, overseas standards or any export interval before using SWITCH. |

| | |
|---|---|
| General Instructions: | GENERAL INSTRUCTIONS<br>SWITCH fungicide is a water dispersible granule containing cyprodinil and fludioxonil. Cyprodinil is systemic compound that is taken up into the cuticle and waxy layers of leaves and fruit. From there it is distributed and penetrates into other parts of the plant. Fludioxonil stays mainly on the leaf and fruit surfaces.<br><br>Application<br>Ground Application Only: Ensure thorough coverage of foliage and fruit (or bunches) using equipment delivering a MEDIUM spray quality (except for vineyard and orchard airblast sprayers).<br><br>Dilute Spraying<br>Use a sprayer designed to apply high volumes of water up to the point of runoff and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of runoff. Avoid excessive runoff. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Directions for Use table for each 100 L of water. Spray to the point of runoff. The required dilute spray volume will change, and the sprayer set up and operation may also need to be changed, as the crop grows.<br>Concentrate Spraying<br>Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of runoff) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute<br>spray volume (see Dilute Spraying) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way:<br>Example Only<br>1. Dilute spray volume as determined above: for example, 1000L/ha<br><br>2. Your chosen concentrate spray volume: for example, 400 L/ha<br>3. The concentration factor in this example is: 2.5 X (ie 1000 L # 400 L = 2.5) |

| | |
|---|---|
| | 4. If the dilute label rate is 80 g/100 L, then the concentrate rate becomes 2.5 x 80, that is 200 g/100 L of concentrate spray.<br>The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training, and follow industry Best Practices.<br><br>Mixing<br>SWITCH is a water dispersible granule (WG fungicide that mixes readily with water. Partly fill the spray tank with water. Start the agitation and add the correct amount of product to the spray tank with the agitation system running. Continue agitation while topping up the spray tank with water. Add a non-ionic surfactant at registered rates (where required at the completion of filling of the spray tank. Continue agitation while spraying.<br><br>Incompatibility/Tank Mixing:<br>DO NOT mix SWITCH fungicide with DELAN* 700 WG FUNGICIDE or any other products containing DITHIANON as the active ingredient. As formulations of other manufacturers' products are beyond the control of Syngenta, and the quality of water may vary with location, all other mixtures should be tested prior to mixing commercial quantities. For further information on product compatibility, consult your local Syngenta representative.<br><br>When mixing SWITCH and other water dispersible granule (WG) or wettable powder (WP) formulations, ensure they are added and mixed well prior to adding emulsifiable concentrate (EC) or suspension concentrate (SC) products. Wettable powder (WP) formulations should be pre-mixed separately and then added to the spray tank. |

| | |
|---|---|
| Resistance Warning: | SWITCH Fungicide is a combination of an anilinopyrimidine and a phenylpyrrole fungicide. For fungicide resistance management, SWITCH is both a Group 9 and a Group 12 fungicide. Some naturally occurring individual fungi resistant to SWITCH and other Group 9 and Group 12 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or other Group 9 and Group 12 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use Syngenta Australia Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.<br>RESISTANCE MANAGEMENT<br>In order to minimise the likelihood of the development of fungicide resistance, it is recommended that SWITCH be used in a program incorporating all available forms of crop protection. SWITCH should be applied as part of a protective spray program using fungicides from different mode of action group/s. Alternating or mixing fungicides with different modes of actions is a recognised disease resistance management strategy. DO NOT wait until disease levels have built up before commencing applications as this reduces the effectiveness of control and increases the risk of resistance development. Disease control may be reduced if strains of pathogens less sensitive to SWITCH Fungicide develop.<br>SWITCH Fungicide should be applied as specified in the Directions for Use in association with the following CropLife Fungicide Resistance Management Strategies:<br>Grapes: If three or fewer bunch rot sprays are applied in a season, use only one spray containing a Group 9 fungicide. If four or more bunch rot sprays are applied in a season, use a maximum of two sprays containing a Group 9 fungicide.<br>Capsicum, cucumbers, and strawberries: If three or fewer Botrytis fungicide sprays are applied per crop, use only one spray containing a Group 9 fungicide. If four to six sprays are applied per crop use a maximum of two sprays containing a Group 9 fungicide. If seven or more sprays are applied per crop use a maximum of three sprays containing a Group 9 fungicide.<br>Note: In the absence of an established resistance management strategy for a particular crop/disease situation, it is recommended that:<br>• The use of fungicides from any given activity group (excluding Group M) be limited to a maximum of 1/3 of the total number of fungicide applications per crop.<br>• A maximum of two consecutive applications of SWITCH are to be applied, including from one season/crop to the next. |

| | |
|---|---|
| Precautions: | PRECAUTION<br>RE-ENTRY PERIOD:<br>After application, DO NOT enter treated areas until the spray has dried unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use. |

| | |
|---|---|
| Protections: | INTEGRATED PEST MANAGEMENT<br>Based on the available information it cannot be ruled out that SWITCH may have an adverse effect on beneficial invertebrates where integrated pest management is practiced.<br><br>PROTECTION OF HONEY BEES AND OTHER INSECT POLLINATORS<br>DO NOT spray apple trees while bees are actively foraging. DO NOT allow spray to drift from apple trees onto flowering weeds or flowering crops in the vicinity of the apple orchard.<br><br>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT<br>Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.<br><br>TO AVOID CROP DAMAGE<br>Nursery stock and cut flowers: This product has demonstrated toxicity under certain situations, application methods, application rates and weather conditions to nursery stocks including Impatiens, New Guinea Impatiens, Geraniums, Ferns, Anemone, Begonia, Kalanchoe, Saint Paulia and Exacum affine. This product may also leave undesirable spray residue on foliage and flowers. It is recommended that users treat a sample number of plants or a small area prior to whole crop treatment and monitor any phytotoxic effects that will compromise production goals. This action cannot guarantee crop safety as application, environmental and crop conditions may vary from test treatment to whole of crop treatment. |

| | |
|---|---|
| Storage and Disposal: | STORAGE AND DISPOSAL<br>Store in the closed, original container in a cool, dry, well ventilated area, out of direct sunlight.<br>Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation, and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product. |

| | |
|---|---|
| Safety Directions: | SAFETY DIRECTIONS<br>Harmful if inhaled or swallowed. May irritate the nose and throat. Will irritate the eyes. Repeated exposure may cause allergic disorders. Avoid contact with eyes. DO NOT inhale dust or spray mist. When opening the container and preparing spray, and using the prepared spray wear:<br>• cotton overalls buttoned to the neck and wrist (or equivalent clothing)<br>• elbow length chemical resistant gloves<br>• face shield or goggles<br>If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves and contaminated clothing. |

| | |
|---|---|
| First Aid Instructions: | FIRST AID<br>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 11 26. |

| | |
|---|---|
| First Aid Warnings: | |

## DIRECTIONS FOR USE

| Crop/ Situation | Disease | Rate | WHP | Critical Comments |
|---|---|---|---|---|
| **Apples** | **Suppression of** Alternaria leaf and fruit blotch (*Alternaria* spp.) | 80 g/100 L | 14 days (H)<br><br>DO NOT graze treated orchards or cut pasture within treated orchards for stock food (G) | Apply SWITCH fungicide prior to the onset of disease. Repeat applications at 7 to 14 day intervals if conditions continue to remain favourable for disease development.<br><br>DO NOT apply more than 3 applications of SWITCH per crop.<br><br>DO NOT apply any post-harvest treatment which contains fludioxonil such as SCHOLAR fungicide. |
| **Capsicums** | Grey mould (*Botrytis cinerea*)<br><br>Sclerotinia rot (*Sclerotinia minor*, *S. sclerotiorum*) | 100 g/100L or 1 kg/ha | 7 days (H) | Apply prior to disease development.<br><br>Do not apply more than 2 applications per crop with an interval of 7 to 14 days between applications. Use the shorter spray interval under high disease pressure or when continual wet, cloudy conditions favour disease development. |
| **Cucumbers** | Grey mould (*Botrytis cinerea*) | 800 g to 1 kg/ha | 3 days (H) | Apply SWITCH fungicide prior to the onset of disease. Repeat applications at 7 to 10 day intervals if conditions continue to remain favourable for disease development. Use a 7 day spray interval and high rate under high disease pressure.<br><br>DO NOT apply more than 3 applications of SWITCH per crop. |
| **Cut Flowers** | Grey mould (*Botrytis* spp.) | 80 g/100 L or 800 g/ha plus 0.02% non-ionic surfactant | Not required when used as directed | Apply before disease appears when conditions favour infection.<br><br>Apply by low or high volume spray.<br><br>Apply a maximum of 2 applications within an annual production cycle.<br><br>For resistance management this product should form part of a disease management strategy.<br><br>See **To Avoid Crop Damage** |
| **Grapes except grapes grown for dried fruit production** | Grey mould (*Botrytis cinerea*) | **Dilute spraying** 80 g/100 L | 4 weeks (H) | DO NOT apply more than 2 applications of SWITCH fungicide per season. Applications can be made at 5% capfall, 80% capfall, pre-bunch closure or veraison.<br><br>Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.<br><br>DO NOT use at rates greater than 200 g/100 L of water ie at concentration factors above 2.5X.<br><br>This use is subject to a CropLife fungicide resistance management strategy (see **Resistance Management**). |

| Crop/ Situation | Disease | Rate | WHP | Critical Comments |
|---|---|---|---|---|
| **Green Beans** | Grey mould (*Botrytis cinerea*)<br><br>Sclerotinia (*Sclerotinia minor* and *Sclerotinia sclerotiorium*) | 800 g to 1 kg/ha | 7 days (H)<br><br>DO NOT feed forage or fodder to lactating dairy animals producing milk for human consumption (G) | Apply SWITCH fungicide prior to the onset of disease. Apply a second application 7 to 10 days later if conditions continue to remain favourable for disease development. Use a 7 day spray interval and high rate under high disease pressure.<br><br>DO NOT apply more than 2 consecutive applications of SWITCH, with no more than 3 applications of SWITCH per crop. |
| **Green Peas** including garden peas, snow peas and sugar snap peas | | | 14 days (H)<br><br>DO NOT feed forage or fodder to lactating dairy animals producing milk for human consumption. DO NOT process into stock food for 14 days after application. (G) | Apply SWITCH fungicide prior to the onset of disease. Apply a second application 7 to 10 days later if conditions continue to remain favourable for disease development. Use a 7 day spray interval and high rate under high disease pressure.<br><br>DO NOT apply more than 2 applications of SWITCH per crop. |
| **Leafy Vegetables** including Asian leafy greens, chard, Chinese cabbage, cress, endive, kale, mustard, rocket, spinach and silverbeet | Sclerotinia (*Sclerotinia minor*, *S. sclerotiorium*) | 800 g to 1 kg/ha | 7 days (H) | Apply SWITCH fungicide prior to the onset of disease. Apply a second application 7 to 14 days later if conditions continue to remain favourable for disease development. Use a 7 day spray interval and high rate under high disease pressure.<br><br>DO NOT apply more than 2 applications of SWITCH per crop. |
| **Lettuce** | Grey mould (*Botrytis cinerea*)<br><br>Sclerotinia rot (*Sclerotinia minor*, *S. sclerotiorium*) | 800 g to 1 kg/ha | 7 days (H) | Apply SWITCH fungicide prior to the onset of disease. Best results are achieved with the first spray commencing soon after transplanting on a 7 to 14 spray interval. Use a 7 day spray interval and high rate under high disease pressure or where conditions are conducive for disease development.<br><br>DO NOT apply more than 2 applications of SWITCH per crop. |
| | Anthracnose (*Microdochium panattonianum*) | 1 kg/ha in a spray volume of 250 to 500 L water/ha plus a registered rate of non-ionic surfactant | | Apply prior to the onset of disease.<br><br>DO NOT apply more than 2 applications per crop per season with an interval of 7 to 14 days between applications. Use the shorter spray interval under high disease pressure or when continual wet, cloudy conditions favour disease development.<br><br>Apply using boom-spray or similar equipment to deliver medium spray quality. Ensure spray application provides good coverage of plant foliage.<br><br>Add a non-ionic surfactant at recommended rates. |

| Crop/ Situation | Disease | Rate | WHP | Critical Comments |
|---|---|---|---|---|
| | | | | Apply in rotation with fungicide chemicals from other mode of action groups as part of a resistance management program. |
| **Nursery Stock (non-food) and Ornamentals (non-food)** including seedlings, plugs, potted colour, trees, shrubs, foliage plants, palms, grasses and fruit trees (non-bearing) | Anthracnose (*Colletotrichum* spp.)<br><br>Grey mould (*Botrytis* spp.)<br><br>Rhizoctonia rot (*Rhizoctonia* spp.)<br><br>Sclerotinia rot (*Sclerotinia* spp.) | 80 g/100 L or 800 g/ha | Not required when used as directed | Apply via calibrated boom or handheld sprayer when conditions favour disease development, prior to the first symptoms.<br><br>Do not apply more than 4 applications per crop at 10 to 14 day intervals.<br><br>DO NOT apply more than 2 consecutive applications before applying at least the same number of sprays from a different mode of action group.<br><br>See **To Avoid Crop Damage** |
| **Onions** | Neck rot (*Botrytis alli*)<br><br>**Suppression of** Black mould (*Aspergillus niger*) | 800 g/ha to 1 kg/ha plus a registered rate of non-ionic surfactant | 7 days (H) | Apply SWITCH fungicide at key timings for botrytis control prior to the onset of disease. Apply a second application 7 to 14 days later if conditions continue to remain favourable for disease development. Use a 7 day spray interval and high rate under high disease pressure.<br><br>Apply in sufficient water to ensure thorough and uniform coverage.<br><br>Applications MUST be made with the addition of a non-ionic surfactant.<br><br>DO NOT apply more than 2 applications of SWITCH per crop. |
| **Alliums** including bulb onions, spring onions, shallots and garlic | **Suppression of** Black mould (*Aspergillus niger*)<br><br>**Suppression of** Grey mould (*Botrytis cinerea*) | | | Apply prior to the onset of disease by foliar application with a boom spray or equivalent. Spray in sufficient volume to ensure thorough and uniform coverage.<br><br>DO NOT apply more than 2 applications per crop with an interval of 7 to 14 days between applications. Use the shorter spray interval under high disease pressure or when continual wet, cloudy conditions favour disease development.<br><br>To avoid resistance, the product should be rotated with other approved fungicides from different chemical groups. |
| **Pyrethrum** | Winter dieback complex:<br>Ray blight (*Phoma ligulicola*) and<br>Tan spot (*Microsphaeropsis tanaceti*) | 1 kg/ha | Not required when used as directed | Apply in a total spray volume of 400 L/ha.<br><br>Do not apply more than 2 applications per crop with an interval of 7 to 14 days between applications. |

| Crop/ Situation | Disease | Rate | WHP | Critical Comments |
|---|---|---|---|---|
| **Strawberries** | Crown and petiole rots (*Colletotrichum gloeosporioides*) | 80 g/100L water | 3 days (H) | Apply SWITCH fungicide prior to the onset of disease. Repeat applications at 7 to 14 day intervals. Use the shorter interval when conditions favour disease development (wet and cloudy conditions).<br><br>The rate suggested is for high volume dilute spraying to the point of "run-off". For concentrate spraying, adjust dilution rate accordingly.<br><br>DO NOT exceed 800 g/ha.<br><br>Apply in sufficient water to ensure thorough and uniform coverage of foliage and plant crowns.<br><br>DO NOT apply more than 3 applications of SWITCH per crop. |
| | Grey mould (*Botrytis cinerea*) | | | Commence during flowering when conditions favour disease. Repeat applications of SWITCH fungicide on a 7 to 10 day interval. Use the shorter interval when conditions favour disease development.<br><br>The rate suggested is for high volume spraying to the point of "run-off". For concentrate spraying, adjust dilution rate accordingly.<br><br>Apply in sufficient water to ensure thorough and uniform coverage of foliage and fruit. Reduce background levels of disease by removing infected plant debris and rotten fruit.<br><br>This use is subject to a CropLife fungicide resistance management strategy. DO NOT apply more than 3 applications of SWITCH per crop (see **Resistance Management**). |

(H) = harvest; (G) = grazing

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**Restraints:**

DO NOT apply by aircraft.

DO NOT apply by equipment carried on the back of the user.

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of field runoff for at least 3 days after application.

## SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between three and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

• Spray droplets are not smaller than a MEDIUM spray droplet size category.

• Minimum distances between the application site and downwind sensitive areas are observed (see the following table titled ‘Buffer zones for boom sprayers’).

**Buffer zone for Boom sprayers**

| Application rate | Boom height above the target canopy | Mandatory downwind buffer zones | | | | |
|---|---|---|---|---|---|---|
| Up to 1 kg/ha | | Bystander areas | Natural aquatic areas | Pollinator areas | Vegetation areas | Livestock areas |
| | 0.5 m or lower | 0 metres | 15 metres | 0 metres | 0 metres | 0 metres |
| | 1.0 m or lower | 0 metres | 50 metres | 0 metres | 0 metres | 0 metres |

DO NOT apply by a vertical sprayer unless the following requirements are met:

• Spray is not directed above the target canopy.

• The outside of the sprayer is turned off when turning at the end of rows and when spraying the outer row on each side of the application site.

• For dilute water rates up to the maximum listed for each type of canopy specified, minimum distances between the application site and downwind sensitive areas are observed (see the following table titled ‘Buffer zones for vertical sprayers’).

### Buffer zones for vertical sprayers

| **Type of target canopy and dilute water rate** | **Mandatory downwind buffer zones** | | | | |
|---|---|---|---|---|---|
| | **Bystander areas** | **Natural aquatic areas** | **Pollinator areas** | **Vegetation areas** | **Livestock areas** |
| 2 metres tall and shorter, maximum dilute water rate of 1000 L/ha | 0 metres | 10 metres | 0 metres | 0 metres | 0 metres |
| Taller than 2 metres (not fully-foliated), maximum dilute water rate of 2000 L/ha | 0 metres | 30 metres | 0 metres | 0 metres | 0 metres |
| Taller than 2 metres (fully-foliated), maximum dilute water rate of 2000 L/ha | 0 metres | 20 metres | 0 metres | 0 metres | 0 metres |