Product Name: THIOVIT JET MICROGRANULE FUNGICIDE/MITICIDE

APVMA Approval No: 53904/135412

RLP
APPROVED

Label Name:	THIOVIT JET MICROGRANULE FUNGICIDE/MITICIDE

Signal Headings:	

Constituent Statements:	800 g/kg SULFUR (S) present as the Elemental Sulphur

Mode of Action:	GROUP **M2** FUNGICIDE

Statement of Claims:	For the control of Powdery Mildew, Rust and Mites in Pome and Stone Fruit, Citrus, Grapevines, Kiwifruit, Mangosteens, Rambutans, Strawberries, Tomatoes, Ornamentals, and some Vegetables as specified in the Directions for Use

Net Contents:	15 kg - 25 kg

Restraints:	DO NOT apply product in combination with spraying oils except to deciduous fruit in dormant period. DO NOT use on sulphur sensitive crops when temperature exceeds 24°C.

Directions for Use:	This section contains file attachment.

Other Limitations:	

Withholding Periods:	WITHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED

Trade Advice:	

General Instructions:	This section contains file attachment.

Resistance Warning:	Fungicide Resistance Warning GROUP M2 FUNGICIDE For fungicide resistance management THIOVIT® Jet is a Group M2 fungicide. Some naturally occurring individual fungi resistant to THIOVIT® Jet and other Group M2 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by THIOVIT® Jet or other Group M2 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Syngenta Australia Pty Ltd accepts no liability for any losses that may result from failure of THIOVIT® Jet to control resistant fungi.

Precautions:	

Protections:	PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT contaminate streams, rivers, or waterways with the chemical or used containers.

Storage and Disposal:	STORAGE AND DISPOSAL Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight. DO NOT store near foodstuffs. Keep out of reach of children. Shake and empty contents into spray tank. DO NOT dispose of undiluted chemicals on site. Break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

Safety Directions:	

First Aid Instructions:	FIRST AID If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.

First Aid Warnings:	

DIRECTIONS FOR USE

Rate			Critical Comments
In the following table all rates given are for dilute spraying. For concentrate spraying, refer to the Application Section.			For all uses in in this table: Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
Crop	**Disease/Pest**	**Rate**	
Tree and Vine Crops			
Citrus	Citrus Rust Mite, Brown Citrus Rust Mite, Bud Mite, Moss, Lichens	400 to 500 g/100 L	Apply during July-August. Leave at least 3 weeks between sprays of oil and product.
	Melanose		Apply in late August for best results.
	White Louse Scale	200 to 300 g/100 L	May be added to copper spray at petal fall or applied separately during Spring or Autumn.
Grapes table grapes, fruit destined for drying	Vine Mite, Grapeleaf Blister Mite	500 g/100 L	Apply before sprouting. Ensure thorough coverage.
	Powdery Mildew	100 to 200 g/100 L	Apply immediately after budburst, then every 2 to 3 weeks or as required. Ensure thorough coverage.
Grapevines wine grapes only	Vine Mite, Grapeleaf Blister Mite	500 g/100 L	Apply before sprouting. Ensure thorough coverage.
	Powdery Mildew	200 to 600 g/100 L	Use rates to the upper end of the rate range when disease pressure is high and/or a higher degree of control is required. Apply immediately after bud burst, then every 14 to 21 days or as required. Crop phototoxicity may occur when application occurs during hot or humid conditions or when the target crop is suffering from moisture stress. DO NOT apply during the heat of the day. Use rates to the lower end of the rate range when warm conditions are present at and immediately after application. Ensure thorough coverage.
Kiwifruit	Two-spotted Mite	150 to 200 g/100 L	Spray only when the pest is present. Repeat application only when necessary. **Caution:** Avoid spraying during the heat of the day.
Mangosteens	Powdery Mildew, Mites	400 g/100 L	Apply as a foliar spray. Spray when pests are present. Repeat applications only when necessary. DO NOT apply during hot weather.
Peaches, Nectarines, Plums	Rust	200 g/100 L	Apply as required during November to January.
		350 g/100 L	Apply at petal fall only.
	Brown Rot Blossom Blight	200 g/100 L	Apply at petal fall and then as a cover spray at regular intervals beginning 3 to 4 weeks after petal fall.
		350 g/100 L	Apply at petal fall only.
Pome Fruit	Powdery Mildew	200 to 400 g/100 L	Start spraying early. Apply at least 2 sprays before blossom. Apply every 2 weeks after blossom or as required. Surfactant may be added at 12 mL/100 L. Regular sprays will also aid in the control of Black Spot (Scab).
	Black Spot (Scab)	350 g/100 L	
Rambutans	Powdery Mildew, Mites	400 g/100 L	Apply as a foliar spray. Spray when pests are present. Repeat applications only when necessary. DO NOT apply during hot weather.

Other Crops			
Crop	**Disease/Pest**	**Rate**	**Critical Comments**
Ornamentals	Powdery Mildew, Two-spotted Mite	200 to 350 g/100 L	Apply when pest or disease is first seen and repeat every 10 to 21 days.
Strawberries	Powdery Mildew	200 to 300 g/100 L	Apply when the disease is first noticed then at 10 to 14 day intervals during humid weather.
Tomatoes for processing	Powdery Mildew, Bean Rust, Tomato Russet Mite, Bean Spider Mite, Two-spotted Mite	2 kg/ha	Apply when pest or disease is first seen and repeat as necessary. Crop phytotoxicity may occur when applications occur during hot conditions. DO NOT apply during the heat of the day.
Vegetables (NOT Cucumber or Rockmelon)	Powdery Mildew, Bean Rust, Tomato Russet Mite, Bean Spider Mite, Two-spotted Mite	200 to 300 g/100 L	Apply when pest or disease is first seen and repeat as necessary. Crop phytotoxicity may occur when applications occur during hot conditions. DO NOT apply during the heat of the day.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

GENERAL INSTRUCTIONS

This product is a micronised formulation which mixes easily with water to form a spray with good spreading and sticking properties. It gives a quick fumigant action and long residual effect.

Mixing
This product must be mixed with water and applied by suitable equipment.

1	2	3
• Fill vat no more than 25% full with clean water BEFORE adding THIOVIT® Jet. • Begin agitating vat contents vigorously and continue agitation during entire mixing and spraying operations.	• Pour required amount of THIOVIT® Jet STEADILY into vat. Allow vigorous bypass agitation to completely disperse product. DO NOT dump product into vat all at once.	• After adding required quantity of THIOVIT® Jet and obtaining COMPLETE DISPERSION, continue to fill vat to desired level for spraying. • Thorough agitation of the spray liquid should continue during the entire spraying operation. • Note: Spray solution should NOT be left standing in the vat overnight.

Tank Mixing
When mixing THIOVIT® Jet and other granule or powder formulations, ensure they are added and mixed well prior to adding emulsifiable concentrate (EC) or suspension concentrate (SC) products. Wettable powder (WP) formulations should be pre-mixed separately and then added to the spray tank.

Application in Tree Crops and Vines
Dilute Spraying: Use a sprayer designed to apply high volumes of water up to the point of runoff and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of runoff. Avoid excessive runoff. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Direction for Use table for each 100 L of water. Spray to the point of runoff. The required dilute spray volume will change, and the sprayer set up and operation may also need to be changed as the crop grows.
Concentrate Spraying: Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of runoff) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (see *Dilute Spraying* above) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way:
Example Only

1. Dilute spray volume as determined above: for example 2000 L/ha
2. Your chosen concentrate spray volume: for example 500 L/ha
3. The concentration factor in this example is: 4 x (ie 2000 L ÷ 500 L = 4)
4. If the dilute label rate is 200 g/100 L, then the concentrate rate becomes 4 x 200, that is 800 g/100 L of concentrate spray.

The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training, and follow industry Best Practices.