Product Name: VERTENTO PLINAZOLIN technology Insecticide

APVMA Approval No: 93995/141169

RLP APPROVED

Label Name:	VERTENTO PLINAZOLIN technology Insecticide

Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	400 g/L ISOCYCLOSERAM

Mode of Action:	GROUP 30 INSECTICIDE

Statement of Claims:	For control of certain insect pests in almonds, macadamia nut and various fruit crops as per the Directions for Use

Net Contents:	1 – 1000 Litres

Restraints:	This section contains file attachment.

Directions for Use:	This section contains file attachment.

Other Limitations:	

Withholding Periods:	WITHHOLDING PERIODS: Avocado, Mango Harvest: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION

	Papaya Harvest: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION Almonds, Citrus, Macadamia: Harvest: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION Grazing: DO NOT GRAZE TREATED ORCHARDS

Trade Advice:	EXPORT OF TREATED PRODUCE Crops treated with VERTENTO® insecticide may contain finite (measurable) residues of isocycloseram and may pose a risk to trade in situations where no residue tolerance (import tolerance) is established in the importing country or where residues in Australian commodities are likely to exceed a residue tolerance (import tolerance) established in the importing country. For further information, please contact your export organisation or Syngenta Australia for the latest information on MRLs and import tolerances before using VERTENTO®.

General Instructions:	GENERAL INSTRUCTIONS Mixing VERTENTO® insecticide is a Suspension Concentrate (SC) formulation. Shake the container well before using. Partially fill the spray tank with clean water and add the required volume of VERTENTO® to the water whilst agitating. Always use an appropriate adjuvant, then top up the tank with clean water to the required volume. VERTENTO® should be agitated constantly before and during application and applied as soon as possible after mixing. Adjuvant Citrus: Apply VERTENTO® insecticide with a non-ionic spray adjuvant such as AGRAL® Spray Adjuvant at 10 mL per 100 L of spray mixture. For Kelly's citrus thrips, apply VERTENTO® with a non-ionic spray adjuvant such as AGRAL® Spray Adjuvant at 10 mL per 100 L of spray mixture or a petroleum oil such as Vicol Summer Oil (825 g/L petroleum oil) at 500 mL per 100 L of spray mixture. Refer to the petroleum oil label for Restraints and Critical Comments for safe use in citrus. For other specified adjuvant - refer to Syngenta Australia Pty Ltd representative for information Almond, Avocado, Macadamia, Mango and Papaya: Apply VERTENTO® with a non-ionic adjuvant such as AGRAL® Spray Adjuvant at 10 mL per 100 L of spray mixture For other specified adjuvant - refer to Syngenta Australia Pty Ltd representative for information. Application Thorough coverage of the target area is essential. Apply in sufficient water and adjust water volumes according to the crop growth stage. Use suitable application parameters (nozzles, pressure, speed, etc.) to ensure thorough and even coverage. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Can be used at up to 2.5X concentrate unless specified in Critical Comments. Dilute application using rate per 100 L: • Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed • Set up and operate the sprayer to achieve even coverage throughout the crop canopy • Apply sufficient water to cover the crop to the point of run-off • Avoid excessive run-off • The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice

	• Add the amount of product specified in the Directions for Use table for each 100 L of water • Spray to the point of run-off, to a maximum of 4000 L/ha for citrus crops or 3000 L/ha for all other crops Concentrate spraying: • Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies spray volumes less than those required to reach the point of run-off) and matched to the crop being sprayed • Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen spray volume • Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy • This is needed to calculate the concentrate mixing rate The mixing rate for concentrate spraying can then be calculated in the following way: Example only • Dilute spray volume as determined above: for example 1500 L/ha • Your chosen concentrate spray volume: for example 600 L/ha • The concentration factor in this example is 2.5X (eg 1500 L ÷ 600 L = 2.5) • If the dilute label rate is 5 mL/100 L then the concentrate rate becomes 2.5x5, which is 12.5 mL/100 L of concentrate spray • The chosen spray volume, amount of product per 100 L of water, and the sprayer set up and operation may need to be changed as the crop grows • DO NOT use at a concentration factor greater than 2.5X (eg at a rate higher than 12.5 mL/100 L where a dilute spraying rate of 5 mL/100 L is specified) • Note that the concentration mixing rate is applicable only to VERTENTO®. The adjuvant remains unchanged (ie no concentration factor applies) • For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry best practice

Resistance Warning:	For insecticide resistance management, VERTENTO® Insecticide is a Group 30 insecticide. Some naturally occurring insect biotypes resistant to VERTENTO® and other Group 30 insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if VERTENTO® or other Group 30 insecticides are used repeatedly. The effectiveness of VERTENTO® on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Syngenta Australia Pty Ltd accepts no liability for any losses that may result from the failure VERTENTO® to control resistant insects. For further information contact your local supplier, Syngenta representative or local agricultural department agronomist. In order to avoid or delay the selection of resistant insects, Group 30 insecticides should be used as part of an insecticide resistance management (IRM) strategy which incorporates the following: • Insecticides from the same mode of action group should not be used to treat successive generations of the target pest. • Multiple applications of VERTENTO® and other products containing Group 30 insecticides may be applied successively but only when targeting a single generation of the target insect. • DO NOT apply sequential applications of VERTENTO® if mites are present. If retreatment is required, apply an insecticide from a different chemical group before using VERTENTO® again. • If more than one application of an insecticide is required to control successive generations of the target pest then alternative insecticides with different modes of action should be utilised in rotation with VERTENTO®. • Where possible incorporate alternative methods of pest control as part of an integrated pest management (IPM) approach.

Precautions:	Re-entry Period: DO NOT enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

Protections:	PROTECTION OF HONEY BEES AND OTHER INSECT POLLINATORS Toxic to bees and harmful to bee brood. DO NOT apply to crops from the onset of flowering until flowering is complete unless the application is made when bees are not foraging. DO NOT allow spray drift to flowering weeds or flowering crops in the vicinity of the treatment area. Before spraying, notify beekeepers to move hives to a safe location with an untreated source of nectar and pollen, if there is potential for managed hives to be affected by the spray or spray drift. INTEGRATED PEST MANAGEMENT Toxic to beneficial arthropods. Not compatible with integrated pest management (IPM) programs utilising beneficial arthropods. Minimise spray drift to reduce harmful effects on beneficial arthropods in non-crop areas. PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

Storage and Disposal:	STORAGE AND DISPOSAL Store in the closed original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to treatment tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

Safety Directions:	May irritate the eyes. Repeated exposure may cause allergic skin disorders. Avoid contact with eyes and skin. When using together with other products, consult their label safety directions. When opening the container and preparing spray and using the prepared spray, wear • cotton overalls (or equivalent clothing) buttoned to the neck and wrist and • a washable hat and • elbow length chemical resistant gloves Wash hands after use. After each day's use, wash gloves and contaminated clothing.

First Aid Instructions:	FIRST AID If poisoning occurs contact a doctor or Poisons Information Centre. Phone 131 126.

First Aid Warnings:	WARNING - Suspected of damaging the unborn child.

Crop	Pest	Rate	Critical Comments
Almonds	Carpophilus beetle (*Carpophilus* spp.)	**Dilute spraying** 5 mL/100 L plus non-ionic adjuvant [Δ] **Concentrate spraying** Can be used at up to 2.5X concentrate. Refer to the Application section	**Carpophilus beetle:** Monitor orchard before and during hull split for the presence of Carpophilus beetles. If numbers have the potential to cause economic loss, apply VERTENTO® insecticide. Continue to monitor orchard and where pest pressure continues from an influx of Carpophilus beetles from surrounding areas apply a second application no less than 7 days after the first VERTENTO® insecticide spray. Kernel damage can still occur if Carpophilus beetles enter the orchard just prior to harvest, when the shell is open, and feed directly on the kernel. If further spray applications are required rotate to an alternative mode of action. **DO NOT apply more than 3 applications per cropping season.** **DO NOT exceed 2 spray applications from hull-split until harvest.** **DO NOT exceed 150 mL of VERTENTO® insecticide per hectare per spray application.** Ensure thorough coverage of the target crop including with concentrate spraying – refer to the Application section in General Instructions. [Δ] Always add a specified adjuvant – refer to the Adjuvant section in General Instructions. VERTENTO® insecticide should form part of an integrated pest program to manage carpophilus beetle populations with a focus on orchard hygiene.

Crop	Pest	Rate	Critical Comments
Citrus	Fuller's rose weevil (*Naupactus godmanni*) Kelly's citrus thrips (*Pezothrips kellyanus*) Light brown apple moth (*Epiphyas postvittana*)	**Dilute spraying** 5 mL/100 L plus adjuvant[Δ] **Concentrate spraying** Can be used at up to 2.5X concentrate. Refer to the Application section	**Kelly's citrus thrips:** Monitor orchard and commence VERTENTO® insecticide applications after flowering when local economic spray thresholds are reached and prior to calyx closure. A single application may be suitable where pest pressure is low. Under high thrips insect pressure, apply a second spray application prior to calyx closure. There must be a minimum of 14 days between each VERTENTO® insecticide application. **Fuller's rose weevil:** Monitor orchard for weevil emergence. Apply VERTENTO® insecticide prior to the start of egg lay which usually occurs in late summer and through autumn. **Light brown apple moth:** Monitor orchard for insect presence and apply VERTENTO® insecticide when local economic spray thresholds are reached. Continue to monitor orchard and where pest pressure continues apply a second application. There must be a minimum of 14 days between each VERTENTO® insecticide application. **DO NOT apply more than 2 applications per cropping season.** **DO NOT exceed 200 mL of VERTENTO® insecticide per hectare per spray application.** Ensure thorough coverage of the target crop including with concentrate spraying – refer to the Application section in General Instructions. [Δ] Always add a specified adjuvant – refer to the Adjuvant section in General Instructions.

Crop	Pest	Rate	Critical Comments
Macadamia	Fruit spotting bug (*Amblypelta nitida*) Macadamia seed weevil (*Kuschelorhynchus macadamiae*)	**Dilute spraying** 5 mL/100 L plus non-ionic adjuvant [Δ] **Concentrate spraying** Can be used at up to 2.5X concentrate. Refer to the Application section	Monitor orchard and commence VERTENTO® insecticide applications after flowering once local economic spray thresholds are reached. Continue to monitor orchard and where pest pressure continues, additional VERTENTO® insecticide applications may be applied. There must be a minimum of 14 days between each VERTENTO® insecticide application. If further spray applications are required, rotate to an alternative mode of action. **DO NOT apply more than 3 applications per cropping season.** **DO NOT exceed 150 mL of VERTENTO® insecticide per hectare per spray application.** Ensure thorough coverage of the target crop, including with concentrate spraying – refer to the Application section in General Instructions. [Δ] Always add a specified adjuvant – refer to the Adjuvant section in General Instructions. VERTENTO® insecticide should be used as part of an integrated pest management approach which should include the use of other measures for control.
Avocado, Mango	Banana spotting bug (*Amblypelta lutescens*) Fruit spotting bug (*Amblypelta nitida*)	**Dilute spraying** 5 mL/100 L plus non-ionic adjuvant [Δ] **Concentrate spraying** Can be used at up to 2.5X concentrate. Refer to the Application section	Monitor orchard for spotting bug presence and spray when local economic spray thresholds are reached. Continue to monitor orchard and where pest pressure continues, additional VERTENTO® insecticide applications may be applied. There must be a minimum of 14 days between each VERTENTO® insecticide application. **DO NOT apply more than 3 applications per cropping season.** **DO NOT exceed 150 mL of VERTENTO® insecticide per hectare per spray application.** Ensure thorough coverage of the target crop, including with concentrate spraying – refer to the Application section in General Instructions. [Δ] Always add a specified adjuvant – refer to the Adjuvant section in General Instructions.

Crop	Pest	Rate	Critical Comments
Papaya	Suppression of: Banana spotting bug (*Amblypelta lutescens*)	75 mL/ha plus non-ionic adjuvant [Δ]	Monitor orchard for spotting bug presence and spray when local economic thresholds are reached. Continue to monitor orchard and where pest pressure continues, additional VERTENTO® insecticide applications may be applied. There must be a minimum of 14 days between each VERTENTO® insecticide application. **DO NOT apply more than 3 applications per cropping season.** Ensure thorough coverage of the target crop – refer to the Application section in General Instructions. **Concentrate spraying is not recommended for this use pattern.** [Δ]Always add a specified adjuvant – refer to the Adjuvant section in General Instructions.

Restraints

DO NOT apply by aerial spraying

DO NOT apply if heavy rains or storms that are likely to cause runoff are forecast within 3 days

DO NOT irrigate to the point of water runoff for at least 3 days after application

DO NOT apply to avocados, macadamias, mangoes or papayas in the Mackay/Whitsunday region

DO NOT apply sequential sprays of VERTENTO® insecticide if mites are present, in order to manage insecticide resistance

Spray Drift Restraints

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the buffer zone table below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a vertical, airblast or oscillating boom sprayer unless the following requirements are met:

- Spray is not directed above the target canopy
- The outside of the sprayer is turned off when turning at the end of rows and when spraying the outer row on each side of the application site
- For dilute and concentrate water rates up to the maximum listed for each type of canopy specified, minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for vertical sprayers') are observed.

Buffer zones for vertical, airblast or oscillating boom sprayers

Crop and Application Rate	Maximum dilute water rate	Type of target canopy	Mandatory Downwind Buffer Zones (metres)				
			Natural Aquatic Areas	Vegetation Areas	Bystander Areas	Pollinator Areas	Livestock Areas
Citrus 5 mL/100 L (max. 200 mL/ha)	1000 L/ha	2 metres tall and shorter	60	0	0	0	0
	4000 L/ha	Taller than 2 metres (fully foliated)	170	0	0	0	0
		Taller than 2 metres (not fully foliated)	150				5
Almond, Avocado, Macadamia, Mango 5 mL/100 L (max. 150 mL/ha)	1000 L/ha	2 metres tall or shorter	75	0	0	0	0
	3000 L/ha	Taller than 2 metres (not fully foliated)	140	0	0	0	0
		Taller than 2 metres (fully foliated)	150				
Papaya 75 mL/ha	1000 L/ha (7.5 mL/100 L)	2 metres tall and shorter	75	0	0	0	0
		Taller than 2 metres (fully foliated)	100	0	0	0	0
	2000 L/ha (3.75 mL/100 L)	Taller than 2 metres (fully foliated)	100	0	0	0	0