Product Name: CAMPBELL TRIDIM 250 EC FUNGICIDE
APVMA Approval No: 51580/122652

RLP
APPROVED

Label Name:	CAMPBELL TRIDIM 250 EC FUNGICIDE

Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	250 g/L TRIADIMENOL SOLVENT :600 g/L N-METHYL-2-PYRROLIDONE

Mode of Action:	GROUP 3 FUNGICIDE

Statement of Claims:	For systemic control of ring spot on cabbage, cauliflower and broccoli, powdery mildew on cucurbits, tomatoes, grapevines, parsnip, radish, swede, turnips and papaws, pineapple disease of sugar cane, Myrtle rust in Riberries, lemon myrtle, anise myrtle, nursery stock (non-food and forestry), non-bearing fruit trees, ornamentals, cut flowers/foliage and oil tea tree, and various diseases of turf as indicated in the directions for use table.

Net Contents:	1L 5L

Restraints:	Do not apply to turf earlier than 5 weeks after sowing

Directions for Use:	This section contains file attachment.

Other Limitations:	

Withholding Periods:	WITHHOLDING PERIODS:

	Harvest: Cucurbits, eggplant, peppers (capsicum and chilli), tomatoes: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION. Brassica vegetables, grapevines, pawpaws, parsnips, radish, swede, turnip : DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION. Riberries,: DO NOT HARVEST FRUIT FOR 4 WEEKS AFTER APPLICATION. Anise Myrtle, Lemon myrtle: DO NOT HARVEST LEAVES FOR 4 MONTHS AFTER APPLICATION. Grazing: Brassica vegetables: DO NOT GRAZE ANY TREATED AREAS OR CUT FOR STOCK FOOD. Parsnips, radish, swede, turnip : DO NOT graze or cut parsnip, radish, swede and turnip crops for animal feed Anise myrtle, Lemon myrtle, Riberries : DO NOT allow livestock to graze treated area during the season of application. Non-bearing fruit trees, tea tree: DO NOT graze during the season of application..

Trade Advice:	Export of treated produce Growers should note that MRLs or import tolerances do not exist in all markets for edible produce treated with Tridim 250EC. If you are growing edible produce for export, please check with Colin Campbell (Chemicals) Ply. Ltd. for the latest information on MRLs and import tolerances before using Tridim 250EC.

General Instructions:	GENERAL INSTRUCTIONS Tridim 250 EC is a broad spectrum, systemic, foliar fungicide that is quickly taken up by plants and translocated within the plant. Because of its systemic activity, Tridim is less affected by rainfall or irrigation than contact type fungicides. MIXING Half fill the spray tank or vat with clean water. Then add the required amount of Tridim with agitator going and top up the tank with water. Agitate thoroughly prior to and during spraying. APPLICATION Dilute Spraying (Grapevines, Pawpaws)- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.-The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.- Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run-off.- The required dilute spray volume will change the sprayer set up and operation may also need to be changed, as the crop grows. Concentrate Spraying (Grapevines, Pawpaws)- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (See dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.- The mixing rate for concentrate spraying can be calculated in the following way: Example only 1. Dilute spray volume as determined above. For example 1500L/ha 2. Your chosen concentrate spray volume: For example 500L/ha 3. The concentration factor in this example is 3x (i.e. 1500L /500L = 3) 4. As the dilute label rate is 10mL/100L for grapevines, then the concentrate rate becomes 3 x 10, that is 30ml/100L of concentrate spray. - The chosen spray volume, amount of product per 100L of water, and the spray set up and operation may need to be changed as the crop grows.

	- For grapevines - DO NOT use a concentrate rate higher than that specified in the Critical Comments because a rate higher than that has not been tested. - For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry best practices.

Resistance Warning:	FUNGICIDE RESISTANCE WARNING GROUP 3 FUNGICIDE Tridim 250EC is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product and other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Colin Campbell (Chemicals) Pty Ltd accepts no liability for any losses that result from failure of this product to control resistant fungi. Resistance management : DO NOT apply more than 2 consecutive applications of any Group 3 fungicide. Rotate approved products from different chemical mode of action groups at regular intervals within a structured disease management plan.

Precautions:	

Protections:	PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT Harmful to fish. Do not contaminate streams, rivers or waterways with this product or used container.

Storage and Disposal:	Store in the closed, original container in a dry, cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Safety Directions:	SAFETY DIRECTIONS Will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length butyl rubber gloves and goggles.. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day’s use, wash gloves, goggles and contaminated clothing.

First Aid Instructions:	FIRST AID

	If poisoning occurs, contact a doctor of Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. If swallowed, do NOT induce vomiting. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

First Aid Warnings:	

DIRECTIONS FOR USE

Crop	Disease	Rate	WHP	Critical Comments
Brassica vegetables (e.g. Cabbage, Cauliflower, Broccoli, Brussels sprouts	Ring spot	200 or 400 mL/ha plus a non-ionic wetter 200 mL/ha	7 days	Apply every 10 days. Start the program immediately following the first two consecutive days of rain after transplanting. Use higher rate under heavy disease pressure or when weather conditions favour development of the disease.
Cucurbits	Powdery mildew	40 mL/100L or 400 mL/ha	1 day	Apply a non-group 3 fungicide on a schedule commencing as soon as possible after emergence until fruit set. Then either tank mix or alternate Tridim with a non-group 3 fungicide on a 5-10 day schedule. Use short spray intervals when disease pressure is higher plants are growing rapidly. Good coverage is essential for good results.
Tomatoes, Peppers (Capsicums and Chillies), Eggplants				Apply Tridim 250EC prior to, or at first sign of disease. Best control is achieved when applied as part of a well-planned preventative spray program, with early applications before heavy disease pressure develops. Tridim 250EC should be either tank mixed or alternated with fungicides from different activity groups (NOT Group 3 Fungicides). Intervals between sprays should be 5-10 days with short spray intervals appropriate when disease pressure is high, or plants are growing rapidly, or weather conditions favour disease development. Good coverage is essential for goodresults.DO NOT apply more than 4 sprays of Tridim 250EC per crop. Application: Apply in sufficient volume to ensure thorough coverage of all foliage. For applications with spray volumes above 1000L/ha apply as a dilute spray to the point of incipient run-off on a per 100L basis.

Crop	Disease	Rate	WHP	Critical Comments
Grapevines	Powdery mildew	**Dilute Spraying** 10mL/100L or 100mL/ha **Concentrate Spraying** Refer to the Application section	7 days	This use pattern is subject to a CropLife Resistance Management strategy (www.croplife.org.au) Apply a maximum of 3 sprays per season, thoroughly and only as part of a spray program. Typically: 1.When shoots are 10-20cm long. 2.Pre flowering. 3.At flowering. 4.After fruit set 5.Before bunches close. A non-Group 3 fungicide should be used after a maximum of 2 consecutive Tridim 250EC sprays as part of this program. DO NOT allow spray intervals to exceed 21 days. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. DO NOT use in equipment that requires rates greater than 50mL/100Lwater. DO NOT apply in water volumes less than 250L/ha
Papaws (QLD, NSW, WA only)	Powdery mildew	**Dilute Spraying** 20mL/100L or 200 mL/ha **Concentrate Spraying** Refer to the Application section	7 days	Spray when powdery mildew infection begins or when conditions favour the disease. Repeat application every 14 days while conditions favour the disease. Ensure thorough coverage of foliage particularly the crown region. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.
Sugar cane (QLD, NSW only)	Pineapple disease	20 mL/100L	-	Apply to setts by dipping or spraying. Ensure thorough wetting of cut ends.
Parsnips Radish Swede Turnip	Powdery mildew	400 mL/ha or 40 mL/100 L	H : 7 days G: Do not graze or cut for feed	Apply foliar spray when first sign of infection is evident or when conditions are highly conducive to disease development. Apply a maximum of 2 foliar applications with a minimum re-treatment interval of 10 days between applications using boom sprayer, hand wands, side-mounted boom or equivalent.

Crop	Disease	Rate	WHP	Critical Comments
Riberries *(Syzygium luehmannii* and *S. fibrosum*) Anise Myrtle (S. anisatum) Lemon myrtle (*Backhousia citriodora)*	Myrtle rust	100-150 mL/ha	Fruit: 4 weeks Leaf: 4 months Grazing: No grazing during the season of application	Apply 2 sprays per crop 4 weeks apart. Minimum re-treatment interval 14-21 days. Apply via ground based equipment on appearance of myrtle rust in a plantation or when conditions favour development of the disease. Use a maximum spray volume of 400 L/ha.
Nursery stock and ornamentals, including nursery stock (non-food, forestry and non-bearing fruit trees *), ornamentals, crops for cut flowers/foliage *DO NOT apply to non-bearing fruit trees less than 6 months prior to first harvest.	Myrtle rust	50mL/100L	Non-bearing fruit trees: DO NOT graze during the season of application.	Apply by ground application only e.g. knapsack, powered hand-gun, boom, air-assisted. Apply in sufficient volume to ensure thorough coverage of all plant surfaces. Treat a sample area and assess appropriately prior to whole crop treatment to help minimise potential for phytotoxic damage. This is particularly important for crops in bloom. Minimum re-treatment interval between consecutive applications : 14-21 days. **For interstate quarantine requirements:** Apply in accordance with the procedures outlined in the appropriate State Import Requirement or Interstate Certification Assurance (ICA) Document. DO NOT apply the same chemical (or chemical class) on despatch that will be used by an importing authority for decontamination on receipt.
Tea tree		100-150 mL/ha	DO NOT graze during the season of application.	Apply 2 sprays per crop 4 weeks apart. Apply via ground based equipment on appearance of myrtle rust in a plantation or when conditions favour development of the disease. Use a maximum spray volume of 400L/ha.

Crop	Disease	Rate	WHP	Critical Comments
Turf	Dollar spot (NSW, Vic, Qld, SA, WA only)	**Preventative** 30mL/100m ^{2}	-	Apply thoroughly on a monthly programme starting immediately when the disease appears.
		Curative 60mL/100m^2		Apply thoroughly under conditions of heavy disease pressure then treat as necessary.
	Black Helminthosporium (QLD, NSW, TAS only)	60mL/100m^2		Use high volume equipment. Low volume may be used provided turf is irrigated thoroughly after spraying. Apply during period of turf growth on a monthly programme.
	Take – all patch (NSW, Vic, Qld, WA only)			Preventive only after soil fumigation with methyl bromide. Apply thoroughly , using high volume equipment, 6-8 weeks after seeding. Repeat applications (every 8 weeks in the first two years) for continued protection.
	Winter Fusarium (NSW, VIC, TAS only)			Apply thoroughly on a monthly programme from end of March until danger of infection has passed.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.