Product Name: RANMAN 400 SC FUNGICIDE
APVMA Approval No: 66411/142290

RLP
APPROVED

Label Name:	RANMAN 400 SC FUNGICIDE

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	ACTIVE CONSTITUENT: 400 g/L CYAZOFAMID

Mode of Action:	GROUP 21 FUNGICIDE

Statement of Claims:	For the control of various fungal diseases in certain crops as specified in the DIRECTIONS FOR USE table

Net Contents:	500 mL - 200 L

Restraints:	This section contains file attachment.

Directions for Use:	This section contains file attachment.

Other Limitations:	

Withholding Periods:	WITHHOLDING PERIODS: Potatoes: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION Spinach and silverbeet: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION Basil: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION

	Brassica vegetables: Nil Brassica vegetable seedlings and brassica leafy vegetable seedlings: Nil Poppies: Harvest: NOT REQUIRED WHEN USED AS DIRECTED. Poppies: Grazing: DO NOT GRAZE OR CUT FOR STOCK FOOD.

Trade Advice:	Trade advice Export of treated produce Growers should note that maximum residue limits (MRLs) or import tolerances may not exist in all markets for edible produce treated with Ranman 400 SC Fungicide. If you are growing edible produce for export, please check with UPL Australia Limited for the latest information on MRLs and import tolerances before using this product.

General Instructions:	GENERAL INSTRUCTIONS Mixing Fill minimum 50% of the required water into the spray tank, and agitate when adding the required amount of Ranman 400 SC Fungicide. Finally add the rest of the required water volume. Keep the spray solution agitated until all product is applied. Never prepare more spray solution than required. Concentrate Ground and Aerial Sprays Apply in 50 to 100 litres water per hectare. Compatibility For information on the compatibility of Ranman 400 SC Fungicide with other products, contact your local UPL Australia Limited representative.

Resistance Warning:	Fungicide Resistance Warning GROUP 21 FUNGICIDE For fungicide resistance management Ranman 400 SC Fungicide is a group 21 fungicide. Some naturally occurring individual fungi resistant to the product and other group 21 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by the product or other Group 21 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, UPL Australia Limited accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

Precautions:	

Protections:	PROTECTION OF LIVESTOCK DO NOT graze or feed treated crops to animals. PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers. PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS DO NOT apply under weather conditions, or from spraying equipment that may cause drift onto nearby plants/crops, cropping lands or pastures.

Storage and Disposal:	STORAGE AND DISPOSAL

	Store in the closed, original container in a cool, well-ventilated area. Do NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to the spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

Safety Directions:	SAFETY DIRECTIONS When preparing the product for use and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). Wear elbow length chemical resistant gloves when opening the container and preparing the spray for aerial spraying equipment. If applying by spraying equipment carried on the back of the user, when preparing the product for use wear elbow length chemical resistant gloves. Protected cropping situations (e.g. greenhouses and plastic tunnels): wear cotton overalls buttoned to the neck and wrist, chemical resistance gloves and a disposable mist mask. Wash hands after use. After each day's use wash gloves and contaminated clothing.

First Aid Instructions:	FIRST AID If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131 126; New Zealand 0800 764 766.

First Aid Warnings:	

RESTRAINTS

DO NOT apply through any type of irrigation equipment.

DO NOT use on forage brassicas.

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

• Spray droplets not smaller than a MEDIUM spray droplet size category.

• Minimum distances between the application site and downwind sensitive areas (see 'Mandatory downwind buffer zones' section of the following table titled 'Buffer zones for boom sprayers') are observed.

Buffer zones for boom sprayers

Application rate	Mandatory downwind buffer zone (metres)
	Natural aquatic areas
Up to maximum label rate	5

DO NOT apply by aircraft unless the following requirements are met:

• Spray droplets not smaller than a MEDIUM spray droplet size category.

• For maximum release height above the target canopy of 3 m or 25 per cent of wingspan or 25 per cent of rotor diameter, whichever is the greatest, minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section) are observed.

Buffer zones for aircraft

Type of aircraft	Wind speed at time of application	Mandatory downwind buffer zones (metres)
		Natural aquatic areas
Fixed-wing	From 3 to 8 kilometres per hour	20
	From 8 to 14 kilometres per hour	25
	From 14 to 20 kilometres per hour	30
Helicopter	From 3 to 8 kilometres per hour	20
	From 8 to 14 kilometres per hour	30
	From 14 to 20 kilometres per hour	40

Vertical sprayers

DO NOT apply by a vertical sprayer unless the following requirements are met:

- Spray is not directed above the target canopy.
- The outside of the sprayer is turned off when turning at the end of rows and when spraying the outer row on each side of the application site.

DIRECTIONS FOR USE

Crop/ Situation	Disease	Rate	WHP	Critical Comments
Potatoes	Late blight *(Phytophthora infestans)*	150 – 200 mL/ha	(H) 7 days	When conditions favour disease development apply consecutive sprays of Ranman 400 SC at 7 to 10 day intervals. DO NOT wait for disease to appear. Use the shorter interval when disease pressure is severe. Ensure thorough coverage of plants. The total number of Ranman 400 SC applications per season should not exceed 6. When applying Ranman 400 SC consecutively, apply a maximum of 3 consecutive sprays, then switch to a fungicide of another chemical group. Apply Ranman 400 SC in 250 L to 600 L water per hectare for dilute sprays. For concentrate ground and aerial sprays see General Instructions.
Broccoli	White Blister (*Albugo candida*)	150 – 200 mL/ha	-	When conditions favour disease development apply consecutive sprays of Ranman 400 SC at 7 to 10 day intervals. DO NOT wait for disease to appear. Use the shorter interval when disease pressure is severe. Ensure thorough coverage of plants. The total number of Ranman 400 SC applications per season should not exceed 6. When applying Ranman 400 SC consecutively, apply a maximum of 3 consecutive sprays, then switch to a fungicide of another chemical group. Apply Ranman 400 SC in 250 L to 800 L water per hectare for dilute sprays. For concentrate ground and aerial sprays see General Instructions.
Brassicas including Cabbages, Broccoli	Downy mildew (*Hyaloperonospora* spp)	150 – 200 mL/ha	Nil	Make fungicide applications on a 7 to 10 day schedule beginning when disease is first seen or weather and downy mildew disease pressures are expected to initiate a disease outbreak. Use the longest interval for preventative applications or very low disease pressure. Shorten the interval as disease

Brassica vegetable seedlings and brassica leafy vegetable seedlings				pressure and/or fast crop development increases, down to the shortest interval. RANMAN 400 SC FUNGICIDE can be tank-mixed with an organosilicone surfactant at the manufacturer's label recommendation. The total number of Ranman 400 SC applications per season should not exceed 6. When applying Ranman 400 SC consecutively, apply a maximum of 3 consecutive sprays, then switch to a fungicide of another chemical group. Apply Ranman 400 SC in 250 L to 800 L water per hectare for dilute sprays. For concentrate ground and aerial sprays see General Instructions.
Spinach	Downy mildew (P*eronospora* spp)		(H) 3 days	

Crop/ Situation	Disease	Rate	WHP	Critical Comments
Poppies (*Papaver somniferum*)	Systemic downy mildew (*Peronospora somniferi*)	200 mL/ha	-	Apply from 4-leaf stage to row closure. DO NOT apply after the full flowering crop growth stage (BBCH 65). Apply by foliar application using calibrated boom sprayer or similar equipment. Use a spray volume of 100 L/h to achieve thorough and uniform foliar coverage. DO NOT apply less than 7 days after the initial treatment. DO NOT apply with aircraft. Apply with Activator Surfactant at a rate of 100 mL product/100 L. DO NOT apply more than 4 applications per crop. When applying cyazofamid consecutively, apply a maximum of three (3) consecutive sprays, then switch to a fungicide of another chemical (mode of action) group. The sensitivity of the crop to be treated has not been fully evaluated. Treat a small area to ascertain the reaction before treating the whole crop.
Nursery stock (non-food) seedling and plugs, potted colour, trees and shrubs, foliage plants, palms and grasses and fruit trees (non-bearing)	Pythium crown and root rots and damping off (*Pythium* spp.) Phytophthora crown and root rots and foliar blights (*Phytophthora* spp.) Downy mildew (*Peronospora* spp., *Plasmopara* spp., *Bremia* spp.)	<u>Foliar application</u> 15 - 20 mL /100 L OR 150 - 200 mL/ha <u>Soil surface spray application</u> 15 - 40 mL /100 L	-	Apply as a protective treatment when conditions first favour disease development. Apply a maximum two (2) treatments per crop, with 14-21 days between successive sprays. Use the higher rate and shorter respray interval when disease pressure is high. The sensitivity of all the plants to be treated has not been fully evaluated. Treat a small area or small number of plants to ascertain the reaction before treating the whole crop **Foliar application** Apply to the point of run-off using sufficient water to ensure thorough coverage of all plant surfaces. A spray volume range of 250-800 L water per hectare is recommended. Apply using calibrated air-blast sprayer, air-shear sprayer, boom sprayer, back-pack mister, hand-lance or equivalent equipment. Addition of a suitable organosilicone surfactant may assist with the control of downy mildew diseases. A wetter is not required for the control of Phytophthora foliar blights. **Soil Surface Spray Application** Use sufficient water to ensure thorough coverage of the soil surface around plants; irrigate after application to wet root zone only. Apply using suitable and correctly calibrated ground application equipment.

Crop/ Situation	Disease	Rate	WHP	Critical Comments
Spinach and silverbeet (field grown)	Pythium damping off (*Pythium* spp.)	200 mL/ha	(H) 3 days	Do not wait for disease to appear. Make the first application to the soil as a directed, post-transplant or post planting application. The directed application should be made as a band 10 to 15 cm wide over the seed line or transplants. Use a spray volume of 400 – 500 L water/ha Apply via ground rig boom sprayers applying sufficient water to obtain thorough and uniform coverage of foliage. Apply a maximum of six (6) foliar applications per crop at a minimum re-treatment interval of 7-10 days. Apply a maximum of three (3) consecutive sprays, then switch to a fungicide of another chemical (mode of action) group. The sensitivity of some species and varieties of the crops to be treated has not been fully evaluated under all growing conditions. Treat a small number of plants to ascertain their reaction before treating the whole crop.
Basil (Field and protected crops)	Downy mildew (*Peronospora belbahrii*)	175 – 200 mL /ha OR 17.5 – 20 mL / 100 L	(H) 1 day	Apply as a protective treatment when conditions favour disease development and before the disease is evident. DO NOT wait for the disease to appear. Apply up to six (6) foliar applications per crop. Apply only three (3) consecutive applications of cyazofamid (Group 21), with an interval of 7-10 days between sprays, before changing to a fungicide product from a different Mode of Action group. Use the higher application rate and shorter interval under conditions that are highly conducive to disease development, or where there is rapid vegetative growth. Apply in sufficient volume of water to achieve thoroughly coverage of all foliage and stems. A spray volume of 250-500 L/ha is recommended. Apply via air-blast sprayer, air-shear sprayer or equivalent. Use a non-ionic surfactant at approved label rates.

(H) Harvest

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.