RLP
APPROVED

Product Name: Fascinate Dry Herbicide
APVMA Approval No: 85197/124905

Label Name:	Fascinate Dry Herbicide

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	500 g/kg GLUFOSINATE-AMMONIUM

Mode of Action:	GROUP **N** HERBICIDE

Statement of Claims:	For non-residual control of broadleaf and grass weeds in various situations as specified in the DIRECTIONS FOR USE table.

Net Contents:	1 kg-20 kg

Restraints:	Restraints DO NOT apply by aircraft. DO NOT apply when rain is expected within 6 hours. DO NOT apply to weeds under stress due to, for example, very dry, very wet, frosty or diseased conditions. DO NOT apply under hot dry conditions (temperatures above 33ºC with a relative humidity below 50%).

Directions for Use:	This section contains file attachment.

Other Limitations:	

Withholding Periods:	WITHHOLDING PERIODS (WHP): Harvest (H) Avocado, banana, blackberry, boysenberry, citrus fruit, feijoa, grapes, guava, kiwifruit, litchi, loganberry, mango, olives, passionfruit, pawpaw, pineapple, rambutan, raspberry, strawberries, tomatoes, tree nuts: NOT REQUIRED WHEN USED AS DIRECTED. Pome and stone fruit: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION. Grazing (G) DO NOT GRAZE OR CUT TREATED AREAS FOR STOCK FOOD FOR 8 WEEKS AFTER APPLICATION. Summer fallow: DO NOT GRAZE OR CUT FOR STOCKFOOD A CROP SOWN FOLLOWING A FALLOW SPRAY FOR 6 WEEKS AFTER SOWING.

Trade Advice:	Export of Treated Produce Growers should note that suitable MRLs or import tolerances may not be established in all markets for produce treated with Fascinate Dry Herbicide. If you are growing produce for export, please check with UPL Australia Pty Ltd for the latest information on MRLs and import tolerances BEFORE using Fascinate Dry Herbicide.

General Instructions:	GENERAL INSTRUCTIONS Fascinate Dry Herbicide is a non-volatile herbicide with non-selective activity against many annual and perennial broadleaf weeds and grasses. Fascinate Dry Herbicide is absorbed by plant foliage and green stems. It is not significantly translocated as an active herbicide throughout the plant, and therefore will only kill that part of a green plant that is contacted by spray. Fascinate Dry Herbicide does not provide residual weed control. Visible symptoms of control appear in 3 to 7 days, but complete desiccation may take 20 to 30 days under cool conditions. Best results are achieved when application is made under good growing conditions. Application to weeds under stress (e.g. due to continuous severe frosts, dry or waterlogged conditions) should be avoided. Soil fumigation / sterilisation Fascinate Dry Herbicide is metabolised (broken down) by microorganisms in the soil to become inactive. Soil fumigation or sterilisation will reduce the number of microorganisms present, thus slowing the breakdown of Fascinate Dry Herbicide. As damage to transplants or seedlings may occur, it is not advisable to apply Fascinate Dry Herbicide in conjunction with soil fumigation or sterilisation. Plastic mulches Fascinate Dry Herbicide will remain active on inert surfaces such as plastic. Special care should be taken when applying Fascinate Dry Herbicide over plastic mulches, as plant contact with the mulch after spraying may result in crop damage. Compatibility Fascinate Dry Herbicide is compatible with most residual herbicides e.g. simazine, diuron, oxyfluorfen (Goal®), norfluazuron (Solicam®) and oryzalin (Surflan®). The addition of a wetting agent or other adjuvant is generally not considered necessary. However, benefit has been obtained using a wetting agent on hard-to-wet weeds when using water rates in excess of 500 L/ha. The rate is 25 mL/100 L of a 1000 g/L non-ionic wetting agent, or equivalent. Mixing Fascinate Dry Herbicide mixes easily with water. Clean water should always be used for mixing with Fascinate Dry Herbicide. Ensure that the spray tank is free of any residues of previous spray materials. Two-thirds fill the spray tank with clean water, and with agitator operating add the required amount of Fascinate Dry Herbicide. Add other relevant compatible products. Top the tank up to the required volume with clean water with agitator running.

	Application Equipment Ground Sprayers Aim to apply a thorough and even coverage of spray to the target plant. Dense stands of weeds should be thoroughly wetted with spray. Incomplete coverage may result in poor control. Equipment should be such that adequate coverage, penetration and volume of spray liquid can be achieved. Boom or Directed Sprayer Equipment Fascinate Dry Herbicide should be applied at label rates (refer to specific column in the lists of weeds controlled) in sufficient water to give thorough coverage of weeds. It has been found that 300 to 500 L/ha has given good results under most weed conditions. Special care must be taken when using sprayer/slasher combination units not to cause dust and turbulence, which can carry spray into non-target areas. Knapsack and Handgun Equipment Fascinate Dry Herbicide should be applied at label rates (refer to specific columns in the lists of weeds controlled) in adequate water to thoroughly wet the weeds being sprayed, i.e. 500 to 1000 L/ha. Dense stands will require up to 1000 L/ha of spray mixture, whereas less dense stands will require less water. High volume application using hollow-cone nozzles for hand spraying is recommended. Controlled Droplet Application (CDA) Equipment Fascinate Dry Herbicide may be applied through CDA row spraying equipment fitted with a solid (impermeable) shroud or skirt, at rates as recommended for boom or directed sprayers (refer to specific column in the lists of weeds controlled), provided thorough spray coverage of weeds can be achieved. Apply preferably when weeds are less than 15 cm in height, with the equipment set up so that the spray dome only just touches the tops of the weeds. A total spray volume of 20 to 30 L/ha has been found to give good results. Do not mix residual herbicides or any spray adjuvants with Fascinate Dry Herbicide when using CDA equipment. Warning: Because the spray solution is highly concentrated particular care must be taken when using Fascinate Dry Herbicide through CDA equipment to avoid contact of the spray solution with any part of the crop trunk or canopy. DO NOT apply Fascinate Dry Herbicide through equipment fitted with bristle skirts. Particular care should be taken when using CDA equipment around green or uncalloused bark. Please refer to PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS. CDA equipment must not be used for application in cherry orchards. Sprayer cleanup Clean all equipment after use by thoroughly flushing with water. Aircraft Do not apply by aircraft.

Resistance Warning:	Resistant Weeds Warning GROUP N HERBICIDE Fascinate Dry Herbicide is a member of the phosphinic acid group of herbicides. Fascinate Dry Herbicide is an inhibitor of glutamine synthetase. For weed resistance management Fascinate Dry Herbicide is a Group N herbicide. Some naturally occurring weed biotypes resistant to Fascinate Dry Herbicide and other Group N herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Fascinate Dry Herbicide or other Group N herbicides. Since occurrence of resistant weeds is difficult to detect prior to use, UPL Australia Pty Ltd

	accepts no liability for any losses that may result from the failure of Fascinate Dry Herbicide to control resistant weeds.

Precautions:	PRECAUTIONS DO NOT allow bystanders to come into contact with the spray cloud. Re-entry period Do not allow entry into treated areas until the spray has dried.

Protections:	PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT contaminate streams, rivers or waterways with this product or the used container. PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. DO NOT apply on desirable foliage or allow spray to drift onto the foliage of desirable plants, trees or vines, as damage will occur. DO NOT allow product to contact green or uncalloused bark (such as on desirable young trees and vines) or cut, cracked, damaged or wounded tissue, where the affected surface is not adequately healed. Fascinate Dry Herbicide may be used around desirable trees/vines less than two years old provided they are effectively shielded from spray and spray drift. DO NOT allow desirable plant foliage to contact any inert surface, such as plastic mulches, which have been treated with Fascinate Dry Herbicide. DO NOT apply Fascinate Dry Herbicide to recently fumigated or sterilised soil.

Storage and Disposal:	STORAGE AND DISPOSAL Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Plastic packaging only Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product. Laminated bag packaging only Single-rinse or shake remainder into spray tank/water/dip/drench, etc. Do not dispose of undiluted chemicals on site. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Safety Directions:	SAFETY DIRECTIONS May irritate the eyes and skin. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes and skin. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). If applying by low pressure hand wand, wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat, elbow length chemical resistant gloves, and half facepiece respirator

	Wash hands after use. After each day's use, wash gloves, contaminated clothing and respirator and if rubber wash with detergent and warm water.

First Aid Instructions:	FIRST AID If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

First Aid Warnings:	

DIRECTIONS FOR USE

A. ORCHARDS, PLANTATIONS, VINEYARDS and OTHER ROW CROPS

CROP/ SITUATION	WEEDS	STATE	RATE	WHP	CRITICAL COMMENTS
Blackberry, boysenberry, loganberry, raspberry	Primocane and sucker control	NSW, ACT, Vic, Tas only	200 g /100 L water	NR (H) 8 weeks (G)	Apply as a directed spray to suckers and primocanes. Contact with flowers, developing fruit or desirable foliage will cause damage. Ensure complete coverage of primocanes / suckers by spraying to the point of runoff, preferably when they are less than 15 cm high. A non-ionic wetting agent (1000 g/L) may be added at a rate of 25 mL/100 L or equivalent.
Avocado, banana, feijoa, guava, kiwifruit, litchi, mango, pawpaw, passionfruit, pineapple, rambutan plantations	See lists of weeds controlled in Tables 1 and 2.	Qld, NSW, ACT, Vic, SA, WA, NT only	400 g to 2 kg/ha	NR (H) 8 weeks (G)	Apply as a directed or shielded spray. Refer to the label section **Application Equipment** for specific information on application methods. Controlled Droplet Application equipment must **not** be used for application in cherry orchards. **Warnings:** Do not allow spray or spray drift to contact desirable foliage or green (uncalloused) bark. To avoid potential crop damage, refer to the label sections on **Application Equipment** and **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS.** Fascinate Dry may be used around trees/vines less than two years old provided they are effectively shielded from spray and spray drift. The recommended rate of use is determined by the following criteria: **WEED SPECIES** **WEED STAGE OF GROWTH** **WEED DENSITY** **CLIMATIC CONDITIONS** **WEED SPECIES** Apply the appropriate rate to control the least susceptible weed present as per the lists of weeds controlled in the accompanying tables. W**EED STAGE OF GROWTH** Use the lower rate when weeds are young and succulent (grasses: pre-tillering; broadleaves: cotyledons to 4-leaf) or the population is very sparse. A median rate should be used for medium sized plants (grasses: tillering; broadleaves: 4 leaf to advanced vegetative) and the high rate should be used when weeds are mature (grasses: noding to flowering; broadleaves: budding to flowering). **WEED DENSITY** Use the higher rates when the weed population is dense. Thorough coverage of weeds is essential for good control. **CLIMATIC CONDITIONS** Best results are achieved when applied under warm humid conditions. Control will be reduced and/or slower under cold conditions
Citrus orchards		All States			
Olive plantations					
Pome and stone fruit orchards				21 days (H) 8 weeks (G)	
Tree nut plantations				NR (H) 8 weeks (G)	
Vineyards					

					and/or overcast conditions. Good results will be achieved under most other conditions, however poor results may occur under hot, dry conditions (temperatures above 33ºC with a relative humidity below 50%). Weeds that have been hardened or stunted in growth due to stressed conditions should be treated at the maximum rate. **COVERAGE:** Complete coverage of weeds is essential for good control. Poor coverage may result in re-growth. **PERENNIAL WEEDS:** Apply when weeds are actively growing. Follow up treatments will be necessary to control re-growth of perennial weeds in most cases.
Strawberries, cane berry fruits (inter-row) Tomatoes (inter-row)	See lists of weeds controlled in Tables 1 and 2.	All States	400 g to 2 kg/ha	NR (H) 8 weeks (G)	Apply as a directed or shielded spray to the inter-row area. Take care not to allow spray or spray drift to contact the crop, including strawberry runners. Refer to **GENERAL INSTRUCTIONS** for warnings concerning plastic mulch and fumigated/sterilised soil. Determine the recommended rate of use by considering the criteria WEED SPECIES, WEED STAGE OF GROWTH, WEED DENSITY and CLIMATIC CONDITIONS, as described above.

B. COMMERCIAL, INDUSTRIAL, NON-AGRICULTURAL AREAS and FORESTRY PLANTATIONS

CROP/ SITUATION	WEEDS	STATE	RATE	WHP	CRITICAL COMMENTS
Commercial & Industrial areas, rights-of-way and other non-agricultural areas	See lists of weeds controlled in Tables 1 and 2.	All States	400 g to 2.4 kg /ha	-	Determine the recommended rate of use by considering the criteria WEED SPECIES, WEED STAGE OF GROWTH, WEED DENSITY and CLIMATIC CONDITIONS as described above. Warnings: Do not allow spray or spray drift to contact desirable plants. To avoid potential crop damage, refer to the label sections on **Application Equipment** and **PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS.**
Commercial & Industrial areas, rights-of-way and other non-agricultural areas	Volunteer or wildling *Pinus* spp.		Handgun and Knapsack application 200 g/ 100 L water		Fascinate Dry is a non-selective herbicide and will affect most weeds. Its forestry use is designed to improve the control of *Pinus* spp. wildings when pre-plant weed control is carried out. To broaden the weed spectrum, mixing with other herbicides such as glyphosate and metsulfuron-methyl at labelled rates may be necessary. **APPLICATION** Apply with an adjuvant. The addition of an adjuvant e.g. Nu-Film® P or Exit® may assist in improving performance. High water volumes or nozzle systems should be used to achieve complete coverage of weeds, which is essential for good control. Handgun and knapsack rates are based on the application of 1000 L of spray mixture per sprayed hectare. This is usually adequate to thoroughly wet dense stands of weeds. Less dense stands will require lower water rates. Fascinate Dry does not provide residual weed control. Refer also to comments in the General Instructions which relate to application. **WEED GROWTH STAGE AND CONDITION** Use on *Pinus* is recommended to maximise efficacy. Apply when weeds are actively growing. Results will be reduced if treated plant is under stress due to very dry, very wet, frosty or diseased conditions. **COVERAGE** Complete coverage of target is essential for good control. Poor coverage may result in re-growth. **CLIMATIC CONDITIONS** **Best results are achieved when applied under warm, humid conditions (**temperatures below 33°C with a relative humidity above 50%)**.** Good results will be achieved under most other conditions, however poor results may occur under hot, dry conditions. Trials have shown better results from autumn and winter applications than from spring and summer applications. **SYMPTOMS**
Forestry plantations (preplant plantation establishment)			2 kg/ha		

					Visible symptoms will appear within 3 weeks; tree death may take several months depending on initial coverage and size of tree. Follow up treatments may be necessary to control re-growth in some cases.
Line-marking on sports grounds	Turf grasses and any weeds	All States	100 g to 200 g /100 L water	-	Refer to General Instructions. Fascinate Dry is a non-selective, non-residual herbicide with limited translocation potential. It is therefore ideally suited for line-marking on sports fields where precise weed control is required. Apply at 6 – 8 week intervals depending on growth of turf. Apply using single boom or hand wand.

C. SUMMER FALLOW SITUATIONS

CROP/ SITUATION	WEEDS	WEED STAGE	RATE	WHP	CRITICAL COMMENTS
Maintenance of summer fallow prior to planting; **Cereal grains** (including wheat, barley, oats, maize and sorghum) **Pulses** (including chickpeas, faba beans, field peas, lentils, lupins and mungbeans), **Oilseed**s (including canola, cotton, soybeans and sunflowers) Do not sow crops until 14 days or more have elapsed after the final application.	**Control of:** Annual polymeria Bellvine Bladder ketmia Caltrop Dwarf amaranth Field bindweed (European bindweed) Flax-leaf fleabane Paddy melon Peach vine Red pigweed Rhyncho (Rhyncosia) Sesbania pea Sowthistle (Milk thistle) Volunteer cotton (other than Liberty Link cotton) Yellow vine **Suppression of:** Chinese lantern (Wild gooseberry) Noogoora burr complex	2-6 leaf	1.5 kg / ha in a minimum of 100 L water	6 weeks (G)	Apply to actively growing weeds. Good coverage is essential. Refer '**Application'** section for details. Do not apply more than three applications per season. Fascinate Dry will have an effect on weeds that are larger than the recommended leaf stage, but speed of activity and level of control may be reduced. **CLIMATIC CONDITIONS** **Best results are achieved when Fascinate Dry is applied under warm humid conditions (temperatures below 33°C with a relative humidity above 50%).** Under any other conditions efficacy and speed of action may be reduced. Do not apply onto weeds when dew, fog or mist is present.

NR- Not Required; H-Harvest; G-Grazing

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

DIRECTIONS FOR USE (continued)
Table 1: Recommendations for weed control (except when referred to Table 2).

COMMON NAME	SCIENTIFIC NAME	APPLICATION RATE		
		Boom or directed sprayer kg/ha	Handgun g/100 L	Knapsack g/15 L
ANNUAL WEEDS				
Amaranthus spp.	*Amaranthus* spp.	0.8 to 2.0	200	30
Apple of Peru	*Nicandra physalodes*	0.6 to 1.2	120	18
Argentine peppercress	*Lepidium bonariense*	0.8 to 1.2	120	18
Awnless barnyard grass	*Echinochloa colona*	1.0 to 1.4	140	21
Barley grass	*Hordeum leporinum*	0.8 to 1.2	120	18
Barnyard grass	*Echinochloa crus galli*	0.8 to 2.0	200	30
Billy goat weed	*Ageratum conyzoides*	0.8 to 2.0	200	30
Bitter cress	*Cardamine hirsuta*	0.8 to 2.0	200	30
Black bindweed (buckwheat) (refer Note 2)	*Fallopia convolvulus*	0.7 to 2.0	200	30
Bladder ketmia	*Hibiscus trionum*	1.2 to 2.0	200	30
Bordered panic	*Entolasia marginata*	0.8 to 1.6	160	24
Brome grasses (refer Note 1)	*Bromus* spp.	0.8 to 1.2	120	18
Calopo	*Calopogonium mucunoides*	0.8 to 2.0	200	30
Caltrop burr (refer also Table 2)	*Tribulus terrestris*	1.2 to 2.0	200	30
Cape weed	*Arctotheca calendula*	0.6 to 2.0	200	30
Clover (subterranean)	*Trifolium subterraneum*	0.7 to 1.2	120	18
Cobbler's peg	*Bidens pilosa*	0.8 to 2.0	200	30
Common storksbill	*Erodium cicutarium*	0.6 to 1.6	160	24
Crowsfoot grass	*Eleusine indica*	1.2 to 2.0	200	30
Dead nettle (refer also Table 2)	*Lamium amplexicaule*	0.8 to 2.0	200	30
Dwarf crumbweed	*Chenopodium pumilo*	1.2 to 2.0	200	30
Fat hen	*Chenopodium album*	1.2 to 2.0	200	30
Fumitory	*Fumaria officinalis*	0.7 to 2.0	200	30

DIRECTIONS FOR USE (continued)

COMMON NAME	SCIENTIFIC NAME	APPLICATION RATE		
		Boom or directed sprayer kg/ha	Handgun g/100 L	Knapsack g/15 L
ANNUAL WEEDS (continued)				
Green crumbweed	*Chenopodium carinatum*	0.8 to 2.0	200	30
Lesser canary grass (refer also Table 2)	*Phalaris minor*	1.2 to 2.0	200	30
Liverseed grass (refer also Table 2)	*Urochloa panicoides*	0.6 to 2.0	200	30
Medics (annual)	*Medicago* spp.	0.4 to 2.0	200	30
Milk thistle	*Sonchus oleraceus*	0.8 to 2.0	200	30
Mint weed	*Salvia reflexa*	1.2 to 2.0	200	30
New Zealand spinach	*Tetragonia tetragoniodes*	0.8 to 2.0	200	30
Patterson's curse	*Echium plantagineum*	0.4 to 1.2	120	18
Peanuts	*Arachis hypogaea*	0.6 to 1.2	120	18
Pigweed	*Portulaca oleracea*	1.2 to 2.0	200	30
Pinkburr	*Urena lobata*	0.8 to 2.0	200	30
Potato weed	*Galinsoga parviflora*	0.8 to 2.0	200	30
Prairie grass (refer Note 1)	*Bromus unioloides*[1]	1.6 to 2.0	200	30
Prickly lettuce	*Lactuca serriola*	1.2 to 2.0	200	30
Red natal grass	*Rhynchelytrum repens*	0.8 to 2.0	200	30
Ryegrass (annual)	*Lolium rigidum*	0.8 to 2.0	200	30
Saffron thistle	*Carthamus lanatus*	0.6 to 2.0	200	30
St. Barnaby's thistle	*Centaurea solstitialis*	0.6 to 2.0	200	30
Sago weed	*Plantago cunninghamii*	0.8 to 1.2	120	18
Scarlet pimpernel	*Anagallis arvensis*	0.8 to 2.0	200	30
Setaria	*Setaria italica*	0.8 to 2.0	200	30
Sheep thistle	*Carduus tenuiflorus*	1.0 to 2.0	200	30
Silver grass	*Vulpia myuros*	0.8 to 2.0	200	30
Sorghum/sudax	*Sorghum bicolor*	0.8 to 2.0	200	30
Square weed	*Spermacoce latifolia*	0.8 to 2.0	200	30
Stagger weed	*Stachys arvensis*	0.8 to 2.0	200	30
Star of Bethlehem	*Ipomoea quamoclit*	0.8 to 2.0	200	30
Summer grass	*Digitaria ciliaris*	0.8 to 2.0	200	30
Thickhead	*Crassocephalum crepidioides*	1.2 to 2.0	200	30
Three cornered jack	*Emex australis*	0.8 to 2.0	200	30
Tomato	*Lycopersicon esculentum*	0.8 to 2.0	200	30
Townsville stylo	*Stylosanthes humilis*	0.4 to 1.2	120	18
Turnip weed	*Rapistrum rugosum*	1.2 to 2.0	200	30
Variegated thistle (refer also Table 2)	*Silybum marianum*	1.0 to 2.0	200	30
Wheat	*Triticum aestivum*	1.6 to 2.0	200	30
Wild carrot	*Daucus glochidiatus*	0.8 to 2.0	200	30

Wild gooseberry	*Physalis minima*	0.8 to 2.0	200	30
Wild mustard	*Sysimbrium orientale*	0.8 to 2.0	200	30
Wild oats (refer also Table 2)	*Avena* spp.	1.2 to 2.0	200	30
Wild radish	*Raphanus raphanistrum*	2.0	200	30
Wire weed (refer also Table 2)	*Polygonum aviculare*	0.6 to 2.0	200	30
PERENNIAL WEEDS				
Blady grass	*Imperata cylindrica*	1.2 to 1.6	160	24
Cape tulip	*Homeria* spp.	0.8 to 1.2	120	18
Centro	*Centrosema pubescens*	0.4 to 2.0	200	30
Clover glycine	*Glycine latrobeana*	0.4 to 1.2	120	18
Couch grass	*Cynodon dactylon*	1.0 to 2.0	200	30
Cow pea	*Vigna unguiculata*	0.4 to 1.2	120	18
Giant sensitive plant	*Mimosa invisa*	0.8 to 2.0	200	30
Greenleaf desmodium	*Desmodium intortum*	0.4 to 1.2	120	18
Johnson grass	*Sorghum halepense*	1.2 to 2.0	200	30
Panicum spp.	*Panicum* spp.	0.8 to 2.0	200	30
Paspalum spp.	*Paspalum* spp.	1.2 to 2.0	200	30
Perennial bindweed	*Convolvulus arvensis*	0.8 to 1.2	120	18
Shamrock	*Oxalis corymbosa*	1.2	120	18
Sida weed (refer also Table 2)	*Sida retusa*	1.2 to 2.0	200	30
Silver leaf desmodium	*Desmodium uncinatum*	1.6 to 2.0	200	30
Siratro	*Macroptilium atropurpureum*	0.4 to 1.2	120	18
Stink grass	*Eragrostis cilianensis*	1.2 to 2.0	200	30
White clover	*Trifolium repens*	1.2 to 2.0	200	30
White eye	*Richardia brasiliensis*	1.2 to 2.0	200	30
Willow herb	*Epilobium* spp.	1.6 to 2.0	200	30

Notes:
1. Well established clumps of Prairie grass and Brome grasses may only be suppressed at these rates. Follow-up treatments may be necessary to control regrowth.
2. Good control will be achieved on small and medium sized plants only in non-crop situation.

Table 2: For control of weeds in Commercial and Industrial areas, rights of way and other non-agricultural areas (when referred from Table 1).

COMMON NAME	SCIENTIFIC NAME	APPLICATION RATE		
		Boom or directed sprayer (kg/ha)	Handgun (g/100 L)	Knapsack (g/15 L)
ANNUAL WEEDS				
Caltrop burr	*Tribulus terrestris*	1.6 to 2.0	200	30
Dead nettle	*Lamium amplexicaule*	2.4	240	36
Lesser canary grass	*Phalaris minor*	1.6 to 2.4	240	36
Liverseed grass	*Urochloa panicoides*	0.6	60	9
Variegated thistle	*Silybum marianum*	2.4	240	36
Wild oats	*Avena* spp.	2.0 to 2.4	240	36
Wire weed	*Polygonum aviculare*	0.8 to 2.0	200	30
PERENNIAL WEEDS				
Sida weed	*Sida retusa*	1.6 to 2.0	200	30