Product Name: SHENZI INSECTICIDE
APVMA Approval No: 92656/136242

RLP
APPROVED

Label Name:	SHENZI INSECTICIDE

Signal Headings:	READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Constituent Statements:	ACTIVE CONSTITUENT: 400 g/L CHLORANTRANILIPROLE

Mode of Action:	GROUP 28 INSECTICIDE

Statement of Claims:	For the control of Lepidopteran species of insect pests in certain Vegetable, Fruit, Cotton and Pulse crops, as per the Directions for Use.

Net Contents:	500 mL - 10 L

Restraints:	This section contains file attachment.

Directions for Use:	This section contains file attachment.

Other Limitations:	

Withholding Periods:	WITHHOLDING PERIODS HARVEST POTATOES: WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED. FRUITING VEGETABLES (CUCURBITS INCLUDING CUCUMBERS, MELONS, PUMPKIN, SQUASH, ZUCCHINI), LEGUME VEGETABLES, STRAWBERRIES: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

	BRASSICA LEAFY VEGETABLES, FRUITING VEGETABLES (excluding CUCURBITS), LEAFY VEGETABLES (including LETTUCE), CELERY and RHUBARB: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION. BRASSICA VEGETABLES (INCLUDING BROCCOLI, BRUSSELS SPROUT, CABBAGE, CAULIFLOWER), SWEET CORN: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION. ALMONDS, POME AND STONE FRUIT: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION. GRAPES: DO NOT HARVEST FOR 8 WEEKS AFTER APPLICATION. COTTON: DO NOT HARVEST FOR 28 DAYS AFTER APPLICATION. CHICKPEA, MUNG BEAN, SOYBEAN: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION. PULSE CROPS (EXCEPT CHICKPEA, MUNG BEAN, SOYBEAN): DO NOT HARVEST FOR 28 DAYS AFTER APPLICATION. GRAZING LEGUME VEGETABLES: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 1 DAY AFTER APPLICATION. SWEET CORN: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION. COTTON: DO NOT ALLOW LIVESTOCK TO GRAZE CROPS, COTTON STUBBLE ORGIN TRASH TREATED WITH SHENZI INSECTICIDE. PULSE CROPS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION. ALMONDS, POME AND STONE FRUIT, GRAPES: DO NOT GRAZE OR CUT FOR STOCK FOOD. OTHER CROPS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

Trade Advice:	Growers should note that maximum residue limits (MRLs) or import tolerances may not exist in all markets for produce treated with Shenzi. If you are growing produce for export, please check with UPL Australia Pty Ltd, your exporter or industry body for the latest information on MRLs and import tolerances before using Shenzi.

General Instructions:	GENERAL INSTRUCTIONS Shenzi is an anthranilic diamide insecticide in the form of a suspension concentrate. Shenzi is particularly active on Lepidopteran insect pests, primarily as a larvicide. Shenzi should be applied after careful field monitoring of pest populations of eggs and larvae to determine the need for application, the correct timing of th e initial application and of any subsequent applications. Subsequent applications are dependent on locally determined economic thresholds, as well as the growth rate of new unprotected plant material. More than one treatment of Shenzi Insecticide may be required to control a population of pests. For Helicoverpa species, spray applications should be timed to coincide with egg hatching and before larvae are entrenched in protected feeding sites. Shenzi Insecticide has been specifically designed for use in Integrated Pest Management (IPM) schemes. Shenzi does not give traditional larval ‘knockdown’ control. It enters the larvae primarily by ingestion of treated foliage, or through penetration of the insect cuticle. After ingesting Shenzi, the larvae cease feeding and die four to five days later. Shenzi provides square, flower and boll protection in cotton and flower and pod protection in pulse crops. MIXING Fill spray tank to ¼ to ½ full of water. Measure the amount of Shenzi Insecticide required for the area to be sprayed. Add Shenzi Insecticide directly to the spray tank with the agitation engaged. Mix thoroughly to disperse the insecticide. Once dispersed, the material must be kept in suspension at all times by continuous agitation. Use mechanical or hydraulic means, DO NOT use air agitation, premix or slurry.

	If spray solution is left standing, ensure thorough re-agitation of the spray mix until fully resuspended. DO NOT allow spray mix to sit overnight, as resuspension may be difficult. SURFACTANT/WETTING AGENT Use a non ionic surfactant/wetting agent as per the directions for use table. DO NOT use BS1000* or Activator* as it may cause crop phytotoxicity. DO NOT add a non-ionic surfactant/wetting agent if: • mixing with another product which already contains a surfactant and/or the product label advises not to add a surfactant. • mixing with a liquid fertiliser. APPLICATION Application equipment should be calibrated to apply at least sixty (60) droplets per cm^2 of target foliage. Droplet VMD should be of medium spray quality. Ground application Use a boom sprayer fitted with high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets. Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size, DOES NOT improve canopy penetration and may increase drift potential. WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE. Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. When applying Shenzi by ground application, keep the boom low to avoid spray drift. For orchard/vineyard sprayers avoid directing spray above trees and always turn-off outward pointing nozzles at row ends and outer ends. For band spraying, increase the number of fan nozzles per crop row as the plant size increases. Aerial application (by fixed wing aircraft or helicopter) Shenzi must only be applied with aircraft fitted with accurately calibrated equipment. Apply a minimum total spray volume of 30 L/ha with nozzles (e.g. Micronaire rotary atomisers, CP nozzles or conventional hydraulic nozzles) set to medium spray quality. A spray drift minimisation strategy should be employed at all times when applying this product. DO NOT apply Shenzi using Ultra Low Volume (ULV) methods. COMPATIBILITY It is recommended that users premix a small quantity of the desired tank mix and observe possible adverse changes (settling out, flocculation etc). Physical compatibility does not guarantee biological compatibility. Not all tank mix partners have been tested. Please consult with a UPL representative for further information. SPRAY EQUIPMENT CLEANOUT Prior to application, start with clean, well-maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove. Drain spray equipment. Thoroughly rinse sprayer and flush hoses, boom, and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. DO NOT clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

Resistance Warning:	For insecticide resistance management Shenzi Insecticide is a Group 28 insecticide. Some naturally occurring insect biotypes resistant to Shenzi Insecticide and other Group 28 insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Shenzi and other Group 28 insecticides are used repeatedly. The effectiveness of Shenzi Insecticide on resistant individuals could be significantly reduced. Since the occurrence of

	resistant individuals is difficult to detect prior to use UPL accepts no liability for any losses that may result from the failure of Shenzi Insecticide to control resistant insects. Shenzi may be subject to specific resistance management strategies. For further information refer to the Insecticide Resistance Management (IRM) section of this label, contact your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, or local UPL Representative. INSECTICIDE RESISTANCE MANAGEMENT (IRM) Shenzi Insecticide contains chlorantraniliprole, a Group 28 Insecticide. Unless directed otherwise in the specific crop/insect sections of the label, the following practices are recommended to prevent or delay the development of insecticide resistance to Shenzi Insecticide and to Group 28 insecticides: • Apply Shenzi Insecticide or other Group 28 insecticides using a "window" approach to avoid exposure of consecutive insect pest generations to the same mode of action. Multiple successive applications of Shenzi or other Group 28 insecticides are acceptable if they are used to treat a single insect generation. • Following a "window" of Shenzi Insecticide or other Group 28 insecticides, rotate to a "window" of applications of effective insecticides with a different mode of action. • The total exposure period of all "Group 28-active windows" applied throughout the crop cycle (from seedling to harvest) should not exceed 50% of the crop cycle. • Incorporate IPM techniques into the overall pest management program. • Monitor insect populations for loss of field efficacy. • Cultivate all cotton and pulse crop fields as soon as possible after picking/harvest to destroy over-wintering pupae of Helicoverpa armigera. For further information contact your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, or local UPL representative. For additional information on insect resistance, modes of action and monitoring visit the Insecticide Resistance Action Committee (IRAC) on the web at http://www.irac-online.org

Precautions:	PRECAUTIONS DO NOT use human flaggers/markers unless they are protected by engineering controls such as vehicles with enclosed cabs.

Protections:	PROTECTIONS PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers. PROTECTION OF HONEY BEES AND OTHER INSECT POLLINATORS Based on Good Agricultural Practices (GAP), DO NOT apply when bees are actively foraging. PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto near-by non-target plants/crops, cropping lands or pastures. IMPORTANT: Not all crops within a crop group, and not all varieties, cultivars or hybrids of crops, have been individually tested for crop safety. To test for crop safety, apply the product in accordance with the label instructions to a small area of the target crop to ensure that a phytotoxic response will not occur, especially where the application is a new use of the product by the applicator.

Storage and Disposal:	Keep out of reach of children. Store in the closed, original container in a cool, well-ventilated area out of direct sunlight. Drummuster eligible containers This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank. Do not dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be taken separately. Plastic containers Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Safety Directions:	May irritate the eyes. Avoid contact with eyes. Wash hands after use.

First Aid Instructions:	First aid is not generally required. If in doubt, contact a Poisons Information Centre (e.g. phone Australia 13 11 26; New Zealand 0800 764 766 or a doctor).

First Aid Warnings:	

RESTRAINTS
DO NOT make more than 3 applications per cotton crop per season, and no more than 2 consecutive sprays per field per season.
DO NOT make more than 1 application per pulse crop per season (except chickpea, soybean and mung bean where a maximum of 2 applications can be made).
DO NOT apply if heavy rains or storms are forecast within 3 days.
DO NOT irrigate to the point of field runoff for at least 3 days after application.
DO NOT allow effluent or run-off from protected cropping systems containing this product to enter dams, streams, ponds or other waterways.
DO NOT apply by aircraft on almonds, pome fruit, stone fruit and grapes.

SPRAY DRIFT RESTRAINTS
Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.

DO NOT allow bystanders to come into contact with the spray cloud.
DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. The buffer zones in the relevant buffer zone table/s below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.
DO NOT apply unless the wind speed is between three and 20 kilometres per hour at the application site during the time of application.
DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.
DO NOT apply by a boom sprayer unless the following requirements are met:
• Spray droplets are not smaller than a MEDIUM spray droplet size category.
• Minimum distances between the application site and downwind sensitive areas are observed (see the following table titled ‘Buffer zones for boom sprayers’).

Buffer zones for boom sprayers

Application rate	Boom height above target canopy	Mandatory downwind buffer zones				
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
Up to 130 mL/ha	0.5 m or lower	0 metres	20 metres	0 metres	0 metres	0 metres
	1.0 m or lower	0 metres	55 metres	0 metres	0 metres	0 metres
Up to 75 mL/ha	0.5 m or lower	0 metres	10 metres	0 metres	0 metres	0 metres
	1.0 m or lower	0 metres	35 metres	0 metres	0 metres	0 metres

DO NOT apply by a vertical sprayer unless the following requirements are met:
• Spray is not directed above the target canopy.
• The outside of the sprayer is turned off when turning at the end of rows and when spraying the outer row on each side of the application site.
• For dilute water rates up to the maximum listed for each type of canopy specified, minimum distances between the application site and downwind sensitive areas are observed (see the following table titled 'Buffer zones for vertical sprayers').

Buffer zones for vertical sprayers

Application rate	Maximum dilute water rate	Type of target canopy	Mandatory downwind buffer zones				
			Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
Up to 16 mL/100mL	1000 L/ha	2 metres tall and shorter	0 metres	10 metres	0 metres	0 metres	0 metres
	1500 L/ha	Taller than 2 metres (not fully-foliated)	0 metres	30 metres	0 metres	0 metres	0 metres
		Taller than 2 metres (fully-foliated)	0 metres	20 metres	0 metres	0 metres	0 metres
Up to 8 mL/100 mL	1000 L/ha	2 metres tall and shorter	0 metres	5 metres	0 metres	0 metres	0 metres
	1500 L/ha	Taller than 2 metres (not fully-foliated)	0 metres	25 metres	0 metres	0 metres	0 metres
		Taller than 2 metres (fully-foliated)	0 metres	15 metres	0 metres	0 metres	0 metres

DO NOT apply by aircraft unless the following requirements are met:
• Spray droplets are not smaller than a MEDIUM spray droplet size category.
• For maximum release height above the target canopy of 3 metres or 25 per cent of wingspan or 25 per cent of rotor diameter, whichever is the greatest, minimum distances between the application site and downwind sensitive areas are observed (see the following table titled 'Buffer zones for aircraft')

Buffer zones for aircraft

Application rate	Type of aircraft	Mandatory downwind buffer zones				
		Bystander areas	Natural aquatic areas	Pollinator areas	Vegetation areas	Livestock areas
Up to 240 mL/ha	Fixed-wing	0 metres	375 metres	0 metres	0 metres	0 metres
	Helicopter	0 metres	230 metres	0 metres	0 metres	0 metres
Up to 130 mL/ha	Fixed-wing	0 metres	220 metres	0 metres	0 metres	0 metres
	Helicopter	0 metres	160 metres	0 metres	0 metres	0 metres

DIRECTIONS FOR USE

For use in all States where appropriate for the crop and/or insect pest.

CROP	PEST	RATE/HA	WHP	CRITICAL COMMENTS
CRITICAL COMMENTS - ALL CROPS Regularly scout crops to monitor for eggs and larvae. Target sprays against eggs and newly hatched larvae before they become entrenched. Apply as egg and larvae reach threshold numbers. A maximum of three (3) applications are to be applied to any one crop. No more than two (2) consecutive sprays per crop, with a minimum spray interval of 7 days (unless stated otherwise). Further treatments should be made with alternative mode of action insecticides. Use enough water to ensure thorough coverage of the crop. Adjust water volumes to crop stage (200 – 1000 L/ha). For dilute spraying- Apply sufficient water to cover the crop to the point of run-off but avoid excessive run-off. The required dilute spray volume will change as the crop grows therefore sprayer set up needs to be adjusted accordingly. Refer to Surfactant/Wetting agent section. Use in accordance with AIRAC Insecticide Resistance Management Strategy guidelines. As part of an Insecticide Resistance Management programme for Cotton bollworm, it is important to plough crops immediately after harvest.				
Brassica vegetables including: Broccoli Brussels sprout Cabbage Cauliflower	Cabbage-centre grub (*Hellula hydralis*) Cabbage cluster caterpillar (*Crocidolomia pavonana*) Cabbage leafminer (*Liriomyza brassicae*) Cabbage white butterfly (*Pieris rapae*) Cluster caterpillar (*Spodoptera litura*) Cotton bollworm (*Helicoverpa armigera*) Diamondback moth (*Plutella xylostella*) Native budworm (*Helicoverpa punctigera*) Soybean looper (*Thysanoplusia orichalcea*)	50 mL + 15 gai/100 L of non-ionic surfactant or 5 mL/100 L (dilute) + 15 gai/100 L of non-ionic surfactant	7 days	
Brassica leafy vegetables including: Buk choy Chinese broccoli (Gai lum/Gai lan/Kai lan), Chinese cabbage (Pet sai/Wombok/Haksukai) Choy sum Gai choy/Am soy Kai choy Kale Mibuna Leafy mustard including Indian mustard and Mustard spinach (Komatsuma) Pak choy Tat soy (for field and protected cropping systems)			3 days	
Stalk & Stem vegetables: Celery Rhubarb	Cotton bollworm (*Helicoverpa armigera*) Native budworm (*Helicoverpa*	50 mL + 15 gai/100 L of non-ionic		

Leafy vegetables (excluding lettuce), including: Cress Endive Silverbeet Spinach (for field and protected cropping systems)	*punctigera*)	surfactant or 5 mL/100 L (dilute) + 15 gai/100 L of non-ionic surfactant		
Lettuce (leaf and closed head varieties) (for field and protected cropping systems)	Cotton bollworm (*Helicoverpa armigera*) Native budworm (*Helicoverpa punctigera*)	75 mL + 15 gai/100 L of non-ionic surfactant or 7.5 mL/100 L (dilute) + 15 gai/100 L of non-ionic surfactant	3 days	
Fruiting vegetables (excluding Cucurbits), including: Capsicum Egg plant Peppers Tomato (trellis and field) (for field and protected cropping systems)	Cotton bollworm (*Helicoverpa armigera*) Native budworm (*Helicoverpa punctigera*) Tomato leaf miner (*Phthorimaea operculella*)	50 mL or 5 mL/100 L (dilute)	3 days	
	Eggfruit caterpillar (*Sceliodes cordalis*)			Ensure spray timing coincides with egg laying/hatching. The pest hatches from the egg and burrows directly into fruit. Larvae entrenched in the fruit at the time of spraying will not be controlled.
Fruiting vegetables (Cucurbits), including: Cucumbers Melons Pumpkin Squash Zucchini (for field and protected cropping systems)	Cotton bollworm (*Helicoverpa armigera*) Native budworm (*Helicoverpa punctigera*) Cucumber moth (*Diaphania indica*)	50 mL + 15 gai/100 L of non-ionic surfactant or 5 mL/100 L (dilute) + 15 gai/100 L of non-ionic surfactant	1 day	Apply with a minimum spray interval of 5 days.
Legume vegetables including: Green beans Green peas Processing peas Snow peas Sugar snap peas (for field and protected cropping systems)	Cotton bollworm (*Helicoverpa armigera*) Native budworm (*Helicoverpa punctigera*)	50 mL + 15 gai/100 L of non-ionic surfactant) or 5 mL/100 L (dilute) + 15 gai/100 L of non-ionic surfactant	1 day	
Potatoes	Cotton bollworm (*Helicoverpa armigera*)	50 mL	Not required	Only target foliar infestations of Potato moth. Moth larvae in the soil or within stems will not be controlled. Apply with

	Native budworm (*Helicoverpa punctigera*) Potato moth (*Phthorimaea operculella*)			a spray interval of 10 - 14 days.
Strawberries (for field and protected cropping systems)	Cluster caterpillar (*Spodoptera litura*) Cotton bollworm (*Helicoverpa armigera*) Native budworm (*Helicoverpa punctigera*)	50 mL + 15 gai/100 L of non-ionic surfactant or 5 mL/100 L (dilute) + 15 gai/100 L of non-ionic surfactant	1 day	
Sweet corn	Cotton bollworm (*Helicoverpa armigera*)	50 mL + 15 gai/100 L of non-ionic surfactant	7 days	Ensure spray timing coincides with egg laying/hatching. Larvae entrenched in cobs at the time of spraying will not be controlled.

CROP	PEST	RATE/100 L	WHP	CRITICAL COMMENTS
ALL CROPS Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. Refer to Application section of the label. Thorough fruit and nut coverage is essential. Use in accordance with AIRAC Insecticide Resistance Management Strategy guidelines.				
Almonds	Carob moth (*Ectomyelois ceratoniae)*	**Dilute spraying:** 16 mL + non ionic surfactant @ 15 gai/100 L **Concentrate spraying:** Refer to **Mixing/ Application** section	14 days	Monitor moth flights through trapping. Applications can be aimed at reducing moth number by targeting mummies. Alternatively, applications should be made when 1 - 5% of the almond hull sutures are opening. If required, retreat at minimum interval of 7 days. Apply by airblast sprayer or equivalent and ensure thorough coverage of all leaf and nut surfaces. **DO NOT** apply more than 2 applications of Shenzi per season. **DO NOT** treat successive generations with Shenzi, alternate with a different mode of action insecticide.
Pome fruit including	Codling moth	**Dilute spraying:** 8 mL + non ionic surfactant @15 gai/100L **Concentrate spraying:** Refer to **Mixing/**	14 days	**DO NOT** make more than three (3) applications per crop per season. Codling moth: A maximum of three (3) applications of Shenzi are to be applied at 14 – 21 day intervals commencing at petal fall (or before 110 Degree Days after Codling moth are detected in traps) until late December. Further treatments should be made with an alternate mode of action insecticide. **Or** a maximum of three (3) applications can be applied commencing from the end of December at 14 - 21 day intervals following treatments with an alternate mode of action product. Oriental fruit moth: When treating the first

Apples Nashi Pears Pears	(*Cydia pomonella*) Budworms (*Helicoverpa* spp). Oriental fruit moth (*Grapholita molesta*)	**Application** section		generation, apply the initial treatment before 110 Degree Days after Oriental fruit moths are detected in traps. The above programme, when commenced at petal fall, will also control Budworms.
	Lightbrown apple moth (*Epiphyas postvittana)*			Lightbrown apple moth: A maximum of three (3) applications of Shenzi Insecticide are to be applied at 14 - 21 day intervals commencing at petal fall or apply at 140 Degree Days after Lightbrown apple moths are detected in traps. Further treatments should be made with alternative mode of action insecticides.
Stone fruit including Apricots Cherries Nectarines Peaches Plums	Oriental fruit moth (*Grapholita molesta*)	**Dilute spraying:** 11 mL + non ionic surfactant @ 15 gai/100 L **Concentrate spraying:** Refer to **Mixing/ Application** section		**DO NOT** make more than two (2) applications per crop per season with a minimum 14 days between applications. When treating the first generation, apply the initial treatment before 110 Degree Days after Oriental fruit moths are detected in traps. A maximum of two (2) applications of Shenzi (minimum of 14 days between applications) to each crop. Target sprays against eggs and newly hatched larvae before they become entrenched. Further treatments should be made with alternative mode of action insecticides.

CROP	PEST	RATE/100 L	WHP	CRITICAL COMMENTS
	Lightbrown apple moth (*E. postvittana)*	**Dilute spraying:** 8mL + non ionic surfactant @ 15 gai/100 L **Concentrate spraying:** Refer to **Mixing/ Application** section		A maximum of two (2) applications of Shenzi Insecticide are to be applied with a minimum spray interval of 14 days commencing at 140 Degree Days after Lightbrown apple moths are detected in traps. Further treatments should be made with alternative mode of action insecticides.
Grapes	Lightbrown apple moth (*E. postvittana*) Grapevine moth (*Phalaenoides glycinae*)	**Dilute spraying:** 8mL + non ionic surfactant @ 15 gai/100 L **Concentrate spraying:** Refer to **Mixing/ Application** section	8 weeks	**DO NOT** make more than two (2) applications per crop per season. Applications to be timed for egg hatch (140 Degree Days after a detected moth flight). **DO NOT** re-treat within fourteen (14) days. A final application may be applied up to bunch closure. **DO NOT** apply after bunch closure. Concentrated spray: **DO NOT** apply in volumes less than 250 L/ha. This low water volume is dependent on the suitability of concentrated spray application equipment. More reliable application may be gained through increased water volumes.

For use in all States where appropriate for the crop and/or insect pest.

CROP	PEST	RATE/HA	WHP	CRITICAL COMMENTS
Cotton	Cotton bollworm (*Helicoverpa armigera*) Native budworm (*Helicoverpa punctigera*) Cluster caterpillar (*Spodoptera litura*)	80 mL or 130mL + non ionic surfactant @ 125 gai/100 L	28 days	Target brown eggs and hatchling (neonates or 1st instar) to small larvae (2nd instar) when they reach the economic spray threshold and before they become entrenched in squares, flowers and bolls. Use the **low** rate on **threshold larvae pressure** (2 larvae per metre row) and low egg pressure. Use the **high** rate with high egg and/or larvae pressure (where potential for >2 larvae per metre row produced) and so as to achieve longer residual control of *Helicoverpa* spp.
	Northern rough bollworm (*Earias vittella*) Rough bollworm (*Earias huegeliana*)	130 mL + non ionic surfactant @ 125 gai/100 L		Target eggs and hatchling (neonates or 1st instar) to small larvae (2nd instar) when they reach the economic spray threshold and before they become entrenched in terminals or bolls.
Chickpea	Cotton bollworm (*Helicoverpa armigera*) Native budworm (*Helicoverpa punctigera*)	60 mL + non-ionic surfactant @ 125 gai/100 L	14 days	A maximum of one application is to be applied to any one pulse crop per season (except Chickpea, Soybean and Mung bean where a maximum of 2 applications can be applied per season). Further treatments should be made with alternative mode of action insecticides. Regularly scout crops to monitor for larvae. Target sprays against larvae. Apply as larvae reach threshold numbers. Larvae in entrenched feeding sites will not be controlled. Use enough water to ensure thorough coverage of the crop. Target a minimum of 100 L/ha by ground rig and a minimum of 30 L/ha by aircraft.
Mung bean, Soybean	Bean podborer (*Maruca vitrata*) Cotton bollworm (*Helicoverpa armigera*)			

	Native budworm (*Helicoverpa punctigera*) Soybean looper (*Thysanoplusia orichalcea*) Bean looper (*Mocis alterna*) Irrorated tabby (*Anticarsia irrorata*)			Use in accordance with Crop Life Insecticide Resistance Management Strategy guidelines. Target brown eggs and hatchlings (neonates or first instar) to small larvae (second instar) when they reach the economic spray threshold and before they become entrenched in flowers or pods.
Winter pulse crops (except Chickpea) including; Faba/Broad bean, Field pea, Lentil, Lupin, Vetch	Cotton bollworm (*Helicoverpa armigera*) Native budworm (*Helicoverpa punctigera*)	60 mL + non-ionic surfactant @ 125 gai/100 L	28 days	
Summer pulse crops (except Mung bean and Soybean) including; Azuki/Adzuki bean, Cow pea, Navy bean, Pigeon pea,	Bean podborer (*Maruca vitrata*) Cotton bollworm (*Helicoverpa armigera*) Native budworm (*Helicoverpa punctigera*) Soybean looper (*Thysanoplusia orichalcea*) Bean looper (*Mocis alterna*) Irrorated tabby (*Anticarsia irrorata*)			

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION