

Textual

In this chapter, we will go through a number of R functions for basic statistics. The focus will be on the results that are presented in form of numbers in text or tables (textual). We will mostly use the builtin functions (from R standard library). Extra packages will be introduced whenever necessary.

Basic descriptive statistics

In this part, we are going to use the functions as applied to a variable. For this purpose, we are going to use builtin datasets in R. You can view the available datasets by

```
data()

## Data sets in package 'datasets':

## AirPassengers      Monthly Airline Passenger Numbers 1949-1960
## BJsales            Sales Data with Leading Indicator
## BJsales.lead (BJsales) Sales Data with Leading Indicator
## BOD                Biochemical Oxygen Demand
## CO2                Carbon Dioxide Uptake in Grass Plants
## ...
```

We can view any dataset description by appending “?” to the dataset name. For example,

```
?chickwts
```

We will start by using `chickwts` dataset that contains both numerical (`weight`) and categorical (`feed`) variables. We can view the first six observations,

```
head(chickwts)

##   weight    feed
## 1    179 horsebean
## 2    160 horsebean
## 3    136 horsebean
## 4    227 horsebean
## 5    217 horsebean
## 6    168 horsebean
```

the last six observations,

```
tail(chickwts)

##   weight    feed
## 66    352 casein
## 67    359 casein
## 68    216 casein
## 69    222 casein
## 70    283 casein
## 71    332 casein
```

and the dimension of the data (row and column).

```
dim(chickwts)
```

```
## [1] 71  2
```

Here we have 71 rows (71 subjects) and two columns (two variables).

Next, view the names of the variables,

```
names(chickwts)
```

```
## [1] "weight" "feed"
```

and view the details of the data,

```
str(chickwts)
```

```
## 'data.frame': 71 obs. of 2 variables:
## $ weight: num 179 160 136 227 217 168 108 124 143 140 ...
## $ feed : Factor w/ 6 levels "casein","horsebean",...: 2 2 2 2 2 2 2 2 2 2 ...
```

which shows that **weight** is a numerical variable and **feed** is a factor, i.e. a categorical variable. **feed** consists of six categories or levels.

We can view the levels in **feed**,

```
levels(chickwts$feed)
```

```
## [1] "casein" "horsebean" "linseed" "meatmeal" "soybean" "sunflower"
```

Describing a numerical variable

A numerical variable is described by a number of descriptive statistics below.

To judge the central tendency of the **weight** variable, we obtain its mean,

```
mean(chickwts$weight)
```

```
## [1] 261.3099
```

and median,

```
median(chickwts$weight)
```

```
## [1] 258
```

To judge its spread and variability, we can view its minimum, maximum and range

```
min(chickwts$weight)
```

```
## [1] 108
```

```
max(chickwts$weight)
```

```
## [1] 423
```

```
range(chickwts$weight)
```

```
## [1] 108 423
```

and obtain its standard deviation (SD)

```
sd(chickwts$weight)
```

```
## [1] 78.0737
```

variance,

```
var(chickwts$weight)
```

```
## [1] 6095.503
```

quantile,

```
quantile(chickwts$weight)
```

```
##      0%   25%   50%   75%  100%  
## 108.0 204.5 258.0 323.5 423.0
```

and interquartile range (IQR)

```
IQR(chickwts$weight)
```

```
## [1] 119
```

There are nine types of quantile algorithms in R (for `quantile` and `IQR`), the default being type 7. You may change this to type 6 (Minitab and SPSS),

```
quantile(chickwts$weight, type = 6)
```

```
##      0%   25%   50%   75%  100%  
##   108   203   258   325   423
```

```
IQR(chickwts$weight, type = 6)
```

```
## [1] 122
```

In addition to SD and IQR, we can obtain its median absolute deviation (MAD),

```
mad(chickwts$weight)
```

```
## [1] 91.9212
```

It is actually simpler to obtain most these in a single command,

```
summary(chickwts$weight)
```

```
##      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.  
##   108.0   204.5   258.0   261.3   323.5   423.0
```

even simpler, obtain all of the statistics using `describe` in the `psych` package

```
install.packages("psych")
```

```
library(psych)
```

```
describe(chickwts$weight)
```

```
##      vars  n   mean    sd median trimmed   mad min max range  skew kurtosis  
## X1      1 71 261.31 78.07   258      261 91.92 108 423   315 -0.01   -0.97  
##      se  
## X1 9.27
```

Describing a categorical variable

A categorical variable is described by its count, proportion and percentage by categories.

We obtain the count of the `feed` variable,

```
summary(chickwts$feed)
```

```
##      casein horsebean  linseed  meatmeal  soybean sunflower  
##          12         10         12         11         14         12
```

```
table(chickwts$feed)
```

```
##
##   casein horsebean   linseed meatmeal   soybean sunflower
##      12      10      12      11      14      12
```

both `summary` and `table` give the same result.

`prop.table` gives the proportion of the result from the count.

```
prop.table(table(chickwts$feed))
```

```
##
##   casein horsebean   linseed meatmeal   soybean sunflower
## 0.1690141 0.1408451 0.1690141 0.1549296 0.1971831 0.1690141
```

the result can be easily turned into percentage,

```
prop.table(table(chickwts$feed))*100
```

```
##
##   casein horsebean   linseed meatmeal   soybean sunflower
## 16.90141 14.08451 16.90141 15.49296 19.71831 16.90141
```

To view the count and the percentage together, we can use `cbind`,

```
cbind(n = table(chickwts$feed), "%" = prop.table(table(chickwts$feed))*100)
```

```
##           n           %
## casein    12 16.90141
## horsebean 10 14.08451
## linseed   12 16.90141
## meatmeal  11 15.49296
## soybean   14 19.71831
## sunflower 12 16.90141
```

We need the quotation marks " " around the percentage sign %, because % also serves as a mathematical operator in R.

More on descriptive statistics

Just now, we viewed all the statistics as applied to a variable. In this part, we are going to view the statistics on a number of variables. This includes viewing a group of numerical variables or categorical variables, or a mixture of numerical and categorical variables. This is relevant in a sense that, most of the time, we want to view everything in one go (e.g. the statistics of all items in a questionnaire), compare the means of several groups and obtain cross-tabulation of categorical variables.

Describing numerical variables

Let us use `women` dataset,

```
head(women)
```

```
##   height weight
## 1     58    115
## 2     59    117
## 3     60    120
## 4     61    123
## 5     62    126
## 6     63    129
```

```
names(women)
```

```
## [1] "height" "weight"
```

```
str(women)
```

```
## 'data.frame': 15 obs. of 2 variables:
## $ height: num 58 59 60 61 62 63 64 65 66 67 ...
## $ weight: num 115 117 120 123 126 129 132 135 139 142 ...
```

which consists of `weight` and `height` numerical variables.

The variables can be easily viewed together by `summary`,

```
summary(women)
```

```
##      height      weight
## Min.   :58.0   Min.   :115.0
## 1st Qu.:61.5   1st Qu.:124.5
## Median :65.0   Median :135.0
## Mean   :65.0   Mean   :136.7
## 3rd Qu.:68.5   3rd Qu.:148.0
## Max.   :72.0   Max.   :164.0
```

even better using `describe` (`psych`),

```
describe(women)
```

```
##      vars  n  mean    sd median trimmed  mad min max range skew
## height   1 15 65.00  4.47    65   65.00  5.93  58  72    14 0.00
## weight   2 15 136.73 15.50   135  136.31 17.79 115 164    49 0.23
##      kurtosis  se
## height   -1.44 1.15
## weight   -1.34 4.00
```

Describing categorical variables

Let us use `infert` dataset,

```
head(infert)
```

```
##   education age parity induced case spontaneous stratum pooled.stratum
## 1    0-5yrs  26     6      1     1           2       1           3
## 2    0-5yrs  42     1      1     1           0       2           1
## 3    0-5yrs  39     6      2     1           0       3           4
## 4    0-5yrs  34     4      2     1           0       4           2
## 5    6-11yrs 35     3      1     1           1       5          32
## 6    6-11yrs 36     4      2     1           1       6          36
```

```
names(infert)
```

```
## [1] "education" "age" "parity" "induced"
## [5] "case" "spontaneous" "stratum" "pooled.stratum"
```

```
str(infert)
```

```
## 'data.frame': 248 obs. of 8 variables:
## $ education : Factor w/ 3 levels "0-5yrs","6-11yrs",...: 1 1 1 1 2 2 2 2 2 2 ...
## $ age : num 26 42 39 34 35 36 23 32 21 28 ...
## $ parity : num 6 1 6 4 3 4 1 2 1 2 ...
```

```
## $ induced      : num  1 1 2 2 1 2 0 0 0 0 ...
## $ case         : num  1 1 1 1 1 1 1 1 1 1 ...
## $ spontaneous  : num  2 0 0 0 1 1 0 0 1 0 ...
## $ stratum      : int   1 2 3 4 5 6 7 8 9 10 ...
## $ pooled.stratum: num   3 1 4 2 32 36 6 22 5 19 ...
```

We notice that `induced`, `case` and `spontaneous` are not yet set as categorical variables, thus we need to factor the variables. We view the value labels in the dataset description,

```
?infert
```

We label the values in the variables according to the description as

```
infert$induced = factor(infert$induced, levels = 0:2, labels = c("0", "1", "2 or more"))
infert$case = factor(infert$case, levels = 0:1, labels = c("control", "case"))
infert$spontaneous = factor(infert$spontaneous, levels = 0:2, labels = c("0", "1", "2 or more"))
str(infert)
```

```
## 'data.frame': 248 obs. of 8 variables:
## $ education : Factor w/ 3 levels "0-5yrs","6-11yrs",...: 1 1 1 1 2 2 2 2 2 ...
## $ age : num 26 42 39 34 35 36 23 32 21 28 ...
## $ parity : num 6 1 6 4 3 4 1 2 1 2 ...
## $ induced : Factor w/ 3 levels "0","1","2 or more": 2 2 3 3 2 3 1 1 1 1 ...
## $ case : Factor w/ 2 levels "control","case": 2 2 2 2 2 2 2 2 2 2 ...
## $ spontaneous : Factor w/ 3 levels "0","1","2 or more": 3 1 1 1 2 2 1 1 2 1 ...
## $ stratum : int 1 2 3 4 5 6 7 8 9 10 ...
## $ pooled.stratum: num 3 1 4 2 32 36 6 22 5 19 ...
```

and we now all these variables are turned into factors.

Again, the variables can be easily viewed together by `summary`,

```
summary(infert[c("education", "induced", "case", "spontaneous")])
```

```
##      education      induced      case      spontaneous
## 0-5yrs : 12    0          :143  control:165    0          :141
## 6-11yrs:120    1          : 68   case  : 83    1          : 71
## 12+ yrs:116    2 or more: 37                2 or more: 36
```

We do not use `table` here in form of `table(infert[c("education", "induced", "case", "spontaneous")])` because `table` used in this form will give us 3-way cross-tabulation instead of count per categories. Cross-tabulation of categorical variables will be covered later.

To obtain the proportion and percentage results, we have to use `lapply`,

```
lapply(infert[c("education", "induced", "case", "spontaneous")],
       function(x) summary(x)/length(x))
```

```
## $education
##    0-5yrs  6-11yrs 12+ yrs
## 0.0483871 0.4838710 0.4677419
##
## $induced
##      0      1 2 or more
## 0.5766129 0.2741935 0.1491935
##
## $case
##   control      case
## 0.6653226 0.3346774
##
```

```
## $spontaneous
##      0      1 2 or more
## 0.5685484 0.2862903 0.1451613

lapply(infert[c("education", "induced", "case", "spontaneous")],
       function(x) summary(x)/length(x)*100)
```

```
## $education
##  0-5yrs 6-11yrs 12+ yrs
## 4.83871 48.38710 46.77419
##
## $induced
##      0      1 2 or more
## 57.66129 27.41935 14.91935
##
## $case
## control    case
## 66.53226 33.46774
##
## $spontaneous
##      0      1 2 or more
## 56.85484 28.62903 14.51613
```

because we need `lapply` to obtain the values for each of the variables. `lapply` goes through each variable and performs this particular part,

```
function(x) summary(x)/length(x)
```

`function(x)` is needed to specify some extra operations to any basic function in R, in our case `summary(x)` divided by `length(x)`, in which the summary results (the counts) are divided by the number of subjects (`length(x)` gives us the “length” of our dataset).

Now, since we already learned about `lapply`, we may also obtain the same results by using `summary` (within `lapply`), `table` and `prop.table`.

```
lapply(infert[c("education", "induced", "case", "spontaneous")], summary)
```

```
## $education
##  0-5yrs 6-11yrs 12+ yrs
##      12      120      116
##
## $induced
##      0      1 2 or more
##     143      68      37
##
## $case
## control    case
##     165      83
##
## $spontaneous
##      0      1 2 or more
##     141      71      36
```

```
lapply(infert[c("education", "induced", "case", "spontaneous")], table)
```

```
## $education
##
##  0-5yrs 6-11yrs 12+ yrs
```

```
##      12      120      116
##
## $induced
##
##      0      1 2 or more
##     143      68      37
##
## $case
##
## control      case
##     165      83
##
## $spontaneous
##
##      0      1 2 or more
##     141      71      36
```

```
lapply(infert[c("education", "induced", "case", "spontaneous")],
       function(x) prop.table(table(x)))
```

```
## $education
## x
##  0-5yrs  6-11yrs  12+ yrs
## 0.0483871 0.4838710 0.4677419
##
## $induced
## x
##      0      1 2 or more
## 0.5766129 0.2741935 0.1491935
##
## $case
## x
## control      case
## 0.6653226 0.3346774
##
## $spontaneous
## x
##      0      1 2 or more
## 0.5685484 0.2862903 0.1451613
```

```
lapply(infert[c("education", "induced", "case", "spontaneous")],
       function(x) prop.table(table(x))*100)
```

```
## $education
## x
##  0-5yrs  6-11yrs  12+ yrs
##  4.83871 48.38710 46.77419
##
## $induced
## x
##      0      1 2 or more
## 57.66129 27.41935 14.91935
##
## $case
## x
```



```
## control      case
## 66.53226 33.46774
##
## $spontaneous
## x
##          0          1 2 or more
## 56.85484 28.62903 14.51613
```

Notice here, whenever we do not need to specify extra operations on a basic function, e.g. `summary` and `table`, all we need to write after the comma in `lapply` is the basic function without `function(x)` and `(x)`.

Describing the variables together

We intentionally went through the descriptive statistics of a variable, followed by a number of variables of the same type. This will give you the basics in dealing with the variables. Most commonly, the variables are described by groups or in form cross-tabulated counts/percentages.

By groups

To obtain all the descriptive statistics by group, we can use `by` with the relevant functions. Let say we want to obtain the statistics by case and control (`case`). We start with numerical variables

```
by(infert[c("age", "parity")], infert$case, summary)
```

```
## infert$case: control
##      age      parity
## Min.   :21.00   Min.   :1.000
## 1st Qu.:28.00   1st Qu.:1.000
## Median :31.00   Median :2.000
## Mean   :31.49   Mean   :2.085
## 3rd Qu.:35.00   3rd Qu.:3.000
## Max.   :44.00   Max.   :6.000
## -----
## infert$case: case
##      age      parity
## Min.   :21.00   Min.   :1.000
## 1st Qu.:28.00   1st Qu.:1.000
## Median :31.00   Median :2.000
## Mean   :31.53   Mean   :2.108
## 3rd Qu.:35.50   3rd Qu.:3.000
## Max.   :44.00   Max.   :6.000
```

```
by(infert[c("age", "parity")], infert$case, describe)
```

```
## infert$case: control
##      vars  n mean  sd median trimmed mad min max range skew kurtosis
## age      1 165 31.49 5.25    31   31.34 5.93  21  44    23 0.23    -0.72
## parity   2 165  2.08 1.24     2    1.88 1.48   1   6     5 1.32     1.42
##      se
## age    0.41
## parity 0.10
## -----
## infert$case: case
##      vars  n mean  sd median trimmed mad min max range skew kurtosis
## age      1  83 31.53 5.28    31   31.39 5.93  21  44    23 0.21    -0.77
```

```
## parity    2 83  2.11 1.28      2    1.90 1.48    1  6    5 1.32    1.34
##          se
## age      0.58
## parity  0.14
```

We can also use `describeBy`, which is an extension of `describe` in the `psych` package.

```
describeBy(infert[c("age", "parity")], group = infert$case)
```

```
##
## Descriptive statistics by group
## group: control
##      vars  n mean  sd median trimmed  mad min max range skew kurtosis
## age      1 165 31.49 5.25    31   31.34 5.93  21 44   23 0.23   -0.72
## parity   2 165  2.08 1.24     2    1.88 1.48   1  6    5 1.32    1.42
##          se
## age      0.41
## parity  0.10
## -----
## group: case
##      vars  n mean  sd median trimmed  mad min max range skew kurtosis
## age      1 83 31.53 5.28    31   31.39 5.93  21 44   23 0.21   -0.77
## parity   2 83  2.11 1.28     2    1.90 1.48   1  6    5 1.32    1.34
##          se
## age      0.58
## parity  0.14
```

which gives us an identical result.

For categorical variables, using `summary`

```
by(infert[c("education", "induced", "spontaneous")], infert$case, summary)
```

```
## infert$case: control
##      education      induced      spontaneous
## 0-5yrs : 8    0          :96    0          :113
## 6-11yrs:80    1          :45    1          : 40
## 12+ yrs:77    2 or more:24    2 or more: 12
## -----
## infert$case: case
##      education      induced      spontaneous
## 0-5yrs : 4    0          :47    0          :28
## 6-11yrs:40    1          :23    1          :31
## 12+ yrs:39    2 or more:13    2 or more:24
```

```
by(infert[c("education", "induced", "spontaneous")], infert$case,
   function(x) lapply(x, function(x) summary(x)/length(x)))
```

```
## infert$case: control
## $education
##    0-5yrs    6-11yrs    12+ yrs
## 0.04848485 0.48484848 0.46666667
##
## $induced
##      0      1 2 or more
## 0.5818182 0.2727273 0.1454545
##
## $spontaneous
```

```
##           0           1 2 or more
## 0.68484848 0.24242424 0.07272727
##
## -----
## infert$case: case
## $education
##    0-5yrs    6-11yrs    12+ yrs
## 0.04819277 0.48192771 0.46987952
##
## $induced
##           0           1 2 or more
## 0.5662651 0.2771084 0.1566265
##
## $spontaneous
##           0           1 2 or more
## 0.3373494 0.3734940 0.2891566
by(infert[c("education", "induced", "spontaneous")], infert$case,
   function(x) lapply(x, function(x) summary(x)/length(x)*100))
```

```
## infert$case: control
## $education
##    0-5yrs    6-11yrs    12+ yrs
## 4.848485 48.484848 46.666667
##
## $induced
##           0           1 2 or more
## 58.18182 27.27273 14.54545
##
## $spontaneous
##           0           1 2 or more
## 68.484848 24.242424 7.272727
##
```

```
## -----
## infert$case: case
## $education
##    0-5yrs    6-11yrs    12+ yrs
## 4.819277 48.192771 46.987952
##
## $induced
##           0           1 2 or more
## 56.62651 27.71084 15.66265
##
## $spontaneous
##           0           1 2 or more
## 33.73494 37.34940 28.91566
```

or by using table

```
by(infert[c("education", "induced", "spontaneous")], infert$case,
   function(x) lapply(x, table))
```

```
## infert$case: control
## $education
##
##    0-5yrs 6-11yrs 12+ yrs
```

```

##      8      80      77
##
## $induced
##
##      0      1 2 or more
##     96     45      24
##
## $spontaneous
##
##      0      1 2 or more
##    113     40      12
##
## -----
## infert$case: case
## $education
##
## 0-5yrs 6-11yrs 12+ yrs
##      4      40      39
##
## $induced
##
##      0      1 2 or more
##     47     23      13
##
## $spontaneous
##
##      0      1 2 or more
##     28     31      24
##
by(infert[c("education", "induced", "spontaneous")], infert$case,
   function(x) lapply(x, function(x) prop.table(table(x))))

## infert$case: control
## $education
## x
##      0-5yrs      6-11yrs      12+ yrs
## 0.04848485 0.48484848 0.46666667
##
## $induced
## x
##      0      1 2 or more
## 0.5818182 0.2727273 0.1454545
##
## $spontaneous
## x
##      0      1 2 or more
## 0.68484848 0.24242424 0.07272727
##
## -----
## infert$case: case
## $education
## x
##      0-5yrs      6-11yrs      12+ yrs
## 0.04819277 0.48192771 0.46987952
##

```

```
## $induced
## x
##      0      1 2 or more
## 0.5662651 0.2771084 0.1566265
##
## $spontaneous
## x
##      0      1 2 or more
## 0.3373494 0.3734940 0.2891566
by(infert[c("education", "induced", "spontaneous")], infert$case,
  function(x) lapply(x, function(x) prop.table(table(x))*100))
```

```
## infert$case: control
## $education
## x
##      0-5yrs      6-11yrs      12+ yrs
## 4.848485 48.484848 46.666667
##
## $induced
## x
##      0      1 2 or more
## 58.18182 27.27273 14.54545
##
## $spontaneous
## x
##      0      1 2 or more
## 68.484848 24.242424 7.272727
## -----
## infert$case: case
## $education
## x
##      0-5yrs      6-11yrs      12+ yrs
## 4.819277 48.192771 46.987952
##
## $induced
## x
##      0      1 2 or more
## 56.62651 27.71084 15.66265
##
## $spontaneous
## x
##      0      1 2 or more
## 33.73494 37.34940 28.91566
```

Please note that simply replacing `table` for `summary` as in `by(infert[c("education", "induced", "spontaneous")], infert$case, table)` will not work as intended. `education` will be nested in `induced`, which is nested in `spontaneous`, listed by `case` instead. And yes, to obtain the proportions and percentages, it gets slightly more complicated as we have to specify `function` twice in `by`.

Simple cross-tabulation

As long as the categorical variables are already **factored** properly, there should not be a problem to obtain the cross-tabulation tables. For example between `education` and `case`,

```
table(infert$education, infert$case)
```

```
##
##           control case
## 0-5yrs         8    4
## 6-11yrs        80   40
## 12+ yrs        77   39
```

We may also include row and column headers, just like `cbind`,

```
table(education = infert$education, case = infert$case)
```

```
##           case
## education control case
## 0-5yrs         8    4
## 6-11yrs        80   40
## 12+ yrs        77   39
```

Since we are familiar with the powerful `lapply`, we can use it to get cross-tabulation of all of the factors with case status,

```
lapply(infert[c("education", "induced", "spontaneous")], function(x) table(x, infert$case))
```

```
## $education
##
## x           control case
## 0-5yrs         8    4
## 6-11yrs        80   40
## 12+ yrs        77   39
##
## $induced
##
## x           control case
## 0             96   47
## 1             45   23
## 2 or more     24   13
##
## $spontaneous
##
## x           control case
## 0             113  28
## 1             40   31
## 2 or more     12   24
```

We may also view subgroup counts (nesting). Here, the cross-tabulation of `education` and `case` is nested within `induced`

```
table(infert$education, infert$case, infert$induced)
```

```
## , , = 0
##
##
##           control case
## 0-5yrs         4    0
## 6-11yrs        57   21
## 12+ yrs        35   26
##
## , , = 1
```

```
##
##
##           control case
## 0-5yrs         0    2
## 6-11yrs        16   11
## 12+ yrs        29   10
##
## , , = 2 or more
##
##           control case
## 0-5yrs         4    2
## 6-11yrs        7    8
## 12+ yrs        13    3
```

which will look nicer if we apply by

```
by(infert[c("education", "case")], infert$induced, table)
```

```
## infert$induced: 0
##           case
## education control case
## 0-5yrs         4    0
## 6-11yrs        57   21
## 12+ yrs        35   26
## -----
## infert$induced: 1
##           case
## education control case
## 0-5yrs         0    2
## 6-11yrs        16   11
## 12+ yrs        29   10
## -----
## infert$induced: 2 or more
##           case
## education control case
## 0-5yrs         4    2
## 6-11yrs        7    8
## 12+ yrs        13    3
```

Summary

In this chapter, we learned about how to handle numerical and categorical variables and obtain the basic and relevant statistics. In the next chapter, we are going to learn about how to explore the variables in visually in form of the relevant graphs and plots.