

**Results**

**Descriptives**

Descriptives


	Dose	Happiness
<b>N</b>	<b>Control</b>	5
	<b>15 mins</b>	5
	<b>30 mins</b>	5
<b>Missing</b>	<b>Control</b>	0
	<b>15 mins</b>	0
	<b>30 mins</b>	0
<b>Mean</b>	<b>Control</b>	2.20
	<b>15 mins</b>	3.20
	<b>30 mins</b>	5.00
<b>Median</b>	<b>Control</b>	2.00
	<b>15 mins</b>	3.00
	<b>30 mins</b>	5.00
<b>Standard deviation</b>	<b>Control</b>	1.30
	<b>15 mins</b>	1.30
	<b>30 mins</b>	1.58
<b>Minimum</b>	<b>Control</b>	1.00
	<b>15 mins</b>	2.00
	<b>30 mins</b>	3.00
<b>Maximum</b>	<b>Control</b>	4.00
	<b>15 mins</b>	5.00
	<b>30 mins</b>	7.00
<b>Skewness</b>	<b>Control</b>	0.541
	<b>15 mins</b>	0.541
	<b>30 mins</b>	0.00
<b>Std. error skewness</b>	<b>Control</b>	0.913
	<b>15 mins</b>	0.913
	<b>30 mins</b>	0.913
<b>Kurtosis</b>	<b>Control</b>	-1.49
	<b>15 mins</b>	-1.49
	<b>30 mins</b>	-1.20
<b>Std. error kurtosis</b>	<b>Control</b>	2.00
	<b>15 mins</b>	2.00
	<b>30 mins</b>	2.00
<b>Shapiro-Wilk W</b>	<b>Control</b>	0.902
	<b>15 mins</b>	0.902
	<b>30 mins</b>	0.987

Descriptives

Shapiro-Wilk p	Control	0.421
	15 mins	0.421
	30 mins	0.967

Plots

Happiness




Relationships, Prediction, and Group Comparisons

You have entered a numeric variable for Variable 1 / Dependent Variable and a nominal variable for Variable 2 / Independent Variables. Hence, a [one way ANOVA](#), which is a test for the difference between several population means, seems to be a good option for you! In order to run this analysis in jamovi, go to: ANOVA > ANOVA

- Drop your dependent (numeric) variable in the box below Dependent Variable and your independent (grouping) variable in the box below Fixed Factors

If the normality or homoscedasticity assumption is violated, you could use the non-parametric [Kruskal-Wallis test](#). Click on the links to learn more about these tests!

## Scatter Plots of Bivariate Relationships - Dependent/Independent Variables



## One-Way ANOVA

One-Way ANOVA

		F	df1	df2	p
<b>Happiness</b>	<b>Welch's</b>	4.32	2	7.94	0.054
	<b>Fisher's</b>	5.12	2	12	0.025

Group Descriptives

	Dose	N	Mean	SD	SE
<b>Happiness</b>	<b>Control</b>	5	2.20	1.30	0.583
	<b>15 mins</b>	5	3.20	1.30	0.583
	<b>30 mins</b>	5	5.00	1.58	0.707

## Assumption Checks

Homogeneity of Variances Tests

		Statistic	df	df2	p
Happiness	Levene's	0.0917	2	12	0.913
	Bartlett's	0.185	2		0.912

Note. Additional results provided by moretests


Normality Tests

		statistic	p
Happiness	Shapiro-Wilk	0.917	0.171
	Kolmogorov-Smirnov	0.179	0.720
	Anderson-Darling	0.517	0.159

Note. Additional results provided by moretests

Plots

Happiness



Post Hoc Tests

Tukey Post-Hoc Test – Happiness

		Control	15 mins	30 mins
Control	Mean difference	—	-1.00	-2.80
	p-value	—	0.516	0.021
15 mins	Mean difference		—	-1.80
	p-value		—	0.147
30 mins	Mean difference			—
	p-value			—

ANOVA

ANOVA - Happiness

	Sum of Squares	df	Mean Square	F	p	$\omega^2$
Dose	20.1	2	10.07	5.12	0.025	0.354
Residuals	23.6	12	1.97			

[3]

Assumption Checks

Homogeneity of Variances Tests

	Statistic	df	df2	p
Levene's	0.0917	2	12	0.913
Bartlett's	0.185	2		0.912


Note. Additional results provided by *moretests*

Normality tests

	statistic	p
Shapiro-Wilk	0.917	0.171
Kolmogorov-Smirnov	0.179	0.720
Anderson-Darling	0.517	0.159

Note. Additional results provided by *moretests*

Q-Q Plot



### Post Hoc Tests

Post Hoc Comparisons - Dose


Comparison							
Dose	Dose	Mean Difference	SE	df	t	P <sub>Tukey</sub>	Cohen's d
Control	15 mins	-1.00	0.887	12.0	-1.13	0.516	-0.713
	30 mins	-2.80	0.887	12.0	-3.16	0.021	-1.997
15 mins	30 mins	-1.80	0.887	12.0	-2.03	0.147	-1.284

*Note.* Comparisons are based on estimated marginal means

[4]

### Estimated Marginal Means

Dose



[4]

## Robust ANOVA

Robust ANOVA

	F	df1	df2	p	ES	Bootstrap CI	
						Lower	Upper
<b>Dose</b>	3.00	2.00	4.00	0.160	0.789	0.421	1.43

Note. Method of trimmed means (level 0.2).

Note. For effect size CI computation (samples 599)

## Post Hoc Tests

Post Hoc Tests - Dose

				95% Confidence interval	
				Lower	Upper
Control	15 mins	-1.00	0.435	-5.32 <sup>a</sup>	3.32 <sup>a</sup>
	30 mins	-3.00	0.181	-7.32 <sup>a</sup>	1.32 <sup>a</sup>
15 mins	30 mins	-2.00	0.317	-6.32 <sup>a</sup>	2.32 <sup>a</sup>

<sup>a</sup> CI are adjusted to control FWE, but not p-values.

## References

[1] The jamovi project (2024). *jamovi*. (Version 2.6) [Computer Software]. Retrieved from <https://www.jamovi.org>.



**[2]** R Core Team (2024). *R: A Language and environment for statistical computing*. (Version 4.4) [Computer software]. Retrieved from <https://cran.r-project.org>. (R packages retrieved from CRAN snapshot 2024-08-07).

**[3]** Fox, J., & Weisberg, S. (2023). *car: Companion to Applied Regression*. [R package]. Retrieved from <https://cran.r-project.org/package=car>.

**[4]** Lenth, R. (2023). *emmeans: Estimated Marginal Means, aka Least-Squares Means*. [R package]. Retrieved from <https://cran.r-project.org/package=emmeans>.